

NOTE

ULTRAVIOLET ABSORPTION SPECTRA OF PERETHYLPOLYGERMANES AND -POLYSTANNANES

W. DRENTH, J. G. NOLTES, E. J. BULTEN AND H. M. J. C. CREEMERS

Institute for Organic Chemistry TNO, Utrecht (The Netherlands)

(Received January 17th, 1969)

As shown by Gilman¹ and Kumada² silanes of general structure $R(SiR_2)_nR$ show ultraviolet absorption. The first absorption maximum is shifted to longer wave length with increasing number of metal atoms in the chain. Shorygin *et al.*² observed the same phenomenon in polygermanes. They also give the spectra of $(CH_3)_6Sn_2$ and $(CH_3)_{14}Sn_6$. The phenomenon of bathochromic shift with increasing chain length is not typical for metal chains. It also occurs in normal hydrocarbons for which Raymonda and Simpson⁴ give an explanation in terms of their independent systems model. Pitt, Jones and Ramsey^{5,6} use molecular orbital theory to account for the shift in metal chains.

According to the latter theory a plot of transition energy versus $2 \cos[\pi/(n+1)]$ gives a straight line the slope of which is equal to $\beta_g + \beta_e$ ⁶. These β 's are exchange integrals between orbitals of neighbouring metal atoms. The subscript g refers to the ground state and the subscript e to the first excited state.

TABLE 1

ULTRAVIOLET ABSORPTION SPECTRA OF POLYMETAL COMPOUNDS^a

<i>n</i>	$CH_3[Si(CH_3)_2]_nCH_3$		$C_2H_5[Ge(C_2H_5)_2]_n C_2H_5$		$C_2H_5[Sn(C_2H_5)_2]_n C_2H_5$	
	λ_{max} (nm)	<i>hν</i> (eV)	λ_{max} (nm)	<i>hν</i> (eV)	λ_{max} (nm)	<i>hν</i> (eV)
2	192	6.45	202 sh	6.14	232 sh ^b	5.34
3	215	5.77	217.5 sh	5.70	250 sh	4.96
4	235	5.28	233.5	5.31	290 sh	4.28
5	250	4.96	248	5.00	310	4.00
6	260	4.77	258	4.81	325 ^b	3.82
8	272.5	4.55				

^a Spectra were recorded in cyclohexane solution using a Cary Model 15 spectrometer. ^b Cf. 210 nm for *n*=2 and 246 nm for *n*=6 in the permethyl compounds³.

Several perethylpolygermanes and -polystannanes have been prepared in this Institute^{7,8} and their ultraviolet absorption spectra measured (Table 1). The wave lengths of maximum absorption of the perethylpolygermanes are slightly higher than the data given by Shorygin *et al.*³ for permethylpolygermanes, but the tin data are




Fig. 1. Plot of $h\nu$ vs. $2 \cos[\pi/(n+1)]$.

appreciably different.

Plots of transition energies corresponding with the data in Table 1, are given in Fig. 1. For comparison, transition energies of permethylsilanes¹ have been included. The silicon, germanium and tin lines are parallel. Thus, $\beta_g + \beta_e$ has equal value for Si-Si, Ge-Ge and Sn-Sn bonds. It amounts to -2.7 eV, compared to -1.7 eV for normal hydrocarbons⁶.

In the series Si_2 , Ge_2 , Sn_2 metal-metal bond energies are approximately 3.0, 2.8 and 2.2 eV⁹. Assuming that β_g will parallel this change and, therefore, will increase to a less negative value, β_e has to decrease to a more negative value. This result indicates an increasing overlap of atomic orbitals which are the basis of the excited state molecular orbitals, in the series $\text{Si} \leq \text{Ge} < \text{Sn}$.

ACKNOWLEDGEMENTS

The authors are indebted to Dr. C. G. PITT and to Professor B. G. RAMSEY for making available their results prior to publication. They wish to thank Professor G. J. M. VAN DER KERK for his interest in this study.

REFERENCES

- 1 H. GILMAN, W. H. ATWELL AND G. L. SCHWEBKE, *J. Organometal. Chem.*, 2 (1964) 369.
- 2 H. SAKURAI AND M. KUMADA, *Bull. Chem. Soc. Jap.*, 37 (1964) 1894.
- 3 P. P. SHORYGIN, V. A. PETUKHOV, O. M. NEFEDOV, S. P. KOLESNIKOV AND V. I. SHIRYAEV, *Teor. Eksp. Khim.*, 2 (1966) 190; *Chem. Abstr.*, 65 (1966) 14660.
- 4 J. W. RAYMONDA AND W. T. SIMPSON, *J. Chem. Phys.*, 47 (1967) 430.
- 5 C. G. PITT, L. L. JONES AND B. G. RAMSEY, *J. Amer. Chem. Soc.*, 89 (1967) 5471.
- 6 B. G. RAMSEY, *Electronic Transitions in Organometalloids*, Academic Press, New York, in the press; C. G. PITT, private communication.
- 7 E. J. BULTEN AND J. G. NOLTES, *J. Organometal. Chem.*, 16 (1969) P8.
- 8 H. M. J. C. CREEMERS AND J. G. NOLTES, *Rec. Trav. Chim. Pays-Bas*, 84 (1965) 382; H. M. J. C. CREEMERS, Ph.D. Thesis, Utrecht, 1967.
- 9 V. I. TELNOI AND I. R. RABINOWITZ, *Zh. Fiz. Khim.*, (1966) 1556; *Chem. Abstr.*, 65 (1966) 15209.