

# Hyperkähler Cones and Orthogonal Wolf Spaces

Lilia Anguelova\*, Martin Roček\* and Stefan Vandoren†

\**C.N. Yang Institute for Theoretical Physics  
SUNY, Stony Brook, NY 11794-3840, USA*

anguelov@grad.physics.sunysb.edu

Roček@insti.physics.sunysb.edu

†*Institute for Theoretical Physics and Spinoza Institute  
Utrecht University, Utrecht, The Netherlands*

vandoren@phys.uu.nl

## ABSTRACT

We construct the hyperkähler cones corresponding to the quaternion-Kähler orthogonal Wolf spaces  $\frac{SO(n+4)}{SO(n) \times SO(4)}$  and their non-compact versions, which appear in hypermultiplet couplings to  $N = 2$  supergravity. The geometry is completely encoded by a single function, the hyperkähler potential, which we compute from an  $SU(2)$  hyperkähler quotient of flat space. We derive the Killing vectors and moment maps for the  $SO(n+4)$  isometry group on the hyperkähler cone. For the non-compact case, the isometry group  $SO(n,4)$  contains  $n+2$  abelian isometries which can be used to find a dual description in terms of  $n$  tensor multiplets and one double-tensor multiplet. Finally, using a representation of the hyperkähler quotient via quiver diagrams, we deduce the existence of a new eight dimensional ALE space.

# 1 Introduction

Quaternion-Kähler (QK) manifolds have recently attracted a lot of attention in the context of  $N = 2$  gauged supergravity, both in four and five spacetime dimensions. In particular, those with negative scalar curvature appear as target spaces for  $N = 2$  hypermultiplets coupled to supergravity [1]. The *homogeneous* QK spaces  $G/H$  are classified by Wolf and Alekseevskii [2]; the compact ones are given by the three infinite series

$$\mathbb{HP}(n) = \frac{Sp(n+1)}{Sp(n) \times Sp(1)}, \quad X(n) = \frac{SU(n+2)}{SU(n) \times U(2)}, \quad Y(n) = \frac{SO(n+4)}{SO(n) \times SO(4)}, \quad (1.1)$$

of dimension  $4n$ , and the five exceptional cases

$$\frac{G_2}{SO(4)}, \quad \frac{F_4}{Sp(3) \times Sp(1)}, \quad \frac{E_6}{SU(6) \times Sp(1)}, \quad \frac{E_7}{Spin(12) \times Sp(1)}, \quad \frac{E_8}{E_7 \times Sp(1)}, \quad (1.2)$$

of dimensions 8, 28, 40, 64 and 112 respectively. Their non-compact versions have negative scalar curvature and hence are relevant for supergravity applications. For low dimensions, there are relations between these spaces: the four-sphere  $Y(1) \cong \mathbb{HP}(1)$ , and  $Y(2) \cong X(2)$ . These isomorphisms are discussed in more detail below.

A complete classification of QK spaces does not exist. Because their  $Sp(1)$  curvature does not vanish, and their quaternionic two-forms are not closed, QK spaces are rather difficult to deal with. This complicates the study of hypermultiplet couplings to supergravity, which arise in the low energy effective action of type II superstring compactification on Calabi-Yau threefolds [3]. In this paper, we focus on the  $Y(n)$  spaces. Their noncompact versions

$$\tilde{Y}(n) = \frac{SO(n,4)}{SO(n) \times SO(4)}, \quad (1.3)$$

describe the classical moduli spaces of type II Calabi-Yau compactifications down to four dimensions, either as dual quaternionic spaces from the  $\mathbf{c}$ -map [3], or from dualizing  $n$  tensor multiplets and a double-tensor multiplet [4].

Vector multiplet couplings to  $N = 2$  supergravity are geometrically much simpler. The scalar fields of the vector multiplets parametrize a special Kähler geometry in  $d=4$  [5] and a very special real geometry in  $d=5$  [6]. Both these geometries are completely determined in terms of a single function from which the target space metric, and other geometrical quantities, can easily be computed. It is desirable to have a similar description for QK manifolds, such that the whole geometry is encoded by a single function of the hypermultiplet scalars. Such a description was first given in the mathematics literature, where it was shown [7], that to every  $4n$  dimensional QK manifold one can associate a  $4(n+1)$  dimensional hyperkähler manifold which admits a conformal homothety and an

isometric  $\mathrm{Sp}(1)$ -action that rotates the three complex structures into each other. The conformal homothety  $\chi^A$  satisfies

$$D_A \chi^B = \delta_A^B, \quad (A, B = 1, \dots, 4n), \quad (1.4)$$

such that locally  $\chi_A = \partial_A \chi$ . The function  $\chi$  is called the hyperkähler potential and the metric is

$$g_{AB} = D_A \partial_B \chi. \quad (1.5)$$

Such a manifold is a hyperkähler cone (HKC) over a base which is an  $\mathrm{Sp}(1)$  fibration over a QK space [7, 8].

It was later anticipated in [9], and shown explicitly in [10, 11] how this connection is precisely realised in the conformal calculus for hypermultiplets. The transition from the HKC to the QK space is called the  $N = 2$  *superconformal quotient*, and is associated with eliminating the compensating hypermultiplet by an  $SU(2)$  quotient and gauge-fixing combined with a reduction along the homothety. The simplest QK manifolds are the quaternionic projective spaces  $\mathbb{H}\mathbb{P}(n)$ . Galicki has shown that the  $X$  and  $Y$  series can be obtained from these by performing quaternionic  $U(1)$  and  $SU(2)$  quotients, respectively [12]. It is interesting to study these spaces in terms of the corresponding HKC's. The HKC of  $\mathbb{H}\mathbb{P}(n-1)$  is simply  $\mathbb{H}^n \cong \mathbb{C}^{2n}$ , and the explicit construction of the HKC of  $X(n-2)$  is a  $U(1)$  hyperkähler quotient of  $\mathbb{H}^n$  [11]. In Section 2 of this paper, we derive the hyperkähler potential  $\chi$  for the orthogonal Wolf spaces  $Y(n-4)$  by performing an  $SU(2)$  hyperkähler quotient on  $\mathbb{H}^n$ . The action of the  $SU(2)$  on the coordinates of  $\mathbb{H}^n$  can be deduced from Galicki's quaternionic quotient [12] because the diagram in Figure 1 is commutative [7]. Hence quaternionic quotients between the QK manifolds lift to hyperkähler quotients between the corresponding HKC's. We also write down the moment maps and Killing vectors for the  $SO(n)$  isometry group on the HKC. These appear in the scalar potential of gauged  $N = 2$  supergravity.

In Section 3 we consider the three special cases for which the hyperkähler potential of the HKC of  $Y(n-4)$  is known:  $n = 4$ , where the HKC of  $Y(0)$  is  $\mathbb{R}^4$ ,  $n = 5$ , where the HKC of  $Y(1)$  is  $\mathbb{R}^8$  and  $n = 6$ , where the HKC of  $Y(2)$  is the HKC of  $X(2)$ . We check our calculation by showing that each of these special cases agrees with the known results.

In Section 4 we construct the dual description of the HKC of the non-compact version  $\tilde{Y}(n)$  in terms of  $n$  tensor multiplets and a double tensor multiplet. This tensor multiplet description naturally appears in type IIB compactifications.

In Section 5 we show how the hyperkähler quotient can be lifted to  $N = 2$  superspace. Once we have this description, we can use the techniques of [14] to write down the quiver diagrams for the HKC's of  $Y(n)$ . This point of view allows us to infer the existence of a new 8-dimensional ALE space.

$$\begin{array}{ccccc}
\text{HKC}(X) & \xleftarrow[-4]{\text{u}(1)} & \mathbb{C}^{2n} = \mathbb{H}^n & \xrightarrow[-12]{\text{su}(2)} & \text{HKC}(Y) \\
\downarrow -4 & & \downarrow & & \downarrow -4 \\
X(n-2) & \xleftarrow[-4]{\text{u}(1)} & \text{HP}(n-1) & \xrightarrow[-12]{\text{su}(2)} & Y(n-4)
\end{array}$$

Figure 1: The horizontal arrows stand for hyperkähler quotients (upper), and quaternionic quotients (lower). The vertical arrows denote the superconformal quotients. The numerical label of each arrow denotes the change of the number of real dimensions accompanying the corresponding quotient.

Our long term goal is to understand the nonperturbative corrections to quaternionic geometries; we expect that these are most easily computed in terms of the hyperkähler potential  $\chi$ . We also note that HKC's appear as moduli spaces of Yang-Mills instantons [15, 16], *e.g.*, the homogeneous QK  $G/H$  manifolds determine the one-instanton moduli spaces with gauge group  $G$  and stability group  $H$ . This suggests a connection between the moduli spaces of Calabi-Yau manifolds and Yang-Mills instantons [16].

## 2 The $SU(2)$ Quotient

In this section, we explicitly perform the non-abelian hyperkähler quotient from  $\mathbb{H}^n = \mathbb{R}^{4n}$  to the HKC of  $Y(n-4)$ ,  $n \geq 4$ . For simplicity, we do the analysis for the compact case; the non-compact case is very similar and we give the final result at the end of this section. The space  $\mathbb{H}^n$  is flat, and has an isometry group  $SO(n)$  which is linearly realized on the  $n$  quaternions

$$q^M = \begin{pmatrix} z_+^M & -\bar{z}_-^M \\ z_-^M & \bar{z}_+^M \end{pmatrix}, \quad M = 1, \dots, n. \quad (2.1)$$

Clearly,  $\mathbb{H}^n$  is itself an HKC with hyperkähler potential (or equivalently,  $N = 1$  superspace Lagrangian)

$$\chi(\mathbb{H}^n) = \frac{1}{2} \text{Tr}(q^M q^{\dagger M}) = z_+^M \bar{z}_+^M + z_-^M \bar{z}_-^M, \quad (2.2)$$

where a sum over repeated indices is understood. The closed hyperkähler two-forms on  $\mathbb{H}^n$  are

$$\Omega^3 = -i \left( dz_+^M \wedge d\bar{z}_+^M + dz_-^M \wedge d\bar{z}_-^M \right), \quad \Omega^+ = dz_+^M \wedge dz_-^M, \quad \Omega^- = (\Omega^+)^*, \quad (2.3)$$

such that the  $SO(n)$  isometries  $q^M \rightarrow O^M_N q^N$  are triholomorphic.

The infinitesimal  $SU(2)$  action that we quotient by can be deduced from the quaternionic quotient in [12]; it acts by left multiplication on each quaternion, with corresponding Killing vector fields on  $\mathbb{H}^n$

$$\begin{aligned} k_3 &= iz_+^M \frac{\partial}{\partial z_+^M} - iz_-^M \frac{\partial}{\partial z_-^M} - i\bar{z}_+^M \frac{\partial}{\partial \bar{z}_+^M} + i\bar{z}_-^M \frac{\partial}{\partial \bar{z}_-^M} , \\ k_+ &= -z_-^M \frac{\partial}{\partial z_+^M} + \bar{z}_+^M \frac{\partial}{\partial \bar{z}_-^M} , \\ k_- &= z_+^M \frac{\partial}{\partial z_-^M} - \bar{z}_-^M \frac{\partial}{\partial \bar{z}_+^M} . \end{aligned} \quad (2.4)$$

Hence, for each  $M$ ,  $(z_+, z_-)$  transforms as an  $SU(2)$  doublet<sup>1</sup>:

$$\begin{aligned} \delta z_\pm &\equiv \lambda^i k_i z_\pm , & i &= 3, +, - \\ \delta z_+ &= i\lambda^3 z_+ - \lambda^+ z_- , & \delta \bar{z}_+ &= -i\lambda^3 \bar{z}_+ - \lambda^- \bar{z}_- , \\ \delta z_- &= -i\lambda^3 z_- + \lambda^- z_+ , & \delta \bar{z}_- &= i\lambda^3 \bar{z}_- + \lambda^+ \bar{z}_+ . \end{aligned} \quad (2.5)$$

This isometry group is also triholomorphic<sup>2</sup> and commutes with  $SO(n)$ .

The components of the corresponding triplets of moment maps<sup>3</sup> on  $\mathbb{H}^n$  are, for  $SU(2)$ ,

$$\begin{aligned} \mu_i^3 &= -\left( z_+ \cdot \bar{z}_+ - z_- \cdot \bar{z}_- , i z_- \cdot \bar{z}_+ , -i z_+ \cdot \bar{z}_- \right) , \\ \mu_i^+ &= \left( -i z_+ \cdot z_- , \frac{1}{2} z_- \cdot z_- , \frac{1}{2} z_+ \cdot z_+ \right) , & \mu_i^- &= (\mu_i^+)^* , \end{aligned} \quad (2.6)$$

and for  $SO(n)$  (with antihermitian generators  $T_I$ )

$$\mu_I^3 = -i \left( (z_+ T_I \bar{z}_+) + (z_- T_I \bar{z}_-) \right) , \quad \mu_I^+ = (z_+ T_I z_-) , \quad \mu_I^- = (\mu_I^+)^* . \quad (2.7)$$

As indicated by these formulae, we sometimes suppress indices in inner products and bilinear forms on  $\mathbb{H}^n$  to simplify the notation.

The  $SU(2)$  hyperkähler quotient requires setting the moment maps (2.6) to zero. Hence the independent set of constraints are the non-holomorphic ones

$$\bar{z}_+ \cdot z_+ = \bar{z}_- \cdot z_- , \quad \bar{z}_+ \cdot z_- = 0 , \quad \bar{z}_- \cdot z_+ = 0 , \quad (2.8)$$

---

<sup>1</sup>Ordinarily, in a hypermultiplet the  $N = 1$  chiral superfields  $z_+$  and  $z_-$  transform in conjugate representations of the gauge group; however, since the spinor representation of  $SU(2)$  is pseudoreal, one may consider an action that mixes them [20]. In the literature, this is sometimes called a ‘‘half-hypermultiplet’’. In section 5 we discuss a formulation in terms of full hypermultiplets that lifts to an  $N = 2$  superspace description.

<sup>2</sup>This  $SU(2)$  should not be confused with the  $Sp(1) = SU(2)$  that rotates the complex structure and is used to perform the  $N = 2$  superconformal quotient. Under its action  $(z_+, \bar{z}_-)$  transforms as a doublet, which corresponds to right multiplication on each quaternion.

<sup>3</sup>We follow the conventions and notations of [11, 17]; for each triholomorphic Killing vector field  $k_I$ , there is a triplet of moment maps  $\vec{\mu}_I = (\mu_I^3, \mu_I^+, \mu_I^-)$ .

and the holomorphic ones

$$z_+ \cdot z_- = 0, \quad z_+ \cdot z_+ = 0, \quad z_- \cdot z_- = 0. \quad (2.9)$$

To perform the quotient one has to solve (2.8) and (2.9), reducing the number of real coordinates by  $3 + 6 = 9$ , and choose a gauge to fix the  $SU(2)$  gauge freedom, further reducing the number of real coordinates by 3; thus the dimension is reduced by 12 real or 6 complex coordinates, consistent with the dimension of the HKC of  $Y(n-4)$ , namely,  $4(n-3)$ . In practice it is hard to solve all the constraints; as explained in [18], one may complexify<sup>4</sup> the gauge group (in our case  $SU(2) \rightarrow SL(2, \mathbb{C})$ ) and solve just the holomorphic constraints, which are preserved by the complexified gauge group. This construction is completely natural in  $N = 1$  superspace, where the non-holomorphic constraints are solved for the  $N = 1$  gauge fields. If we choose 3 *complex* gauge conditions and impose the holomorphic constraints, we see that we still reduce the dimension by 12. The hyperkähler potential on the quotient space is found by gauging the  $N = 1$  Lagrangian for  $\mathbb{H}^n$  and substituting the solution for the gauge fields.

Recall that the hyperkähler potential for the flat space  $\mathbb{H}^n$  is (2.2). For each value of the index  $M$  the pair  $(z_+^M, z_-^M)$  is a doublet under the  $SU(2)$  which we quotient by (see (2.5)). Hence we have  $n$  copies of the 2-dimensional representation of  $SU(2)$  in (2.2).

We introduce a gauge field  $e^V$ , such that the gauged hyperkähler potential of  $\mathbb{H}^n$  is ([20, 19, 18], see also section five and appendix B of [11]):

$$\hat{\chi} = \sum_{M=1}^n (\bar{z}_+^M \ \bar{z}_-^M) \begin{pmatrix} e^V \end{pmatrix} \begin{pmatrix} z_+^M \\ z_-^M \end{pmatrix}, \quad (2.10)$$

where  $\begin{pmatrix} e^V \end{pmatrix}$  is the  $2 \times 2$  matrix of  $N = 1$   $SU(2)$  gauge fields and hence is hermitian with determinant 1. The nonholomorphic constraints (2.8) (*c.f.* (2.6)) can be written as

$$\mu_i^3 = (\bar{z}_+^M \ \bar{z}_-^M) T_i \begin{pmatrix} z_+^M \\ z_-^M \end{pmatrix} = 0, \quad i = 3, +, - \quad (2.11)$$

where  $T_i$  are the  $SU(2)$  generators normalized as

$$T_3 = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}, \quad T_+ = \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}, \quad T_- = \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix}. \quad (2.12)$$

The gauged nonholomorphic moment maps are [20, 19, 18]:

$$\hat{\mu}_i^3 = (\bar{z}_+^M \ \bar{z}_-^M) T_i \begin{pmatrix} e^V \end{pmatrix} \begin{pmatrix} z_+^M \\ z_-^M \end{pmatrix} = 0. \quad (2.13)$$

---

<sup>4</sup>The complexification of the gauge group means doubling of the number of generators (to the set of initial Killing vectors  $k_I$  one adds the set  $J^3 k_I$ , where  $J^3$  is the complex structure determined by  $\Omega^3(X, Y) = g(J^3 X, Y)$ ).

Solving (2.13) for the gauge fields we find

$$\left( e^V \right) = \frac{1}{\sqrt{(\bar{z}_+ \cdot z_+)(\bar{z}_- \cdot z_-) - (\bar{z}_+ \cdot z_-)(\bar{z}_- \cdot z_+)}} \begin{pmatrix} \bar{z}_- \cdot z_- & -\bar{z}_- \cdot z_+ \\ -\bar{z}_+ \cdot z_- & \bar{z}_+ \cdot z_+ \end{pmatrix}, \quad (2.14)$$

and hence the hyperkähler potential of the quotient is

$$\chi_{\text{quotient}} = 2\sqrt{\bar{z}_+^M z_+^M \bar{z}_-^N z_-^N - \bar{z}_+^M z_-^M \bar{z}_-^N z_+^N}, \quad (2.15)$$

with  $M, N$  still running from 1 to  $n$ . To find coordinates on the quotient we gauge-fix and solve the holomorphic constraints; we can choose

$$z_+^n = 0, \quad z_-^{n-1} = 0, \quad z_+^{n-2} = 1. \quad (2.16)$$

Now the holomorphic constraints (2.9) can be easily solved:

$$z_-^{n-2} = -z_+^a z_-^a, \quad z_+^{n-1} = i\sqrt{1 + z_+^a z_+^a}, \quad z_-^n = i\sqrt{z_+^a z_-^a z_+^b z_-^b + z_-^a z_-^a}, \quad (2.17)$$

where  $\{z_+^a, z_-^b\}$ ,  $a, b = 1, \dots, n-3$  are the coordinates on the HKC of  $Y(n-4)$ . Thus the hyperkähler potential on the HKC of  $Y(n-4)$  is given by (2.15), subject to (2.16) and (2.17). Furthermore the dilatations act on  $z_+$  and  $z_-$  with scaling weights 0 and 2 respectively, and the holomorphic two-form on the HKC is simply

$$\Omega^+ = dz_+^a \wedge dz_-^a, \quad a = 1, \dots, n-3. \quad (2.18)$$

The gauge choice (2.16) is legitimate, but not always the most convenient, as we see in the next section.

The moment maps of the  $SO(n)$  isometry on the HKC can easily be computed from the  $SU(2)$  quotient of (2.7). The  $SU(2)$  gauged  $SO(n)$  moment maps take a similar form to (2.13), and eliminating the gauge fields yields

$$\begin{aligned} \mu_I^3 &= \frac{2i}{\chi} \left( (\bar{z}_- \cdot z_-)(\bar{z}_+ T_I z_+) - (\bar{z}_+ \cdot z_-)(\bar{z}_- T_I z_+) + (+ \leftrightarrow -) \right), \\ \mu_I^+ &= (z_+ T_I z_-), \quad \mu_I^- = (\mu_I^+)^*, \end{aligned} \quad (2.19)$$

subject to the constraints (2.16) and (2.17).

The  $SO(n)$  Killing vectors on the quotient space can also be easily computed. On  $\mathbb{H}^n$ , the Killing vectors are simply  $k^M_{I\pm} = (T_I)^M_N z_{\pm}^N$ , but after the quotient we have to add local compensating  $SU(2)$  transformations to preserve the gauge conditions (2.16). We find

$$\begin{aligned} k^M_{I+} &= (T_I)^M_N z_+^N + (T_I)^n_N z_+^N \frac{z_-^{n-2} z_+^M - z_-^M}{z_-^n} - (T_I)^{n-2}_N z_+^N z_+^M, \\ k^M_{I-} &= (T_I)^M_N z_-^N - (T_I)^n_N \frac{z_+^N z_-^{n-2} z_-^M}{z_-^n} - (T_I)^{n-1}_N \frac{z_-^N z_+^M}{z_+^{n-1}} + (T_I)^{n-2}_N z_+^N z_-^M. \end{aligned} \quad (2.20)$$

One can verify that they preserve (2.16) as well as (2.17).

These Killing vectors and moment maps form the main ingredients of the scalar potential on the HKC. Performing the superconformal quotient leads to the Killing vectors and moment maps on  $Y(n-4)$ , which can be compared to those given in [21].

To end this section, we discuss the non-compact case; the QK space is now

$$\tilde{Y}(n-4) \equiv \frac{SO(n-4, 4)}{SO(n-4) \times SO(4)} . \quad (2.21)$$

The above formulae are modified by putting in a pseudo-Riemannian metric in the inner products: one starts with a metric  $\eta_{M\bar{N}}$  on  $\mathbb{H}^{n-4,4}$  of signature  $\eta_{M\bar{N}} = \text{diag}(- - - - + \cdots +)$ . The metric then used in the products  $z^a \eta_{a\bar{b}} \bar{z}^b$  on the HKC of  $\tilde{Y}(n-4)$  has signature  $\eta_{a\bar{b}} = \text{diag}(- + \cdots +)$ , such that, in our conventions, the metric on the QK space is positive definite.<sup>5</sup>

### 3 The Special Cases

In this section, we give more details about the  $SU(2)$  gauge fixing, and compare our results with the known cases for  $n = 4, 5, 6$ . It is convenient to introduce the  $SU(2)$  gauge invariant variables (for all  $n$ ),

$$z^{MN} \equiv z_+^M z_-^N - z_+^N z_-^M . \quad (3.1)$$

The hyperkähler potential on the quotient (2.15) is then, for all  $n$ ,

$$\chi_{\text{quotient}} = \sqrt{2 z^{MN} \bar{z}^{MN}} , \quad (3.2)$$

and is manifestly  $SU(2)$  gauge invariant. The variables (3.1) satisfy a Bianchi-like identity,

$$z^{MN} z^{PQ} + z^{MQ} z^{NP} + z^{MP} z^{QN} = 0 , \quad (3.3)$$

and the holomorphic constraints (2.9) imply

$$z^{MN} z^{NP} = 0 . \quad (3.4)$$

We show below how these constraints can be solved for  $n = 4, 5, 6$ .

#### 3.1 $n = 4$

In this case the quaternion-Kähler manifold is just one point, and so the hyperkähler cone above it must be  $\mathbb{R}^4$ . The gauge choice (2.16) does not give the canonical quadratic hyperkähler potential on  $\mathbb{C}^2$ , so we look for a different gauge. For  $n = 4$ , any antisymmetric

---

<sup>5</sup>This differs by an overall sign from the conventions of [11, 17].



matrix can be decomposed into its selfdual and anti-selfdual part. Notice then that (3.3) and (3.4) are solved by

$$z^{MN} = \pm \epsilon^{MNPQ} z^{PQ} , \quad (3.5)$$

so that we can parametrize

$$z^{MN} = u^i (\bar{\sigma})_{ij}^{MN} u^j . \quad (3.6)$$

Here,  $u^1, u^2$  are two complex coordinates and  $(\bar{\sigma})_{ij}^{MN}$  are the standard selfdual matrices<sup>6</sup>.

Using sigma-matrix algebra, it is easy to check that the constraints (3.3) and (3.4) are satisfied and the hyperkähler potential becomes simply

$$\chi = 4 \sum_{i=1}^2 u^i \bar{u}^i , \quad (3.7)$$

which is the Kähler potential of  $\mathbb{C}^2$  with flat metric.

The gauge choice (3.6) for the bilinears  $z^{MN}$  implies

$$z_+ = (u, v, iu, iv) , \quad z_- = (iv, -iu, v, -u) . \quad (3.8)$$

with  $u^1 = u$  and  $u^2 = v$ . Notice that this  $SU(2)$  gauge solves *all* the constraints (2.8) and (2.9).

### 3.2 $n = 5$

It is well-known that

$$Y(1) = \frac{SO(5)}{SO(4)} = \frac{Sp(2)}{Sp(1) \times Sp(1)} = \mathbb{HP}(1) , \quad (3.9)$$

and hence the HKC above it must be  $\mathbb{H}^2 = \mathbb{R}^8$ . The gauge (2.16) again does not give the canonical, quadratic hyperkähler potential on  $\mathbb{C}^4$ , so we search for a more suitable gauge.

The coordinates  $z_{\pm}^M$  transform in the vector representation of  $SO(5)$ :

$$z_{\pm}^M \rightarrow O^M{}_N z_{\pm}^N , \quad M, N = 1, \dots, 5 , \quad (3.10)$$

We rewrite this in  $Sp(2)$  notation by using the Clifford algebra in four Euclidean dimensions, where all gamma matrices can be taken hermitian. We define  $4 \times 4$  matrices

$$(v_{\pm})_{ij} \equiv -\frac{1}{2} z_{\pm}^M \gamma^M{}_i{}^k \mathcal{C}_{kj} \quad i, j = 1, \dots, 4 , \quad (3.11)$$

---

<sup>6</sup>We use the conventions  $\bar{\sigma}^{MN} = \frac{1}{2}(\bar{\sigma}^M \sigma^N - \bar{\sigma}^N \sigma^M)$  with  $\sigma^M = (\vec{\tau}, i)$  and  $\bar{\sigma}^M = (\vec{\tau}, -i)$ . The  $(\vec{\tau})$  are the three Pauli matrices and  $i, j$  indices are raised and lowered with  $\epsilon_{ij}$  such that  $(\bar{\sigma})_{ij}^{MN}$  is symmetric. We could also choose the anti-selfdual matrices  $\sigma^{MN} = \frac{1}{2}(\sigma^M \bar{\sigma}^N - \sigma^N \bar{\sigma}^M)$ , but this leads to equivalent results.

where  $\gamma^5 = \gamma^1\gamma^2\gamma^3\gamma^4$  and  $\mathcal{C}$  is the charge conjugation matrix, which can be taken antisymmetric and with square  $-1$ , such that  $(\gamma^M\mathcal{C})_{ij}$  is antisymmetric.  $\mathcal{C}$  serves as an  $Sp(2)$  metric and can be used to lower and raise  $(i, j)$  indices. Hence the matrices  $v_{\pm}$  are antisymmetric; the relation  $(\mathcal{C}^{-1})^{ij}v_{\pm ij} = 0$ , then leaves five independent components. We can invert (3.11),

$$z_{\pm}^M = -\frac{1}{2}\text{Tr}\left(\gamma^M v_{\pm}\mathcal{C}^{-1}\right). \quad (3.12)$$

Now we introduce  $SU(2)$  gauge invariant variables analogous to (3.1):

$$w_i{}^j \equiv \left(v_+\mathcal{C}^{-1}v_-\mathcal{C}^{-1} - v_-\mathcal{C}^{-1}v_+\mathcal{C}^{-1}\right). \quad (3.13)$$

The matrix  $w_{ij} \equiv (w\mathcal{C})_{ij}$  is symmetric; the relation to (3.1) is

$$w = \frac{1}{4}z^{MN}\gamma^{MN}. \quad (3.14)$$

It is straightforward to show that the hyperkähler potential (2.15) becomes

$$\chi = \sqrt{\text{Tr}(ww^\dagger)}, \quad (3.15)$$

up to an irrelevant overall factor. In these variables, the moment map constraints (2.9) are

$$z_{\pm} \cdot z_{\pm} = \text{Tr}(v_{\pm}\mathcal{C}^{-1}v_{\pm}\mathcal{C}^{-1}) = 0, \quad z_+ \cdot z_- = \text{Tr}(v_+\mathcal{C}^{-1}v_-\mathcal{C}^{-1}) = 0. \quad (3.16)$$

Imposing them, one can check that the matrix  $ww$  is zero as it should be according to (3.4).

Since we know that the HKC is flat, there should be coordinates  $u_i; i = 1, \dots, 4$ , in which the hyperkähler potential is

$$\chi = \sum_{i=1}^4 u_i \bar{u}_i. \quad (3.17)$$

If we can find a gauge for  $v_{\pm}$ , satisfying (3.16), such that

$$w_{ij} = u_i u_j, \quad (3.18)$$

then (3.15) reduces to (3.17) and we have proven that (2.15) gives the correct result.

This can be achieved by choosing the three complex  $SU(2)$  gauge conditions

$$v_{+14} = v_{-13}, \quad v_{+13} = v_{-14} = 0, \quad (3.19)$$

and identifying

$$v_{+14} = v_{-13} = u_1, \quad v_{+12} = -u_3, \quad v_{-12} = u_4. \quad (3.20)$$

Then we solve the moment map constraints (3.16) for  $v_{+23}$ ,  $v_{-24}$  and  $v_{-23}$ , and  $w_{ij} = u_i u_j$  for  $v_{+24}$ . This gives

$$v_{+23} = \frac{u_3^2}{u_1}, \quad v_{+24} = \frac{u_3 u_4}{u_1} + u_2, \quad v_{-23} = -\frac{u_3 u_4}{u_1} + u_2, \quad v_{-24} = -\frac{u_4^2}{u_1}. \quad (3.21)$$

Using (3.12), (3.19), (3.20) and (3.21) it is straightforward to find the coordinates  $z_{\pm}$  in terms of the flat coordinates  $u_i$  for any explicit representation of the Clifford algebra. For example,

$$z_+ = \left( -2u_3, \quad -\frac{i u_3^2}{u_1} - i u_1, \quad -\frac{u_3^2}{u_1} + u_1, \quad \frac{u_3 u_4}{u_1} + u_2, \quad -\frac{i u_3 u_4}{u_1} - i u_2 \right),$$

$$z_- = \left( 2u_4, \quad \frac{i u_3 u_4}{u_1} - i u_2, \quad \frac{u_3 u_4}{u_1} - u_2, \quad -\frac{u_4^2}{u_1} + u_1, \quad \frac{i u_4^2}{u_1} + i u_1 \right).$$

One can easily check that these  $z_{\pm}^M$ 's satisfy (2.9) and give  $\chi = \sum_{i=1}^4 u_i \bar{u}_i$ .

### 3.3 $n = 6$

Recall that

$$X(2) = \frac{SU(4)}{SU(2) \times U(2)}, \quad Y(2) = \frac{SO(6)}{SO(2) \times SO(4)}, \quad (3.22)$$

and hence  $X(2) = Y(2)$  and their HKC's must coincide. The hyperkähler potential of the HKC of  $X(n)$  was obtained in [11] by a  $U(1)$  hyperkähler quotient of  $\mathbb{H}^{n+2}$ . Denoting complex coordinates of  $\mathbb{H}^4 = \mathbb{C}^8$  by  $w_+^i$  and  $w_-^i$ ,  $i = 1, \dots, 4$ , transforming in conjugate representations of  $SU(4)$ , we have

$$\chi(X) = 2\sqrt{w_+^i \bar{w}_+^i w_-^j \bar{w}_-^j}, \quad \text{where} \quad w_-^4 = 1 \quad w_+^4 = -w_+^a w_-^a \quad a = 1, 2, 3. \quad (3.23)$$

To compare the HKC of  $Y(2)$  with (3.23), we follow the same strategy as for  $n = 5$ , namely we make use of the local isomorphism  $SU(4) = SO(6)$ . To this end we define  $U(1)$  gauge invariant (and traceless) matrix

$$w^i_j = w_+^i w_-^j \quad i, j = 1, \dots, 4, \quad (3.24)$$

and write (3.23) in matrix notation as

$$\chi(X) = 2\sqrt{\text{Tr}(w w^\dagger)}. \quad (3.25)$$

On the  $Y(2)$ -side, the coordinates  $z_{\pm}^M$ ,  $M = 1, \dots, 6$  are in the vector representation of  $SO(6)$ , similarly to (3.10). As in the  $n = 5$  case, we rewrite the hyperkähler potential (2.15) in  $SU(4)$  notation. Then we make gauge choices and solve the holomorphic moment map constraints such that (2.15) coincides with (3.25).

We introduce a basis of selfdual and anti-selfdual four by four matrices  $\eta, \bar{\eta}$  and write

$$v_{\pm ij} \equiv z_{\pm}^a \eta_{ij}^a + i z_{\pm}^{a+3} \bar{\eta}_{ij}^a z, \quad (3.26)$$

where  $a = 1, 2, 3$  and  $\eta^a$  and  $\bar{\eta}^a$  are selfdual and anti-selfdual respectively, see *e.g.*, [22]. We can raise and lower pairs of indices using  $\epsilon^{ijkl}$ ,

$$*v_{\pm}^{ij} \equiv \frac{1}{2} \epsilon^{ijkl} v_{\pm kl}, \quad (3.27)$$

such that

$$z_{\pm}^a = \frac{1}{4} \eta_{ij}^a *v_{\pm}^{ij}, \quad z_{\pm}^{a+3} = \frac{i}{4} \bar{\eta}_{ij}^a *v_{\pm}^{ij}. \quad (3.28)$$

The  $SU(2)$  gauge invariant variables can be put in the traceless matrix

$$v^i_j \equiv *v_+^{ik} v_{-kj} - *v_-^{ik} v_{+kj}, \quad (3.29)$$

and the hyperkähler potential is then, up to a numerical factor,

$$\chi(Y(2)) = \sqrt{\text{Tr}(v v^\dagger)}. \quad (3.30)$$

The holomorphic moment map constraints (2.9) take the form

$$\text{Tr}(v_+ * v_+) = 0 \quad \text{Tr}(v_- * v_-) = 0 \quad \text{Tr}(v_+ * v_-) = 0. \quad (3.31)$$

We must fix a gauge in which the matrices  $v^i_j$  and  $w^i_j$  of (3.24) coincide. This can be done by setting

$$v_{+13} = v_{-12} = 0, \quad v_{-13} = -\frac{1}{2}, \quad (3.32)$$

and identifying

$$v_{+12} = w_+^1, \quad v_{+23} = -w_+^3, \quad v_{-14} = \frac{1}{2} w_-^3. \quad (3.33)$$

We then solve (3.31) in such a way that the matrices  $w$  and  $v$  become equal. The answer is

$$\begin{aligned} v_{+14} &= -w_+^1 w_-^2, & v_{+24} &= w_+^1 w_-^2 + w_+^3 w_-^3, & v_{+34} &= -w_-^2 w_+^3, \\ v_{-23} &= -\frac{1}{2} \frac{w_+^2}{w_+^1}, & v_{-24} &= -\frac{1}{2} \frac{w_+^3 w_+^2}{w_+^1}, & v_{-34} &= -\frac{1}{2} w_-^2 - \frac{1}{2} \frac{w_+^2 w_-^2}{w_+^1}. \end{aligned} \quad (3.34)$$

Using (3.28) it is straightforward to find  $z_{\pm}^M$ ,  $M = 1, \dots, 6$  in terms of  $w_{\pm}^N$   $N = 1, \dots, 4$ .

## 4 The Dual Description of the HKC

As is well known, whenever there is an abelian isometry group one can find a dual description in terms of  $N = 2$  tensor multiplets. In this section, we consider the noncompact case, which is the one relevant for the low energy limit of string compactifications [3]. The isometry group of  $\tilde{Y}(n)$  is  $SO(n, 4)$  (see (1.3)). Since the coordinates  $z_+^M, z_-^M, M = 1, \dots, n+4$  transform in the vector representation of this group, from (2.15) it follows that the hyperkähler potential of the HKC of  $\tilde{Y}(n)$  is also invariant, as are the two-forms (2.3).

To find the generators of a maximal abelian subgroup of  $SO(n, 4)$  note that any two matrices  $T, T'$  of the form

$$T = \begin{pmatrix} 0 & 0 & a \\ 0 & 0 & a \\ a^t & -a^t & 0 \end{pmatrix}, \quad (4.1)$$

where  $a$  is an arbitrary  $(n+2)$ -dimensional row vector, commute with each other for any  $a, a'$ , and are nilpotent of third order. Such  $T$ 's are generators in the Lie algebra of the pseudo-orthogonal group that preserves the metric  $\eta = \text{diag}(- + \hat{\eta})$ , for any  $\hat{\eta}$ . If, for example, we arrange the coordinates as follows:  $(1, 5, 2, 3, \dots, n+4)$ , then the flat metric on  $\mathbb{H}^{n,4}$  has this form, and we can write the generators of an  $n+2$  dimensional abelian subgroup of  $SO(n, 4)$  as

$$(T_I)_M^N = (\delta_M^1 + \delta_M^5)\delta_I^N \sigma_I + \delta_{MI}(\delta_1^N - \delta_5^N), \quad I \neq 1, 5, \quad (4.2)$$

where  $\sigma_I$  is defined by  $\eta_{MN} = \delta_{MN}\sigma_M$ . It is easy to check that

$$(T_I)_M^L \eta_{LN} + (T_I)_N^L \eta_{LM} = 0. \quad (4.3)$$

We dualize with respect to the subgroup  $U(1)^{n+2}$  generated by the set  $\{T_I\}$  by starting with the flat space Lagrangian for  $\mathbb{H}^{n,4}$ , gauging both the  $n+2$  dimensional abelian symmetry found above as well as the  $SU(2)$  of Section 2, and constraining the  $U(1)$  gauge fields with  $N = 1$  tensor multiplet Lagrange multipliers  $G_I$ . We also have to solve the holomorphic moment map constraints for the action of the  $SU(2)$  (2.9) and those for the action of the  $U(1)^{n+2}$ :

$$z_{+M} z_{-N} \eta^{MK} (T_I)_K^N = v_I, \quad (4.4)$$

where the  $v_I$  are  $N = 1$  chiral superfields that together with  $G_I$  make up an  $N = 2$  tensor multiplet. Finally, we gauge fix both  $SU(2)$  and  $U(1)^{n+2}$ . The dual description is given by a Lagrangian which is a function of  $v, \bar{v}$  and the real coordinates  $G$ .

Let us see that all of the above requirements can be fulfilled consistently. Using (4.2), (4.4) becomes

$$(z_{+5} - z_{+1})z_{-I}\sigma_I + (z_{-1} - z_{-5})z_{+I}\sigma_I = v_I. \quad (4.5)$$

The  $SU(2)$  action on each pair  $(z_+^M, z_-^M)$  (2.5) is fixed completely by setting

$$z_{+1} = z_{-5} \equiv \phi, \quad z_{-1} = z_{+5} = 0. \quad (4.6)$$

The  $U(1)^{n+2}$  transformation rules on  $\mathbb{H}^{n,4}$ ,  $\delta z_{\pm M} = \lambda_I (T_I)_M^N z_{\pm N}$ , or explicitly

$$\delta z_{\pm 1} = \lambda_I z_{\pm I} \sigma_I, \quad \delta z_{\pm 5} = \lambda_I z_{\pm I} \sigma_I, \quad \delta z_{\pm I} = \lambda_I (z_{\pm 1} - z_{\pm 5}), \quad (4.7)$$

acquire, on the HKC of  $\tilde{Y}$ , compensating  $SU(2)$  transformations in order to preserve the gauge (4.6), similar to (2.20). They become

$$\begin{aligned} \delta z_{+1} &= \delta z_{-5} = \delta \phi = \frac{1}{2} \sigma_I \lambda_I (z_{+I} + z_{-I}), \\ \delta z_{+I} &= \lambda_I \phi - \frac{1}{2} \sigma_J \frac{z_{+I} \lambda_J (z_{+J} - z_{-J})}{\phi} - \sigma_J \frac{z_{-I} \lambda_J z_{+J}}{\phi}, \\ \delta z_{-I} &= -\lambda_I \phi - \frac{1}{2} \sigma_J \frac{z_{-I} \lambda_J (z_{+J} - z_{-J})}{\phi} - \sigma_J \frac{z_{+I} \lambda_J z_{-J}}{\phi}. \end{aligned} \quad (4.8)$$

The holomorphic  $SU(2)$  moment map constraints (2.9) reduce to

$$\phi^2 = z_{+I} z_{+J} \eta^{IJ}, \quad \phi^2 = -z_{-I} z_{-J} \eta^{IJ}, \quad 0 = z_{+I} z_{-J} \eta^{IJ}. \quad (4.9)$$

We choose a  $U(1)^{n+2}$  gauge

$$z_{+I} = 1 \quad \text{for all } I \neq 1, 5. \quad (4.10)$$

Then (4.9) become

$$\phi^2 = n - 4, \quad \phi^2 = -z_{-I}^2 \sigma_I, \quad z_{-I} \sigma_I = 0, \quad (4.11)$$

which we can solve for  $\phi$  and two of the  $z_{-I}$ 's. The  $U(1)^{n+2}$  holomorphic moment map (4.5) after substituting (4.6), (4.10) and the solution for  $\phi$  is

$$-\sqrt{n-4} (z_{-I} + 1) \sigma_I = v_I. \quad (4.12)$$

As a consequence of (4.11), there are  $n$  independent  $v$ 's.

The Lagrangian of the dual theory is

$$\sum_{M,N=1}^{n+4} \sigma_N (e^{i \sum_I T_I V_I})_{MN} (\bar{z}_+^M \bar{z}_-^M) \left( e^V \right) \begin{pmatrix} z_+^N \\ z_-^N \end{pmatrix} - \sum_{M \neq 1,5} \sigma_M G_M V_M, \quad (4.13)$$

after we eliminate all gauge fields in (4.13) via their equations of motion and impose the conditions (4.6), (4.10) as well as the solutions to (4.11), (4.12). Thus the dual description is in terms of  $n$   $N = 1$  chiral superfields  $v$  and  $n + 2$   $N = 1$  tensor multiplets  $G$ .

Unfortunately we haven't been able to solve for all gauge fields ( $V$  and  $V_I$ ) explicitly; and thus we have not been able to construct the dual Lagrangian. However the analysis that we made above shows that there is a consistent solution of all requirements (constraints and allowed gauge choices) for the existence of a dual description. In  $N = 2$  language,  $n$  of the  $G$ 's combine with the  $n$   $v$ 's into  $n$   $N = 2$  tensor multiplets and the remaining two  $G$ 's combine into a double-tensor multiplet. This is consistent with the expectations from type IIB compactifications [4]. There the double-tensor multiplet appears naturally: it contains the dilaton, axion and the RR and NS two-forms.

Though we cannot solve for all gauge fields simultaneously, we can easily solve for either the  $SU(2)$  prepotential  $V$  or all  $V_I$ 's that gauge the  $U(1)^{n+2}$ . Eliminating the first gives the same answer for the dual Lagrangian as (2.15), but with  $U(1)^{n+2}$  gauged as above. On the other hand, integrating out the  $U(1)^{n+2}$  gauge fields gives us a dual description, although the result is still a function of the  $SU(2)$  gauge fields, which for convenience we now write as  $U = e^V$ . Using the nilpotency properties of the generators (and arranging the indices as  $1, 5, 2, \dots, n + 4$ ),

$$e^{i\sum_{I\neq 1,5} T_I V_I} = 1_{(n+4)\times(n+4)} + i \begin{pmatrix} 0 & 0 & \sigma_2 V_2 & \sigma_3 V_3 & \dots \\ 0 & 0 & \sigma_2 V_2 & \sigma_3 V_3 & \dots \\ V_2 & -V_2 & 0 & 0 & \dots \\ V_3 & -V_3 & 0 & 0 & \dots \\ \vdots & \vdots & \vdots & \vdots & \vdots \end{pmatrix} + \frac{1}{2} \begin{pmatrix} \sum_{I\neq 1,5} \sigma_I V_I^2 & -\sum_{I\neq 1,5} \sigma_I V_I^2 & 0 & 0 & \dots \\ \sum_{I\neq 1,5} \sigma_I V_I^2 & -\sum_{I\neq 1,5} \sigma_I V_I^2 & 0 & 0 & \dots \\ 0 & 0 & 0 & 0 & \dots \\ 0 & 0 & 0 & 0 & \dots \\ \vdots & \vdots & \vdots & \vdots & \vdots \end{pmatrix}, \quad (4.14)$$

we can rewrite (4.13) as

$$\bar{Z}^\alpha M_{\alpha\beta} \sigma_\beta Z^\beta - \sigma_I G_I V_I, \quad \alpha, \beta = 1, \dots, 2(n+4). \quad (4.15)$$

We have denoted  $\bar{Z} = (\bar{z}_+^1, \bar{z}_-^1, \bar{z}_+^5, \bar{z}_-^5, \bar{z}_+^2, \bar{z}_-^2, \dots, \bar{z}_+^{n+4}, \bar{z}_-^{n+4})$  and

$$M = \begin{pmatrix} (1 - \frac{1}{2} \sum_I \sigma_I V_I^2) U & (\frac{1}{2} \sum_I \sigma_I V_I^2) U & i\sigma_2 V_2 U & i\sigma_3 V_3 U & \dots \\ (-\frac{1}{2} \sum_I \sigma_I V_I^2) U & (1 + \frac{1}{2} \sum_I \sigma_I V_I^2) U & i\sigma_2 V_2 U & i\sigma_3 V_3 U & \dots \\ iV_2 U & -iV_2 U & U & 0 & \dots \\ iV_3 U & -iV_3 U & 0 & U & \dots \\ \vdots & \vdots & \vdots & \vdots & \vdots \end{pmatrix}, \quad (4.16)$$

where  $M$  is a  $(n + 4) \times (n + 4)$  matrix of  $2 \times 2$  matrices. The solution for the  $U(1)^{n+2}$  gauge fields is

$$V_I = \frac{G_I - i(\bar{z}_+^1 + \bar{z}_+^5, \bar{z}_-^1 + \bar{z}_-^5) U \begin{pmatrix} z_+^I \\ z_-^I \end{pmatrix} + i(\bar{z}_+^I, \bar{z}_-^I) U \begin{pmatrix} z_+^1 + z_+^5 \\ z_-^1 + z_-^5 \end{pmatrix}}{(\bar{z}_+^1 - \bar{z}_+^5, \bar{z}_-^1 - \bar{z}_-^5) U \begin{pmatrix} z_+^1 - z_+^5 \\ z_-^1 - z_-^5 \end{pmatrix}}. \quad (4.17)$$

Finally we note that if we dualize only  $n + 1$  instead of all  $n + 2$  generators of  $U(1)^{n+2}$ , we find  $n + 1$   $N = 1$  tensor multiplets and  $n + 1$   $N = 1$  chiral superfields, or equivalently,  $n + 1$   $N = 2$  tensor multiplets.

## 5 Half-hypermultiplets and $N = 2$ Superspace

We close with a few comments on other representations of the HKC. The half-hypermultiplet formulation that we have used so far is not easily described in  $N = 2$  superspace; however, a half-hypermultiplet has an alternative formulation as the double cover of an  $A_1$  singularity, that is, a  $U(1)$  hyperkähler quotient of a full hypermultiplet doublet with vanishing Fayet-Iliopoulos terms.

Explicitly, consider a hypermultiplet doublet  $z_\pm^{(1)}, z_\pm^{(2)}$ , with  $z_+$  and  $z_-$  in conjugate representations of a  $U(1)$  gauge group with an  $N = 1$  gauge prepotential  $V_0$  as well as the  $SU(2)$  gauge group of the previous sections with an  $N = 1$  gauge prepotential  $V$ :

$$\chi = e^{V_0} \begin{pmatrix} \bar{z}_+^{(1)} & \bar{z}_+^{(2)} \end{pmatrix} \begin{pmatrix} e^V \end{pmatrix} \begin{pmatrix} z_+^{(1)} \\ z_+^{(2)} \end{pmatrix} + e^{-V_0} \begin{pmatrix} z_-^{(1)} & z_-^{(2)} \end{pmatrix} \begin{pmatrix} e^{-V} \end{pmatrix} \begin{pmatrix} \bar{z}_-^{(1)} \\ \bar{z}_-^{(2)} \end{pmatrix}. \quad (5.1)$$

This must be supplemented by the holomorphic moment map constraints; in particular, the holomorphic  $U(1)$  moment map is [11]

$$z_+^{(1)} \bar{z}_-^{(1)} + z_+^{(2)} \bar{z}_-^{(2)} = 0. \quad (5.2)$$

We shall see that integrating out  $V_0$ , imposing (5.2), and choosing a convenient gauge for the  $U(1)$  symmetry gives us back the half-hypermultiplet formulation.

Integrating out the  $U(1)$  gauge field  $V_0$  gives [11]

$$2 \sqrt{\begin{pmatrix} \bar{z}_+^{(1)} & \bar{z}_+^{(2)} \end{pmatrix} \begin{pmatrix} e^V \end{pmatrix} \begin{pmatrix} z_+^{(1)} \\ z_+^{(2)} \end{pmatrix} \begin{pmatrix} z_-^{(1)} & z_-^{(2)} \end{pmatrix} \begin{pmatrix} e^{-V} \end{pmatrix} \begin{pmatrix} \bar{z}_-^{(1)} \\ \bar{z}_-^{(2)} \end{pmatrix}} \quad (5.3)$$

We can choose a  $U(1)$  gauge

$$z_+^{(2)} = z_-^{(1)} \equiv z_-. \quad (5.4)$$



Then the holomorphic  $U(1)$  moment map (5.2) implies

$$z_-^{(2)} = -z_+^{(1)} \equiv -z_+ . \quad (5.5)$$

Using (5.4) and (5.5) we express everything in (5.3) in terms of  $z_\pm$ :

$$2 \sqrt{(\bar{z}_+ \ \bar{z}_-) \begin{pmatrix} e^V \\ \end{pmatrix} \begin{pmatrix} z_+ \\ z_- \end{pmatrix} (z_- \ -z_+) \begin{pmatrix} e^{-V} \\ \end{pmatrix} \begin{pmatrix} \bar{z}_- \\ -\bar{z}_+ \end{pmatrix}} . \quad (5.6)$$

The second factor in the square root can be rewritten as

$$\begin{aligned} (z_- \ -z_+) \begin{pmatrix} e^{-V} \\ \end{pmatrix} \begin{pmatrix} \bar{z}_- \\ -\bar{z}_+ \end{pmatrix} &= (\bar{z}_- \ -\bar{z}_+) \begin{pmatrix} e^{-V^T} \\ \end{pmatrix} \begin{pmatrix} z_- \\ -z_+ \end{pmatrix} \\ &= (\bar{z}_+ \ \bar{z}_-) \sigma_2 \begin{pmatrix} e^{-V^T} \\ \end{pmatrix} \sigma_2 \begin{pmatrix} z_+ \\ z_- \end{pmatrix} \\ &= (\bar{z}_+ \ \bar{z}_-) e \begin{pmatrix} -\sigma_2 V^T \sigma_2 \\ \end{pmatrix} \begin{pmatrix} z_+ \\ z_- \end{pmatrix} . \end{aligned} \quad (5.7)$$

Since  $V$  is an  $SU(2)$  gauge prepotential, we write it as  $V = V^i \sigma_i$ ,  $i = 1, 2, 3$ , where  $\sigma_i$  are the Pauli matrices, and

$$\sigma_2 \sigma_i^T \sigma_2 = -\sigma_i . \quad (5.8)$$

Hence (5.7) is equal to the first factor under the square root in (5.6) and we confirm that (5.1) is the same as (2.10) (up to an insignificant factor of 2).

To complete the proof of the equivalence of (5.1) and (2.10), we must also compare the  $SU(2)$  holomorphic moment maps. For (5.1) the  $SU(2)$  moment map is [18]

$$\begin{pmatrix} z_-^{(1)} & z_-^{(2)} \end{pmatrix} T_i \begin{pmatrix} z_+^{(1)} \\ z_+^{(2)} \end{pmatrix} = 0 , \quad (5.9)$$

where  $T_i$  are the same generators as (2.12). Using (5.4) and (5.5) to express  $z_\pm^{(1,2)}$  in terms of  $z_\pm$ , we get the same constraints as (2.9).

Having this description in hand, we can now give the  $N = 2$  projective superspace formulation; we use the polar multiplet formulation (for a pedagogical summary as well as a definition of our notation, see the appendix B of [11]; earlier references are contained therein and include [13]). The HKC for  $Y(n-4)$  has the projective superspace Lagrangian

$$L_{N=2} = \oint \frac{d\xi}{2\pi i} \sum_{i=1}^n e^{\mathbf{V}_i \tilde{\Upsilon}_i} e^{\mathbf{V}} \cdot \Upsilon_i , \quad (5.10)$$

where each  $\Upsilon_i$  is a doublet of polar multiplets, each  $e^{\mathbf{V}_i}$  is a tropical multiplet describing a  $U(1)$  gauge prepotential (analogous to the  $N = 1$  gauge prepotential), and  $e^{\mathbf{V}}$  is a hermitian unimodular tropical multiplet describing the  $SU(2)$  gauge prepotential. Formally, we may integrate out the latter and obtain:

$$L_{N=2} = \oint \frac{d\xi}{2\pi i} 2 \left( \det \left[ \sum_{i=1}^n \Upsilon_i \otimes e^{\mathbf{V}_i} \bar{\Upsilon}_i \right] \right)^{\frac{1}{2}} . \quad (5.11)$$

The full hypermultiplet description can also be written as a quiver; indeed, the quiver for the HKC of  $Y(1)$  was explored in quite some detail in the last section of [14]. The general quiver is a higher dimensional extension of (a cover of) the orbifold limit of the  $D_4$  ALE space, and is shown in Figure 2: The observation that the HKC of  $Y(1)$  is flat




Figure 2: The quiver describing the HKC of  $Y(n-4)$ . Each node with a label  $k$  represents a factor  $U(k)$  in the quotient group and each link is a hypermultiplet transforming in the bifundamental of the groups at the nodes it connects. An overall  $U(1)$  acts trivially, and in our particular example, can conveniently be factored out by reducing the  $U(2)$  at the central node to  $SU(2)$ .

implies that the specific quiver discussed in [14], when the orbifold singularity is blown up (rather than removed by going to the cover) gives rise to a genuine 8-dimensional ALE manifold that has not been studied before. However, the higher dimensional analogs cannot be ALE, as the limit without any blowing up is not flat.

## Acknowledgements

SV thanks B. de Wit and M. Trigiante for stimulating discussions. MR thanks the NSF for partial support under grant NSF PHY-0098527.

## References

- [1] J. Bagger, E. Witten, *Matter couplings in  $N = 2$  supergravity*, Nucl. Phys. **B222** (1983) 1.
- [2] A. Wolf, J. Math. Mech. **14** (1965) 1033;  
D. Alekseevskii, Funct. Anal. Appl. **2** (1968) 11, Math. USSR-Izv. **9** (1975) 297.
- [3] S. Cecotti, S. Ferrara, L. Girardello, *Geometry of type II superstrings and the moduli of superconformal field theories*, Int. J. Mod. Phys. **A4** (1989) 2475;  
S. Ferrara and S. Sabharwal, *Quaternionic manifolds for type II superstring vacua of Calabi-Yau spaces*, Nucl. Phys. **B332** (1990) 317.
- [4] R. Böhm, H. Günther, C. Herrmann, J. Louis, *Compactification of type IIB string theory on Calabi-Yau threefolds*, Nucl. Phys. **B569** (2000) 229; hep-th/9908007.
- [5] B. de Wit, P. G. Lauwers, R. Philippe, S. Q. Su and A. Van Proeyen, *Gauge and matter fields coupled to  $N=2$  supergravity*, Phys. Lett. **B134** (1984) 37;  
B. de Wit and A. Van Proeyen, *Potentials and symmetries of general gauged  $N=2$  supergravity - Yang-Mills models*, Nucl. Phys. **B245** (1984) 89.
- [6] M. Gunaydin, G. Sierra and P. K. Townsend, *The geometry of  $N=2$  Maxwell-Einstein supergravity and Jordan algebras*, Nucl. Phys. **B242** (1984) 244.
- [7] A. Swann, *Hyperkähler and quaternionic Kähler geometry*, Math. Ann. **289** (1991) 421.
- [8] G. W. Gibbons, P. Rychenkova, *Cones, tri-Sasakian structures and superconformal invariance*, Phys. Lett. **B443** (1998) 138, hep-th/9809158.
- [9] K. Galicki, *Geometry of the scalar couplings in  $N=2$  supergravity models*, Class. Quantum Grav. **9** (1992) 27.
- [10] B. de Wit, B. Kleijn and S. Vandoren, *Superconformal Hypermultiplets*, Nucl. Phys. **B568** (2000) 475, hep-th/9909228.
- [11] B. de Wit, M. Roček, S. Vandoren, *Hypermultiplets, hyperkähler cones and quaternion-Kähler geometry*, JHEP 0102 (2001) 039, hep-th/0101161.
- [12] K. Galicki, *A Generalization Of The Momentum Mapping Construction For Quaternionic Kahler Manifolds*, Commun. Math. Phys. **108** (1987) 117.

- [13] U. Lindström and M. Roček, *New Hyperkahler Metrics And New Supermultiplets*, Commun. Math. Phys. **115**, 21 (1988).  
 F. Gonzalez-Rey, M. Roček, S. Wiles, U. Lindström and R. von Unge, *Feynman rules in  $N = 2$  projective superspace. I: Massless hypermultiplets*, Nucl. Phys. B **516**, 426 (1998) [arXiv:hep-th/9710250].
- [14] U. Lindström, M. Roček and R. von Unge, *Hyperkahler quotients and algebraic curves*, JHEP **0001**, (2000) 022, hep-th/9908082.
- [15] A. Maciocia, *Metrics On The Moduli Spaces Of Instantons Over Euclidean Four Space*. Commun. Math. Phys. **135** (1991) 467;  
 C. Boyer, B. Mann, Proc. of Symp. in Pure Math. **54** part 2 (1993) 45;
- [16] S. Vandoren, *Instantons and quaternions*, Proc. of the TMR conference “Nonperturbative Quantum Effects”, Paris 2000, eds. D. Bernard et al., hep-th/0009150.
- [17] B. de Wit, M. Roček, S. Vandoren, *Gauging isometries on hyperkahler cones and quaternion-Kähler manifolds*, Phys. Lett. **B511** (2001) 302, hep-th/0104215.
- [18] N. Hitchin, A. Karlhede, U. Lindström, M. Roček, *Hyperkahler metrics and supersymmetry*, Commun. Math. Phys. **108** (1987) 535.
- [19] U. Lindström and M. Roček, *Scalar Tensor Duality And  $N=1$ ,  $N=2$  Nonlinear Sigma Models*, Nucl. Phys. B **222**, 285 (1983).
- [20] P. Breitenlohner, M. Sohnius, *Matter couplings and non-linear sigma models in  $N=2$  supergravity*, Nucl. Phys. **B187** (1981), 409.
- [21] P. Fre, L. Girardello, I. Pesando and M. Trigiante, *Spontaneous  $N=2 \rightarrow N=1$  local supersymmetry breaking with surviving compact gauge groups*, Nucl. Phys. **B493** (1997) 231, hep-th/9607032.
- [22] G. 't Hooft, *Computation of the quantum effects due to a four-dimensional pseudoparticle*, Phys. Rev. **D14** (1976) 3432; *ibid.* (E) **D18** (1978) 2199.