

DEL PEZZO MODULI VIA ROOT SYSTEMS

ELISABETTA COLOMBO, BERT VAN GEEMEN, AND EDUARD LOOIJENGA

ABSTRACT. Coble defined in his 1929 treatise invariants for cubic surfaces and quartic curves. We reinterpret these in terms of the root systems of type E_6 and E_7 that are naturally associated to these varieties, thereby giving some of his results a more intrinsic treatment. Our discussion is uniform for all Del Pezzo surfaces of degree 2,3,4 and 5.

To Professor Yuri Manin, for his 70th birthday.

INTRODUCTION

A *Del Pezzo surface of degree d* is a smooth projective surface with semi-ample anticanonical bundle whose class has self-intersection d . The degree is always between 1 and 9 and the surface is either a quadric ($d = 8$ in that case) or is obtained from blowing up $9 - d$ points in the projective plane that satisfy a mild genericity condition. So moduli only occur for $1 \leq d \leq 4$. The anticanonical system is d -dimensional and when $d \neq 1$, it is also base point free. For $d = 4$, the resulting morphism is birational onto a complete intersection of two quadrics in \mathbb{P}^4 , for $d = 3$ it is birational onto a cubic surface in \mathbb{P}^3 and for $d = 2$ we get a degree two map onto \mathbb{P}^2 whose discriminant curve is a quartic (we will ignore the case $d = 1$ here). This image surface (resp. discriminant) is smooth in case the anticanonical bundle is ample; we then call the Del Pezzo surface a *Fano surface*. Otherwise it might have simple singularities in the sense of Arnol'd (that have a root system label A , D or E). Conversely, every complete intersection of two quadrics in \mathbb{P}^4 , cubic surface in \mathbb{P}^3 or quartic curve in \mathbb{P}^2 with such singularities thus arises.

We mentioned that a degree d Del Pezzo surface, $d \neq 8$, is obtained from blowing up $9 - d$ points in \mathbb{P}^2 in general position. A more precise statement is that if we are given as many disjoint exceptional curves E_1, \dots, E_{9-d} on the Del Pezzo surface S , then these can be simultaneously contracted to produce a projective plane. So the images of these curves yield $9 - d$ numbered points p_1, \dots, p_{9-d} in \mathbb{P}^2 given up to projective equivalence. Hence every polynomial expression in terms of the projective coordinates of these points that is invariant under $SL(3, \mathbb{C})$ is a ‘covariant’ for the tuple $(S; E_1, \dots, E_{9-d})$. Coble exhibited such covariants for the important cases $d = 2$ and $d = 3$. These are in general not covariants of S itself, since the surface may have many exceptional systems (E_1, \dots, E_{9-d}) . Indeed, if we assume that S is Fano, then, as Manin observed, these systems are simply transitively permuted by a Weyl group W , which acts here as a group of Cremona transformations.

Therefore, this group will act on the space of such covariants. Coble's covariants span a W -invariant subspace and Coble was able to identify the W -action as a Cremona group (although the Weyl group interpretation was not available to him). For $d = 3$ he found an irreducible representation of degree 10 of a E_6 -Weyl group and for $d = 2$ he obtained an irreducible representation of degree 15 of a E_7 -Weyl group.

The present paper purports to couch Coble's results in terms of a moduli space of tuples $(S; E_1, \dots, E_{9-d})$ as above, for which S is semistable in the sense of Geometric Invariant Theory. This moduli space comes with an action of the Weyl group W . It also carries a natural line bundle, called the *determinant bundle*, to which the W -action lifts: this line bundle assigns to a Del Pezzo surface the line that is the dual of the top exterior power of the space of sections of its anticanonical bundle. It turns out that this bundle is proportional to the one that we use to do geometric invariant theory with (and from which our notion of semistability originates). We show that the Coble covariants can be quite naturally understood as sections of this bundle and we reprove the fact known to Coble that these sections span an irreducible representation of W . We also show that these sections separate the points of the above moduli space so that one might say that Coble's covariants of a stable tuple $(S; E_1, \dots, E_{9-d})$ make up a complete set of invariants. This approach not only covers the cases Coble considered (degree 2 and 3), but also the degree 4 case and, somewhat amusingly, even the degree 5 case, for which there are no moduli at all. For the case of degree 3 we also make the connection with earlier work of Naruki and Yoshida. This allows us to conclude that the Coble covariants define a complete linear system and define a closed immersion of the GIT-compactification of the moduli space of marked cubic surfaces in a 9-dimensional projective space. Our results are less complete when the degree is 2; for instance, we did not manage to establish that the Coble covariants define a complete linear system.

We end up with a description of the GIT moduli space that is entirely in terms of the corresponding root system. Our results lead to us to some remarkable integrability properties of the module of W -invariant vector fields on the vector space that underlies the defining (reflection) representation of W and we raise the question of whether this is a special case of a general phenomenon.

Since the appearance of Coble's book a great deal of work on Del Pezzo moduli has seen the day. As its sheer volume makes it impossible to give our predecessors their fair due, any singling out of contributions will be biased. While keeping that in mind we nevertheless wish to mention the influential book by Manin [18], the Astérisque volume by Dolgachev-Ortland [10], Naruki's construction of a smooth compactification of the moduli space of marked cubic surfaces [16], the determination of its Chow groups in [5], the Lecture Note by Hunt [15] and Yoshida's revisit of the Coble covariants [19]. The ball quotient description of the moduli space of cubic surfaces by

Allcock, Carlson and Toledo [2], combined with Borchers' theory of modular forms, led Allcock and Freitag [1] to construct an embedding of the moduli space of marked cubic surfaces, which coincides with the map given by the Coble invariants [13], [14].

We now briefly review the organization of the paper. The first section introduces a moduli space for marked Fano surfaces of degree $d \geq 2$ as well as the line bundle over that space that is central to this paper, the determinant line bundle. This assigns to a Fano surface the determinant line of the dual of the space of sections of its anticanonical sheaf (this is also the determinant of the cohomology of its structure sheaf). We show that this line bundle can be used to obtain in a uniform manner a compactification (by means of GIT) so the determinant bundle extends over this compactification as an ample bundle. In Section 2 we introduce the Coble covariants and show that they can be identified with sections of the determinant bundle. The next section expresses these covariants purely in terms of the associated root system. In Section 4 we identify (and discuss) the Weyl group representation spanned by the Coble covariants. The final section investigates the separating properties of the Coble covariants, where the emphasis is on the degree 3 case.

As we indicated, Manin's work on Del Pezzo surfaces has steered this beautiful subject in a new direction. Although this represents only a small part of his many influential contributions to mathematics, we find it therefore quite appropriate to dedicate this paper to him on the occasion of his 70th birthday.

CONTENTS

Introduction	1
1. Moduli spaces for marked Del Pezzo surfaces	4
Completion of the moduli spaces by means of GIT	6
Degree 4 surfaces in projective 4-space	7
Cubic surfaces	9
Quartic curves	9
Completion of the moduli space of marked Fano surfaces	11
2. Coble's covariants	11
Action of the Weyl group on the Coble factors	13
The covariants	16
3. Anticanonical divisors with a cusp	18
Anticanonical cuspidal cubics on Del Pezzo surfaces	18
Anticanonical cuspidal cubics on Fano surfaces	20
The universal parabolic curve of a cubic surface	23
Moduli of degree 4 Del Pezzo surfaces	25
4. Coble's representations	26
Macdonald's irreducibility theorem	26
The Coble representation of a Weyl group of type E_6	27

The Coble representation of a Weyl group of type E_7	31
5. The Coble linear system	36
The Coble system in the degree four case	37
The Coble system in the degree three case	38
The Coble system in the degree two case	45
References	45

1. MODULI SPACES FOR MARKED DEL PEZZO SURFACES

We call a smooth complete surface S a *Del Pezzo surface of degree d* if its anticanonical bundle ω_S^{-1} is semi-ample and $\omega_S \cdot \omega_S = d$. It is known that then $1 \leq d \leq 9$ and that S is isomorphic either to a smooth quadric or to a surface obtained from successively blowing up $9 - d$ points of \mathbb{P}^2 . In order that a successive blowing up of $9 - d$ points of \mathbb{P}^2 yields a Del Pezzo surface it is necessary and sufficient that we blow up on (i.e., over the strict transform of) a smooth cubic curve (which is an anticanonical divisor of \mathbb{P}^2). This is equivalent to the apparently weaker condition that we blow up at most 3 times on a line and at most 6 times on a conic. It is also equivalent to the apparently stronger condition that the anticanonical system on this surface is nonempty and has dimension d .

Let S be a Del Pezzo surface of degree d . The vector space $V(S) := H^0(\omega_S^{-1})^*$ (which we will usually abbreviate by V) has dimension $d + 1$. The anticanonical system defines the (rational) *anticanonical map* $S \dashrightarrow \mathbb{P}(V)$. When $d \geq 2$, it has no base points, so that the anticanonical map is a morphism. For $d = 1$, it has a single base point; if we blow up this point, then the anticanonical map lifts to a morphism $\tilde{S} \rightarrow \mathbb{P}(V)$ which makes \tilde{S} a rational elliptic surface (with a section defined by the exceptional curve of the blowup).

Let $S \rightarrow \tilde{S}$ contract the (-2) -curves on S (we recall that a curve on a smooth surface is called a (-2) -*curve* if it is a smooth rational curve with self-intersection -2 .) Then \tilde{S} has rational double point singularities only and its dualizing sheaf $\omega_{\tilde{S}}$ is invertible and anti-ample. We shall call such a surface an *anticanonical surface*. If $d \geq 2$, then the anticanonical morphism factors through \tilde{S} . For $d \geq 3$ the second factor is an embedding of \tilde{S} in a projective space of dimension d ; for $d = 3$ this yields a cubic surface and for $d = 4$ a complete intersection of two quadrics. When $d = 2$, the second factor realizes \tilde{S} as a double cover of a projective plane ramified along a quartic curve with only simple singularities in the sense of Arnol'd (accounting for the rational double points on \tilde{S}).

Adopting the terminology in [11], we say that S is a *Fano surface* of degree d if ω_S^{-1} is ample (but beware that other authors call this a Del Pezzo surface). If S is given as a projective plane blown up in $9 - d$ points, then it is Fano precisely when the points in question are distinct, no three lie on

a line, no six lie on a conic and no eight lie on a cubic which has a singular point at one of them. This is equivalent to requiring that S contains no (-2) -curves.


From now on we assume that S is not isomorphic to a smooth quadric. We denote the canonical class of S by $k \in \text{Pic}(S)$ and its orthogonal complement in $\text{Pic}(S)$ by $\text{Pic}_0(S)$. An element $e \in \text{Pic}(S)$ is called an *exceptional class* of S if $e \cdot e = e \cdot k = -1$. Every exceptional class is representable by a unique effective divisor. A *marking* of S is an ordered $(9 - d)$ -tuple of exceptional classes (e_1, \dots, e_{9-d}) on $\text{Pic}(S)$ with $e_i \cdot e_j = -\delta_{ij}$. Given a marking, there is a unique class $\ell \in \text{Pic}(S)$ characterized by the property that $3\ell = -k + e_1 + \dots + e_{9-d}$ and $(\ell, e_1, \dots, e_{9-d})$ will be basis of $\text{Pic}(S)$. The marking is said to be *geometric* if S can be obtained by $(9 - d)$ - successive blowups of a projective plane in such a manner that e_i is the class of the total transform E_i of the exceptional curve of the i th blowup. An ℓ -marking of S consists of merely giving the class ℓ . So if \mathcal{L} is a representative line bundle, then \mathcal{L} is base point free and defines a birational morphism from S to a projective plane and the anticanonical system on S projects onto a d -dimensional linear system of cubic curves on this plane.

Since we are interested here in the moduli of Fano surfaces, we usually restrict to the case $d \leq 4$: if S is given as a blown up projective plane then four of the $9 - d$ points to be blown up can be used to fix a coordinate system, from which it follows that we have a fine moduli space $\mathcal{M}_{m,d}^o$ of marked Fano surfaces of degree d that is isomorphic to an affine open subset of $(\mathbb{P}^2)^{5-d}$.

From now on we assume that $d \leq 6$. With Manin we observe that then the classes $e_i - e_{i+1}$, $i = 1, \dots, 8 - d$ and $\ell - e_1 - e_2 - e_3$ make up a basis of $\text{Pic}_0(S)$ and can be thought of as a system of simple roots of a root system R_{9-d} . This root system is of type E_8, E_7, E_6, D_5, A_4 and $A_2 + A_1$ respectively. The roots that have fixed inner product with ℓ make up a single S_{9-d} -orbit and we label them accordingly:

- (0) $h_{ij} := e_i - e_j$, ($i \neq j$),
- (1) $h_{ijk} := \ell - e_i - e_j - e_k$ with i, j, k pairwise distinct,
- (2) $(2\ell - e_1 - e_2 - e_3 - e_4 - e_5 - e_6 - e_7) + e_i$, denoted h_i when $d = 2$,
For $d = 3$, this is only makes sense for $i = 7$ and we then may write h instead.
- (3) $-k - e_i$ ($d = 1$ only).

Notice that $h_{ij} = -h_{ji}$, but that in h_{ijk} the order of the subscripts is irrelevant.



The marking defined by (e_1, \dots, e_{9-d}) is geometric if and only if for every (-2) -curve C its intersection product with each of the simple roots is not positive.

The Weyl group $W(R_{9-d})$ is precisely the group of orthogonal transformations of $\text{Pic}(S)$ that fix k . It acts simply transitively on the markings. In particular it acts on $\mathcal{M}_{m,d}^\circ$ and the quotient variety $\mathcal{M}_d^\circ := W(R_{9-d}) \backslash \mathcal{M}_{m,d}^\circ$ can be interpreted as the coarse moduli space of Fano surfaces of degree d . The orbit space of $\mathcal{M}_{m,d}^\circ$ relative to the permutation group of the e_1, \dots, e_{9-d} (a Weyl subgroup of type A_{8-d}) is the moduli space $\mathcal{M}_{\ell,d}^\circ$ of ℓ -marked Fano surfaces of degree d .

Completion of the moduli spaces by means of GIT. Fix a 3-dimensional complex vector space A and a generator $\alpha \in \det(A)$. We think of α as a translation invariant 3-vector field on A . If $f \in \text{Sym}^3 A^*$ is a cubic form on A , then the contraction of α with df , $\iota_{df}\alpha$, is a 2-vector field on A that is invariant under scalar multiplication and hence defines a 2-vector field on $\mathbb{P}(A)$. We thus obtain an isomorphism between $\text{Sym}^3 A^*$ and $H^0(\omega_{\mathbb{P}(A)}^{-1})$.

Let $d \in \{2, 3, 4\}$. A $(d+1)$ -dimensional linear quotient V of $\text{Sym}^3 A$ defines a linear subspace $V^* \subset \text{Sym}^3 A^*$, i.e., a linear system of cubics on $\mathbb{P}(A)$ of dimension d . If we suppose that this system does not have a fixed component, then its base locus consists of $9-d$ points (multiplicities counted) and blowing up this base locus produces an ℓ -marked Del Pezzo surface S with the property that $H^0(\omega_S^{-1})^*$ can be identified with V (we excluded $d = 1$ here because then the base locus has 9 points and we get a rational elliptic surface). If we specify an order for the blowing up, then S is even geometrically marked. The quotient surface \bar{S} obtained from contracting (-2) -curves is more canonically defined as it can be described in terms of the rational map $\mathbb{P}(A) \dashrightarrow \mathbb{P}(V)$: for $d = 3, 4$ it is the image of this map and for $d = 2$ the Stein factorization realizes \bar{S} as a double cover $\mathbb{P}(V)$ ramified over a quartic curve.

The condition that the linear system has no fixed component defines an open subset $\Omega_d \subset G_{d+1}(\text{Sym}^3 A^*)$ whose complement has codimension > 1 . Over Ω_d we have a well-defined ℓ -marked family \bar{S}_d/Ω_d to which the $\text{SL}(A)$ -action lifts. Any ℓ -marked anticanonical surface is thus obtained so that we have a bijection between the set of isomorphism classes of ℓ -marked Del Pezzo surfaces and the set of $\text{SL}(A)$ -orbits in Ω_d . It is unlikely that this can be lifted to the level of varieties and we therefore we invoke geometric invariant theory. We begin with defining the line bundle that is central to this paper.

Definition 1.1. If $f : S \rightarrow B$ is a family of anticanonical surfaces of degree d , then its *determinant bundle* $\text{Det}(S/B)$ is the line bundle over B that is the dual of the determinant of the rank $9-d$ vector bundle $R^1 f_* \omega_{S/B}^{-1}$ (so this assigns to a Del Pezzo surface S , the line $\det H^0(\omega_S^{-1})^*$).

Thus we have a line bundle $\text{Det}(\mathcal{S}_d/\Omega_d)$. Its fiber over the $(d+1)$ -dimensional subspace $V^* \subset \text{Sym}^3 A^*$ is the line $\det(V)$ and hence the fiber of the ample bundle $\mathcal{O}_{G_{d+1}(\text{Sym}^3 A^*)}(1)$. A section of $\mathcal{O}_{G_{d+1}(\text{Sym}^3 A^*)}(k)$ determines a section of $\text{Det}^{\otimes k}(\mathcal{S}_d/\Omega_d)$ and any section of the latter so occurs because the complement of Ω_d in $G_{d+1}(\text{Sym}^3 A^*)$ has codimension > 1 . Since the action of $\text{SL}(A)$ on $\text{Sym}^3 A^*$ is via $\text{PGL}(A)$ (the center μ_3 of $\text{SL}(A)$ acts trivially), we shall regard this as a representation of the latter. Consider the subalgebra of $\text{PGL}(A)$ -invariants in homogeneous coordinate ring of $G_{d+1}(\text{Sym}^3 A^*)$,

$$R_d^\bullet := \left(\bigoplus_{k=0}^{\infty} H^0(\mathcal{O}_{G_{d+1}(\text{Sym}^3 A^*)}(k)) \right)^{\text{PGL}(A)},$$

The affine cone $\text{Spec}(R_d^\bullet)$ has the interpretation as the categorical $\text{PGL}(A)$ -quotient of the (affine) Plücker cone over the Grassmannian. It may be thought of as the affine hull of the moduli space of triples (S, ℓ, δ) with (S, ℓ) an ℓ -marked Del Pezzo surface of degree d and δ a generator of $\det H^0(\omega_S^{-1})$. Since we shall find that the base of this cone, $\text{Proj}(R_d^\bullet)$, defines a projective completion of $\mathcal{M}_{\ell,d}^\circ$, we denote it by $\mathcal{M}_{\ell,d}^*$. The asserted interpretation of $\mathcal{M}_{\ell,d}^*$ of course requires that we verify that the orbits defined by Fano surfaces are stable. We will do that in a case by case discussion that relates this to GIT completions that are obtained in a different manner. In fact, for each of the three cases $d = 2, 3, 4$ we shall construct a GIT completion \mathcal{M}_d^* of \mathcal{M}_d° in such a way that the forgetful morphism $\mathcal{M}_{\ell,d}^\circ \rightarrow \mathcal{M}_d^\circ$ extends to a finite morphism of GIT completions $\mathcal{M}_{\ell,d}^* \rightarrow \mathcal{M}_d^*$. This description will also help us to identify (and interpret) the boundary strata.

We recall that the proj construction endows $\mathcal{M}_{\ell,d}^*$ for every $k \geq 0$ with a coherent sheaf $\mathcal{O}_{\mathcal{M}_{\ell,d}^*}(k)$ of rank one whose space of sections is R_d^k . We call $\mathcal{O}_{\mathcal{M}_{\ell,d}^*}(1)$ the *determinant sheaf*; it is a line bundle in the orbifold setting.

In what follows, V_{d+1} is a fixed complex vector space of dimension $d+1$ endowed with a generator μ of $\det(V_{d+1})$. We often regard μ as a translation invariant $(d+1)$ -polyvector field on V_{d+1} .

Degree 4 surfaces in projective 4-space. If a pencil of quadrics in $\mathbb{P}(V_5)$ contains a smooth quadric, then the number of singular members of this pencil (counted with multiplicity) is 5. According to Wall [20] the geometric invariant theory for intersections of quadrics is as follows: for a plane $P \subset \text{Sym}^2 V_5^*$, $[\wedge^2 P] \in \mathbb{P}(\wedge^2(\text{Sym}^2 V_5^*))$ is $\text{SL}(V_5)$ -stable (resp. $\text{SL}(V_5)$ -semistable) if and only if the divisor on $\mathbb{P}(P)$ parameterizing singular members is reduced (resp. has all its multiplicities ≤ 2). A semistable pencil belongs to minimal orbit if and only if its members can be simultaneously diagonalized. So a stable pencil is represented by a pair $\langle Z_0^2 + Z_1^2 + Z_2^2 + Z_3^2 + Z_4^2, a_0 Z_0^2 + a_1 Z_1^2 + a_2 Z_2^2 + a_3 Z_3^2 + a_4 Z_4^2 \rangle$ with a_0, \dots, a_4 distinct. This is equivalent to the corresponding surface S_P in $\mathbb{P}(V_5)$ being smooth. The minimal strictly semistable orbits allow at most two pairs of coefficients to

be equal. In case we have only one pair of equal coefficients, S_P has two A_1 -singularities and in case we have two such pairs, four. The fact that these singularities come in pairs can be ‘explained’ in terms of the D_5 -root system in the Picard group of a Del Pezzo surface of degree 4: a A_1 -singularity is resolved by a single blowup with a (-2) -curve as exceptional curve whose class is a root in the Picard root system. The roots perpendicular to this root make up a root system of type $D_4 + A_1$ and the class of the companion (-2) -curve will sit in the A_1 -summand. Besides, a minimal strictly semistable orbit with 2 resp. 4 A_1 -singularities is adjacent to a semistable orbit without such A_1 -pairs and represented by a pair of quadrics one of which is defined by $Z_0^2 + Z_1^2 + Z_2^2 + Z_3^2 + Z_4^2$ and the other by $Z_0Z_1 + a_1Z_1^2 + a_2Z_2^2 + a_3Z_3^2 + a_4Z_4^2$ resp. $Z_0Z_1 + Z_2Z_3 + a_3Z_3^2 + a_4Z_4^2$.

The center μ_5 of $SL(V_5)$ acts faithfully by scalars on $\wedge^2(\mathrm{Sym}^2 V_5^*)$ and for that reason the $SL(V_5)$ -invariant part of the homogeneous coordinate ring of $\mathrm{Gr}_2(\mathrm{Sym}^2 V_5^*)$ lives in degrees that are multiples of 5:

$$S_4^\bullet := \bigoplus_{k=0}^{\infty} S_4^k, \quad S_4^k := H^0(\mathcal{O}_{\mathrm{Gr}_2(\mathrm{Sym}^2 V_5^*)}(5k))^{\mathrm{SL}(V_5)}.$$

We obtain a projective completion $\mathcal{M}_4^* := \mathrm{Proj} S_4^\bullet$ of \mathcal{M}_4° with twisting sheaves $\mathcal{O}_{\mathcal{M}_4^*}(k)$ such that $S_4^k = H^0(\mathcal{O}_{\mathcal{M}_4^*}(k))$. The singular complete intersections are parameterized by a hypersurface in $\mathrm{Gr}_2(\mathrm{Sym}^2 V_5^*)$. Since the Picard group of this Grassmannian is generated by $\mathcal{O}_{\mathrm{Gr}_2(\mathrm{Sym}^2 V_5^*)}(1)$, this discriminant is defined by a section of some $\mathcal{O}_{\mathrm{Gr}_2(\mathrm{Sym}^2 V_5^*)}(20)$ and so $B_4 := \mathcal{M}_4^* - \mathcal{M}_4^\circ$ is defined by a section of $\mathcal{O}_{\mathcal{M}_4^*}(4)$.

Suppose we are given a surface $S \subset \mathbb{P}(V_5)$ defined by a pencil of quadrics. So S determines a line Φ_S in $\wedge^2(\mathrm{Sym}^2 V_5^*)$. Any generator $F_1 \wedge F_2 \in \Phi_S$ and $u \in V_5^*$ determine a 2-vector field on V by $\iota_{du} \wedge dF_1 \wedge dF_2 \mu$. This 2-vector field is invariant under scalar multiplication and tangent to the cone over S . Hence it defines a 2-vector field on S , or equivalently, an element of $H^0(\omega_S^{-1})$. The map thus defined is an isomorphism

$$V_5^* \otimes \Phi_S \cong H^0(\omega_S^{-1}).$$

By taking determinants we get an identification of $\Phi_S^5 \cong \det H^0(\omega_S^{-1})$. We may think of Φ_S^5 as the quotient of the line Φ_S by the center μ_5 of $SL(V_5)$. Thus $\mathrm{Spec}(S_4^\bullet)$ may be regarded as the affine hull of the moduli space of pairs (S, δ) with S a Del Pezzo surface of degree 4 and δ a generator of $\det(H^0(\omega_S^{-1}))$. We conclude:

Proposition 1.2. *We have a natural finite embedding $S_4^\bullet \subset R_4^\bullet$ of graded \mathbb{C} -algebras so that the forgetful morphism $\mathcal{M}_{\ell,4}^\circ \rightarrow \mathcal{M}_4^\circ$ extends to a finite morphism of GIT completions $\mathcal{M}_{\ell,4}^* \rightarrow \mathcal{M}_4^*$ (and the notions of semistability coincide in the two cases) and $\mathcal{O}_{\mathcal{M}_4^*}(1)$ is the determinant sheaf.*

Cubic surfaces. Following Hilbert the cubic surfaces in $\mathbb{P}(V_4)$ that are stable (resp. semistable) relative to the $\mathrm{SL}(V_4)$ -action are those that have an A_1 -singularity (resp. A_2 -singularity) at worst. There is only one strictly semistable minimal orbit and that is the one that has three A_2 -singularities.

The center μ_4 of $\mathrm{SL}(V_4)$ acts faithfully by scalars on $\mathrm{Sym}^3 V_4^*$ and so the $\mathrm{SL}(V_4)$ -invariant part of the homogeneous coordinate ring of $\mathrm{Sym}^3 V_4^*$ lives in degrees that are multiples of 4:

$$S_3^\bullet := \bigoplus_{k=0}^{\infty} S_3^k, \quad S_3^k := H^0(\mathcal{O}_{\mathbb{P}(\mathrm{Sym}^3 V_4^*)}(4k))^{\mathrm{SL}(V_4)}.$$

We thus find the projective completion $\mathcal{M}_3^* := \mathrm{Proj} S_3^\bullet$ of \mathcal{M}_3° with twisting sheaves $\mathcal{O}_{\mathcal{M}_3^*}(k)$ such that $S_3^k = H^0(\mathcal{O}_{\mathcal{M}_3^*}(k))$. The discriminant hypersurface in the linear system of degree d hypersurfaces in \mathbb{P}^n has degree $(n+1)(d-1)^n$. So the singular cubic surfaces are parameterized by a hypersurface in $\mathbb{P}(\mathrm{Sym}^3 V_4^*)$ of degree 32. The stable locus $\mathcal{M}_3^\circ \subset \mathcal{M}_3 \subset \mathcal{M}_3^*$ is the complement of a single point. Furthermore, $B_3 := \mathcal{M}_3^* - \mathcal{M}_3^\circ$ is defined by a section of $\mathcal{O}_{\mathcal{M}_3^*}(8)$.

Let $S \subset \mathbb{P}(V_4)$ be a cubic surface defined by a line Φ_S in $\mathrm{Sym}^3 V_4^*$. Proceeding as in the degree 4 case we find for that a generator $F \in \Phi_S$ and an $u \in V^*$ the expression $\iota_{du} \wedge dF \mu$ defines a 2-vector field on S and that we thus get an isomorphism

$$V_4^* \otimes \Phi_S \cong H^0(\omega_S^{-1}).$$

By taking determinants we get an identification of $\Phi_S^4 \cong \det H^0(\omega_S^{-1})$. We think of Φ_S^4 as the quotient of the line Φ_S by the center μ_4 of $\mathrm{SL}(V_4)$ and conclude as before:

Proposition 1.3. *We have a natural finite embedding $S_3^\bullet \subset R_3^\bullet$ of graded \mathbb{C} -algebras so that the forgetful morphism $\mathcal{M}_{\ell,3}^\circ \rightarrow \mathcal{M}_3^\circ$ extends to a finite morphism of GIT completions $\mathcal{M}_{\ell,3}^* \rightarrow \mathcal{M}_3^*$ (and the notions of semistability coincide in the two cases) and $\mathcal{O}_{\mathcal{M}_3^*}(1)$ is the determinant sheaf.*

Quartic curves. The case of degree 2 is a bit special because $W(E_7)$ has a nontrivial center (of order two). The center leaves invariant the (isomorphism type of the) surface: it acts as an involution and only changes the marking. The latter even disappears if we only remember the fixed point set of this involution, the quartic curve. Van Geemen [10] observed that the marking of the Del Pezzo surface then amounts to a principal level two structure on the quartic curve (this is based on the fact that $W(E_7)$ modulo its center is isomorphic to the symplectic group $\mathrm{Sp}(6, \mathbb{Z}/2)$). Since a smooth quartic curve is a canonically embedded genus three curve, \mathcal{M}_2° can also be interpreted as the moduli space of nonhyperelliptic genus three curves with principal level two structure (here we ignore the orbifold structure).

The projective space $\mathbb{P}(\mathrm{Sym}^4 V_3^*)$ parameterizes the quartic curves in the projective plane $\mathbb{P}(V_3)$. The geometric invariant theory relative its $\mathrm{SL}(V_3)$ -action is as follows: a quartic curve is stable if and only if it has singularities

no worse than of type A_2 . A quartic is unstable if and only if it has a point of multiplicity ≥ 3 (or equivalently, a D_4 -singularity or worse) or consists of a cubic plus an inflectional tangent. The latter gives generically a A_5 -singularity, but such a singularity may also appear on a semistable quartic, for instance on the union of two conics having a point in common where they intersect with multiplicity 3. Let us, in order to understand the incidence relations, review (and redo) the classification of nonstable quartics.

A plane quartic curve C that is not stable has a singularity of type A_3 or worse. So it has an equation of the form $cy^2z^2 + yzf_2(x, y) + f_4(x, y)$ with f_2 and f_4 homogeneous. Consider its orbit under the \mathbb{C}^\times -copy in $SL(V_3)$ for which $t \in \mathbb{C}^\times$ sends (x, y, z) to $(x, ty, t^{-1}z)$. If we let $t \rightarrow 0$, then the equation tends to $cy^2z^2 + ax^2yz + bx^4$, where $f_2(x, 0) = ax^2$ and $f_4(x, 0) = bx^4$. We go through the possibilities.

If $c = 0$, then C has a triple point and the equation $ax^2yz + bx^4$ is easily seen to be unstable. We therefore assume that $c = 1$ and we denote the limit curve by C_0 .

If $a^2 - 4b \neq 0 \neq b$, then C_0 is made up of two nonsingular conics meeting in two distinct points with a common tangent (having therefore a A_3 -singularity at each) and the original singularity was of type A_3 .

If $a \neq 0 = b$, then we have the same situation except that one of the conics has now degenerated into a union of two lines.

The most interesting case is when $a^2 - 4b = 0 \neq b$. Then C_0 is a double nonsingular conic and in case $C \neq C_0$, C has a singularity of type A_k for some $4 \leq k \leq 7$. The case of an A_7 -singularity occurs for the curve C_1 given by $(yz + x^2)(yz + x^2 + y^2)$: it consists of two nonsingular conics meeting in a single point with multiplicity 4. This is also the most degenerate case next after C_0 : any $SL(V_3)$ -orbit that has C_0 in its closure is either the orbit of C_0 or has C_1 in its closure. So although a double conic does not yield a Del Pezzo surface, the corresponding point of \mathcal{M}_2^* is uniquely represented by a geometrically marked Del Pezzo surface with a A_7 -singularity.

On the other hand, the condition $a = b = 0$ (which means that f_2 and f_4 are divisible by y so that we have a cubic plus an inflectional tangent or worse), gives the limiting curve defined by $y^2z^2 = 0$, which is clearly unstable.

We shall later find that the ambiguous behaviour of a A_5 -singularity reflects a feature of the E_7 -root system: this system contains two Weyl group equivalence classes of subsystems of type A_5 : one type is always contained in a A_7 -subsystem (the semistable case) and the other is not (the unstable case).

Since the center μ_3 of $SL(V_3)$ acts faithfully by scalars on $\text{Sym}^4 V_3^*$ we have as algebra of invariants

$$S_2^\bullet := \bigoplus_{k=0}^{\infty} S_2^k, \quad S_2^k := H^0(\mathcal{O}_{\mathbb{P}(\text{Sym}^4 V_3^*)}(3k))^{\text{SL}(V_3)}.$$

Thus $\mathcal{M}_2^* := \text{Proj } \mathbb{C}[\text{Sym}^4 V_3^*]^{\text{SL}(V_3)}$ is a projective completion of \mathcal{M}_2° . It comes with twisting sheaves $\mathcal{O}_{\mathcal{M}_2^*}(k)$ such that $S_2^k = H^0(\mathcal{O}_{\mathcal{M}_2^*}(k))$. Let us write $\mathcal{M}_2^\circ \subset \mathcal{M}_2 \subset \mathcal{M}_2^*$ for the stable locus; this can be interpreted as the moduli space of marked Del Pezzo surfaces of degree 2 with A_2 -singularities at worst. Its complement in \mathcal{M}_2^* is of dimension one. Since the singular quartics make up a hypersurface of degree 27 in $\mathbb{P}(\text{Sym}^4 V_3^*)$, $B_2 := \mathcal{M}_2^* - \mathcal{M}_2^\circ$ is defined by a section of $\mathcal{O}_{\mathcal{M}_2^*}(9)$. In particular, B_2 is a Cartier divisor.

Let C be a quartic curve in $\mathbb{P}(V_3)$ defined by the line $\Phi_C \subset \text{Sym}^4 V_3^*$. If $F \in \Phi_C$ is a generator, then a double cover S of $\mathbb{P}(V_3)$ totally ramified along C is defined by $w^2 = F$ in $V_3 \times \mathbb{C}$ (more precisely, it is Proj of the graded algebra obtained from $\mathbb{C}[V_3]$ by adjoining to it a root of F). Then for every $u \in V_3^*$, the 2-vector field $w^{-1} \iota_{du} \wedge dF \mu$ defines a section of ω_S^{-1} . We thus get an isomorphism $V_3^* \otimes w^{-1} dF \cong H^0(\omega_S^{-1})$. If we take the determinants of both sides, we find that $(w^{-1} dF)^3$ determines a generator of $\det H^0(\omega_S^{-1})$. So $F^{-3} (dF)^6$ gives one of $(\det H^0(\omega_S^{-1}))^2$, in other words, we have a natural isomorphism $\Phi_C^3 \cong (\det H^0(\omega_S^{-1}))^2$. That the square of the determinant appears here reflects the fact that the central element -1 of $W(E_7)$ induces an involution in S which acts as the scalar -1 on $\det H^0(\omega_S^{-1})$. We obtain:

Proposition 1.4. *We have a natural finite embedding $S_2^\bullet \subset R_2^\bullet$ of graded \mathbb{C} -algebras so that the forgetful morphism $\mathcal{M}_{\ell,2}^\circ \rightarrow \mathcal{M}_2^\circ$ extends to a finite morphism of GIT completions $\mathcal{M}_{\ell,2}^* \rightarrow \mathcal{M}_2^*$ (and the notions of semistability coincide in the two cases) and $\mathcal{O}_{\mathcal{M}_2^*}(1)$ is the square of the determinant sheaf.*

Completion of the moduli space of marked Fano surfaces. We have produced for $d = 4, 3, 2$, a GIT completion \mathcal{M}_d^* of \mathcal{M}_d° that we were able to identify with a finite quotient of $\mathcal{M}_{\ell,d}^*$. This implies that $\mathcal{M}_{\ell,d}^*$ contains $\mathcal{M}_{\ell,d}^\circ$ as an open dense subset and proves that every point of $\mathcal{M}_{\ell,d}^*$ can be represented by an ℓ -marked Fano surface.

We define a completion $\mathcal{M}_{m,d}^*$ of the moduli space $\mathcal{M}_{m,d}^\circ$ of marked Fano surfaces of degree d simply as the normalization of \mathcal{M}_d^* in $\mathcal{M}_{m,d}^\circ$. This comes with an action of $W(R_{9-d})$ and the preceding discussion shows that $\mathcal{M}_{\ell,d}^*$ can be identified with the orbit space of $\mathcal{M}_{m,d}^\circ$ by the permutation group of the e_1, \dots, e_{9-d} (a Weyl subgroup of type A_{8-d}).

2. COBLE'S COVARIANTS

In this section we assume that the degree d of a Del Pezzo surface is at most 6 (we later make further restrictions).

Let $(S; e_1, \dots, e_{9-d})$ be a geometrically marked Del Pezzo surface. Recall that we have a class $\ell \in \text{Pic}(S)$ characterized by the property that $-3\ell + e_1 + \dots + e_{9-d}$ equals the canonical class k . Let us choose a line bundle \mathcal{L} on S which represents ℓ : $H^0(\mathcal{L})$ is then of dimension 3 and if we denote its dual by A , then the associated linear system defines a birational morphism

$S \rightarrow \mathbb{P}(A)$ which has $E = E_1 + \dots + E_{9-d}$ as its exceptional divisor. The direct image of \mathcal{L} on $\mathbb{P}(A)$ is still a line bundle (namely $\mathcal{O}_{\mathbb{P}(A)}(1)$, but we continue to denote this bundle by \mathcal{L}).

We claim that there is a natural identification

$$\omega_S^{-1} \cong \mathcal{L}^3(-E) \otimes \det A.$$

To see this, we note that if $p \in \mathbb{P}(A)$ and $\lambda \subset A$ is the line defined by p , then the tangent space of $\mathbb{P}(A)$ at p appears in the familiar exact sequence

$$0 \rightarrow \mathbb{C} \rightarrow \text{Hom}(\lambda, A) \rightarrow T_p \mathbb{P}(A) \rightarrow 0,$$

from which it follows that $\det T_p \mathbb{P}(A) = \lambda^{-3} \det(A)$ (we often omit the \otimes -symbol when lines or line bundles are involved). So the anticanonical bundle $\omega_{\mathbb{P}(A)}^{-1}$ of $\mathbb{P}(A)$ is naturally identified with $\mathcal{L}^3 \otimes \det(A)$. Since $S \rightarrow \mathbb{P}(A)$ is the blowup with exceptional divisor E , we see that the above identification makes ω_S^{-1} correspond to $\mathcal{L}^3(-E) \otimes \det(A)$.

The following simple lemma will help us to understand Coble's covariants.

Lemma 2.1. *For a Del Pezzo surface S , the determinant lines of the vector spaces $H^0(\mathcal{O}_E \otimes \mathcal{L}^3) \otimes \det(A)$ and $V(S) := H^0(\omega_S^{-1})^*$ are canonically isomorphic.*

Proof. The identification $\omega_S^{-1} \cong \mathcal{L}^3(-E) \otimes \det A$ above gives rise to the short exact sequence

$$0 \rightarrow \omega_S^{-1} \rightarrow \mathcal{L}^3 \otimes \det A \rightarrow \mathcal{O}_E \otimes \mathcal{L}^3 \otimes \det A \rightarrow 0.$$

This yields an exact sequence on H^0 because $H^1(\omega_S^{-1}) = 0$. If we take into account that $H^0(\mathcal{L}^3) = \text{Sym}^3 H^0(\mathcal{L}) = \text{Sym}^3 A^*$, then we find the exact sequence

$$0 \rightarrow V^* \rightarrow \text{Sym}^3 A^* \otimes \det A \rightarrow H^0(\mathcal{O}_E \otimes \mathcal{L}^3) \otimes \det A \rightarrow 0.$$

Since $\dim(\text{Sym}^3 A^*) = 10$ and $\det(\text{Sym}^3 A^*) = (\det A)^{-10}$, the determinant of the middle term has a canonical generator. This identifies the determinant of V with the one of the right hand side. \square

It will be convenient to have a notation for the one dimensional vector space appearing in the preceding lemma: we denote

$$L(S, E) := \det(H^0(\mathcal{O}_E \otimes \mathcal{L}^3 \otimes \det A)) = \det(H^0(\mathcal{O}_E \otimes \mathcal{L}^3)) \otimes (\det A)^{9-d},$$

so that the lemma asserts that $L(S, E)$ may be identified with $\det V(S)$.

We continue with $(S; e_1, \dots, e_{9-d})$ and \mathcal{L} . If p_i denotes the image point of E_i , then the geometric fiber of \mathcal{L} over p_i is $\lambda_i := H^0(\mathcal{L} \otimes \mathcal{O}_{E_i})^*$ (a one dimensional vector space). So $L(S, E) = (\lambda_1 \cdots \lambda_{9-d})^{-3} \otimes \det(A)^{9-d}$. For e_i, e_j, e_k distinct, the map defined by componentwise inclusion

$$\lambda_i \oplus \lambda_j \oplus \lambda_k \rightarrow A$$

is a linear map between 3-dimensional vector spaces. It is an isomorphism if p_i, p_j, p_k are not collinear. Hence the corresponding map on the third exterior powers, yields an element

$$|ijk| \in \lambda_i^{-1} \lambda_j^{-1} \lambda_k^{-1} \det A$$

that is nonzero in the Fano case (recall that we usually omit the \otimes -sign when lines are involved). Notice that the line $\lambda_i^{-1} \lambda_j^{-1} \lambda_k^{-1} \det A$ attached to \mathcal{L} only depends on the marked surface $(S; e_1, \dots, e_{9-d})$ and not on the choice of the \mathcal{L} : as said above \mathcal{L} is unique up to isomorphism and the only possible ambiguity therefore originates from the action of \mathbb{C}^\times in the fibers of \mathcal{L} . But it is clear that this \mathbb{C}^\times -action is trivial on this line. For e_{i_1}, \dots, e_{i_6} distinct, we also have a linear map between 6-dimensional vector spaces

$$\lambda_{i_1}^2 \oplus \dots \oplus \lambda_{i_6}^2 \rightarrow \text{Sym}^2 A.$$

Since $\det(\text{Sym}^2 A) = (\det A)^4$, this defines a determinant

$$|i_1 \dots i_6| \in \lambda_{i_1}^{-2} \dots \lambda_{i_6}^{-2} (\det A)^4.$$

It is nonzero if and only if p_1, \dots, p_6 do not lie on a conic, which is the case when S is Fano. The elements $|ijk|$ and $|i_1 \dots i_6|$ just introduced will be referred to as *Coble factors*.

Action of the Weyl group on the Coble factors. We now assume that S is a Fano surface of degree ≤ 6 . Another marking of S yields another ℓ' and hence another line $L(S, E')$. Nevertheless they are canonically isomorphic to each other since both have been identified with $\det V$. For what follows it is important to make this isomorphism concrete. We will do this for the case that the new marking is the image of the former under the reflection in h_{123} . So $\ell' = 2\ell - e_1 - e_2 - e_3$, $e'_1 = \ell - e_2 - e_3$ (E'_1 is the strict transform of $\overline{p_2 p_3}$), e'_2 and e'_3 are expressed in a likewise manner and $e'_i = e_i$ for $i > 3$. We represent ℓ' by

$$\mathcal{L}' := \mathcal{L}^2(-E_1 - E_2 - E_3)$$

so that $A' = H^0(\mathcal{L}')^*$. Before proceeding, let us see what happens if we do this twice, that is, if we apply h_{123} once more:

$$\mathcal{L}'' = \mathcal{L}'^2(-E'_1 - E'_2 - E'_3) = \mathcal{L}^4(-2E_1 - 2E_2 - 2E_3 - E'_1 - E'_2 - E'_3).$$

The line bundle $\mathcal{L}^3(-2E_1 - 2E_2 - 2E_3 - E'_1 - E'_2 - E'_3)$ is trivial (a generator is given by a section of \mathcal{L}^3 whose divisor is the triangle spanned by p_1, p_2, p_3) and so if I denotes its (one dimensional) space of sections, then \mathcal{L}'' is identified with $\mathcal{L} \otimes I$. We note that for $i > 3$, the restriction map $I \rightarrow \lambda_i^{-3}$ is an isomorphism of lines and that we also have a natural isomorphism $I \rightarrow (\lambda_1 \lambda_2 \lambda_3)^{-1}$, which, after composition with the inverse of $|123|$ yields an isomorphism $I \rightarrow \det(A)^{-1}$.

The space of sections of \mathcal{L}' is the space of quadratic forms on A that are zero on λ_1, λ_2 and λ_3 . This leads to an exact sequence

$$0 \rightarrow \lambda_1^2 \oplus \lambda_2^2 \oplus \lambda_3^2 \rightarrow \text{Sym}^2 A \rightarrow A' \rightarrow 0.$$

The exactness implies that

$$\det A' = (\det A)^4 (\lambda_1 \lambda_2 \lambda_3)^{-2}.$$

We have $(\lambda'_i)^{-1} = H^0(\mathcal{L}^2(-E_1 - E_2 - E_3) \otimes \mathcal{O}_{E'_i})$ by definition. For $i > 3$, this is just the space of quadratic forms on the line λ_i , i.e., λ_i^{-2} and so $\lambda'_i = \lambda_i^2$ in that case. For $i = 1$,

$$(\lambda'_1)^{-1} = H^0(\mathcal{L}^2(-E_1 - E_2 - E_3) \otimes \mathcal{O}_{E'_1}) = H^0(\mathcal{L}^2 \otimes \mathcal{O}_{\overline{p_2 p_3}}(-((p_2) - (p_3))))$$

is the space of quadratic forms on $\lambda_2 + \lambda_3$ that vanish on each summand, i.e., $\lambda_2^{-1} \lambda_3^{-1}$. Thus $\lambda'_1 = \lambda_2 \lambda_3$ and likewise $\lambda'_2 = \lambda_3 \lambda_1$, $\lambda'_3 = \lambda_1 \lambda_3$. Notice that λ''_i is naturally identified with $\lambda_i \otimes I^{-1}$. Thus

$$\begin{aligned} L(S, E') &= (\lambda'_1 \cdots \lambda'_{9-d})^{-3} (\det A')^{9-d} = \\ &= (\lambda_1 \cdots \lambda_{9-d})^{-6} (\lambda_1 \lambda_2 \lambda_3)^{-2(9-d)} (\det A)^{4(9-d)} = \\ &= L(S, E) (\lambda_1^{-1} \lambda_2^{-1} \lambda_3^{-1} \det A)^{21-2d} \prod_{i=4}^{9-d} (\lambda_i^{-3} \det A). \end{aligned}$$

The identifications above of $\lambda_1 \lambda_2 \lambda_3$, λ_i^3 ($i > 3$) and $\det(A)$ with I^{-1} show that the twisting line $(\lambda_1^{-1} \lambda_2^{-1} \lambda_3^{-1} \det A)^{21-2d} \prod_{i=4}^{9-d} (\lambda_i^{-3} \det A)$ has a canonical generator δ . This generator can be expressed in terms of our Coble factors as

$$\delta := |123|^9 \prod_{i=4}^{9-d} (|12i||23i||31i|)$$

(which indeed lies in $(\lambda_1^{-1} \lambda_2^{-1} \lambda_3^{-1} \det A)^{21-2d} \prod_{i=4}^{9-d} (\lambda_i^{-3} \det A)$).

Proposition 2.2. *The isomorphism $L(S; E) \cong L(S', E')$ defined above coincides with the isomorphism that we obtain from the identification of domain and range with $\det V$.*

Proof. Choose generators $a_i \in \lambda_i$ and write x_1, x_2, x_3 for the basis of A^* dual to a_1, a_2, a_3 . The basis (a_1, a_2, a_3) of A defines a generator $a_1 \wedge a_2 \wedge a_3$ of $\det A$. This determines an isomorphism $\phi : \omega_S^{-1} \cong \mathcal{L}^3(-E)$. That isomorphism fits in the exact sequence

$$0 \rightarrow V^* \rightarrow \text{Sym}^3 A^* \rightarrow \bigoplus_{i=1}^{9-d} \lambda_i^{-3} \rightarrow 0.$$

The middle space has the cubic monomials in x_1, x_2, x_3 as a basis. The triple (x_1^3, x_2^3, x_3^3) defines a basis dual to $(a_1^3, a_2^3, a_3^3) \in \lambda_1^3 \oplus \lambda_2^3 \oplus \lambda_3^3$. It follows that we have an exact subsequence

$$(1) \quad 0 \rightarrow V^* \rightarrow K \rightarrow \bigoplus_{i=4}^{9-d} \lambda_i^{-3} \rightarrow 0,$$

where $K \subset \text{Sym}^3 A^*$ is the span of the cubic monomials that are not a third power. This yields an identification

$$(\lambda_4 \cdots \lambda_{9-d})^{-3} \cong \det V \det(\langle x_1^2 x_2, \dots, x_2 x_3^2, x_1 x_2 x_3 \rangle).$$

We now do the same for $\mathcal{L}' = \mathcal{L}^2(-E_1 - E_2 - E_3)$. The space A' comes with a basis $(a'_1 = a_2 a_3, a'_2 = a_3 a_1, a'_3 = a_1 a_2)$ which is dual to the basis $(x_2 x_3, x_3 x_1, x_1 x_2)$ of $H^0(\mathcal{L}^2(-E_1 - E_2 - E_3))$. The monomial $x_1 x_2 x_3$ is the obvious generator of $I = H^0(\mathcal{L}^3(-E'_1 - E'_2 - E'_3 - 2E_1 - 2E_2 - 2E_3))$ so that have an associated isomorphism

$$\phi' := \phi \otimes x_1 x_2 x_3 : \omega_S^{-1} \rightarrow (\mathcal{L}')^3(-E').$$

We fit this in the exact sequence

$$0 \rightarrow H^0(\omega_S^{-1}) \rightarrow H^0((\mathcal{L}')^3) \rightarrow \bigoplus_{i=1}^{9-d} \lambda_i'^{-3} \rightarrow 0.$$

The vector space of sections of the middle term has the cubic monomials in $x_2 x_3, x_3 x_1, x_1 x_2$ as a basis. The lines $\lambda'_1, \lambda'_2, \lambda'_3$ are spanned by $a_2 a_3, a_3 a_1$ and $a_1 a_2$ respectively and λ'_i is for $i = 4, 5, \dots, 9-d$ spanned by a_i^2 . So $(x_2 x_3)^3$ spans $\lambda_1'^{-3}$ and similarly for $\lambda_2'^{-3}$ and $\lambda_3'^{-3}$. The space spanned by cubic monomials in $x_2 x_3, x_3 x_1, x_1 x_2$ that are not pure powers is just $K' := x_1 x_2 x_3 K$. It follows that we get an exact subsequence analogous to the sequence (1):

$$(2) \quad 0 \rightarrow V^* \rightarrow K' \rightarrow \bigoplus_{i=4}^{9-d} \lambda_i'^{-3} \rightarrow 0,$$

where the embedding $V^* \rightarrow K'$ is the composite of $V^* \rightarrow K$ and the isomorphism $K \cong K'$ given by multiplication by $x_1 x_2 x_3$. This identifies $(a_i')^{-3} \in \lambda_i'^{-3}$ with $(x_1 x_2 x_3)(a_i) \cdot a_i^{-3} \in \lambda_i^{-3}$. From this we deduce that the generator $(a'_1 \cdots a'_{9-d})^{-3} (a'_1 \wedge a'_2 \wedge a'_3)^{9-d}$ of $(\lambda'_1 \cdots \lambda'_{9-d})^{-3} (\det A')^{9-d}$ corresponds to $\prod_{i=4}^{9-d} (x_1 x_2 x_3)(a_i)$ times the generator $(a_1 \cdots a_{9-d})^{-3} (a_1 \wedge a_2 \wedge a_3)^{9-d}$ of $(\lambda_1 \cdots \lambda_{9-d})^{-3} (\det A)^{9-d}$. Since $\delta = \prod_{i=4}^{9-d} (x_1 x_2 x_3)(a_i)$, this proves that the isomorphism $L(S; E) \cong \det V \cong L(S, E')$ sends the generator of the former to δ times the generator of the latter. \square

We next determine how the Coble factors for ℓ' are expressed in terms of those of ℓ . We retain our $0 \neq a_i \in \lambda_i$ and write x_1, x_2, x_3 for the basis of A^* dual to a_1, a_2, a_3 as before. We identify $|ijk|$ resp. $|i_1 \cdots i_6|$ with the $a_1 \wedge a_2 \wedge a_3$ -coefficient of $a_i \wedge a_j \wedge a_k$ resp. the $a_1^2 \wedge a_2^2 \wedge a_3^2 \wedge a_2 a_3 \wedge a_3 a_1 \wedge a_1 a_2$ -coefficient of $a_{i_1}^2 \wedge \cdots \wedge a_{i_6}^2$. Here are some typical cases, where it is assumed that the free indices are distinct and > 3 :

$$\begin{aligned} |123| &= 1, \\ |12k| &= \langle x_3 | a_k \rangle, \\ |1JFK| &= \langle x_2 \wedge x_3 | a_j \wedge a_k \rangle, \\ |ijk| &= \langle x_1 \wedge x_2 \wedge x_3 | a_i \wedge a_j \wedge a_k \rangle, \\ |123ijk| &= \langle x_2 x_3 \wedge x_3 x_1 \wedge x_1 x_2 | a_i^2 \wedge a_j^2 \wedge a_k^2 \rangle, \\ |12ijkl| &= \langle x_2^2 \wedge x_2 x_3 \wedge x_3 x_1 \wedge x_1 x_2 | a_i^2 \wedge a_j^2 \wedge a_k^2 \wedge a_l^2 \rangle, \\ |1ijklm| &= \langle x_2^2 \wedge x_3^2 \wedge x_2 x_3 \wedge x_3 x_1 \wedge x_1 x_2 | a_i^2 \wedge a_j^2 \wedge a_k^2 \wedge a_l^2 \wedge a_m^2 \rangle. \end{aligned}$$

The corresponding expressions for the new marking are converted into the old marking by the substitutions

$$a'_i = \begin{cases} a_2 a_3 & \text{when } i = 1, \\ a_3 a_1 & \text{when } i = 2, \\ a_1 a_2 & \text{when } i = 3, \\ a_i^2 & \text{when } i > 3. \end{cases}, \quad x'_i = \begin{cases} x_2 x_3 & \text{when } i = 1, \\ x_3 x_1 & \text{when } i = 2, \\ x_1 x_2 & \text{when } i = 3, \\ x_i^2 & \text{when } i > 3. \end{cases}.$$

We thus find:

$$\begin{aligned} |123|' &= 1 = |123|, \\ |12k|' &= x_1 x_2(a_k) = |23k||31k|, \\ |1jk|' &= -x_1(a_j)x_1(a_k)|1jk|, \\ |ijk|' &= |123ijk|, \\ |123ijk|' &= x_1 x_2 x_3(a_i).x_1 x_2 x_3(a_j).x_1 x_2 x_3(a_k).|ijk|, \\ |12ijkl|' &= x_1 x_2(a_i).x_1 x_2(a_j).x_1 x_2(a_k).x_1 x_2(a_l).|12ijkl|, \end{aligned}$$

The expression for $|1ijklm|'$ does not appear to have a pleasant form: we find that

$$\begin{aligned} |1ijklm|' &= x_1(a_i)x_1(a_j)x_1(a_k)x_1(a_l)x_1(a_m). \\ &\cdot \langle x_3^2 x_1 \wedge x_1 x_3^2 \wedge x_1 x_2 x_3 \wedge x_2^2 x_3 \wedge x_2 x_3^2 \mid a_i^3 \wedge a_j^3 \wedge a_k^3 \wedge a_l^3 \wedge a_m^3 \rangle. \end{aligned}$$

The covariants. Here is the definition.

Definition 2.3. Let $(S; e_1, \dots, e_{9-d})$ be a marked Fano surface of degree $d \leq 6$. A *Coble covariant* is an element of $L(S, E)$ that is a product of Coble factors $|ijk|$ and $|i_1 \cdots i_6|$ in such a manner that every unordered pair in $\{1, 2, \dots, 9-d\}$ appears in one of these factors.

This notion also makes sense for a marked Del Pezzo surface, and indeed, in case the E_1, \dots, E_{9-d} are irreducible (or equivalently, p_1, \dots, p_{9-d} are distinct), then we adopt this as a definition. But when this is not the case, this is not the ‘right’ definition (see Remark 2.5).

It is easily verified that Coble covariants exist only when $2 \leq d \leq 5$. In these cases they are as follows:

- $d = 5$: There is only one Coble covariant, namely $|123||234||341||412|$. It is nonzero if and only if no three points are collinear, that is, if S is a Fano surface.
- $d = 4$: A typical Coble covariant is $|123||234||345||451||512|$. It depends on a cyclic ordering of $\{1, 2, \dots, 5\}$, with the opposite cycle giving the same element. So the number of Coble covariants up to sign is equal to $4!/2 = 12$. A Coble covariant can be nonzero even if S has (-2) -curves. For instance, if (p_1, p_2, p_4) and (p_2, p_3, p_5) are collinear but are otherwise generic then the given Coble covariant is nonzero and S has a $2A_1$ -configuration (i.e., two disjoint (-2) -curves).

- $d = 3$: We have two typical cases: one is $|134||234||356||456||512||612|$ and another is $|123||456||123456|$. The former type amounts to dividing the 6-element set $\{e_1, \dots, e_6\}$ in three equal parts (of two) and cyclically order the three parts (there are 30 such) and the latter to splitting of $\{e_1, \dots, e_6\}$ into two equal parts (there are 10 of these). So there are 40 Coble covariants up to sign.
- $d = 2$: We have two typical cases: $|351||461||342||562||547||217||367|$ (of which there are 30) and $|123456||127||347||567|$ (105 in number). So up to sign we find 135 cases.

Proposition 2.4. *If S is Fano, then the collection of Coble covariants, when considered as elements of $\det V(S)$, is independent of the marking.*

Proof. It is enough to show that the collection is invariant under the reflection in h_{123} . So in view of Proposition 2.2 we need to verify that if we make the above substitutions for a Coble covariant relative to (e'_1, \dots, e'_{9-d}) , then we get δ times a Coble covariant relative to (e_1, \dots, e_{9-d}) . This is a straightforward check. We do a few examples. For $d = 3$, we find

$$\begin{aligned} |134|'|234|'|356|'|456|'|512|'|612|' &= \\ &= -x_1x_3(a_4).x_2x_3(a_4).x_3(a_5)x_3(a_6)|356|. |123456|.x_1x_2(a_5).x_1x_2(a_6) = \\ &= -\delta|124||356||123456|, \end{aligned}$$

which is indeed the image of $|124||356||123456|$. Similarly,

$$|123|'|456|'|123456|' = |123|. |123456|. \delta |456|,$$

which is the image of $|123||456||123456|$.

The other cases are similar and are left to the reader to verify. \square

Remark 2.5. The notion of a Coble covariant extends to the case of a geometrically marked Del Pezzo surface. There is not much of an issue here as long as the points p_1, \dots, p_{9-d} remain distinct, but when two coalesce the situation becomes a bit delicate, since we wish to land in $\det V(S)$. It is clear that if p_2 approaches p_1 , then any Coble covariant involving these points such as $|123| \in \lambda_1^{-1}\lambda_2^{-1}\lambda_3^{-1} \det A$ tends to 0. But we should regard $|123|$ as an element of $\det(H^0(\mathcal{O}_{E_1+E_2+E_3} \otimes \mathcal{L}^3)) \otimes \det A$ and when p_2 tends to p_1 , then E_1 becomes decomposable and of the form $F + E_2$. The component F is the strict transform of the exceptional curve of the first blowup and hence a (-2) -curve. Let q be the point where F and E_2 meet. This corresponds to tangent direction at p_1 , or equivalently, to a plane $P_q \subset A$ that contains λ_1 . If λ_2 moves in W towards λ_1 , then $H^0(\mathcal{O}_{E_1+E_2})$ becomes $H^0(\mathcal{O}_{F+E_2})$. The exact sequence

$$0 \rightarrow \mathcal{I}_{F+E_2}/\mathcal{I}_{F+2E_2} \rightarrow \mathcal{O}_{F+2E_2} \rightarrow \mathcal{O}_{F+E_2} \rightarrow 0$$

induces an exact sequence on sections. Notice that the first term is a constant sheaf on E_2 . Its fiber over q is $T_q^*F \otimes T_q^*E_2$. This fiber is apparently also the determinant of $H^0(\mathcal{O}_{F+2E_2})$. Since $T_qF = \text{Hom}(\lambda, W_q/\lambda)$ and

$T_q E_2 = \text{Hom}(W_q/\lambda, A/W_q)$, we have $T_q^* F \otimes T_q^* E_2 = (\det A)^{-1} \det P_q \otimes \lambda_1$. It follows that

$$\det(H^0(\mathcal{O}_{F+2E_2+E_3} \otimes \mathcal{L}^3)) \otimes \det A = \lambda_1^{-1} \lambda_3^{-1} \det P_q.$$

It is in this line where |123| should take its value. (This also explains why in the next section we need to divide by a discriminant $\Delta(t_q, \dots, t_{g-d})$.)

Since every point of $\mathcal{M}_{m,d}^*$ is representable by a marked Del Pezzo surface, a Coble covariant can be regarded as a section of the determinant sheaf $\mathcal{O}_{\mathcal{M}_{m,d}^*}(1)$. It follows from Section 2 that $W(R_{g-d})$ permutes these sections transitively.

Definition 2.6. The Coble space \mathcal{C}_d is the subspace of $H^0(\mathcal{O}_{\mathcal{M}_{m,d}^*}(1))$ spanned by the Coble covariants.

We shall prove that for $d = 3, 4$, \mathcal{C}_d is complete, i.e., all of $H^0(\mathcal{O}_{\mathcal{M}_{m,d}^*}(1))$, but we do not know whether that is true when $d = 2$.

Remark 2.7. We shall see that \mathcal{C}_d is an irreducible representation of $W(R_{g-d})$ of dimension 6, 10, 15 for resp. $d = 4, 3, 2$. It follows from the discussion in Section 1 that any $W(R_{g-d})$ -invariant polynomial of degree k in the Coble covariants has an interpretation in terms of classical invariant theory: for $d = 4$ we get a $SL(5)$ -invariant of degree $5k$ for pencils of quadrics, for $d = 3$, a $SL(4)$ -invariant of degree $4k$ for cubic forms and for $d = 2$, a $SL(3)$ -invariant of degree $3k/2$ for quartic forms.

Here is an example that illustrate this. There is only one irreducible representation of $W(E_6)$ of degree 10. This representation is real and has therefore a nonzero $W(E_6)$ -invariant quadratic form. According to the preceding this produces a $SL(4)$ -invariant of degree 8 for cubic forms. This is indeed the lowest degree of a such an invariant.

3. ANTICANONICAL DIVISORS WITH A CUSP

Anticanonical cuspidal cubics on Del Pezzo surfaces. Let S be a Del Pezzo surface of degree d that is not isomorphic to a smooth quadric. Assume that is also given a reduced anticanonical curve K on S isomorphic to a cuspidal cubic (notice that if $d = 1$ such a curve will not always exist). The curve K is given by a hyperplane $V_K \subset V = H^0(\omega_S^{-1})^*$. We write l_K for the line V/V_K so that l_K^* is a line in $V^* = H^0(\omega_S^{-1}) = \text{Hom}(\omega_S, \mathcal{O}_S)$. The image of l_K^* is $\text{Hom}(\omega_S, \mathcal{O}_S(-K)) \subset \text{Hom}(\omega_S, \mathcal{O}_S)$ and hence l_K may be identified with $H^0(\omega_S(K))$. So a nonzero $\kappa \in l_K$ can be understood as a rational 2-form κ on S whose divisor is K . The residue $\text{Res}_K(\kappa)$ of κ on the smooth part of K identifies $\text{Pic}^0(K)$ with \mathbb{C} as an algebraic group: we may represent an element of $\text{Pic}^0(K)$ by a difference $(q) - (p)$ and then $\text{Res}_K(\kappa)$ assigns to this element the integral of $\text{Res}_K(\kappa)$ along any arc in K from p to q . This identifies l_K with $\text{Hom}(\text{Pic}^0(K), \mathbb{C})$ or equivalently, $\text{Pic}^0(K)$ with l_K^* .

Recall that $\text{Pic}_0(S) \subset \text{Pic}(S)$ denotes the orthogonal complement of the class of ω_S^{-1} . It is then clear that restriction defines a homomorphism $r : \text{Pic}_0(S) \rightarrow \text{Pic}^0(K) \rightarrow \mathfrak{l}_K^*(\subset V^*)$. We extend r to an algebra homomorphism

$$r : \text{Sym}^\bullet \text{Pic}_0(S) \rightarrow \text{Sym}^\bullet \mathfrak{l}_K^*(\subset \text{Sym}^\bullet V^*).$$

For $\kappa \in \mathfrak{l}_K$, we compose this map with the evaluation in κ and obtain an algebra homomorphism

$$r_\kappa : \text{Sym}^\bullet \text{Pic}_0(S) \rightarrow \mathbb{C}.$$

Suppose now S geometrically marked by (e_1, \dots, e_{9-d}) as before. With the notation of the previous section, we have a line bundle \mathcal{L} on S and an associated contraction morphism $S \rightarrow \mathbb{P}(A)$, where $A = H^0(\mathcal{L})^*$, with E_i mapping to a singleton. The cuspidal curve K meets E_i in a single point p_i (with intersection number one). It is mapped isomorphically to its image in $\mathbb{P}(A)$.

The Zariski tangent space of K at its cusp is a line (with multiplicity two, but that will be irrelevant here). Let $u \in A^*$ be such that $u = 0$ defines the corresponding line in $\mathbb{P}(A)$. We may then extend u to a coordinate system (u, v, w) for A such that K is given by the equation $u^2w - v^3$ (this makes $[1 : 0 : 0]$ the unique flex point of K and $w = 0$ its tangent line). This coordinate system is for a given u almost unique: if (u, v', w') is any other such coordinate system, then $v' = cv$ and $w' = c^3w$ for some $c \in \mathbb{C}^\times$.

However, a choice of a generator $\kappa \in \mathfrak{l}_K$ singles out a natural choice (v, w) by requiring that the residue of κ on K is the restriction of $d(v/u)$. We put $t := v/u$ (we should write $v_\kappa, w_\kappa, t_\kappa$, but we do not want to overburden the notation, let us just remember that v, w, t are homogeneous of degree 1, 3, 1 in κ). The dependence of v and w on u is clearly homogeneous of degree 1. The smooth part K is then parameterized by $t \in \mathbb{C} \mapsto p(t) := [1 : t : t^3]$ such that dt corresponds to $\text{Res}_K \kappa$ and r_κ sends $(p(t)) - (p(t')) \in \text{Pic}^0(K)$ to $t - t' \in \mathbb{C}$.

Assume for the moment that the p_i 's are distinct (so that the E_i 's are irreducible and the λ_i 's are distinct). Let us denote the restriction of $u \in A^*$ to λ_i by u_i . This is clearly a coordinate for λ_i and hence a generator of λ_i^{-1} . We thus obtain the generator

$$\begin{aligned} \epsilon'_\kappa &:= (u_1 \cdots u_{9-d})^3 (du \wedge dv \wedge dw)^{-(9-d)} \\ &\in \lambda_1^{-3} \cdots \lambda_{9-d}^{-3} (\det A)^{9-d} = L(S, E). \end{aligned}$$

Since v and w are homogeneous of degree one in u , it follows that ϵ'_κ is independent of the choice of u . But they are homogeneous of degree 1 and 3 respectively in κ , and so ϵ'_κ is homogeneous of degree $-4(9-d)$ in κ .

Let us now see which linear forms we get on $\lambda_1^3 \oplus \cdots \oplus \lambda_{9-d}^3$ by restriction of cubic monomials in u, v, w . They will be of the form $(t_1^k u_1^3, \dots, t_{9-d}^k u_{9-d}^3)$ for some $k \geq 0$: for the monomial $u^a v^b w^c$ we have $k = b + 3c$. We thus get all integers $0 \leq k \leq 9$ except 8 (and 3 occurs twice since u^2v and w^3 yield

the same restriction):

$$\begin{array}{ccccccc} u^3 \mapsto 1 & u^2v \mapsto t & uv^2 \mapsto t^2 & u^2w, v^3 \mapsto t^3 & uvw \mapsto t^4 \\ v^2w \mapsto t^5 & uw^2 \mapsto t^6 & vw^2 \mapsto t^7 & w^3 \mapsto t^9. \end{array}$$

If we select $9 - d$ such monomials and compute the determinant of their restrictions to $\lambda_1^3 \oplus \cdots \oplus \lambda_{9-d}^3$, we see that that it is either zero or equal to $(u_1 \cdots u_{9-d})^3 \det((t_i)^{k_j})_{1 \leq i, j \leq 9-d}$ for some $0 \leq k_1 < \cdots < k_{9-d}$. The latter expression lies in $\mathbb{Z}[t_1, \dots, t_{9-d}]$ and is divisible by the discriminant that we get by taking $9 - d$ monomials corresponding to $1, t, \dots, t^{8-d}$, namely

$$\Delta(t_1, \dots, t_{9-d}) := \prod_{1 \leq i < j \leq 9-d} (t_i - t_j).$$

In other words, if we regard t_1, \dots, t_{9-d} as variables, then the $(9 - d)$ th exterior power over $\mathbb{Z}[t_1, \dots, t_{9-d}]$ of the homomorphism

$$\mathbb{Z}[t_1, \dots, t_{9-d}] \otimes \text{Sym}^3 A^* \rightarrow \bigoplus_{i=1}^{9-d} \mathbb{Z}[t_1, \dots, t_{9-d}] u_i^3$$

has its image generated by $\Delta(t_1, \dots, t_{9-d}) u_1^3 \cdots u_{9-d}^3$. That a division by $\Delta(t_1, \dots, t_{9-d})$ is appropriate is suggested by Remark 2.5 and so we use

$$e_\kappa := \Delta(t_1, \dots, t_{9-d}) e'_\kappa$$

as a generator of $\lambda_1^{-3} \cdots \lambda_{9-d}^{-3} (\det A)^{9-d}$ instead. We may rephrase this more sensibly in terms of our exact sequence

$$0 \rightarrow V^* \rightarrow \text{Sym}^3(A^*)(\det A)^{10} \rightarrow H^0(\mathcal{L}^3 \otimes \mathcal{O}_E \otimes \det A) \rightarrow 0.$$

We see that our coordinates define a basis of the middle term in such a manner that they make the sequence split: $9 - d$ cubic monomials that yield $1, t, \dots, t^{8-d}$ define a partial basis of $\text{Sym}^3(A^*)(\det A)^{10}$ whose $(9 - d)$ th exterior power maps onto e_κ . Notice that this remains true if some of the points p_i coalesce, that is, if S is just a geometrically marked Del Pezzo surface—this is in contrast to e'_κ . Since e_κ is homogeneous in κ of degree $-4(9 - d) + \binom{9-d}{2} = \frac{1}{2}d(9 - d)$, we have constructed an isomorphism

$$e : \mathfrak{l}_\kappa^{d(9-d)/2} \cong L(S, E).$$

Anticanonical cuspidal cubics on Fano surfaces. Let us return to the Fano case. If $p_i = p(t_i) = [1 : t_i : t_i^3]$, then the generator of λ_i dual to u_i is clearly $\tilde{p}_i = (1, t_i, t_i^3)$. It is not hard to verify that for i, j, k distinct in $\{1, \dots, 9 - d\}$, we have the following identity in $(\lambda_i \lambda_j \lambda_k)^{-1} \det A$:

$$\begin{aligned} \tilde{p}_i \wedge \tilde{p}_j \wedge \tilde{p}_k &= \Delta(t_i, t_j, t_k) (-t_i - t_j - t_k) (du \wedge dv \wedge dw)^{-1} = \\ &= \Delta(t_i, t_j, t_k) r_\kappa(h_{ijk}) (du \wedge dv \wedge dw)^{-1}. \end{aligned}$$

Similarly we find for i_1, \dots, i_6 distinct in $\{1, \dots, 9-d\}$ the following identity in the line $\lambda_{i_1}^{-2} \dots \lambda_{i_6}^{-2} \det(\text{Sym}^2 A)$:

$$\begin{aligned} \tilde{p}_{i_1}^2 \wedge \dots \wedge \tilde{p}_{i_6}^2 &= \mp(t_{i_1} + \dots + t_{i_6})\Delta(t_{i_1}, \dots, t_{i_6})(du \wedge dv \wedge dw)^{-4} = \\ &= \pm\Delta(t_{i_1}, \dots, t_{i_6})r_\kappa(2\ell - e_{i_1} - \dots - e_{i_6})(du \wedge dv \wedge dw)^{-4}. \end{aligned}$$

In this manner we get for example when $d = 3$:

$$\begin{aligned} \pm|123456||123||456| &= \\ &= \Delta(t_1, \dots, t_6)r_\kappa(\Delta(R'))u_1^3 \dots u_6^3(du \wedge dv \wedge dw)^{-6} = r_\kappa(\Delta')\epsilon_\kappa, \end{aligned}$$

where Δ' is the discriminant of the root subsystem of type $3A_2$ given by $\langle h_{12}, h_{23}, h_{45}, h_{56}, h_{123}, h \rangle$, and similarly

$$\pm|134||234||356||456||512||612| = r_\kappa(\Delta'')\epsilon_\kappa,$$

where Δ'' is the $3A_2$ -discriminant of $\langle h_{12}, h_{134}, h_{34}, h_{356}, h_{56}, h_{125} \rangle$. In this manner to each Coble covariant there is associated the discriminant of a A_2^3 -subsystem of the E_6 root system and vice versa (there are indeed 40 such subsystems). In similar fashion we find that for $d = 2$ a Coble covariant is the discriminant of a A_1^7 -subsystem of the E_7 root system and vice versa (there are 135 such); the two typical cases yield

$$\begin{aligned} \pm|135||146||234||256||457||127||367| &= r_\kappa(h_{135}h_{146}h_{234}h_{256}h_{457}h_{127}h_{367})\epsilon_\kappa, \\ \pm|123456||127||347||567| &= r_\kappa(h_{12}h_{34}h_{56}h_{127}h_{347}h_{567}h_7)\epsilon_\kappa. \end{aligned}$$

For $d = 4$, we do not get the discriminant of a root subsystem. The best way to describe this case is perhaps by just giving a typical case in terms of the standard representation of D_5 in \mathbb{R}^5 as in Bourbaki ([3]), where the roots are $\pm\epsilon_i \pm \epsilon_j$ with $1 \leq i < j \leq 5$. One such case is

$$\prod_{i \in \mathbb{Z}/5} (\epsilon_i - \epsilon_{i+1})(\epsilon_i + \epsilon_{i+1}) = \prod_{i \in \mathbb{Z}/5} (\epsilon_i^2 - \epsilon_{i+1}^2).$$

There are indeed 12 such elements up to sign.

Finally, we observe that for $d = 5$, we get

$$\pm|123||234||341||412| = r_\kappa(\Delta)\epsilon_\kappa,$$

where Δ is the discriminant of the full A_4 -system $\langle h_{12}, h_{23}, h_{34}, h_{123} \rangle$.

This makes it clear that in all these cases $\epsilon : \mathfrak{l}_K^{d(9-d)/2} \cong L(S, E)$ is independent of the marking once we identify $L(S, E)$ with $\det V$. Notice that the polynomials defining Coble covariants indeed have the predicted degree $\frac{1}{2}d(9-d)$.

We can restate this as follows. Consider the Lobatschevki lattice $\Lambda_{1,9-d}$ whose basis elements are denoted $(\ell, e_1, \dots, e_{9-d})$ (the inner product matrix is in diagonal form with $(+1, -1, \dots, -1)$ on the diagonal). Put $k := -3\ell + e_1 + \dots + e_{9-d}$ and let $(h_{123} = 3\ell - e_1 - e_2 - e_3, h_{12} = e_1 - e_2, \dots, h_{8-d,9-d} =$

$e_{8-d} - e_{9-d}$). This is a basis of k^\perp that is at the same time a root basis of a root system R_{9-d} .

If R is a root system, then let us denote its root lattice $Q(R)$, by $W(R)$ its Weyl group, by $\mathfrak{h}(R) := \text{Hom}(Q(R), \mathbb{C})$ the Cartan algebra on which R is defined. Let $\mathfrak{h}(R)^\circ \subset \mathfrak{h}(R)$ stand for the reflection hyperplane complement (which, in the parlance of Lie theory, is the set of its *regular elements*). We abbreviate the projectivizations of these last two spaces by $\mathbb{P}(R)$ and $\mathbb{P}(R)^\circ$. In the presence of a nondegenerate $W(R)$ -invariant symmetric bilinear form on $Q(R)$ we tacitly identify $\mathfrak{h}(R)$ with its dual.

So $Q(R_{9-d}) = k^\perp$. It is clear that a marking of a Del Pezzo surface amounts to an isomorphism $\text{Pic}(S) \cong \Lambda_{1,9-d}$ which sends the canonical class to k (and hence $\text{Pic}_0(S)$ to $Q(R_{9-d})$). These isomorphisms are simply transitively permuted by the Weyl group $W(R_{9-d})$. If we are given a marked Fano surface $(S; e_1, \dots, e_{9-d})$ of degree d and a rational 2-form κ on S whose divisor is a cuspidal curve K , then we can associate to these data an element of $\mathfrak{h}(R_{9-d})$ by

$$Q(R_{9-d}) \cong \text{Pic}_0(S) \rightarrow \text{Pic}^\circ(K) \xrightarrow{\tau_\kappa} \mathbb{C}.$$

It is known that we land in $\mathfrak{h}(R_{9-d})^\circ$ and that we thus obtain a bijection between the set of isomorphism classes of systems $(S; e_1, \dots, e_{9-d}; \kappa)$ and the points of $\mathfrak{h}(R_{9-d})^\circ$. This isomorphism is evidently homogeneous of degree one: replacing κ by $c\kappa$ multiplies the image by a factor c . In other words, the set of isomorphism classes of systems $(S; e_1, \dots, e_{9-d}; K)$, where $(S; e_1, \dots, e_{9-d})$ is a marked Fano surface of degree d and K is a cuspidal anticanonical curve on S can be identified with $\mathbb{P}(\mathfrak{h}_{9-d})^\circ$ (where we have abbreviated $\mathfrak{h}(R_{9-d})$ by \mathfrak{h}_{9-d}) in a such a manner that if l is a line in \mathfrak{h}_{9-d} associated to $(S; e_1, \dots, e_{9-d}; K)$, then l gets identified with the line $H^0(\omega_S(K))$.

We sum up the preceding in terms of the forgetful morphism $\mathbb{P}(\mathfrak{h}_{9-d})^\circ \rightarrow \mathcal{M}_{m,d}^\circ$:

Theorem 3.1. *Assume that $d \in \{2, 3, 4, 5\}$. Then the forgetful morphism from $\mathbb{P}(\mathfrak{h}_{9-d})^\circ$ to $\mathcal{M}_{m,d}^\circ$ is surjective and flat. It is covered by a natural isomorphism between the pull-back of the determinant bundle and $\mathcal{O}_{\mathbb{P}(\mathfrak{h}_{9-d})^\circ}(\frac{1}{2}d(9-d))$ and under this isomorphism the Coble covariants form a single $W(R_{9-d})$ -orbit, which up to a constant common scalar factor is as follows:*

- ($d=5$) the discriminant of the A_4 -system (a polynomial of degree 10),
- ($d=4$) the $W(D_5)$ -orbit of the degree 10 polynomial $\prod_{i \in \mathbb{Z}/5} (e_i^2 - e_{i+1}^2)$,
- ($d=3$) the discriminants of subsystems of type $3A_2$ (a $W(E_6)$ -orbit of polynomials of degree 9),
- ($d=2$) the discriminants of subsystems of type $7A_1$ (a $W(E_7)$ -orbit of polynomials of degree 7).

In particular, the Coble space \mathcal{C}_d can be identified with the linear span of the above orbit of polynomials.

The Weyl group representation \mathcal{C}_d was, at least for $d = 2$ and $d = 3$, already considered by Coble [4], although the notion of a Coxeter group

was not available to him. We shall consider these representations in more detail in Section 4. In that section we also investigate the rational map $\mathbb{P}(\mathfrak{h}_{\mathcal{R}_{d-2}}) \dashrightarrow \mathcal{M}_{m,d}^*$ defined by the morphism $\mathbb{P}(\mathfrak{h}_{\mathcal{R}_{d-2}})^\circ \rightarrow \mathcal{M}_{m,d}^\circ$.

Theorem 3.1 presents (for $d = 2, 3, 4, 5$) the moduli space $\mathcal{M}_{m,d}^\circ$ as a flat $W(\mathcal{R}_{d-2})$ -equivariant quotient of $\mathbb{P}(\mathcal{R}_{d-2})^\circ$ with $(d-2)$ -dimensional fibers. Admittedly that description is somewhat indirect from a geometric point of view. We will here offer in the next two subsections a somewhat more concrete characterisation when fiber and base are positive dimensional (so for $d = 3$ and $d = 4$). A fiber is then irreducible and we show that the $(d-1)$ $W(\mathcal{R}_{d-2})$ -invariant vector fields of lowest degree span in $\mathbb{P}(\mathcal{R}_{d-2})^\circ$ a $(d-2)$ -dimensional foliation whose leaves are the fibers of $\mathbb{P}(\mathcal{R}_{d-2})^\circ \rightarrow \mathcal{M}_{m,d}^\circ$. We first do the case $d = 3$.

The universal parabolic curve of a cubic surface. We begin with recalling a classical definition.

Definition 3.2. Let S be a Fano surface of degree 3. A point of S is said to be *parabolic* if there is a cuspidal anticanonical curve on S that has its cusp singularity at that point.

We may think of the surface S as anticanonically embedded in \mathbb{P}^3 as a smooth cubic surface. If $F(Z_0, Z_1, Z_2, Z_3)$ is a defining equation, then the locus of parabolic points is precisely the part of S where S meets its *Hessian surface* (defined by $\det(\partial^2 F / \partial Z_i \partial Z_j)$) transversally. So this is a nonsingular curve on S that need not be closed (in fact, it isn't: a point of S where its tangent plane intersects S in a union of a conic and a line tangent lies in the Zariski boundary of the parabolic curve). If we fix a marking for S , so that is determined a point of $p \in \mathcal{M}_{m,3}^\circ$, then the parabolic locus can be identified with the fiber of $\mathbb{P}(\mathcal{R}_{d-2})^\circ$ over p .

We now return to the situation of the beginning of this section, where we essentially have a fixed cuspidal cubic curve K in the projective plane \mathbb{P}^2 whose smooth part has in terms of affine coordinates the parameter form $(v, w) = (t, t^3)$ (this puts the cusp at infinity and the unique flex point at the origin). The points $p_1, \dots, p_6 \in \mathbb{P}_\ell^2$ that we blow up in order to produce S lie on K_{reg} ; we denote their $t(=v)$ -coordinates t_1, \dots, t_6 . The system $(S; e_1, \dots, e_6; K)$ defines a point $\tilde{p} \in \mathbb{P}(E_6)^\circ$ and (t_1, \dots, t_6) describes a point of $\mathfrak{h}(E_6)^\circ$ that lies over \tilde{p} . The vector fields X on \mathbb{P}^2 with the property that X be tangent to K at the points p_1, \dots, p_6 make up a vector space of dimension two. It contains the field $X_E = v\partial/\partial v + 3w\partial/\partial w$, which is tangent to K_{reg} everywhere (it generates a \mathbb{C}^\times -action on \mathbb{P}^2 that preserves K). If X is in this vector space, then

$$\hat{X} := \sum_i X(p_i) \frac{\partial}{\partial t_i}$$

is a tangent vector of $\mathfrak{h}(E_6)$ at $(t_1, \dots, t_6) \in \mathfrak{h}(E_6)^\circ$. For X_E this yields $\hat{X}_E = \sum_{i=1}^6 t_i \frac{\partial}{\partial t_i}$, in other words, we get the Euler field of $\mathfrak{h}(E_6)$ at (t_1, \dots, t_6) .

Lemma 3.3. *If X is not proportional to X_E , then the line in $\mathbb{P}(E_6)$ through \tilde{p} that is defined by \hat{X} is tangent to the fiber of $\mathbb{P}(E_6)^\circ \rightarrow \mathcal{M}_{m,3}^\circ$ at \tilde{p} .*

Proof. This is mostly a matter of geometric interpretation. If we view $\hat{X} = (X(p_1), \dots, X(p_6))$ as an infinitesimal displacement of the point configuration (p_1, \dots, p_6) in \mathbb{P}^2 , then \hat{X} does not effectively deform the corresponding Fano surface, because X is an infinitesimal automorphism of \mathbb{P}^2 . But if we view \hat{X} as an infinitesimal displacement (p_1, \dots, p_6) then it will induce a nontrivial line field (a priori with singularities) unless X is tangent to K . This last condition is equivalent to X being proportional to X_E . \square

We now calculate the resulting field on $\mathbb{P}(E_6)^\circ$. A vector field X on \mathbb{P}^2 has in our affine coordinates (v, w) the form

$$X = (a_0 + a_1v + a_2w + c_1v^2 + c_2vw) \frac{\partial}{\partial v} + (b_0 + b_1v + b_2w + c_1vw + c_2w^2) \frac{\partial}{\partial w}$$

Since we may calculate modulo X_E we assume $b_2 = 0$. The condition that X be tangent to C at p_i amounts to: $3t^2(a_0 + a_1t + a_2t^3 + c_1t^2 + c_2t^4) = b_0 + b_1t + c_1t^4 + c_2t^6$ for $t = t_i$, or equivalently, that $2c_2t^6 + 3a_2t^5 + 2c_1t^4 + 3a_1t^3 + 3a_0t^2 - b_1t - b_0$ has (the distinct) zeroes $t = t_i$ for $i = 1, \dots, 6$. This means that

$$\frac{a_2}{c_2} = -\frac{2}{3}\sigma_1, \quad \frac{c_1}{c_2} = \sigma_2, \quad \frac{a_1}{c_2} = -\frac{2}{3}\sigma_3, \quad \frac{a_0}{c_2} = \frac{2}{3}\sigma_4, \quad \frac{b_1}{c_2} = 2\sigma_5, \quad \frac{b_0}{c_2} = -2\sigma_6,$$

where σ_i stands for the i th symmetric function of t_1, \dots, t_6 . So we may normalize X by taking $c_2 = 1$. The value of X in p_i is in terms of the t -coordinate its x -component and hence equal to

$$(a_0 + a_1t_i + c_1t_i^2 + a_2t_i^3 + c_2t_i^4) \frac{\partial}{\partial t} = \left(\frac{2}{3}\sigma_4 - \frac{2}{3}\sigma_3t_i + \sigma_2t_i^2 - \frac{2}{3}\sigma_1t_i^3 + t_i^4 \right) \frac{\partial}{\partial t}.$$

It follows that

$$(*) \quad \hat{X} = \sum_{i=1}^6 \left(\frac{2}{3}\sigma_4 - \frac{2}{3}\sigma_3t_i + \sigma_2t_i^2 - \frac{2}{3}\sigma_1t_i^3 + t_i^4 \right) \frac{\partial}{\partial t_i}.$$

We see in particular that if regard \hat{X} as a vector field $((t_1, \dots, t_6)$ varies), then it is homogeneous of degree 3. The space of homogenous vector fields of degree 3 on $\mathfrak{h}(E_6)$ is as a W_6 -representation space isomorphic to $\text{Sym}^4(\mathfrak{h}^*(E_6)) \otimes \mathfrak{h}(E_6)$; this has a one-dimensional space of invariants and contains no other $W(E_6)$ -invariant one-dimensional subspace. It follows that \hat{X} is $W(E_6)$ -invariant and is characterized by this property up to a constant factor.

Corollary 3.4. *The fibration $\mathbb{P}(E_6)^\circ \rightarrow \mathcal{M}_{m,3}^\circ$ integrates the one dimensional foliation defined by a $W(E_6)$ -invariant vector field that is homogeneous of degree three.*

A natural way to produce such an invariant vector field is to take the nonzero $W(E_6)$ invariants polynomials f_2, f_5 on $\mathfrak{h}(E_6)$ of degree 2 and 5 (these are unique up to a constant factor); since f_2 is nondegenerate we can choose coordinates z_1, \dots, z_6 such that $f_2 = \sum_i z_i^2$. The gradient vector field relative to f_2 ,

$$\nabla f_5 = \sum_{i=1}^6 \frac{\partial f_5}{\partial z_i} \frac{\partial}{\partial z_i}$$

is a $W(E_6)$ -invariant homogeneous vector field of degree 3. So we can restate the preceding corollary as

Theorem 3.5. *Let \mathfrak{h} denote the natural representation space of a Coxeter group W of type E_6 . The natural W -invariant rational dimension-one foliation on $\mathbb{P}(\mathfrak{h})^\circ$ of degree 3 (i.e., the one defined by the gradient of a nonzero invariant quintic form with respect a nonzero (hence nondegenerate) invariant quadratic form on \mathfrak{h}) is algebraically integrable and has a leaf space that is in a W -equivariant manner isomorphic to the moduli space of marked smooth cubic surfaces.*

Moduli of degree 4 Del Pezzo surfaces. This is a slight modification of the argument for the degree 3 case. We have one point less and so by letting t_6 move over the affine line we may regard formula (*) as defining a one parameter family of vector fields on $\mathfrak{h}(D_5)$. For $i \geq 1$, we have $\sigma_i(t_1, \dots, t_6) = \sigma_i(t_1, \dots, t_5) + t_6 \sigma_{i-1}(t_1, \dots, t_5)$ and so we immediately see that we this is a linear family spanned by the two vector fields

$$\hat{X}_3 = \sum_{i=1}^5 \left(\frac{2}{3} \sigma_4 - \frac{2}{3} \sigma_3 t_i + \sigma_2 t_i^2 - \frac{2}{3} \sigma_1 t_i^3 + t_i^4 \right) \frac{\partial}{\partial t_i},$$

$$\hat{X}_2 = \sum_{i=1}^5 \left(\frac{2}{3} \sigma_3 - \frac{2}{3} \sigma_2 t_i + \sigma_1 t_i^2 - \frac{2}{3} t_i^3 \right) \frac{\partial}{\partial t_i}.$$

The subscript indicated of course the degree. Since \hat{X} is $W(E_6)$ -invariant, \hat{X}_3 and \hat{X}_2 will be invariant under the $W(E_6)$ -stabilizer of e_6 , that is, $W(D_5)$. The $W(D_5)$ -invariant vector fields on $\mathfrak{h}(D_5)$ form a free module on the polynomial algebra of $W(D_5)$ -invariant functions. The latter algebra has its generators f_2, f_4, f_5, f_6, f_8 in degrees indicated by the subscript. The generator f_2 is a nondegenerate quadratic form and the module of invariant vector fields is freely generated by the gradients of the f_i relative to f_2 , $X_{i-2} := \nabla f_i$ (these have the degree indicated by the subscript; X_0 is the Euler field). We conclude that the plane distribution on $\mathbb{P}(D_5)^\circ$ spanned by the vector fields X_2 and X_3 is also defined by \hat{X}_2 and \hat{X}_3 . We conclude :

Theorem 3.6. *Let \mathfrak{h} denote the natural representation space of a Coxeter group W of type D_5 . The natural W -invariant rational dimension-two foliation on $\mathbb{P}(\mathfrak{h})^\circ$ defined by the gradients of a nonzero invariant forms of degree 4 and 5 with respect a nonzero (hence nondegenerate) invariant quadratic form on*

\mathfrak{h} is algebraically integrable and has a leaf space that is in a W -equivariant manner isomorphic to the moduli space of marked Fano surfaces of degree 4.

Remark 3.7. The Frobenius integrability is remarkable, because it tells us that the degree four vector field $[X_2, X_3]$ does not involve the degree four generator X_4 . It is of course even more remarkable that it is algebraically so (in the sense that its leaves are the fibers of a morphism). It makes one wonder how often this happens. For instance one can ask: given a Coxeter arrangement complement \mathfrak{h}° and a positive integer k , when is the distribution on \mathfrak{h}° spanned by the subset of homogeneous invariant generating vector fields whose degree is $\leq k$ algebraically integrable?

4. COBLE'S REPRESENTATIONS

This section discusses the main properties of the representations of a Weyl group of type D_5 , E_6 or E_7 that we encountered in Theorem 3.1.

Macdonald's irreducibility theorem. We will use the following beautiful (and easily proved!) theorem of MacDonal [17] which states that the type of representation under consideration is irreducible.

Proposition 4.1 (Macdonald). *Let R be a root system, \mathfrak{h} the complex vector space it spans and $S \subset R$ a reduced root subsystem. Then the $W(R)$ -subrepresentation of $\text{Sym}^{|S|/2} \mathfrak{h}$ generated by the discriminant of S is irreducible. In particular, the Coble representations of type E_6 and E_7 are irreducible. \square*

Proof. Since the proof is short, we reproduce it here. If $L \subset \text{Sym}^{|S|/2} \mathfrak{h}$ denotes the line spanned by the discriminant of S , then $W(S)$ acts on L with the sign character. In fact, L is the entire eigensubspace of $\text{Sym}^{|S|/2} \mathfrak{h}$ defined by that character is, for if $G \in \text{Sym}^{|S|/2} \mathfrak{h}$ is such that $s(G) = -G$ for every reflection s in $W(S)$, then G is zero on each reflection hyperplane of $W(S)$ and hence divisible by the discriminant of S . Since G and the discriminant have the same degree, G must be proportional to it.

Let $V = \mathbb{C}[W(R)]L$ be the $W(R)$ -subrepresentation of $\text{Sym}^{|S|/2} \mathfrak{h}$ generated by L . We must prove that every W -equivariant map $\phi : V \rightarrow V$ is given by a scalar. From the preceding it follows that ϕ preserves L and so is given on L as multiplication by a scalar, λ say. Then $\phi - \lambda \mathbf{1}_V$ is zero on L and hence zero on V . \square

Unfortunately Macdonald's theorem does not come with an effective way to compute the degree of such representations and that is one of several good reasons to have a closer look at them. (The Coble representations had indeed been considered by Coble and presumably by others before him. Their irreducibility and their degrees were known at the time.)

It will be convenient (and of course quite relevant for the application we have in mind) to work for $d = 2, 3, 4$ with the Manin model of the R_{9-d} root system as sitting in the Lobatchevski lattice $\Lambda_{1,9-d}$ so that $\mathfrak{h}_{9-d} := \mathfrak{h}(R_{9-d})$ is

the orthogonal element of $k = -3\ell + e_1 + \cdots + e_{9-d}$. For $d = 2, 3, 4$ we have a corresponding (Coble) representation $\mathcal{C}_d = \mathcal{C}(R_{9-d})$, which in case $d = 3$ resp. $d = 2$ is spanned by the discriminants subsystems of type $3A_2$ resp. $7A_1$. We may regard \mathcal{C}_d as a linear system of hypersurfaces of degree 10 ($d = 4$), 9 ($d = 3$) or 7 ($d = 2$) in $\mathbb{P}(\mathfrak{h}_{9-d})$. Among our goals is to compute the dimension of this system and to investigate its separating properties.

The Manin basis recognizes one particular weight, namely the orthogonal projection of ℓ in \mathfrak{h}_{9-d} . Its $W(R_{9-d})$ -stabilizer is the symmetric group \mathcal{S}_{9-d} of e_1, \dots, e_{9-d} (a Weyl subgroup of type A_{8-d}). We shall denote by $\pi_{9-d} : \mathbb{C} \otimes \Lambda_{1,9-d} \rightarrow \mathfrak{h}_{9-d}$ the orthogonal projection. So $\pi_{9-d}(e_i) = e_i + \frac{1}{3}k$.

The Coble representation of a Weyl group of type E_6 . Here $R = R_6$ and the representation of $W(E_6)$ in question is the subspace $\mathcal{C}_3 \subset \text{Sym}^3 \mathfrak{h}_6$ spanned by the discriminants of subsystems of type $3A_2$ of R . Following Proposition 4.1, this representation is irreducible. We shall prove that $\dim \mathcal{C}_3 = 10$ and that quotients of elements of \mathcal{C}_3 separate the isomorphism types of cubic surfaces.

Lemma 4.2. *The Weyl group $W(R)$ acts transitively on the collection of ordered triples of mutually orthogonal roots. If $(\alpha_1, \alpha_2, \alpha_3)$ is such a triple, then*

- (i) *there is a root $\alpha \in R$ perpendicular to each α_i and this root is unique up to sign,*
- (ii) *the roots $\alpha_1, \alpha_2, \alpha_3, \alpha$ belong to (unique) subsystem of type D_4 ,*
- (iii) *there are precisely two subsystems of type $3A_2$ containing $\{\alpha_1, \alpha_2, \alpha_3\}$ and these two subsystems are interchanged by s_α .*

Proof. The transitivity assertion and the properties (i) and (ii) are known and a proof goes like this: the orthogonal complement of a root $\alpha_1 \in R$ is a subsystem $R' \subset R$ of type A_5 , the orthogonal complement of a root $\alpha_2 \in R'$ in R' is a subsystem $R'' \subset R'$ of type A_3 and the orthogonal complement of a root $\alpha_3 \in R''$ is a subsystem $R''' \subset R''$ of type A_1 . Since all the root systems encountered have the property that their Weyl group acts transitively on the roots, the first assertion follows. Notice that we proved (i) at the same time. The remaining properties now only need to be verified for a particular choice of $(\alpha_1, \alpha_2, \alpha_3)$.

We take this triple to be (h_{12}, h_{34}, h_{56}) . Then we may take $\alpha = h$ and we see that these are roots of the D_4 -system spanned by $h_{12}, h_{34}, h_{56}, h_{135}$. The two $3A_2$ -subsystems containing $\{h_{12}, h_{34}, h_{56}\}$ are then easily seen to be $\langle h_{12}, h_{134} \rangle \perp \langle h_{34}, h_{356} \rangle \perp \langle h_{56}, h_{125} \rangle$ and $\langle h_{12}, h_{156} \rangle \perp \langle h_{34}, h_{123} \rangle \perp \langle h_{56}, h_{345} \rangle$. We observe that s_h interchanges them. \square

The following notion is the root system analogue of its namesake introduced by Allcock and Freitag [1].

Lemma-Definition 4.3. *Let R be a root system of type E_6 and $S \subset R$ a subsystem of type $3A_2$. If $\alpha \in R$ is not orthogonal to any summand of S , then the roots in S orthogonal to α make up a subsystem of type $3A_1$ (which then must*

meet every summand of S). This sets up a bijection between the antipodal pairs $\{\pm\alpha\}$ that are not orthogonal to any summand of S and $3A_1$ -subsystems of S .

For (S, α) as above and S^+ a set of positive roots for S , the degree nine polynomial $(1 - s_\alpha)\Delta(S^+)$ is called a cross of R .

Proof. If we are given a A_2 -subsystem of R , then any root not in that subsystem is orthogonal to some root in that subsystem. This implies that in the above definition we can find in each of the three A_2 -summands a root that is orthogonal to α . Since α is not orthogonal to any summand, this root is unique up to sign and so the roots in S fixed by s form a $3A_1$ -subsystem as asserted.

Conversely, if $R' \subset S$ is a subsystem of type $3A_1$, and $\alpha \in R - R'$ is as in Lemma 4.2, then $s = s_\alpha$ has the desired property. \square

Lemma 4.4. *Let $S^+ \subset R$ and $\alpha \in R - S$ be as in Lemma 4.3. If $\alpha_1, \alpha_2, \alpha_3$ are the roots in S^+ perpendicular to α , then the cross $(1 - s_\alpha)\Delta(S^+)$ is divisible by $\alpha_1\alpha_2\alpha_3\alpha$:*

$$(1 - s_\alpha)\Delta(S^+) = \alpha_1\alpha_2\alpha_3\alpha F_1,$$

and the quotient $F_1 \in \text{Sym}^5 \mathfrak{h}_6$ is invariant under the Weyl group of the D_4 -subsystem that contains $\alpha_1, \alpha_2, \alpha_3, \alpha$.

Proof. It is clear that both $\Delta(S^+)$ and $s_\alpha\Delta(S^+) = \Delta(s_\alpha S^+)$ are divisible by $\alpha_1\alpha_2\alpha_3$. It is also clear that $(1 - s_\alpha)\Delta(S^+)$ is divisible by α . So F_1 is defined as an element of $\text{Sym}^5 \mathfrak{h}$.

We will now prove that there exists a $g \in \text{GL}(\mathbb{Q} \otimes \Lambda_{1,6})$ which centralizes the Weyl group in question and is such that the transform of F_1 under g^{-1} is a $W(R)$ -invariant in $\text{Sym}^5 \mathfrak{h}$. This will clearly suffice.

We may, in view of Lemma 4.2, assume without loss of generality that $(\alpha_1, \alpha_2, \alpha_3, \alpha) = (\mathfrak{h}_{12}, \mathfrak{h}_{34}, \mathfrak{h}_{56}, \mathfrak{h})$ so that the D_4 -subsystem containing these roots is $\langle \mathfrak{h}_{12}, \mathfrak{h}_{34}, \mathfrak{h}_{56}, \mathfrak{h}_{135} \rangle$. Denote by \mathfrak{h}' the subspace of \mathfrak{h} spanned by these roots.

We first recall a remarkable result due to Naruki. The set of exceptional classes that have inner product 1 with α is $\{e_1, \dots, e_6\}$ (this set and its s_α -transform make up what is classically known as a *double six*). Consider the element $(1 - s_\alpha) \prod_{i=1}^6 \pi_*(e_i) \in \text{Sym}^6 \mathfrak{h}$. It is clearly divisible by α and the quotient $F \in \text{Sym}^5 \mathfrak{h}$ will evidently be invariant under a Weyl subgroup of $W(R)$ of type $A_5 + A_1$. But according to Naruki ([15], p. 235) F is even invariant under all of $W(R)$.

The orthogonal complement of \mathfrak{h}' in $\mathbb{C} \otimes \Lambda_{1,6}$ is spanned by the members of the ‘anticanonical triangle’

$$(\epsilon_0 := \ell - e_1 - e_2, \epsilon_1 := \ell - e_3 - e_4, \epsilon_2 := \ell - e_5 - e_6)$$

and the intersection $\mathfrak{h} \cap \mathfrak{h}'^\perp$ is spanned by the differences $\epsilon_0 - \epsilon_1$ and $\epsilon_1 - \epsilon_2$. Let $g \in \text{GL}(\mathbb{Q} \otimes \Lambda_{1,6})$ be the transformation that is the identity on \mathfrak{h}' and takes e_i to $e_i + 2e_{i+1}$ for $i \in \mathbb{Z}/3$. This transformation preserves \mathfrak{h}'^\perp and hence commutes with all the transformations that preserve \mathfrak{h}' and act as the

identity on \mathfrak{h}'^\perp . We also note that $g(k) = 3k$, that g preserves the orthogonal complement of k and hence g commutes with π .

We claim that $g\pi(e_1) = -h_{134} \in R$. One easily checks that $2e_1 + \epsilon_0 + k \in \mathfrak{h}'$ and so

$$\begin{aligned} g(e_1) &= g\left(\frac{1}{2}(2e_1 + \epsilon_0 + k)\right) - \frac{1}{2}g(\epsilon_0) - \frac{1}{2}g(k) = \\ &= \frac{1}{2}(2e_1 + \epsilon_0 + k) - \frac{1}{2}(\epsilon_0 + 2e_1) - \frac{3}{2}k = \\ &= e_1 - \epsilon_1 - k = -\ell + e_1 + e_3 + e_4 - k = -h_{134} + k. \end{aligned}$$

Applying π to this identity yields $g\pi(e_1) = -h_{134}$.

We get similar formulas for the $g\pi(e_i)$ and thus find that

$$g\pi\{e_1, \dots, e_6\} = \langle -h_{134}, -h_{234} \rangle \perp \langle -h_{356}, -h_{456} \rangle \perp \langle -h_{125}, -h_{126} \rangle.$$

Notice that the union of this set with $\{h_{12}, h_{34}, h_{56}\}$ is a system of positive roots of a $3A_2$ -system. This union will be our S^+ . So g_* takes the polynomial $h_{12}h_{34}h_{56} \prod_{i=1}^6 \pi(e_i)$ to $\Delta(S^+)$. Since s_α commutes with g we have

$$\begin{aligned} (1 - s_\alpha)\Delta(S^+) &= h_{12}h_{34}h_{56}g\left((1 - s_\alpha) \prod_{i=1}^6 \pi(e_i)\right) = \\ &= h_{12}h_{34}h_{56}g(\alpha F) = h_{12}h_{34}h_{56}\alpha g(F) \end{aligned}$$

so that $F_1 = g(F)$. This proves the lemma. \square

Corollary 4.5. *For any three pairwise perpendicular roots in R there exists a cross that is divisible by their product. This cross is unique up to sign and is also divisible by a root perpendicular to these three. This yields a bijection between $4A_1$ -subsystems of R and antipodal pairs of crosses.* \square

Corollary 4.6. *Let $R' \subset R$ be a subsystem of type D_4 . The $4A_1$ -subsystems of R' define three crosses up to sign whose sum (up to sign) is zero. These crosses span a $W(R')$ -invariant plane in \mathcal{C}_3 . A quotient of the discriminants of two $4A_1$ -subsystems of R' is a quotient of two crosses.* \square

We now fix a D_5 -subsystem $R_o \subset R$. It has precisely 5 subsystems of type D_4 . As we just observed, each of these defines a plane in \mathcal{C}_3 . Therefore, the crosses associated to the $4A_1$ -subsystems of R_o span a subspace of \mathcal{C}_3 of dimension at most 10.

Lemma 4.7. *For every subsystem $S \subset R$ of type $3A_2$, $S \cap R_o$ is of type $2A_1 + A_2$ and hence contains three subsystems of type $3A_1$; for every such $3A_1$ -subsystem the associated $4A_1$ -subsystem of R is in fact contained in R_o . Moreover $S \mapsto S \cap R_o$ defines a bijection between the $3A_2$ -subsystems of R and the $2A_1 + A_2$ -subsystems of R_o and $W(R_o)$ acts transitively on both sets.*

Proof. It is easy (and left to the reader) to find one subsystem $S \subset R$ of type $3A_2$ such that $R_o \cap S$ has the stated properties. It therefore suffices to prove the transitivity property. This involves a simple count: The $W(R_o)$ -stabilizer of $R_o \cap S$ of R_o contains $W(R_o \cap S)$ as a subgroup of index two

(there is an element in the stabilizer that interchanges the A_1 -summands and is minus the identity on the A_2 -summand) and so the number of systems $W(R)$ -equivalent to $R_o \cap S$ is $|W(D_5)|/2|W(A_2 + 2A_1)| = 40$. That is just as many as there are $3A_2$ -subsystems of R . \square

We continue with the D_5 -subsystem $R_o \subset R$ that we fixed above. Let $S \subset R$ be any subsystem of type $3A_2$. By Lemma 4.7 $R_o \cap S$ is of type $2A_1 + A_2$. Let be s_1, s_2, s_3 be the three reflections in the Weyl group of the A_2 -summand. The two A_1 -summands and the antipodal root pair attached to s_i make up a $3A_1$ -subsystem $R_1^{(i)}$ of $R_o \cap S$. Each of these subsystems is contained in a unique $4A_1$ -subsystem. Let $s^{(i)}$ denote the reflection in the extra A_1 -summand. According to Lemma 4.7, $s^{(i)} \in W(R_o)$, so that $R_2^{(i)} := s^{(i)}S$ has the property that $R^{(i)} \cap R_o = R \cap R_o$.

Lemma 4.8. *The discriminant $\Delta(S^+)$ is fixed under $s' + s'' + s'''$, in other words,*

$$2\Delta(S^+) = (1 - s')\Delta(S^+) + (1 - s'')\Delta(S^+) + (1 - s''')\Delta(S^+),$$

where we note that the right hand side is a sum of three crosses attached to subsystems of R_o of type $4A_1$. In particular, \mathcal{C}_3 is generated by the crosses.

Proof. It is clear that $s_i\Delta(S^+) = -\Delta(S^+)$. Since $s_3 = s_1s_2s_1^{-1}$, we have $R_1''' = s_1(R_1'')$. This implies that $s''' = s_1s''s_1$ and so

$$(s'' + s''')\Delta(S^+) = (s'' + s_1s''s_1)\Delta(S^+) = (1 - s_1)s''\Delta(S^+) = (1 - s_1)\Delta(s''S^+).$$

Since $s_1 \notin W(s''S)$, the right hand side is a cross. We claim that this cross equals the cross $(1 - s')\Delta(S^+)$ up to sign. For this it suffices to show that there exist four perpendicular roots such that each is divisible by three of them. It is clear that $(1 - s_1)s''\Delta(S^+)$ is divisible by a root attached to s_1 and by the roots in the two A_1 -summands of S^+ (for these are unaffected by s'' and s_1). On the other hand, $(1 - s')\Delta(S^+)$ is divisible by a root attached to s' and the roots in the two A_1 -summands of S . It remains to observe that the roots attached to s_1 and s' are perpendicular.

Thus $(s'' + s''')\Delta(S^+) = \pm(1 - s')\Delta(S^+)$. Suppose the minus sign holds, so that $1 - s' + s'' + s'''$ kills $\Delta(S^+)$. The cyclic permutation $1 + s' + s'' - s'''$ then also kills $\Delta(S^+)$ and hence so will $1 + s''$. In other words, $\Delta(S^+)$ will be anti-invariant under s'' . Since $s'' \notin W(S)$, this is a contradiction. Hence the plus sign holds and the lemma follows. \square

Theorem 4.9. *The planes defined by the five subsystems of R_o of type D_4 make up a direct sum decomposition of \mathcal{C}_3 . In particular \mathcal{C}_3 is the irreducible representation of the E_6 -Weyl group of degree 10.*

Proof. Lemma 4.8 shows that \mathcal{C}_3 is spanned by the crosses attached to $4A_1$ -subsystems of R_o . So the five planes in question span \mathcal{C}_3 and $\dim \mathcal{C}_3 \leq 10$. The irreducible representations of $W(R)$ of degree < 10 are the trivial representation, the sign representation (which are both of degree 1) and the

defining representation (of degree 6) and \mathcal{C}_3 is clearly neither of these. The theorem follows. \square

We now determine the common zero set of the Coble covariants. We first make some general remarks that also apply to the D_5 and the E_7 -case. The zero set of a Coble invariant is a union of reflection hyperplanes and hence each irreducible component of their common intersection, $Z_r \subset \mathfrak{h}_r$, is an intersection of reflection hyperplanes. As Z_r is invariant under the Weyl group, so is the collection of its irreducible components. So an irreducible component is always the translate of common zero set of a subset of the given root basis of R_r . (This subset need not be unique.)

Proposition 4.10. *The common zero set $Z_6 \subset \mathfrak{h}$ of the members of \mathcal{C}_3 is the union of the linear subspaces that are pointwise fixed by a Weyl subgroup of type A_3 .*

Proof. We first verify that for any A_3 -subsystem of R , the subspace of \mathfrak{h} perpendicular to it is in the common zero set of the members of \mathcal{C}_3 . Since an A_3 -subsystem is contained in a D_5 -subsystem, it is in view of Lemma 4.7 enough to show that a $A_2 + 2A_1$ -subsystem and a A_3 -subsystem in a D_5 -system always meet. This is easily verified.

We next show that Z_6 is not larger. Any subsystem generated by fundamental roots that does not contain a A_3 -system is contained in a subsystem of type $2A_2 + A_1$. There is a single Weyl group equivalence class of such subsystems and so it suffices to give two subsystems of R , of type $3A_2$ and of type $2A_2 + A_1$ that are disjoint. We take $\langle \mathfrak{h}_{12}, \mathfrak{h}_{23}, \mathfrak{h}_{45}, \mathfrak{h}_{56}, \mathfrak{h}_{123}, \mathfrak{h} \rangle$ and $\langle \mathfrak{h}_{16}, \mathfrak{h}_{125}, \mathfrak{h}_{34}, \mathfrak{h}_{136}, \mathfrak{h}_{25} \rangle$. \square

Question 4.11. Is \mathcal{C}_3 the space of degree 9 polynomials on \mathfrak{h} that vanish on Z_6 ? This is probably equivalent to the completeness of \mathcal{C}_3 as a linear system on $\mathcal{M}_{m,3}^*$ (which is known, though in a rather indirect manner, see Remark 5.10).

The Coble representation of a Weyl group of type E_7 . The Weyl group $W(E_7)$ decomposes as $W_+(E_7) \times \{1, c\}$, where $W_+(E_7) \subset W(E_7)$ is the subgroup of elements that have determinant one in the Coxeter representation and $c \in W(E_7)$ is minus the identity in the Coxeter representation. This implies that every irreducible representation of $W(E_7)$ is obtained as an irreducible representation of $W_+(E_7)$ plus a degree as to whether c acts as 1 or as -1 .

We know that the representation of $W(E_7)$ defined by \mathcal{C}_2 (which we recall, is spanned by products of seven pairwise perpendicular roots of the E_7 root system) is irreducible and we want to prove:

Proposition 4.12. *The representation \mathcal{C}_2 of $W(E_7)$ is of degree 15 and the nontrivial central element of $W(E_7)$ acts as -1 .*

It is known that there is just one isomorphism types of irreducible representations of $W_+(E_7)$ in degree 15 and so Proposition 4.12 identifies the isomorphism type of the representation.

In what follows R stands for the root system R_7 of type E_7 .

Lemma 4.13. *The Weyl group $W(R)$ acts transitively on the collection of $7A_1$ -subsystems of R . If we are given a subsystem R' of type $2A_1$, then the roots perpendicular to R' make up a subsystem of type $A_1 + D_4$. In particular, there is a unique subsystem of type $3A_1$ that contains R' and is orthogonal to a subsystem of type D_4 . Conversely, the roots perpendicular to a given subsystem of R of type D_4 make up a system of type $3A_1$.*

Proof. This lemma is known and the proof is standard. The first assertion follows from the fact that the roots orthogonal to a given root of R form a subsystem of type D_6 and the roots orthogonal to a root of a root system of type D_6 form a subsystem of type $D_4 + A_1$.

Any root subsystem of R of type D_4 is saturated and so a root basis of this subsystem extends to a root basis of R . As the group $W(R)$ acts transitively on the set of root bases, it also acts transitively on the set of subsystems of type D_4 . \square

So if we have a subsystem $R_1 \subset R$ of type $7A_1$, then any two summands of R_1 (making up a subsystem $R' \subset R_1$ of type $2A_1$) determine a third summand and the remaining 4 summands will lie in a D_4 -subsystem. In this way we can construct a 2-dimensional simplicial complex with 7 vertices indexed by the summands of R_1 : three vertices span a 2-simplex if and only if the orthogonal complement of the sum of their associated A_1 -summands is of type D_4 . The preceding lemma tells us that every edge is in exactly one 2-simplex. Probably the $W(E_7)$ -stabilizer of R_1 is the full automorphism group of this complex.

The subsystems of type $7A_1$ make up two S_7 -orbits, represented by

- (A) $\langle h_7, h_{12}, h_{34}, h_{56}, h_{127}, h_{347}, h_{567} \rangle$, 105 in number and
- (B) $\langle h_{123}, h_{145}, h_{167}, h_{256}, h_{247}, h_{357}, h_{346} \rangle$, of which there are 30.

We designate by the same letters (A) and (B) the type of the corresponding product of roots.

Lemma 4.14. *Let F be a product of roots of type (B). Then the S_7 -stabilizer of F acts transitively on its factors and has order $7 \cdot 3 \cdot 2^3$. The subgroup that stabilizes a given factor is isomorphic to S_4 .*

Proof. Let F be of type (B) and let α be a factor of F . Write $\alpha = h_{abc}$. Then the other factors are of the form $h_{axy}, h_{azw}, h_{bxz}, h_{byw}, h_{cxw}, h_{cyz}$, where x, y, z, w are the distinct elements of $\{1, 2, \dots, 7\} - \{a, b, c\}$. So these factors are given by an indexing by a, b, c of the three ways we can split $\{1, 2, \dots, 7\} - \{a, b, c\}$ into two pairs. This description proves that S_7 is transitive of the collection of pairs (F, α) with stabilizer mapping isomorphically onto the permutation group of $\{x, y, z, w\}$. The lemma follows. \square

Lemma 4.15. *The space \mathcal{C}_2 is spanned by the 30 root products of type (B) and is annihilated by $\sum_{w \in \mathcal{S}(i,j,k)} \text{sign}(w)w$ for any 3-element subset $\{i, j, k\}$ of $\{1, \dots, 7\}$.*

For the proof we need:

Lemma 4.16. *A root system S of type D_4 contains exactly 3 subsystems of type $4A_1$ and the discriminants of these three subsystems (relative some choice of positive roots) are such that a signed sum is zero. More precisely, if $S_o \subset S$ is a subsystem of type $4A_1$, then the (8) reflections in $W(S) - W(S_o)$ decompose into two equivalence classes with the property that two reflections s, s' belong to different classes if and only if they do not commute. In that case $S_o, sS_o, s'S_o$ are the distinct $4A_1$ -subsystems of S and if f is the product of 4 pairwise perpendicular roots in S_o , then $f = s(f) + s'(f)$. The plane in the fourth symmetric power of the complex span of the root system generated by these discriminants affords an irreducible representation of the Weyl group of the root system.*

Proof. In terms of the standard model for the D_4 -system, the set of vectors $\pm \epsilon_i \pm \epsilon_j$, $1 \leq i < j \leq 4$ in Euclidean 4-space, the $4A_1$ -subsystems correspond to the three ways of partitioning $\{1, 2, 3, 4\}$ into parts of size 2. For instance, the partition $\{\{1, 2\}, \{3, 4\}\}$ yields $\{\pm \epsilon_1 \pm \epsilon_2, \pm \epsilon_3 \pm \epsilon_4\}$, whose discriminant is (up to sign) equal to $\epsilon_1^2 \epsilon_3^2 + \epsilon_2^2 \epsilon_4^2 - \epsilon_1^2 \epsilon_4^2 + \epsilon_2^2 \epsilon_3^2$. We can verify the lemma for $s = s_{\epsilon_1 - \epsilon_3}$, $s' = s_{\epsilon_1 - \epsilon_4}$ and deduce the general case from that.

The last clause is easily verified. \square

Proof of Lemma 4.15. Consider $F = h_7 h_{12} h_{34} h_{56} h_{127} h_{347} h_{567}$ (a typical root product of type (A)). The four factors that are not of type (1), $h_7, h_{12}, h_{34}, h_{56}$, lie in a subsystem of type D_4 . If we let s resp. s' be the reflection in h_{135} resp. h_{246} , then

$$\begin{aligned} s(h_7 h_{12} h_{34} h_{56}) &= h_{246} h_{235} h_{145} h_{136} \\ s'(h_7 h_{12} h_{34} h_{56}) &= h_{135} (-h_{146}) (-h_{236}) (-h_{245}) = -h_{135} h_{146} h_{236} h_{245}. \end{aligned}$$

Notice that the second product is obtained from the first by applying minus the transposition (34). According to Lemma 4.16 we then have

$$(AB) \quad h_7 h_{12} h_{34} h_{56} = (1 - (34)) h_{246} h_{235} h_{145} h_{136}.$$

After multiplying both sides with $h_{127} h_{347} h_{567}$, we see that F has been written as a difference of two products of type (B): $f = (1 - (34)) G$ with

$$G := h_{127} h_{347} h_{567} h_{246} h_{235} h_{145} h_{136}.$$

In particular, the type (B)-products generate \mathcal{C}_2 . It follows from Lemma 4.14 that the S_7 -stabilizer of G has two orbits in the collection of 3-element subsets $\{i, j, k\} \subset \{1, \dots, 7\}$: those for which h_{ijk} is a factor of G and those for which there exist a factor h_{abc} of G with $\{a, b, c\} \cap \{i, j, k\} = \emptyset$. So there are only two cases to verify.

We first do the case $I = \{3, 4, 5\}$. We are then in the second case because $I \cap \{1, 2, 7\} = \emptyset$ and h_{127} is a factor of G . To this purpose we look at the D_4 -system defined by the pair h_7, h_{127} : it is the system that contains the four roots $h_{34}, h_{56}, h_{347}, h_{567}$. The reflections s resp. s' perpendicular to h_{367} resp. h_{467} lie in this D_4 summand and do not commute. We have

$$\begin{aligned} s(h_{34}h_{56}h_{347}h_{567}) &= h_{467}(-h_{357})h_{64}h_{35} \\ s'(h_{34}h_{56}h_{347}h_{567}) &= (-h_{367})(-h_{457})h_{63}h_{45} \end{aligned}$$

so that

$$h_{34}h_{56}h_{347}h_{567} - h_{467}h_{357}h_{46}h_{35} + h_{367}h_{457}h_{36}h_{45} = 0.$$

The second resp. third term are obtained from the first by applying to it minus the transposition (45) resp. minus the transposition (35), so that

$$(1 - (45) - (35)) h_7 h_{12} h_{127} h_{34} h_{56} h_{347} h_{567} = 0.$$

If we combine this with (AB), and observe that

$$(1 - (45) - (35))(1 - (34)) = \sum_{w \in \mathcal{S}(3,4,5)} \text{sign}(w)w,$$

then we find that the latter kills $G = h_{127}h_{347}h_{567}h_{246}h_{235}h_{145}h_{136}$.

An instance of the first case, namely the assertion that G is also killed by $\sum_{w \in \mathcal{S}(1,2,7)} \text{sign}(w)wG$, follows by exploiting the symmetry properties of G : the transpositions (34) and (35) have the same effect on G as resp. (12) and (17). This implies that

$$\sum_{w \in \mathcal{S}(1,2,7)} \text{sign}(w)wG = \sum_{w \in \mathcal{S}(3,4,5)} \text{sign}(w)wG = 0.$$

□

Corollary 4.17. *We have $\dim \mathcal{C}_2 \leq 15$.*

Proof. Let $f \in \mathcal{C}_2$. Since a monomial of type (B) has a unique factor of the form (12a), $a \in \{3, \dots, 7\}$, we can write f accordingly: $f = (124)f_3 + \dots + (127)f_7$. For every a , we have 6 type (B) monomials corresponding to the ways we index the splittings of the complement of a in $\{3, 4, 5, 6, 7\}$ by $\{1, 2, a\}$. The symmetric group $\mathcal{S}(1, 2, a)$ permutes these six root products simply transitively. These root products satisfy the corresponding alternating sum relation and so we can arrange that each f_a is a linear combination of 6 monomials whose alternating sum of coefficients is zero. If we take as our guiding idea to make a as small as possible, then it turns out that in half of the cases we can do better.

Let us first assume $a \in \{5, 6, 7\}$. We then invoke the relation defined by $\{3, 4, a\}$:

$$\sum_{w \in \mathcal{S}(3,4,a)} \text{sign}(w)wf_a = 0.$$

Four of the six terms have a factor (123) or (124), whereas the other two have a factor (12a) and combine to $(1 - (34))f_\alpha$. So this relation allows us to arrange that f_α and $(34)f_\alpha$ have the same coefficient. We thus make f_α vary in a space of dimension ≤ 3 . If $\alpha \in \{6, 7\}$, then we can repeat this game with $\{4, 5, \alpha\}$. This allows us to assume in addition that f_α and $(45)f_\alpha$ have the same coefficient. But then f_α must have all its coefficients equal. So f_α varies in a space of dimension ≤ 1 for $\alpha = 6, 7$, of dimension ≤ 3 for $\alpha = 5$, and of dimension ≤ 5 for $\alpha = 3, 4$. This proves that $\dim \mathcal{C}_2 \leq 15$. \square

Proof of Proposition 4.12. If we combine Proposition 4.1 and Corollary 4.17, we see that \mathcal{C}_2 is an irreducible representation of $W(R)$ of dimension ≤ 15 . Since $W(R) = W(R)_+ \times \{1, c\}$, it will then also be an irreducible representation of $W(R)_+$. The symmetric bilinear form on the root lattice induces a nondegenerate form on the root lattice modulo two times the weight lattice (this is a \mathbb{F}_2 -vector space of dimension 6). This identifies $W(R)_+$ with the symplectic group $\mathrm{Sp}(6, \mathbb{Z}/2)$ and it is well-known (see for instance [7], where this group is denoted $S_6(2)$) that the irreducible representations of dimension < 15 are the trivial representation, the sign representation and the standard representation of degree 7. It is easy to see that \mathcal{C}_2 is neither of these. Since c acts as -1 in \mathcal{C}_2 , the proposition follows. \square

The roots orthogonal to an A_5 -subsystem of an E_6 -system make up a system of type A_1 or A_2 . In terms of our root basis, they are represented by $\langle h_{23}, h_{34}, h_{45}, h_{56}, h_{67} \rangle$ (with $\langle h_1 \rangle$ as the perpendicular system) and the system $\langle h_{123}, h_{34}, h_{45}, h_{56}, h_{67} \rangle$ (with $\langle h_1, h_{12} \rangle$ as perpendicular system). We shall call an A_5 -subsystem of the second type *special*. Conversely, the roots perpendicular to a A_2 -subsystem form a special A_5 -system. Since the A_2 -subsystems make up a single Weyl group equivalence class, the same is true for the special A_5 -subsystems.

Proposition 4.18. *If we regard \mathcal{C}_2 as a vector space of degree 7 polynomials on \mathfrak{h}_7 , then their common zero set Z_7 is the union of the linear subspaces perpendicular to a system of type D_4 or to a special system of type A_5 .*

Proof. The D_4 -subsystems constitute a single Weyl group equivalence class and so we may take as our system the one spanned by the fundamental roots $\langle h_{23}, h_{34}, h_{45}, h_{123} \rangle$. We must show that every $7A_1$ -subsystem of R meets this D_4 -system. The positive roots of the D_4 -system are $\{h_{ij}\}_{2 \leq i < j \leq 5}$ and $\{h_{1ij}\}_{2 \leq i < j \leq 5}$. It is easy to see from our description that every $7A_1$ -system of type (A) contains a root h_{ij} with $2 \leq i < j \leq 5$. Similarly, we see that every $7A_1$ -system of type (B) contains a root h_{1ij} with $2 \leq i < j \leq 5$.

We argue for the special A_5 -system $\langle h_{123}, h_{34}, h_{45}, h_{56}, h_{67} \rangle$ in a similar fashion. Its positive roots are $\{h_{ij}\}_{3 \leq i < j \leq 7}$ and $\{h_{12i}\}_{3 \leq i \leq 7}$. Every $7A_1$ -system of type (A) contains a root h_{ij} with $3 \leq i < j \leq 7$ and every $7A_1$ -system of type (B) contains a root h_{12i} with $3 \leq i \leq 7$.

It remains to show that this exhausts Z_7 . Every subsystem of R that does not contain a D_4 -subsystem has only components of type A. If in addition

it does not contain a special A_5 -system, then any such a subsystem is Weyl group-equivalent to a proper subsystem in the (nonsaturated) A_7 -system spanned by the fundamental roots $h_{12}, h_{34}, \dots, h_{67}$ and the highest root h_1 . The latter has as its positive roots $\{h_{ij}\}_{1 \leq i < j \leq 7} \cup \{h_i\}_{i=1}^7$ and is therefore disjoint with the $7A_1$ -subsystem $\langle h_{123}, h_{145}, h_{167}, h_{256}, h_{247}, h_{357}, h_{346} \rangle$. This implies that Z_7 is as asserted. \square

Question 4.19. Is \mathcal{C}_2 the space of degree 7 polynomials on \mathfrak{h}_7 that vanish on Z_7 ? We expect this to be equivalent to the question whether \mathcal{C}_2 is complete as a linear system on $\mathcal{M}_{m,2}^*$.

5. THE COBLE LINEAR SYSTEM

Let A be a vector space of dimension three so that $\mathbb{P}(A)$ is a projective plane. Given a numbered set (p_1, \dots, p_N) of $N \geq 5$ points in $\mathbb{P}(A)$ that are in generic position, then for any 5-tuple (i_0, \dots, i_4) with i_0, \dots, i_4 pairwise distinct and taken from $\{1, \dots, N\}$, the four ordered lines $p_{i_0}p_{i_1}, p_{i_0}p_{i_2}, p_{i_0}p_{i_3}, p_{i_0}p_{i_4}$ through p_{i_0} have a cross ratio. The collection of cross ratio's thus obtained make up a complete projective invariant of (p_1, \dots, p_N) : we may choose coordinates such that $p_1 = [1 : 0 : 0]$, $p_2 = [0 : 1 : 0]$, $p_3 = [0 : 0 : 1]$, $p_4 = [1 : 1 : 1]$ and the coordinates $[z_0 : z_1 : z_2]$ for p_i , $i > 4$, are then given by cross ratio's. For instance, $z_1 : z_2 = (p_1p_2 : p_1p_3 : p_1p_4 : p_1p_i)$. If $a_i \in A$ represents p_i , then we can write the this as a cross ratio of 4 lines in the plane $a_1 \wedge A \subset \wedge^2 A$: $(a_1 \wedge a_2 : a_1 \wedge a_3 : a_1 \wedge a_4 : a_1 \wedge a_i)$.

Now let us observe that if (v_1, v_2, v_3, v_4) is a generic ordered 4-tuple in a vector space T of dimension two, then the corresponding points in $\mathbb{P}(T)$ have a cross ratio that can be written as a ratio of two elements of $\det(T)^2$, namely $(v_1 \wedge v_4)(v_2 \wedge v_3) : (v_2 \wedge v_4)(v_1 \wedge v_3)$. If we apply this to to the present case, then we get

$$z_1 : z_2 = (p_1p_2 : p_1p_3 : p_1p_4 : p_1p_i) = |12i||134| : |13i||124|,$$

where we used the Coble notation. Thus the cross ratio's formed in this manner allow us to reconstruct (p_1, \dots, p_5) up to projective equivalence. We can express this in terms of roots as follows.

Lemma 5.1. *Let $(S; e_1, \dots, e_{9-d})$ be a marked Del Pezzo surface of degree $d \leq 4$, $S \rightarrow \check{\mathbb{P}}^2(H^0(S, \ell))$ the contraction morphism defined by the linear system $|\ell|$ (as usual) and p_i the image of E_i . Then p_1, \dots, p_4 are in general position if and only if none of the roots in the A_4 -subsystem generated by $(h_{123}, h_{12}, h_{23}, h_{34})$ is effective in $\text{Pic}(S)$.*

If that is the case and K is a cuspidal anticanonical curve on S , then for $i > 4$ the cross ratio $(p_1p_2 : p_1p_3 : p_1p_4 : p_1p_i)$ equals the ratio of the two elements of the line $\text{Pic}(K)^\circ$ given by $r_K(h_{2i}h_{12i}h_{34}h_{134})$ and $r_K(h_{3i}h_{13i}h_{24}h_{124})$.

Proof. The first part is left to the reader as an exercise. As to the second part, choose affine coordinates (x, y) in $\check{\mathbb{P}}^2(H^0(S, \ell))$ such that the K is given by $y^3 = x^2$. So $p_i = (t_i, t_i^3)$ for some t_i . For $i \neq 1$, the line p_1p_i has tangent

$[t_i - t_1 : t_i^3 - t_1^3] = [1 : t_i^2 + t_i t_1 + t_1^2]$. So the cross ratio of the lines $p_1 p_i$ involves factors of the form

$$(t_j^2 + t_j t_1 + t_1^2) - (t_i^2 + t_i t_1 + t_1^2) = (t_j - t_i)(t_j + t_i + t_1), \quad 2 \leq i < j \leq 5.$$

If we use the x -coordinate to identify $\text{Pic}_0(K)$ with \mathbb{C} , then such a factor can be written $r_K(h_{ij}h_{1ij})$. The last assertion follows. \square

Remark 5.2. Notice that the roots that appear in the numerator resp. the denominator of $h_{25}h_{12i}h_{34}h_{134} : h_{3i}h_{13i}h_{24}h_{124}$ are four pairwise perpendicular roots which all lie in a single D_4 -subsystem.

The Coble system in the degree four case. We first consider a Fano surface of degree 5. We recall that such a surface S can be obtained by blowing up 4 points of a projective plane in general position, and so is unique up to isomorphism. Any automorphism of this surface that acts trivially on its Picard group preserves every exceptional curve (= line) and hence is the identity. It follows that the automorphism group of S is the Weyl group $W(A_4)$. There are 10 lines on S . If five of them make up pentagon, then their sum is an anticanonical divisor. There are 12 such pentagons and they generate the anticanonical system. Let us now fix a marking (e_1, e_2, e_3, e_4) for S . To every $p \in S$ we associate a marked Del Pezzo surface $(S_p; e_1, \dots, e_5)$ of degree 4 by letting S_p be the blowup of S in p and letting e_5 be the class of the exceptional divisor. This defines a rational map $S \dashrightarrow \mathcal{M}_{m,4}^*$. We will see that this is in fact an isomorphism.

Proposition 5.3. *The 12 Coble covariants for D_5 span a complete linear system of dimension 6 and define an embedding of $\mathcal{M}_{m,4}^*$ in a projective space of dimension 5. The image is ‘the’ anticanonically embedded Fano surface of degree 5 (so that the Coble system is anticanonical) with $\mathcal{M}_{m,4}^* - \mathcal{M}_{m,4}^\circ$ mapping onto the union of its ten lines. The divisor of every Coble covariant is a pentagon on this Fano surface and every pentagon thus occurs.*

Proof. Let A be a complex vector space of dimension 3 and let $p_1, \dots, p_5 \in \mathbb{P}(A)$. We first assume that p_1, \dots, p_4 are in general position (i.e., no three collinear). We then adapt our coordinate system accordingly: $p_i = [a_i]$, with $a_1 = (1, 0, 0)$, $a_2 = (0, 1, 0)$, $a_3 = (0, 0, 1)$ and $a_4 = (1, 1, 1)$. If $a_5 = (z_0, z_1, z_2)$, then typical determinants involving a_5 are:

$$|125| = z_2, \quad |145| = z_2 - z_1.$$

So for instance

$$|415||152||523||234||341| = (z_1 - z_2) \cdot (-z_2) \cdot z_0 \cdot 1 \cdot -1 = z_0 z_1 z_2 - z_1^2 z_2.$$

We obtain in this manner the 6 polynomials $z_0 z_1 z_2 - z_i^2 z_j$, $i \neq j$, and it is easily verified that any other Coble covariant is a linear combination of these. They are visibly linearly independent and hence form a basis for \mathcal{C}_5 . It is precisely the linear system of cubic curves that pass through p_1, \dots, p_4 . So the Coble system is anticanonical and defines an embedding of the blowup

S of $\mathbb{P}(A)$ in p_1, \dots, p_4 to \mathbb{P}^5 . The remaining assertions are verified in a straightforward manner. \square

Remark 5.4. This proposition and its proof show that the moduli space $\mathcal{M}_{m,4}^*$ is as a variety simply obtained from \mathbb{P}^2 by blowing up the vertices p_1, \dots, p_4 of the coordinate simplex. This argument then also shows that there is universal semistable marked Del Pezzo surface of degree 4, $\mathcal{S}_{m,4} \rightarrow \mathcal{M}_{m,4}^*$:

over $p_5 \in \mathcal{M}_{m,4}^*$ we put the blowup of the surface $\mathcal{M}_{m,4}^*$ in p_5 so that $\mathcal{S}_{m,4}$ is simply $\mathcal{M}_{m,4}^* \times \mathcal{M}_{m,4}^*$ blown up along the diagonal with one of the projections serving as the structural morphism.

The Coble system in the degree three case. Our discussion starts off with the following lemma.

Lemma 5.5. *The Coble system \mathcal{C}_3 has no base points.*

Proof. The Coble linear system pulled back to $\mathbb{P}(h_6)$ has according to Proposition 4.10 as its base point locus the projective arrangement $\mathbb{P}(Z_6)$, the union of the fixed point hyperplanes of Weyl subgroups of type A_3 . Since \mathcal{M}_3^* is a quotient of $\mathbb{P}(h_6) - \mathbb{P}(Z_6)$ it follows that \mathcal{C}_3 has no base points and hence defines a morphism to a \mathbb{P}^9 . \square

We use what we shall call the *Naruki model* of $\mathcal{M}_{m,3}^*$. This is based on a particular way of getting a degree 3 Fano surface as a blown-up projective plane: we suppose the points in question to be labelled p_i, q_i with $i \in \mathbb{Z}/3$ and to lie on the coordinate lines of \mathbb{P}^2 as follows:

$$\begin{aligned} p_0 &= [0 : 1 : a_0], & p_1 &= [a_1 : 0 : 1], & p_2 &= [1 : a_2 : 0], \\ q_0 &= [0 : 1 : b_0], & q_1 &= [b_1 : 0 : 1], & q_2 &= [1 : b_2 : 0]. \end{aligned}$$

For the moment we assume that blowing up these points gives rise to a Fano surface S so that in particular none of the a_i, b_i is zero and $a_i \neq b_i$. If we blow up these points, the strict transform of the coordinate triangle is an anticanonical curve K (it is a tritangent of the corresponding cubic surface). This ‘partial rigidification’ reduces the projective linear group $\mathrm{PGL}(3, \mathbb{C})$ to its maximal subtorus that leaves the coordinate triangle invariant. The following expressions are invariant under that torus

$$\alpha_i := a_i/b_i \ (i \in \mathbb{Z}/3), \quad \delta := -b_0 b_1 b_2,$$

and together they form a complete projective invariant of the configuration. Notice the formulas

$$a_0 b_1 b_2 = -\alpha_0 \delta, \quad a_0 a_1 b_2 = -\alpha_0 \alpha_1 \delta, \quad a_0 a_1 a_2 = -\alpha_0 \alpha_1 \alpha_2 \delta.$$

As explained in the Appendix of [16], α_i and δ have a simple interpretation in terms of the of pair (S, K) : if we denote the exceptional curves A_i, B_i , and E is the strict transform of the line through q_1 and q_2 , then the cycles $A_i - B_i$, $i \in \mathbb{Z}/3$ and $B_0 - E$ span in $\mathrm{Pic}(S)$ the orthogonal complement to the components of K and the numbers in question can be interpreted as their images in $\mathrm{Pic}^0(K) \cong \mathbb{C}^\times$. The classes themselves make up the basis

of a D_4 root system with the last one representing the central node. We denote the 4-torus for which $\alpha_0, \alpha_1, \alpha_2, \delta$ is a basis of characters by T . This torus comes with an action of $W(D_4)$ and this makes it an adjoint torus of type D_4 . We denote by T° the open set of its regular elements. This is the complement of the union of reflection hypertori, i.e., the locus where none of the D_4 -roots is 1. It has the interpretation as the moduli space of marked nonsingular cubic surfaces with the property that a particular tritangent (which is entirely given by the marking) has not its three lines collinear. If we want to include that case too, we must first blow up the identity element of T , $\text{Bl}_1(T) \rightarrow T$, and then remove the strict transforms of the reflection hypertori. This open subset, $\text{Bl}_1(T)^\circ \subset \text{Bl}_1(T)$, is a model for the moduli space of marked nonsingular cubic surfaces, in other words, it can be identified with $\mathcal{M}_{m,3}^\circ$. The modular interpretation implies that this variety has a $W(E_6)$ -action, although only the action of a Weyl subgroup of type D_4 is manifest. The action of the two missing fundamental reflections was written down by Naruki and Sekiguchi: the one that in the Dynkin diagram is attached to α_i is given by

$$\begin{aligned}\alpha_i &\mapsto -\alpha_0\alpha_1\alpha_2\delta^2 \frac{1 - \alpha_i}{1 - \alpha_0\alpha_1\alpha_2\delta^2} \\ \alpha_{i\pm 1} &\mapsto \frac{(1 - \alpha_0\alpha_{i\pm 1}\delta)(1 - \alpha_0\alpha_1\alpha_2\delta)}{(1 - \alpha_0\delta)(\alpha_{i\pm 1} - \alpha_0\alpha_1\alpha_2\delta)} \\ \delta &\mapsto \delta^{-1} \frac{(1 - \alpha_i\delta)(1 - \alpha_0\alpha_1\alpha_2\delta^2)}{(1 - \alpha_i)(1 - \alpha_0\alpha_1\alpha_2\delta)}.\end{aligned}$$

(We give this formula because of its remarkable form only—we shall not use it.) Subsequently Naruki [16] found a nice $W(E_6)$ -equivariant smooth projective completion of this space with a normal crossing divisor as boundary. What is more relevant here is a (projective) blow-down of his completion that was also introduced by him. We shall take Naruki's construction of the latter as our guide and reprove some of his results in the process.

We identify the complexification of $\text{Hom}(\mathbb{C}^\times, T)$ with the Lie algebra \mathfrak{t} of T , so that the latter has a natural \mathbb{Q} -structure. The decomposition Σ of $\mathfrak{t}(\mathbb{R})$ into its $W(D_4)$ -chambers has its rays spanned by the coweights that lie in the $W(D_4)$ -orbit of a fundamental coweight. One of these is the orbit of coroots and has 24 elements; the other three consist of minuscule weights and are a single orbit under the full automorphism group of the D_4 -system; it has also 24 elements. If we remove the faces that contain a coroot we obtain a coarser decomposition of $\mathfrak{t}(\mathbb{R})$ that we denote by Σ' ; a maximal face of Σ' is now an orbit of a Weyl chamber under the stabilizer of a coroot (a type $3A_1$ -Weyl group).

Let $T \subset T_\Sigma$ be the associated torus embedding. It is a smooth with normal crossing boundary. The boundary divisors are in bijective correspondence with the above coweights. We shall refer to those that correspond to coroots

resp. minuscule weights as *toric coroot divisors* resp. *toric minuscule weight divisors*. So there are 24 of each.

Now blow up successively in T_Σ : the identity element (in other words the fixed point set of $W(D_4)$), the fixed point sets of the Weyl subgroups of type A_3 , the fixed point sets of the Weyl subgroups of type A_2 . We denote the resulting blowup \hat{T}_Σ . In this blowup the exceptional divisors of type A_3 have been separated and each is naturally a product. To be precise, a $W(A_3)$ -Weyl subgroup $G \subset W(D_4)$ has as its fixed point locus in \hat{T}_Σ a copy of a \mathbb{P}^1 whose tangent line at the identity is the G -fixed point set in \mathfrak{t} and the divisor associated to G is then naturally the product of the projective line \hat{T}_Σ^G and the projective plane $\mathbb{P}(\mathfrak{t}/\mathfrak{t}^G)$ blown up in the fixed points of the A_2 -Weyl subgroups of G (there are four such and they are in general position).

The Coble system on Naruki's completion, together with the $W(E_6)$ -action on it, was identified in [14] (5.9 and 4.5). It is the pull-back of

$$H^0(T_\Sigma, \mathcal{O}(D_S + 2D_R) \otimes \mathfrak{m}_e^3)$$

to \hat{T}_Σ , here D_S, D_R are the sum of the 24 divisors corresponding to the rays spanned by the minuscule weight and the coroots respectively, and \mathfrak{m}_e is the ideal sheaf of the identity element $e \in T_\Sigma$.

We now give an explicit description of the Coble covariants in terms of $(\alpha_0, \alpha_1, \alpha_2, \delta)$. For this we begin with observing the following simple identities:

$$|p_0 p_1 p_2| = \alpha_0 \alpha_1 \alpha_2 + 1, \quad |p_i q_i p_{i+1}| = (b_i - a_i) a_{i+1}, \quad |p_i q_i p_{i-1}| = b_i - a_i.$$

A straightforward computation yields

$$|p_0 p_1 p_2 q_0 q_1 q_2| = \pm (b_0 - a_0)(b_1 - a_1)(b_2 - a_2)(1 - \alpha_0 \alpha_1 \alpha_2 b_0 b_1 b_2).$$

We substitute these values in the formulae for the Coble covariants, but for reasons similar as in Section 3 we divide these by $(b_0 - a_0)(b_1 - a_1)(b_2 - a_2)$. For example,

$$\begin{aligned} |p_0 p_1 p_2 q_0 q_1 q_2| \cdot |p_0 q_0 p_1| \cdot |q_1 p_2 q_2| &= \\ &= \pm (1 - \alpha_0 \alpha_1 \alpha_2 b_0 b_1 b_2) \cdot (b_0 - a_0) a_1 \cdot (b_2 - a_2) \\ &= \pm \alpha_1 \delta (1 - \alpha_0 \alpha_1 \alpha_2 \delta) (1 - \alpha_0) (1 - \alpha_2) \end{aligned}$$

and

$$\begin{aligned} |p_0 q_0 p_1| \cdot |p_0 q_0 q_2| \cdot |p_1 q_1 q_2| \cdot |p_1 p_2 q_2| \cdot |p_0 q_1 p_2| \cdot |q_0 q_1 p_2| &= \\ = \pm \frac{(b_0 - a_0) a_1 \cdot (b_0 - a_0) \cdot (b_1 - a_1) b_2 \cdot (b_2 - a_2) (\alpha_0 b_1 \alpha_2 + 1) (b_0 b_1 \alpha_2 + 1)}{(b_0 - a_0)(b_1 - a_1)(b_2 - a_2)} &= \\ = \pm \alpha_0 \alpha_1 \delta (1 - \alpha_0) (1 - \alpha_0 \alpha_2 \delta) (1 - \alpha_2 \delta). \end{aligned}$$

We thus find for the Coble covariants (40 up to sign) the following expressions:

$$\begin{array}{ll}
1 & \delta(1 - \alpha_0)(1 - \alpha_1)(1 - \alpha_2) \\
2 & \alpha_0\alpha_1\alpha_2\delta^2(1 - \alpha_0)(1 - \alpha_1)(1 - \alpha_2) \\
3 & (1 - \alpha_0\delta)(1 - \alpha_1\delta)(1 - \alpha_2\delta) \\
4 & \alpha_0\alpha_1\alpha_2\delta(1 - \alpha_0\delta)(1 - \alpha_1\delta)(1 - \alpha_2\delta) \\
5 & (1 - \alpha_0\alpha_1\delta)(1 - \alpha_1\alpha_2\delta)(1 - \alpha_2\alpha_0\delta) \\
6 & \delta(1 - \alpha_0\alpha_1\delta)(1 - \alpha_1\alpha_2\delta)(1 - \alpha_2\alpha_0\delta) \\
7_i & (1 - \alpha_{i-1}\delta)(1 - \alpha_{i+1}\delta)(1 - \alpha_0\alpha_1\alpha_2\delta) \\
8_i & \alpha_i\delta(1 - \alpha_{i-1})(1 - \alpha_{i+1})(1 - \alpha_0\alpha_1\alpha_2\delta^2) \\
9_i & \delta(1 - \alpha_{i-1})(1 - \alpha_{i+1})(1 - \alpha_0\alpha_1\alpha_2\delta^2) \\
10_i & \alpha_{i-1}\alpha_{i+1}\delta(1 - \delta)(1 - \alpha_i\alpha_{i-1}\delta)(1 - \alpha_i\alpha_{i+1}\delta) \\
11_i & (1 - \delta)(1 - \alpha_i\alpha_{i+1}\delta)(1 - \alpha_i\alpha_{i-1}\delta) \\
12_i & \alpha_i\delta(1 - \alpha_{i+1}\delta)(1 - \alpha_{i-1}\delta)(1 - \alpha_0\alpha_1\alpha_2\delta) \\
\\
13 & (1 - \delta)(1 - \alpha_0\alpha_1\alpha_2\delta)(1 - \alpha_0\alpha_1\alpha_2\delta^2) \\
14_i & \alpha_{i-1}\alpha_{i+1}\delta(1 - \delta)(1 - \alpha_i)(1 - \alpha_i\delta) \\
15_i & (1 - \alpha_i\delta)(1 - \alpha_{i-1}\alpha_{i+1}\delta)(1 - \alpha_0\alpha_1\alpha_2\delta^2) \\
16_i & \delta(1 - \alpha_i)(1 - \alpha_{i-1}\alpha_{i+1}\delta)(1 - \alpha_0\alpha_1\alpha_2\delta) \\
17_{ijk} & \alpha_i\delta(1 - \alpha_j)(1 - \alpha_k\delta)(1 - \alpha_j\alpha_k\delta)
\end{array}$$

Notice that the zero sets in T of any of these expressions is the union of the reflection hypertori of a subsystem of type $3A_1$ (in the first 12 cases) or the A_2 (for the last 5). We now quote from [16] (Proposition 11.3):

Theorem 5.6 (Naruki). *There is a projective contraction*

$$\hat{T}_\Sigma \rightarrow \check{T}_\Sigma$$

which contracts each A_3 -divisor along the projection on its 2-dimensional factor. The contracted variety \check{T}_Σ is nonsingular and the action of $W(E_6)$ on $(\text{Bl}_e T)^\circ$ extends regularly to it. This action is transitive on the collection of 40 divisors that are of toric coroot type or of A_2 -type.

The 40 divisors in question are easily seen to be pairwise disjoint. Naruki also shows (Section 12 of *op. cit.*) that each of these can be contracted to a point. We can see that quickly using the theory of torus embeddings: the 24 toric coroot divisors get contracted if we replace in the above discussion the decomposition Σ of $\mathfrak{t}(\mathbb{R})$ by the coarser one, Σ' , that we obtain by removing the faces that contain a coroot. The $W(E_6)$ -action and the theorem above imply that this contraction is then also possible for the remaining 16 divisors. The singularities thus created, for instance the one defined by the ray spanned by the coroot δ^\vee , can be understood as follows: the natural affine T -invariant neighborhood of the point of $T_{\Sigma'}$ defined by the ray spanned by the coroot δ^\vee is Spec of the algebra generated by the elements of the orbit of δ under the Weyl group of the $3A_1$ -subsystem $\langle \alpha_0, \alpha_1, \alpha_2 \rangle$: $\text{Spec} \mathbb{C}[\delta, \alpha_0\delta, \alpha_1\delta, \dots, \alpha_0\alpha_1\alpha_2\delta]$. This is a cone over the Veronese embedding of $(\mathbb{P}^1)^3$.

The following more precise result is in [14] (Theorem 5.7) and follows also from [19].

Theorem 5.7. *The Coble covariants generate on \check{T}_Σ a linear system without base points that has the property that its restriction each of the 40 divisors of toric coroot type or of A_2 -type is trivial. The resulting morphism to a nine dimensional projective space realizes Naruki's contraction.*

Outline of proof. We have a natural decomposition of T_Σ into strata by type: D_4 (yielding the identity element), A_3 , A_2 and $\{1\}$ (being open in T_Σ). There is a corresponding decomposition of \hat{T}_Σ (and of \check{T}_Σ , but we find it more convenient to work on the former), albeit that strata are then indexed by chains of strata in T_Σ that are totally ordered for incidence. We first check that along every stratum the Coble covariants define, modulo the stated contractions property, an embedding. For this, the $W(D_4)$ -equivariance allows us to concentrate on the open subset $U = \text{Spec}(\mathbb{C}[\alpha_0, \alpha_1, \alpha_2, \delta])$ of T_Σ and its preimage \hat{U} in \hat{T}_Σ .

It is clear from the expressions we found that the Coble covariants generate $\mathbb{C}[\delta, \alpha_0\delta, \dots, \alpha_0\alpha_1\alpha_2\delta]$ after we localize away from the kernels of the roots (that is, we make each expression root -1 invertible). So we have an embedding on the corresponding open subset of \check{T}_Σ (this contains the singular point defined the coroot δ^\vee). A closer look at the equations shows that this is in fact even true if we allow some of the roots to be 1, provided that they are mutually perpendicular. In other words, the linear system defines an embedding on the intersection of the open stratum with U .

Now let Z be the A_3 -stratum that is open in $\alpha_0 = \alpha_1 = \delta = 1$. All Coble covariants vanish on Z and we readily verify that Coble covariants generate the ideal defining Z . Thus the system has no base points on the blowup of Z and the system contracts this exceptional divisor along the Z -direction.

Now let us look at an A_2 -stratum Z' , say the one that is open in $\alpha_0 = \delta = 1$. We observe that the restriction of every covariant to Z' is proportional with $(1 - \alpha_1)(1 - \alpha_2)(1 - \alpha_1\alpha_2)$ (and can be nonzero). So the linear system will define the constant map on Z' (or the exceptional divisor over Z').

We turn to the situation at the identity of T . Every covariant vanishes there with order three and has for initial part a product of three roots, viewed as linear forms on \mathfrak{t} . Up to sign, the roots in such a product are the positive roots of a $3A_1$ -subsystem (in the first 12 cases) or of a A_2 -system (the last 5 cases). The preceding implies that the linear system restricted to a A_2 -line in $\mathbb{P}(\mathfrak{t})$ is constant. With some work we find that the Coble linear system is without base points and generates the ideal defining the identity away from the union of the A_2 -loci.

The remaining strata on \hat{T}_Σ are defined by 'flags': chains of strata above totally ordered by incidence, with $\{1\} < Z' < Z$ as a typical degenerate case. In that situation, one checks that the covariants generate on that stratum the ideal $\mathcal{I}_{\{e\}}\mathcal{I}_{Z'}\mathcal{I}_Z$. We thus see that we have a local embedding along this stratum. The other strata are dealt with in the same way.

This shows that the linear system defines local embeddings modulo the contraction property. So \check{T}_Σ is defined as a projective quotient of \hat{T}_Σ and the linear system maps it as local embedding to a nine dimensional projective space. It remains to see that the images of the strata are disjoint. This is left to the reader. \square

Remark 5.8. The construction of the Naruki quotient comes with a stratification and as one may expect, each of its members has a modular interpretation. We here give that interpretation without proof. As mentioned, the open stratum T° is the moduli space of systems $(S; e_1, \dots, e_6; K)$ with $(S; e_1, \dots, e_6)$ a marked Fano surface of degree 3 (equivalently, a cubic surface) and K an anticanonical divisor made up of three nonconcurrent exceptional curves (so that the isomorphism $S \rightarrow \bar{S}$ maps K onto a tritangent $\bar{K} \subset \bar{S}$). Suppose now that S is merely a Del Pezzo surface whose configuration of (-2) -curves is nonempty, but disjoint with K . Then that configuration is of type rA_1 and we are on a stratum contained in T of type rA_1 ($1 \leq r \leq 4$) or it is of type $rA_1 + sA_1$ with $r \geq 1$ and we are on one of the 24 points that are images of toric coroot divisors (these are the punctual strata of T_Σ). In these cases \bar{K} is a genuine tritangent of \bar{S} (that lies in the smooth part of \bar{S}). The other strata are loci for which \bar{K} is no longer a tritangent: if \bar{K} defines an Eckardt point (so that K consists of three distinct concurrent exceptional curves), then we find ourselves in the stratum that is open in the preimage of the unit element of T . If \bar{K} becomes a union of a double line and another line, then it contains two distinct A_1 -singularities of \bar{S} and K is of the form $(2E + E' + C + C')$, where E, E' are exceptional curves and C, C' are (-2) -curves with E', C, C' pairwise disjoint and meeting E normally. We are then on a stratum that is open in the image of an A_3 -locus in \check{T} . If \bar{K} becomes a triple line, then it contains two distinct A_2 -singularities of \bar{S} and K is of the form $(3E + C + C')$, where E is exceptional curve and C, C' are disjoint A_2 -curves meeting E normally. We are then representing one of the 16 punctual strata that are images of an A_2 -locus in T .

Corollary 5.9. *The GIT completion of the moduli space of marked cubic surfaces, $\mathcal{M}_{m,3}^*$, is $W(E_6)$ -equivariantly isomorphic to the Naruki contraction of \hat{T}_Σ . The Coble linear system embeds $\mathcal{M}_{m,3}^*$ in projective nine space.*

Proof. The Coble linear system on \hat{T}_Σ is without base points and so the resulting morphism $f : \hat{T}_\Sigma \rightarrow \mathbb{P}^9$ realizes the Naruki contraction. Recall that we have an identification of $\mathcal{M}_{m,3}^\circ$ with $(\text{Bl}_e T)^\circ$. This isomorphism clearly extends to a morphism $\mathcal{M}_{m,3}^* \rightarrow f(\hat{T}_\Sigma)$. This morphism is birational and since $f(\hat{T}_\Sigma)$ is normal, it must be a contraction. \square

Remark 5.10. It is known [2] that the moduli space of stable cubic surfaces is Galois covered by the complex 4-ball with an arithmetic group Γ as Galois group. The group Γ has a single cusp and this cusp represents the minimal strictly stable orbit of cubic surfaces (i.e., those having three A_2 -singularities). This gives \mathcal{M}_3 the structure of an arithmetic ball quotient

for which \mathcal{M}_3^* is its Baily-Borel compactification. Allcock and Freitag [1] have used Γ -modular forms to construct an embedding of this Baily-Borel compactification in a 9-dimensional projective space. This is precisely the embedding that appears here (see also Freitag [13] and van Geemen [14]). Via this interpretation it also follows that the Coble system is complete [12].

Remark 5.11. The linear system \mathcal{C}_3 can be also studied by restricting it, as was done in [6], to the exceptional divisor $\mathbb{P}(\mathfrak{h}_5)$ of the blowup of the e_6 -point in $\mathbb{P}(\mathfrak{h}_6)$. The generic point of $\mathbb{P}(\mathfrak{h}_5)$ has a modular interpretation: it parameterizes marked cubic surfaces with a point where the tangent space meets the surface in the union of a conic and a line tangent to that conic. The marking determines the line, but not the conic, for the system of conics on a cubic surface that lie in plane that contains a given line on that surface has two members that are tangent to the line. So we have a natural involution ι on that space. The projective space $\mathbb{P}(\mathfrak{h}_5)$ can be seen as the projective span of the D_5 -subsystem R_5 , spanned by the roots not involving e_6 .

In order to be explicit we also use the standard model for the D_5 root system, i.e., the model for which $\epsilon_i - \epsilon_{i+1} = h_{i,i+1}$ ($i = 1, \dots, 4$) and $\epsilon_4 + \epsilon_5 = h_{123}$. This makes $W(R_5)$ the semidirect product of the group of permutations of the basis elements $\epsilon_1, \dots, \epsilon_5$ and the group of sign changes in the basis elements $(\epsilon_1, \dots, \epsilon_5) \mapsto (\pm\epsilon_1, \dots, \pm\epsilon_5)$ with an even number of minus signs. We denote the basis dual to $(\epsilon_1, \dots, \epsilon_5)$ by (x_1, \dots, x_5) . The Coble covariant we attached to the $3A_2$ -system $\langle h_{ij}, h_{jk}, h_{lm}, h_{m6}, h_{ijk}, h \rangle$ gives, after dividing by a common degree 4 factor, the quintic form on \mathfrak{h}_5 defined by

$$h_{ij}h_{jk}h_{ik}h_{lm}h_{ijk} = (x_i - x_j)(x_j - x_k)(x_i - x_k)(x_l^2 - x_m^2),$$

whereas the Coble covariant attached to $\langle h_{ij}, h_{ilm}, h_{lm}, h_{km6}, h_{k6}, h_{ijk} \rangle$ gives

$$h_{ij}h_{jlm}h_{ilm}h_{lm}h_{ijk} = (x_i - x_j)(x_j + x_k)(x_i + x_k)(x_l^2 - x_m^2).$$

These all lie in a single $W(R_5)$ -orbit as predicted by Lemma 4.7. It can be easily checked that the base locus of the system on $\mathbb{P}(\mathfrak{h}_5)$ is the set of points fixed by a Weyl subgroup of type A_3 . There are two orbits of subroot systems of type A_3 : one has 40 elements and is represented by $\langle \epsilon_1 - \epsilon_2, \epsilon_2 - \epsilon_3, \epsilon_3 - \epsilon_4 \rangle$ and the other has 10 elements and is represented by $\langle \epsilon_1 - \epsilon_2, \epsilon_2 - \epsilon_3, \epsilon_2 + \epsilon_3 \rangle$. So the base locus is a union of 50 lines. The locus where two such lines meet are the (16) fixed points of a Weyl subgroup of type A_4 and the (5) fixed points of a Weyl subgroup of type D_4 . Blowing up first the 21 points and then the strict transforms of the 50 lines we obtain a smooth fourfold $\widetilde{\mathbb{P}(\mathfrak{h}_5)}$ in which the strict transforms of the planes defined by the (40) root subsystems of type A_2 have become disjoint. The Coble system defines a morphism

$$\Psi : \widetilde{\mathbb{P}(\mathfrak{h}_5)} \longrightarrow \mathbb{P}^9$$

which is generically two to one: it identifies the orbits of the involution ι which on $\mathbb{P}(\mathfrak{h}_5)$ is given as the rational map

$$[x_1 : \dots : x_5] \longmapsto [x_1^{-1} : \dots : x_5^{-1}].$$

The morphism Ψ is ramified along the exceptional divisors over the A_4 -points and contracts the exceptional divisors over the A_3 -lines to planes and the 40 planes of type A_2 to points.

The Coble system in the degree two case. An analogue of Lemma 5.5 holds:

Proposition 5.12. *The linear system \mathcal{C}_2 is without base points on $\mathcal{M}_{m,2}^*$. Its restriction to $\mathcal{M}_{m,2}^\circ$ is an embedding.*

Proof. The proof of the first assertion only differs from the one of Lemma 5.12 essentially by replacing the reference to Proposition 4.10 by a reference to Proposition 4.18: the pull-back of \mathcal{C}_2 to $\mathbb{P}(\mathfrak{h}_7)$ has according to Proposition 4.18 as base locus $\mathbb{P}(Z_7)$ the projective arrangement defined by the subsystems of type D_4 and the special subsystems of type A_5 . Since the map from $\mathbb{P}(\mathfrak{h}_7) - \mathbb{P}(Z_7)$ to \mathcal{M}_2^* is a surjective morphism, \mathcal{C}_2 is without base points.

The second assertion follows from Lemma 4.13: if we are given a D_4 -subsystem of a root system of type E_6 , then orthogonal to it we have a $3A_1$ -system. Thus two disjoint $4A_1$ -subsystems of the given D_4 -system have a discriminants whose quotient is a quotient of Coble covariants. According to Lemma 5.1 the quotient of two such $4A_1$ -subsystems is a cross ratio. Hence all (generalized) cross ratio's are recovered from the Coble covariants. If the seven points are in general position, then these cross ratio's determine the points up to projective equivalence. \square

Recall from Proposition 1.4 that we identified \mathcal{C}_2 with a space of sections of a square root of $\mathcal{O}_{\mathcal{M}_2^*}(1)$.

Conjecture 5.13. *The linear system \mathcal{C}_2 is without base points and hence defines an injective morphism from the moduli space \mathcal{M}_2^* of semistable quartic curves with level two structure to a 14-dimensional projective space.*

We also expect that there is an analogue of the results of Naruki and Yoshida with the role of D_4 taken by E_6 .

REFERENCES

- [1] D. Allcock, E. Freitag: *Cubic surfaces and Borchers products*, Comment. Math. Helv. 77 (2002), 270–296.
- [2] D. Allcock, J.A. Carlson, D. Toledo: *The complex hyperbolic geometry of the moduli space of cubic surfaces*, J. Algebraic Geom. 11 (2002), 659–724.
- [3] N. Bourbaki: *Groupes et algèbres de Lie*, Ch. 4,5 et 6, 2^{ième} éd., Masson, Paris 1981.
- [4] A.B. Coble: *Algebraic Geometry and Theta Functions*, AMS New York (1929).
- [5] E. Colombo, B. van Geemen: *The Chow group of the moduli space of marked cubic surfaces*, Ann. Mat. Pura Appl. (4) 183 (2004), 291–316.
- [6] E. Colombo, B. van Geemen: *A family of marked cubic surfaces and the root system D_5* , available at math.AG/0509561, to appear in Internat. J. Math.

- [7] J.H. Conway, R.T. Curtis, S.P. Norton, R.A. Parker, R.A. Wilson: *Atlas of finite groups*, Oxford University Press, Eynsham, 1985.
- [8] F.R. Cossec, I.V. Dolgachev: *Enriques Surfaces I*, Birkhäuser, PM 76 (1989).
- [9] M. Demazure: *Surfaces de Del Pezzo*, in Séminaire sur les Singularités des Surfaces, Lecture Note in Math. 777, Springer, Berlin-New York, 1980.
- [10] I. Dolgachev, D. Ortland: *Point sets in projective spaces and theta functions*, Astérisque 165 (1988), 210 pp.
- [11] I. Dolgachev, B. van Geemen, S. Kondō: *A complex ball uniformization of the moduli space of cubic surfaces via periods of K3 surfaces*, J. Reine Angew. Math. 588 (2005), 99–148.
- [12] E. Freitag: *A graded algebra related to cubic surfaces*, Kyushu J. Math. 56 (2002), no. 2, 299–312.
- [13] E. Freitag: *Comparison of different models of the moduli space of marked cubic surfaces*, Proc. Japanese-German seminar Riushi-do, T. Ibukiyama, W. Kohnen eds., 74-79 (2002), also available from www.rzuser.uni-heidelberg.de/~t91/index4.html.
- [14] B. van Geemen: *A linear system on Naruki's moduli space of marked cubic surfaces*, Internat. J. Math. 13 (2002), 183-208.
- [15] B. Hunt: *The geometry of some special arithmetic quotients*. Lecture Notes in Mathematics, 1637, Springer-Verlag, Berlin (1996).
- [16] I. Naruki: *Cross ratio variety as a moduli space of cubic surfaces*, Proc. London Math. Soc. 45 (1982), 1-30.
- [17] I.G. Macdonald: *Some irreducible representations of Weyl groups*, Bull. London Math. Soc. 4 (1972), 148–150.
- [18] Y. Manin: *Cubic Forms*, 2nd edition, North Holland Math. Library 4, North Holland 1986.
- [19] M. Yoshida: *A $W(E_6)$ -equivariant projective embedding of the moduli space of cubic surfaces*, 15 p. available at [arXiv:math. AG/0002102](https://arxiv.org/abs/math/0002102)
- [20] C.T.C. Wall: *Geometric invariant theory of linear systems*, Math. Proc. Cambridge Philos. Soc. 93 (1983), 57–62.
- [21] C.T.C. Wall: *Nets of quadrics, and theta-characteristics of singular curves*, Philos. Trans. Roy. Soc. London Ser. A 289 (1978), no. 1357, 229–269.

DIPARTIMENTO DI MATEMATICA, UNIVERSITÀ DI MILANO, VIA SALDINI 50, I-20133 MILANO, ITALIA

E-mail address: elisabetta.colombo@mat.unimi.it, geemen@mat.unimi.it

MATHEMATISCH INSTITUUT, UNIVERSITEIT UTRECHT, P.O. BOX 80.010, NL-3508 TA UTRECHT, NEDERLAND

E-mail address: looiijeng@math.uu.nl