

On ws -convergence of product measures

Erik J. Balder, Mathematical Institute, University of Utrecht, Netherlands

20 December, 1999

A canonical redecomposition makes it possible to study the ws -topology for measures on a product space by means of recent techniques, developed for the narrow topology for transition probabilities. Thus a central result is obtained that generalizes both Prohorov's theorem and Komlós' theorem; it is in terms of pointwise w -convergence of averages of transition measures (*Komlós-convergence*). New results for sequential ws -convergence follow; these include two versions of Prohorov's theorem for relative sequential ws -compactness and a complete characterization of sequential ws -convergence in terms of Komlós-convergence. Specializations yield the criterion for relative ws -compactness of Schäl (1975), the refined characterizations of ws -convergence of Galdéano and Truffert (1997, 1998) and a new version of Fatou's lemma in several dimensions. Separately, a non-sequential extension of Prohorov's theorem for relative ws -compactness is presented as well; it generalizes the corresponding relative ws -compactness criterion of Jacod and Mémin (1981).

1 Introduction

Let (Ω, \mathcal{A}) be an abstract measurable space and let S be a topological space which is completely regular and Suslin; we equip S with its Borel σ -algebra $\mathcal{B}(S)$. Recall from definitions III.67, III.79 in Dellacherie and Meyer (1975) that a Suslin space is the image of a Polish space under a continuous mapping. Let $\mathcal{M}(\Omega \times S)$ be the set of all finite nonnegative measures on $(\Omega \times S, \mathcal{A} \otimes \mathcal{B}(S))$. On this set the following *weak-strong* topology (*ws*-topology for short) was introduced by Schäl (1975) (as usual, $\mathcal{C}_b(S)$ stands here for the space of all bounded continuous functions on S).

Definition 1.1 The *ws-topology* on $\mathcal{M}(\Omega \times S)$ is the coarsest topology for which all functionals $\pi \mapsto \int_{A \times S} c(s)\pi(d(\omega, s)), A \in \mathcal{A}, c \in \mathcal{C}_b(S)$, are continuous.

This is one of several equivalent definitions discussed in Theorem 3.7 of Schäl (1975). The *ws*-topology is called the “measurable-continuous topology” by Jacod and Mémin (1981) and the “narrow topology” by Galdéano and Truffert (1997, 1998). If a sequence $(\pi_n) := (\pi_n)_{n \in \mathbb{N}}$ in $\mathcal{M}(\Omega \times S)$ converges in the *ws*-topology to a limit $\pi_\infty \in \mathcal{M}(\Omega \times S)$, then this will be indicated by $\pi_n \xrightarrow{ws} \pi_\infty$. A standard argument shows the *ws*-topology to be Hausdorff. It “straddles” two classical topologies:

Definition 1.2 (i) The *s-topology* on $\mathcal{M}(\Omega)$ is the coarsest topology for which all functionals $\lambda \mapsto \lambda(A), A \in \mathcal{A}$, are continuous.
(ii) The *w-topology* on $\mathcal{M}(S)$ is the coarsest topology for which all functionals $\nu \mapsto \int_S c(s)\nu(ds), c \in \mathcal{C}_b(S)$, are continuous.

Clearly, the *s*-topology is the finest topology on $\mathcal{M}(\Omega)$ for which $\pi \mapsto \pi^\Omega := \pi(\cdot \times S)$, the marginal projection from $\mathcal{M}(\Omega \times S)$ onto $\mathcal{M}(\Omega)$, is continuous. Similarly, the *w*-topology is the finest one on $\mathcal{M}(S)$ for which $\pi \mapsto \pi^S := \pi(\Omega \times \cdot)$, the marginal projection from $\mathcal{M}(\Omega \times S)$ onto $\mathcal{M}(S)$, is continuous. Conversely, it is not possible to describe the *ws*-topology solely in terms of these marginal topologies, because different measures in $\mathcal{M}(\Omega \times S)$ may have the same marginal projections. Important compactness results for the *s*-topology can be found in Gänssler (1971). The *w*-topology

is well-known under the name *weak* (or *narrow*) topology. It has been studied extensively in probability and measure theory; e.g., cf. Ash (1972), Billingsley (1968), Dellacherie and Meyer (1975), Schwartz (1975). Schäl (1975) gave some fundamental results for the *ws*-topology (for S separable and metric). These include Theorem 3.7 of Schäl (1975), which extends the classical portmanteau theorem. In Theorem 3.10 of Schäl (1975) he also gave a criterion for relative *ws*-compactness, but only in terms of *w*-compactness in $\mathcal{M}(\Omega \times S)$ (see Corollary 2.2). For this, he additionally supposed Ω to be topological. As also shown by him, the *ws*-topology leads naturally to a topology for policy-induced measures, the ws^∞ -topology, that is useful for existence in stochastic dynamic programming; see Nowak (1988) and Balder (1989b, 1992) for related subsequent work. Independently, Jacod and Mémin (1981) also studied the *ws*-topology. Their choice for a Polish space S opens up a richer variety of results (the present paper's more frugal choice for a completely regular Suslin space S does the same). While their portmanteau-type Proposition 2.4 is still covered by Theorem 3.7 of Schäl (1975), their Theorem 2.16 goes considerably further. In their Theorem 2.8 Jacod and Mémin (1981) also gave a relative *ws*-compactness result that goes further than the corresponding result of Schäl (1975) in that it addresses the situation where the measurable space (Ω, \mathcal{A}) is abstract (but with S Polish, as already mentioned before). The portmanteau-type results of Jacod and Mémin (1981), cited above, were recently refined by Galdéano (1997) in her doctoral thesis and by Galdéano and Truffert (1998), notably in connection with variational convergence. Like Jacod and Mémin (1981), they use abstract (Ω, \mathcal{A}) and Polish S .

The foundations for the *ws*-topology in Schäl (1975), which lie in statistical decision theory, have much in common with the foundations of what is now often called Young measure theory. The principal object of study there is the so-called narrow topology (alias Young measure topology) for transition probabilities, which we shall now recall. A *transition measure* (alias *kernel*) with respect to (Ω, \mathcal{A}) and $(S, \mathcal{B}(S))$ is a mapping $\tilde{\delta} : \Omega \mapsto \mathcal{M}(S)$ such that $\tilde{\delta}(\cdot)(B) : \omega \mapsto \tilde{\delta}(\omega)(B)$ is \mathcal{A} -measurable for every $B \in \mathcal{B}(S)$. We denote the set of all such transition measures by $\mathcal{T}(\Omega; S)$. A *transition probability* (alias *Markov kernel*) is a transition measure $\tilde{\delta} \in \mathcal{T}(\Omega; S)$ which takes only values in the set $\mathcal{P}(S)$ of all probability measures on S . Let $\mathcal{R}(\Omega; S)$ be the set of all such transition probabilities; thus, for $\tilde{\delta} \in \mathcal{T}(\Omega; S)$ we have $\tilde{\delta} \in \mathcal{R}(\Omega; S)$ if and only if $\tilde{\delta}(\cdot)(S) \equiv 1$. See section 2.6 in Ash (1972), Definition IX.1 of Dellacherie and Meyer (1975) and section III.2 of Neveu (1965) for technical backup. In Young measure theory the measurable space (Ω, \mathcal{A}) is endowed with a *fixed* measure $\mu \in \mathcal{M}(\Omega)$. Now there corresponds to every $\tilde{\delta} \in \mathcal{T}(\Omega; S)$ – and in particular to every $\tilde{\delta} \in \mathcal{R}(\Omega; S)$ – a canonical product measure (possibly infinite) on $(\Omega \times S, \mathcal{A} \otimes \mathcal{B}(S))$; it is given by

$$(\mu \otimes \tilde{\delta})(E) := \int_{\Omega} \tilde{\delta}(\omega)(E_{\omega}) \mu(d\omega), E \in \mathcal{A} \otimes \mathcal{B}(S).$$

E.g., see section 2.6 in Ash (1972) or section III.2 in Neveu (1965). Here E_{ω} denotes the section of E at ω . Observe that $\mu \otimes \tilde{\delta} \in \mathcal{M}(\Omega \times S)$ whenever $\tilde{\delta}(\cdot)(S)$ is μ -integrable. The following definition was given in Balder (1984b):

Definition 1.3 The *narrow topology* on $\mathcal{R}(\Omega; S)$ is the coarsest topology for which all functionals $\delta \mapsto \int_{A \times S} c(s)(\mu \otimes \delta)(d(\omega, s))$, $A \in \mathcal{A}$, $c \in \mathcal{C}_b(S)$, are continuous.

Note that this is one of several equivalent definitions; cf. Theorem 2.2 of Balder (1988). Definitions 1.1 and 1.3 show that the mapping $\delta \mapsto \mu \otimes \delta$ is a homeomorphism between $\mathcal{R}(\Omega; S)$, endowed with the narrow topology, and the subset $\{\pi \in \mathcal{M}(\Omega \times S) : \pi^{\Omega} = \mu\}$ of $\mathcal{M}(\Omega \times S)$, endowed with the relative *ws*-topology. Here a well-known disintegration of product measures is used – cf. section 2. Hence, the *ws*-topology generalizes the narrow topology for transition probabilities. As shown by the following example, the connections are less direct in the opposite direction:

Example 1.1 Let Ω be the unit interval $[0, 1]$, equipped with the Lebesgue σ -algebra \mathcal{A} and the Lebesgue measure λ . Let $r_1(\omega) := 1$ if $\omega \in [0, 1/2]$ and $r_1(\omega) := 0$ if $\omega \in (1/2, 1]$, and extend r_1 to \mathbb{R} by periodicity with period 1. Let $r_n(\omega) := r_1(2^{n-1}\omega)$. Consider the sequence (μ_n) in $\mathcal{M}(\Omega)$, given by $\mu_n(A) := \int_A r_n d\lambda$. Then it follows by standard arguments that $\mu_n(A) \rightarrow \mu_{\infty}(A) := \lambda(A)/2$ for every $A \in \mathcal{A}$. Consider also the sequence (δ_n) in $\mathcal{R}(\Omega; S)$, defined by $\delta_n(\omega) := \epsilon_{r_n(\omega)}$ (we use

$S := \{0, 1\}$). Here ϵ_a is the usual notation for the Dirac point measure at $a \in [0, 1]$. By the same sort of argument (see Example 2.6 in Balder (1988)) it follows that (δ_n) converges narrowly to the constant transition probability $\delta_\infty \in \mathcal{R}(\Omega; S)$, defined by $\delta_\infty(\omega) := (\epsilon_0 + \epsilon_1)/2$. This holds both when Ω is equipped with λ or with $\mu_\infty = \lambda/2$. Now $\mu_n \otimes \delta_n \xrightarrow{ws} \pi_\infty$, with $\pi_\infty := \mu_\infty \otimes \epsilon_1$. To see this, observe that for every $A \in \mathcal{A}$ and $c \in \mathcal{C}_b(S)$ one has $\int_{A \times S} cd(\mu_n \otimes \delta_n) = \int_A r_n(\omega)c(r_n(\omega))\lambda(d\omega) = \mu_n(A)c(1) \rightarrow \mu_\infty(A)c(1)$. Consequently, we do *not* have $\mu_n \otimes \delta_n \xrightarrow{ws} \mu_\infty \otimes \delta_\infty$.

While this example shows that the reverse direction is not without some intricacy, this paper will show that, nevertheless, the reverse route is still a viable one, which leads to many new results for the *ws*-topology. Our principal tool on this route is a canonical redecomposition of the product measures. Namely, relative compactness and related questions for the *ws*-topology, including all questions involving the *ws*-convergence of *sequences*, can essentially be resolved by a rather refined apparatus developed for the study of narrow convergence of transition probabilities, that is to say, by modern Young measure theory. Given the results already obtained within this theory (see Balder (1984b, 1988, 1995, 1998, 1999)), we shall describe the route in detail, but not all the details to which it leads, for this would be unnecessarily repetitive. Instead, we present some major results that have currently no counterpart whatsoever in the cited literature on the *ws*-topology. These include the following: (i) Theorem 2.2, a simultaneous generalization of Prohorov's Theorem 2.1 and Komlós Theorem 2.3, (ii) Theorem 2.6, a complete, useful characterization of sequential *ws*-convergence in terms of Komlós-convergence (i.e., in terms of *pointwise w*-convergence of averages), (iii) Theorem 3.2, an upper semicontinuity result for the pointwise support sets of a *ws*-convergent sequence, and (iv) Theorems 2.4 and 5.1; these form two further extensions of Prohorov's Theorem 2.1 and generalize the above-mentioned compactness criteria of Schäl (1975) and Jacod and Mémin (1981) (see Corollary 2.2 and Theorem 5.2). The usefulness of these results is illustrated by some applications, including a new version of Fatou's lemma in several dimensions (Theorem 4.1). Other applications are given in Balder and Yannelis (1999).

2 Three Prohorov-type theorems

Recall from Theorem 1 of Valadier (1973) that a measure $\pi \in \mathcal{M}(\Omega \times S)$ can be decomposed (or disintegrated) as follows: there exists a transition probability δ_π in $\mathcal{R}(\Omega; S)$ (see section 1) such that

$$\pi(E) = \int_\Omega \delta_\pi(\omega)(E_\omega)\pi^\Omega(d\omega), E \in \mathcal{A} \otimes \mathcal{B}(S). \quad (2.1)$$

In terms of section 1, (2.1) states that π can be decomposed into the product measure $\pi^\Omega \otimes \delta_\pi$. Observe also that the condition in Valadier (1973) that the marginal π^S of π be Radon follows by Theorem III.69 of Dellacherie and Meyer (1975), in view of the fact that S is Suslin. Now suppose that $\Pi \subset \mathcal{M}(\Omega \times S)$ is such that the collection Π^Ω of its Ω -marginals, defined by $\Pi^\Omega := \{\pi^\Omega : \pi \in \Pi\}$, is dominated by some $\mu \in \mathcal{M}(\Omega)$ (from now on, this will be called *marginal domination* of Π by μ). Correspondingly, for any $\pi \in \Pi$ we indicate by $\tilde{\phi}_\pi \in \mathcal{L}_\mathbb{R}^1(\Omega, \mu)$ an arbitrary but fixed version of the Radon-Nikodym density of π^Ω with respect to μ . Then (2.1) can be restated as follows (from now on we call this *redecomposition*):

$$\pi(E) = \int_\Omega \tilde{\phi}_\pi(\omega)(E_\omega)\mu(d\omega), E \in \mathcal{A} \otimes \mathcal{B}(S). \quad (2.2)$$

That is to say, every $\pi \in \Pi$ can also be decomposed as $\mu \otimes \tilde{\delta}_\pi$, where $\tilde{\delta}_\pi \in \mathcal{T}(\Omega; S)$ is now a transition *measure*; it is given by $\tilde{\delta}_\pi(\omega) := \tilde{\phi}_\pi(\omega)\delta_\pi(\omega)$. Observe that this implies $\tilde{\phi}_\pi = \tilde{\delta}_\pi(\cdot)(S)$. Particular examples of marginally dominated sets Π are:

- (i) Any sequence (π_n) in $\mathcal{M}(\Omega \times S)$.
- (ii) Any subset Π of $\mathcal{M}(\Omega \times S)$ for which Π^Ω is relatively *s*-compact.

Here the first case is evident (e.g., $\mu := \sum_n 2^{-n} \pi_n^\Omega / (1 + \pi_n(\Omega \times S))$ marginally dominates (π_n)), and the second case follows by Proposition 2.2 below. To some extent the fact that sequences are always marginally dominated, regardless of relative s -compactness of the marginals, explains the finer results that we shall obtain for sequences. The following definition is classical; see Billingsley (1968), Dellacherie and Meyer (1975) or Schwartz (1975):

Definition 2.1 A set $M \subset \mathcal{M}(S)$ is *tight* if for every $\epsilon > 0$ there is a compact $K_\epsilon \subset S$ such that $\sup_{\nu \in M} \nu(S \setminus K_\epsilon) < \epsilon$.

We recall Prohorov's famous theorem (Theorem 1.12, p. 170 of Prohorov (1956)). It asserts that tightness in the classical sense of Definition 2.1, together with boundedness, constitutes a sufficient condition for both relative sequential w -compactness and relative (topological) w -compactness in $\mathcal{M}(S)$:

Theorem 2.1 (Prohorov) *If $M \subset \mathcal{M}(S)$ is tight and bounded, then*

- (i) *M is relatively sequentially w -compact,*
- (ii) *M is relatively w -compact.*

Recall that M is said to be *bounded* if $\sup_{\nu \in M} \nu(S)$ is finite. Part (i) of this theorem can be found in Theorem 6.1 of Billingsley (1968) and part (ii) in Theorem III.59 of Dellacherie and Meyer (1975). A fine point in part (i) is that Billingsley (1968) requires S to be metrizable. However, our completely regular Suslin space S has a *weak metric*, i.e., a metric d whose topology is not finer than the original topology on S (e.g., see Theorem III.66 of Dellacherie and Meyer (1975)). Indeed, observe that by complete regularity the functions in $\mathcal{C}_b(S)$ separate the points of S . Hence, by the Suslin property and Lemma III.31 of Castaing and Valadier (1977), a countable subcollection (c_i) in $\mathcal{C}_b(S)$ already separates the points. So

$$d(s, z) := \sum_{i=1}^{\infty} 2^{-i} (1 + \|c_i\|_{\infty})^{-1} |c_i(s) - c_i(z)|$$

forms a weak metric on S (here $\|c_i\|_{\infty} := \sup_S |c_i|$). It follows that (S, d) is also Suslin, and on compact sets the two topologies are actually equivalent. Moreover, the corresponding Borel σ -algebras coincide by Corollary 2, p. 101, of Schwartz (1975). From these facts it is not hard to deduce that the above part (i) of the theorem still holds in our setting (cf. Theorems 2.4, 2.5 in Balder (1999)). We now extend tightness as in Definition 2.1 in two versions. The first of these comes from Young measure theory (see Berliocchi and Lasry (1973) and Balder (1979, 1984b)), where it is simply called *tightness*. We shall use it to extend Theorem 2.1(i), i.e., the sequential part of Prohorov's theorem, in two different forms (see Theorem 2.2 and Corollary 2.1 below). The second version of tightness, which we call *ws-tightness*, is more demanding. It serves for extensions to the *ws*-topology of both the sequential part (i) of Prohorov's Theorem 2.1 and the nonsequential part (ii). This is done in Theorems 2.4 and 5.1 respectively.

Definition 2.2 (i) A set $\Pi \subset \mathcal{M}(\Omega \times S)$ is *tight* if there exists a $\mathcal{A} \otimes \mathcal{B}(S)$ -measurable function $h : \Omega \times S \rightarrow [0, +\infty]$ such that the set $\{s \in S : h(\omega, s) \leq \beta\}$ is compact for every $\omega \in \Omega$ and $\beta \in \mathbb{R}_+$ and such that $\sup_{\pi \in \Pi} \int_{\Omega \times S} h \, d\pi < +\infty$.

(ii) A set $\Pi \subset \mathcal{M}(\Omega \times S)$ is *ws-tight* if Π is tight and Π^Ω is relatively s -compact.

Observe that *ws-tightness* of $\Pi \subset \mathcal{M}(\Omega \times S)$ implies that $\sup_{\pi \in \Pi} \pi(\Omega \times S) < +\infty$, i.e., Π is *bounded* (just note that $\lambda \mapsto \lambda(S)$ is s -continuous on the compact s -closure of Π^Ω). To compare the new definition of tightness with the classical one in Definition 2.1, we give an equivalent version of part (i) of Definition 2.2. We do so by means of the following proposition (cf. Jawhar (1984) and Exercise 10 on p. 109 of Bourbaki (1974), Chapter 5).

Proposition 2.1 *For every $\Pi \subset \mathcal{M}(\Omega \times S)$ the following are equivalent:*

- (a) *Π is tight in the sense of Definition 2.2(i).*
- (b) *For every $\epsilon > 0$ there exists a compact-valued multifunction $\Gamma_\epsilon : \Omega \rightarrow 2^S$, with $\mathcal{A} \otimes \mathcal{B}(S)$ -measurable graph $\text{gph } \Gamma_\epsilon$, such that $\sup_{\pi \in \Pi} \pi((\Omega \times S) \setminus \text{gph } \Gamma_\epsilon) < \epsilon$.*

Here $\text{gph } \Gamma_\epsilon := \{(\omega, s) \in \Omega \times S : s \in \Gamma_\epsilon(\omega)\}$.

PROOF. (a) \Rightarrow (b): Let h be as in Definition 2.2(i). Take $\Gamma_\epsilon(\omega) := \{s \in S : h(\omega, s) \leq \sigma/\epsilon\}$, with $\sigma := \sup_{\pi \in \Pi} \int h d\pi$. Then, clearly, Γ_ϵ has a measurable graph and compact values. To see that also the inequality holds, we simply observe that $\sigma \geq \int_{(\Omega \times S) \setminus \text{gph } \Gamma_\epsilon} \sigma/\epsilon d\pi$ holds for all $\pi \in \Pi$.

(b) \Rightarrow (a): Take $\epsilon := 3^{-n}$; rather than taking finite unions of multifunctions, we can suppose without loss of generality that the multifunctions $\Gamma_{1/3^n}$ are pointwise nondecreasing (in n). Now set $h(\omega, s) := 2^n$ if $s \in \Gamma_{1/3^{n+1}}(\omega) \setminus \Gamma_{1/3^n}(\omega)$ and $h(\omega, s) := 0$ if $s \in \Gamma_{1/3^n}(\omega)$. Then it is easy to see that h has the properties required in Definition 2.2(i). QED

It is clear from this proposition that classical tightness as in Definition 2.1 is generalized by tightness in the sense of Definition 2.2 (simply trivialize the space (Ω, \mathcal{A}) by taking Ω equal to a singleton or by setting $\mathcal{A} := \{\emptyset, \Omega\}$).

Proposition 2.2 *For every $\Pi \subset \mathcal{M}(\Omega \times S)$ the following are equivalent:*

- (a) Π^Ω is relatively s -compact.
- (b) Π^Ω is relatively sequentially s -compact.
- (c) Π^Ω is dominated by a measure $\mu \in \mathcal{M}(\Omega)$ and the corresponding collection $\{\tilde{\phi}_\pi : \pi \in \Pi\}$ of densities is uniformly μ -integrable.

PROOF. Each of (a), (b) and (c) implies boundedness of Π^Ω (i.e., $\sup_{\pi \in \Pi} \pi^\Omega(S) < +\infty$). So the equivalences hold by Theorem 2.6 of Gänssler (1971). Observe that 2.6(iii) of Gänssler (1971) states only uniform absolute continuity, but, in combination with $\sup_{\pi \in \Pi} \int_\Omega \tilde{\phi}_\pi d\mu = \sup_{\pi \in \Pi} \pi^\Omega(S) < +\infty$, this yields uniform μ -integrability as stated in (c) (apply Proposition II.5.2 in Neveu (1965)). QED

This shows that the relative s -compactness condition in Definition 2.2(ii) can be stated in several alternative ways. The next result applies in particular when $\Pi \subset \mathcal{M}(\Omega \times S)$ is ws -tight; its version for the narrow topology for transition probabilities is well-known. The proof does not make any use of the Suslin property of S (it only uses the separability and metrizability of S); thus, this proposition extends Remark 3.11 of Schäl (1975).

Proposition 2.3 *Suppose that \mathcal{A} is countably generated and S is metrizable (i.e., S is metrizable Suslin). Then every $\Pi \subset \mathcal{M}(\Omega \times S)$ such that Π^Ω is relatively s -compact is metrizable for the ws -topology.*

PROOF. By Proposition 2.2, there exists a dominating measure $\mu \in \mathcal{M}(\Omega)$ for Π^Ω . By hypothesis, there exists a countable (at most) algebra $\mathcal{A}_0 \subset \mathcal{A}$ which generates the σ -algebra \mathcal{A} . Let us write $\mathcal{A}_0 := \{A_j : j \in \mathbb{N}\}$. By Proposition 7.19 of Bertsekas and Shreve (1978) there exists a countable subset (c'_i) of $\mathcal{C}_b(S)$ such that for any net (ν_γ) in $\mathcal{M}(S)$ and any $\bar{\nu} \in \mathcal{M}(S)$ the following is true: $\lim_\gamma \int_S c'_i d\nu_\gamma = \int_S c'_i d\bar{\nu}$ for every $i \in \mathbb{N}$ implies $\bar{\nu} = w\text{-lim}_\gamma \nu_\gamma$. Now set

$$\rho(\pi, \pi') := \sum_{i,j} 2^{-i-j} (1 + \|c'_i\|_\infty)^{-1} \left| \int_{A_j \times S} c'_i d\pi - \int_{A_j \times S} c'_i d\pi' \right|.$$

First, observe that this defines a metric on $\mathcal{M}(\Omega \times S)$ which is not finer than the ws -topology. It remains to prove that $\bar{\pi} = ws\text{-lim}_\gamma \pi_\gamma$ for any net (π_γ) in Π and any $\bar{\pi} \in \Pi$ such that $\lim_\gamma \rho(\pi_\gamma, \bar{\pi}) = 0$. To this end, let $A \in \mathcal{A}$ and $c \in \mathcal{C}_b(S)$ be arbitrary. Define $\bar{\pi}^A := \bar{\pi}(A \times \cdot)$ and $\pi_\gamma^A := \pi_\gamma(A \times \cdot)$ in $\mathcal{M}(S)$. By the above property of (c'_i) , the hypothesis $\lim_\gamma \rho(\pi_\gamma, \bar{\pi}) = 0$ implies $\bar{\pi}^{A_j} = w\text{-lim}_\gamma \pi_\gamma^{A_j}$ for every j . In particular, this gives $\lim_\gamma \int_{A_j \times S} c d\pi_\gamma = \int_{A_j \times S} c d\bar{\pi}$ for every j . By Theorem 1.3.11 in Ash (1972), there exists for every $\epsilon > 0$ a set $A_j \in \mathcal{A}_0$ such that $\int_\Omega |1_A - 1_{A_j}| d\mu < \epsilon$. Using boundedness of Π , it follows that on Π the functional $\pi \mapsto \int_{A \times S} c d\pi$ is the uniform limit of a certain sequence of functionals $\pi \mapsto \int_{A_{j'} \times S} c d\pi$. Therefore, we conclude that $\lim_\gamma \int_{A \times S} c d\pi_\gamma = \int_{A \times S} c d\bar{\pi}$. QED

Remark 2.1 In their Proposition 2.10 Jacod and Mémin (1981) claim that $\mathcal{M}(\Omega \times S)$ as a whole is metrizable for the ws -topology if \mathcal{A} is countably generated, regardless of any s -compactness of

marginals. The present author does not know a counterexample to this claim, but wishes to point out that the proof of Proposition 2.10 on p. 535 of Jacod and Mémin (1981) is unconvincing. Namely, for $\Omega := [0, 1]$ and trivial S it already breaks down for the sequence $(\epsilon_{1/2^n})$ and ϵ_0 in $\mathcal{M}(\Omega)$. In that situation \mathcal{A}_0 , the algebra of finite disjoint unions of right-open and left-closed intervals with rational endpoints, generates $\mathcal{A} := \mathcal{B}([0, 1])$. But while $\epsilon_{1/2^n}(A) \rightarrow \epsilon_0(A)$ for every $A \in \mathcal{A}_0$, which is in complete accordance with the hypotheses on p. 535 of Jacod and Mémin (1981), we have $\epsilon_{1/2^n}(B) \not\rightarrow \epsilon_0(B)$ for $B := \{1/2^j : j \in \mathbb{N}\}$.

The remainder of this section is devoted to three different extensions of the sequential part (i) of Theorem 2.1 and to an associated characterization of sequential *ws*-convergence. Given this sequential orientation, it should not come as a surprise that it only makes use of the *sequential* compactness of the subsets of S used in Definition 2.2(i) (by using the weak metric mentioned above, it is clear that such sequential compactness is implied by compactness – note that the converse need not be true). In other words, for the sole purpose of extending part (i) of Prohorov's Theorem 2.1, one could phrase Definition 2.2(i) in terms of sequential compactness; this was done in Balder (1989c, 1990, 1995, 1998, 1999). In Balder (1989c, 1990) the following intermediate, nontopological mode of convergence was introduced and studied in a more abstract context. For sequences (π_n) in $\mathcal{M}(\Omega \times S)$ we shall use it to characterize *ws*-convergence completely in terms of the associated sequence $(\tilde{\delta}_{\pi_n})$ in $\mathcal{T}(\Omega; S)$.

Definition 2.3 Given $\mu \in \mathcal{M}(\Omega)$, a sequence $(\tilde{\delta}_n)$ of transition measures in $\mathcal{T}(\Omega; S)$ *K-converges under* μ to $\tilde{\delta}_\infty \in \mathcal{T}(\Omega; S)$ (notation: $\tilde{\delta}_n \xrightarrow{\mu, K} \tilde{\delta}_\infty$) if for every subsequence $(\tilde{\delta}_{n_j})$ of $(\tilde{\delta}_n)$ there is a μ -null set N in \mathcal{A} – possibly depending on that subsequence – such that

$$\frac{1}{m} \sum_{j=1}^m \tilde{\delta}_{n_j}(\omega) \xrightarrow{w} \tilde{\delta}_\infty(\omega) \text{ in } \mathcal{M}(S) \text{ for every } \omega \in \Omega \setminus N. \quad (2.3)$$

Example 2.1 (i) Independent and identically distributed sequences in $\mathcal{R}(\Omega; S)$ provide concrete and interesting examples of *K*-convergence. For instance, let $\Omega := [0, 1]$ be equipped with the Lebesgue measure μ and let (r_n) be the sequence of Rademacher functions $r_n(\omega) := \text{sgn}(\cos(2^n \pi \omega))$. For $S := \{1, -1\}$ we can define $\delta_n(\omega) := \epsilon_{r_n(\omega)}$. Then the random measures $\delta_n : [0, 1] \rightarrow \mathcal{P}(\{1, -1\})$ are independent and identically distributed. By Kolmogorov's strong law of large numbers, which can be applied to every subsequence of (δ_n) (observe that $\mathcal{P}(\{1, -1\})$ has dimension 1), we obtain $\delta_n \xrightarrow{\mu, K} \delta_\infty$ with $\delta_\infty \equiv (\epsilon_1 + \epsilon_{-1})/2$.

(ii) A less interesting illustration of *K*-convergence is as follows. Let $(\tilde{\delta}_n)$ and $\tilde{\delta}_\infty$ be given in $\mathcal{T}(\Omega; S)$ with $\tilde{\delta}_n(\omega) \xrightarrow{w} \tilde{\delta}_\infty(\omega)$ in $\mathcal{M}(S)$ for μ -a.e. ω in Ω . Concretely, for $\Omega := [0, 1]$, equipped with the Lebesgue σ -algebra \mathcal{A} and the Lebesgue measure μ , and for $S := \{0\}$ we could take the following sequence $(\tilde{\delta}_n)$. For $\omega \in [0, 1/n]$ let $\tilde{\delta}_n(\omega)(\{0\}) := n$ and for $\omega \in (1/n, 1]$ let $\tilde{\delta}_n(\omega)(\{0\}) := 0$. Also, let $\tilde{\delta}_\infty(\omega)(\{0\}) = 0$ for all $\omega \in \Omega$. This example also shows that, unlike *ws*-convergence in $\mathcal{M}(\Omega \times S)$ and *K*-convergence in $\mathcal{R}(\Omega; S)$, *K*-convergence in the space of transition measures $\mathcal{T}(\Omega; S)$ need not preserve aggregate measure in the limit. Notably, in the above situation $(\mu \otimes \tilde{\delta}_n)(\Omega \times S)$ equals 1 for all $n \in \mathbb{N}$, but it equals 0 for $n = \infty$.

In Example 2.1(i) Kolmogorov's theorem is actually applied uncountably many times (viz. once for each subsequence). Each such application yields an exceptional μ -null set N (i.e., the null set that figures in Kolmogorov's limit statement). While Definition 2.3 allows for this, it does not mean perforce that the total number of exceptional null sets N involved in Definition 2.3 is uncountably infinite as well. For instance, in Example 2.1(ii) one and the same null set can serve for all subsequences. The following fact, however, is elementary: for any (α_n) and α_∞ in \mathbb{R} :

$$\lim_m \frac{1}{m} \sum_{j=1}^m \alpha_{n_j} = \alpha_\infty \text{ for every subsequence } (\alpha_{n_j}) \text{ of } (\alpha_n) \text{ implies } \alpha_n \rightarrow \alpha_\infty. \quad (2.4)$$

This means that in Example 2.1(i) the uncountable number of applications of Kolmogorov's theorem is indeed matched by an uncountable number of exceptional null sets. This finding underlines the

importance of the null sets in Definition 2.3: their plurality distinguishes stronger from weaker modes of convergence in $\mathcal{T}(\Omega; S)$.

Next, we state a useful lower semicontinuity property of K -convergence. This combines a Fatou- and a Fatou-Vitali-type result. Recall here that a *normal integrand* on $\Omega \times S$ is a $\mathcal{A} \otimes \mathcal{B}(S)$ -measurable function $g : \Omega \times S \rightarrow (-\infty, +\infty]$ such that $g(\omega, \cdot)$ is lower semicontinuous on S for every $\omega \in \Omega$.

Proposition 2.4 *If $\tilde{\delta}_n \xrightarrow{\mu, K} \tilde{\delta}_\infty$ for $(\tilde{\delta}_n)$ and $\tilde{\delta}_\infty$ in $\mathcal{T}(\Omega; S)$ and $\mu \in \mathcal{M}(\Omega)$, then the following hold:*

- (i) $\liminf_n \int_{\Omega \times S} g d(\mu \otimes \tilde{\delta}_n) \geq \int_{\Omega \times S} g d(\mu \otimes \tilde{\delta}_\infty)$ for every nonnegative normal integrand g on $\Omega \times S$.
- (ii) $\liminf_n \int_{\Omega \times S} g d(\mu \otimes \tilde{\delta}_n) \geq \int_{\Omega \times S} g d(\mu \otimes \tilde{\delta}_\infty)$ for every normal integrand g on $\Omega \times S$ that is bounded below, provided that $(\tilde{\delta}_n(\cdot)(S))$ is μ -uniformly integrable.

PROOF. (i) Let $\beta := \liminf_n \int_{\Omega \times S} g d(\mu \otimes \tilde{\delta}_n)$. For elementary reasons, there is a subsequence $(\mu \otimes \tilde{\delta}_{n_j})$ of $(\mu \otimes \tilde{\delta}_n)$ such that $\beta = \lim_j \int_{\Omega \times S} g d(\mu \otimes \tilde{\delta}_{n_j})$. Set $\psi_n(\omega) := \int_S g(\omega, s) \tilde{\delta}_n(\omega)(ds)$ for $n \in \mathbb{N} \cup \{\infty\}$. Then $\beta = \lim_j \int_\Omega \psi_{n_j} d\mu$. Of course, this implies also $\beta = \lim_m \int_\Omega \frac{1}{m} \sum_{j=1}^m \psi_{n_j} d\mu$. Now (2.3) gives $\liminf_m \frac{1}{m} \sum_{j=1}^m \psi_{n_j}(\omega) \geq \psi_\infty(\omega)$ for μ -a.e. ω , because the function $g(\omega, \cdot)$ is lower semicontinuous and nonnegative on S (apply Theorem III.55 of Dellacherie and Meyer (1975)). Hence, an application of Fatou's lemma gives $\beta \geq \int_\Omega \psi_\infty d\mu$. Since $\int_\Omega \psi_\infty d\mu = \int_{\Omega \times S} g d(\mu \otimes \tilde{\delta}_*)$, the proof of (i) is finished.

(ii) Again we set $\beta := \liminf_n \int_{\Omega \times S} g d(\mu \otimes \tilde{\delta}_n)$. As before, there exists a subsequence $(\tilde{\delta}_{n'_j})$ of $(\tilde{\delta}_n)$ for which $\beta = \lim_{n'_j} \int_{\Omega \times S} g d(\mu \otimes \tilde{\delta}_{n'_j})$. By uniform integrability of $(\tilde{\delta}_{n'_j}(\cdot)(S))$ and the Dunford-Pettis theorem, there exist a further subsequence $(\tilde{\delta}_{n'_j})$ of $(\tilde{\delta}_{n'_j})$ and a function $\psi_* \in \mathcal{L}_\mathbb{R}^1(\Omega, \mu)$ such that

$$(\tilde{\delta}_{n'_j}(\cdot)(S)) \text{ converges to } \psi_* \text{ in the weak topology } \sigma(\mathcal{L}_\mathbb{R}^1(\Omega, \mu), \mathcal{L}_\mathbb{R}^\infty(\Omega)). \quad (2.5)$$

By (2.3) we have $\frac{1}{m} \sum_{j=1}^m \tilde{\delta}_{n'_j}(\omega) \xrightarrow{w} \tilde{\delta}_\infty(\omega)$ for μ -a.e. ω , so in particular $\frac{1}{m} \sum_{j=1}^m \tilde{\delta}_{n'_j}(\omega)(S) \rightarrow \tilde{\delta}_\infty(\omega)(S)$. Because of (2.5), the same averages $\frac{1}{m} \sum_{j=1}^m \tilde{\delta}_{n'_j}(\cdot)(S)$ also converge weakly to ψ_* in $\sigma(\mathcal{L}_\mathbb{R}^1, \mathcal{L}_\mathbb{R}^\infty)$. As is well-known, these two facts together imply $\tilde{\delta}_\infty(\omega)(S) = \psi_*(\omega)$ for μ -a.e. ω (use the Lebesgue-Vitali theorem). By hypothesis, there is a constant $\alpha \in \mathbb{R}$ such that $g \geq -\alpha$. So $g + \alpha$ is a nonnegative normal integrand on $\Omega \times S$. By part (i)

$$\beta + \alpha \liminf_j (\mu \otimes \tilde{\delta}_{n'_j})(\Omega \times S) \geq \int_{\Omega \times S} g d(\mu \otimes \tilde{\delta}_\infty) + \alpha (\mu \otimes \tilde{\delta}_\infty)(\Omega \times S).$$

Here $\liminf_j (\mu \otimes \tilde{\delta}_{n'_j})(\Omega \times S) = \lim_j \int_\Omega \tilde{\delta}_{n'_j}(\cdot)(S) d\mu = \int_\Omega \psi_* d\mu$, as follows by (2.5). So, in view of $\tilde{\delta}_\infty(\cdot)(S) = \psi_*$ μ -a.e., the inequality simplifies to $\beta \geq \int_{\Omega \times S} g d(\mu \otimes \tilde{\delta}_\infty)$. QED

For $(\tilde{\delta}_n)$ in $\mathcal{R}(\Omega; S) \subset \mathcal{T}(\Omega; S)$ uniform integrability as in part (ii) of the above proposition holds trivially by $\tilde{\delta}_n(\cdot)(S) \equiv 1$ for all n . Hence, the distinction between parts (i) and (ii) in the above proposition is not encountered in Young measure theory.

Our first and central extension of Theorem 2.1(i) can now be stated. It states that tightness is a sufficient condition for “relative compactness” for K -convergence in $\mathcal{M}(\Omega \times S)$ (as Komlós-convergence is nontopological, parentheses are called for). Recall from section 2 that a sequence (π_n) is *always* marginally dominated by some measure $\mu \in \mathcal{M}(\Omega)$, causing every π_n , $n \in \mathbb{N}$, to be redecosposable as $\mu \otimes \tilde{\delta}_{\pi_n}$, by virtue of (2.2).

Theorem 2.2 *If (π_n) in $\mathcal{M}(\Omega \times S)$ is tight, bounded and marginally dominated by $\mu \in \mathcal{M}(\Omega)$, then there exist a subsequence $(\tilde{\delta}_{\pi_{n'}})$ of $(\tilde{\delta}_{\pi_n})$ and a transition measure $\tilde{\delta}_* \in \mathcal{T}(\Omega; S)$ such that $\tilde{\delta}_*(\cdot)(S)$ is μ -integrable and $\tilde{\delta}_{\pi_{n'}} \xrightarrow{\mu, K} \tilde{\delta}_*$.*

If one trivializes (Ω, \mathcal{A}) , then it is easy to see that Theorem 2.2 reduces to part (i) of Prohorov's Theorem 2.1 (use (2.4)). On the other hand, if one trivializes S , then Theorem 2.2 reduces to Komlós' theorem, which is as follows (see Komlós (1967) or Chatterji (1973)):

Theorem 2.3 (Komlós) Let (ψ_n) be a sequence in $\mathcal{L}_{\mathbb{R}}^1(\Omega, \mu)$ such that $\sup_n \int_{\Omega} |\psi_n| d\mu < +\infty$. Then there exist a subsequence $(\psi_{n'})$ of (ψ_n) and a function $\psi_* \in \mathcal{L}_{\mathbb{R}}^1(\Omega, \mu)$ such that for every further subsequence $(\psi_{n'_j})$ of $(\psi_{n'})$ there is a μ -null set N – possibly depending on that subsequence – such that

$$\lim_{m \rightarrow \infty} \frac{1}{m} \sum_{j=1}^m \psi_{n'_j}(\omega) = \psi_*(\omega) \text{ for every } \omega \text{ in } \Omega \setminus N.$$

Beautiful connections exist between Theorem 2.3 and Kolmogorov's strong law of large numbers; e.g., see Aldous (1977) and Chatterji (1985). Very directly – e.g., see the exercise on p. 217 of Stout (1974) or see Valadier (1991) – Theorem 2.3 implies the SLLN, and hence extends it to sequences of non-independent random variables. Therefore, the SLLN is also extended by Theorem 2.2, which generalizes Theorem 2.3, as we saw above. Theorem 2.2 also extends the very similar Prohorov-Komlós theorem for transition probabilities in Theorem 5.1 of Balder (1990); however, that result does not reduce to Komlós' theorem if S is trivial. Our second extension of Theorem 2.1(i) is as follows:

Corollary 2.1 If (π_n) in $\mathcal{M}(\Omega \times S)$ is tight, bounded and marginally dominated by $\mu \in \mathcal{M}(\Omega)$, then there exist a subsequence $(\pi_{n'})$ of (π_n) , a measure $\pi_* \in \mathcal{M}(\Omega \times S)$, marginally dominated by μ , and a nonincreasing sequence (A_p) of sets in \mathcal{A} such that $\lim_p \mu(A_p) = 0$ and

$$\lim_{n'} \int_{A \times S} c(s) \pi_{n'}(d(\omega, s)) = \int_{A \times S} c(s) \pi_*(d(\omega, s))$$

for every $p \in \mathbb{N}$, $A \in \mathcal{A}$, $A \subset \Omega \setminus A_p$ and $c \in \mathcal{C}_b(S)$.

PROOF. By Theorem 2.2, there exists a subsequence $(\pi_{n'})$ of (π_n) and $\tilde{\delta}_* \in \mathcal{T}(\Omega; S)$ such that $\tilde{\delta}_*(\cdot)(S)$ is μ -integrable and $\tilde{\delta}_{\pi_{n'}} \xrightarrow{\mu, K} \tilde{\delta}_*$. The former means that $\pi_* := \mu \otimes \tilde{\delta}_*$ is well-defined in $\mathcal{M}(\Omega \times S)$ (see section 1). Also, we have $\sup_n \int_{\Omega} \tilde{\phi}_{\pi_n} d\mu = \sup_n \pi_n(\Omega \times S) < +\infty$. Hence, by the biting lemma (see Gaposhkin (1972) or Brooks and Chacon (1977), p. 17) there exists a sequence (A_p) that decreases to a null set such that $(\tilde{\phi}_{\pi_n})$ is uniformly μ -integrable over $\Omega \setminus A_p$ for every fixed $p \in \mathbb{N}$. For $A \subset \Omega \setminus A_p$, $p \in \mathbb{N}$, it remains to invoke Proposition 2.4(ii) twice: set $\Omega := A$ and set first $g(\omega, s) := c(s)$ and then $g(\omega, s) := -c(s)$. This gives $\lim_n \int_{A \times S} c d(\mu \otimes \tilde{\delta}_{\pi_n}) = \int_{A \times S} c d(\mu \otimes \tilde{\delta}_*)$. In view of (2.2) and the definition of π_* , this finishes the proof. QED

We shall now give a quick proof of Theorem 2.2 by means of the abstract generalization of Komlós' Theorem 2.3, given in Theorem 2.1 of Balder (1990) (see Balder and Hess (1996) for further developments in this direction). This proof requires only a slight extension of the demonstration of Theorem 5.1 of Balder (1990), as given in section 5 of that reference. A second, more elaborate proof of Theorem 2.2, starting directly from Theorem 2.3, is given in the appendix.

PROOF OF THEOREM 2.2. In order to apply Theorem 2.1 of Balder (1990) we slightly modify the substitutions made in section 5 of Balder (1990). We now take $E := \mathcal{M}(S)$, equipped with the w -topology, which takes the place of $\mathcal{P}(S)$ in Balder (1990). Consequently, the last line on p. 12 of that reference must be adapted as follows: $h(\omega, x) := \int_S h'(\omega, s)x(ds) + x(S)$, $x \in E$ (here h' plays the same role as h in Definition 2.2). This causes $h(\omega, \cdot)$ to be sequentially w -inf-compact on $\mathcal{M}(S)$ for every $\omega \in \Omega$ by Prohorov's Theorem 2.1. Also, the definition of a^g in p. 13 of that reference must be slightly adapted: we still define $a^g : \Omega \times \mathcal{M}(S) \rightarrow \mathbb{R}$ by $a^g(\omega, x) := \int_{\Omega \times S} g(\omega, s)x(ds)$, but this time we use the *bounded* Carathéodory functions, i.e., bounded $\mathcal{A} \otimes \mathcal{B}(S)$ -measurable $g : \Omega \times S \rightarrow \mathbb{R}$ such that $g(\omega, \cdot)$ is continuous on S for every $\omega \in \Omega$. Let $\|g\|_{\infty} := \sup_{\Omega \times S} |g|$; then the inequality $|a^g(\omega, x)| \leq \|g\|_{\infty} x(S) \leq \|g\|_{\infty} h(\omega, x)$ shows that condition (B) on p. 3 of Balder (1990) continues to hold. The result now follows from Theorem 2.1 of that same reference, as shown in its section 5. QED

Observe that Theorem 2.2 and Corollary 2.1 require tightness, but not ws -tightness. This allows for situations where all marginal projections π_n^{Ω} , $n \in \mathbb{N}$, are absolutely continuous with respect to some given $\mu \in \mathcal{M}(\Omega)$, but where π_*^{Ω} is not absolutely continuous with respect to μ :

Example 2.2 Let $\Omega := [0, 1]$ be equipped with the Lebesgue σ -algebra \mathcal{A} and the Lebesgue measure μ . Let $S := \{0\}$ and define $\pi_n \in \mathcal{M}(\Omega \times S)$ by $\pi_n(A \times S) := n\mu(A \cap [0, 1/n])$. Here all π_n^Ω , $n \in \mathbb{N}$, are absolutely continuous with respect to μ . Now (π_n) is tight (take $\Gamma_\epsilon \equiv S = \{0\}$ in Proposition 2.1), but not *ws*-tight (notice that $\pi_n^\Omega \xrightarrow{w} \epsilon_0$, but not $\pi_n^\Omega \xrightarrow{s} \epsilon_0$). Yet Corollary 2.1 applies, and from the preceding analysis one sees immediately that any nonincreasing sequence (A_p) will do for which $\bigcap_p A_p = \{0\}$. For $(\pi_{n'})$ one can simply take (π_n) itself and for π_* the null measure in $\mathcal{M}(\Omega \times S)$.

Our third generalization of Prohorov's Theorem 2.1(i) is a full-fledged generalization to *ws*-convergence. It requires the full force of *ws*-tightness (to see that it generalizes, one just trivializes (Ω, \mathcal{A}) again). This third generalization also includes the sequential versions of Prohorov's theorem for narrow convergence of transition probabilities in Balder (1989c, 1990, 1999). As we know from section 1, these have for Π^Ω a singleton $\{\mu\}$. A non-sequential companion result is Theorem 5.1, given below. It extends the remaining part (ii) of Prohorov's Theorem 2.1.

Theorem 2.4 *If $\Pi \subset \mathcal{M}(\Omega \times S)$ is *ws*-tight, then Π is relatively sequentially *ws*-compact.*

PROOF. Let (π_n) be any sequence in Π and let $\mu \in \mathcal{M}(\Omega)$ be as in Proposition 2.2. By Theorem 2.2, there exists a subsequence $(\pi_{n'})$ of (π_n) and $\tilde{\delta}_* \in \mathcal{T}(\Omega; S)$ such that $\tilde{\delta}_*(\cdot)(S)$ is μ -integrable and $\tilde{\delta}_{\pi_{n'}} \xrightarrow{\mu, K} \tilde{\delta}_*$. Proposition 2.2, $(\tilde{\phi}_{\pi_{n'}})$ is uniformly μ -integrable. One now proceeds as in the proof of Corollary 2.1 to prove $\pi_{n'} \rightarrow \pi_* := \mu \otimes \tilde{\delta}_*$ by means of Proposition 2.4. QED

Theorem 2.4 can be augmented to deal with situations where S is metrizable or Polish:

Theorem 2.5 *For $\Pi \subset \mathcal{M}(\Omega \times S)$ consider the following statements:*

- (a) $\Pi^\Omega \subset \mathcal{M}(\Omega)$ is relatively *s*-compact and $\Pi^S := \{\pi^S : \pi \in \Pi\} \subset \mathcal{M}(S)$ is tight.
 - (b) Π is *ws*-tight.
 - (c) Every sequence in Π is *ws*-tight.
 - (d) Π is relatively sequentially *ws*-compact.
 - (e) $\Pi^\Omega \subset \mathcal{M}(\Omega)$ is relatively *s*-compact and $\Pi^S \subset \mathcal{M}(S)$ is relatively sequentially *w*-compact.
- The following hold:*
- (i) In general (a) \Rightarrow (b) \Rightarrow (c) \Rightarrow (d) \Rightarrow (e).
 - (ii) If S is metrizable, then (c) \Leftrightarrow (d) \Leftrightarrow (e).
 - (iii) If S is Polish, then (a) \Leftrightarrow (b) \Leftrightarrow (c) \Leftrightarrow (d) \Leftrightarrow (e).

Observe that in parts (ii)-(iii) the *w*-topology on $\mathcal{M}(S)$ is metrizable (apply Theorem III.60 of Dellacherie and Meyer (1975)); hence, in (e) $\Pi^S \subset \mathcal{M}(S)$ is also relatively *w*-compact.

We show now that the criterion for relative *ws*-compactness in Theorem 3.10 of Schäl (1975) follows directly from part (ii) of the above theorem, in combination with the metrizability Proposition 2.3. This result of Schäl (1975) has a metrizable Lusin space Ω , with $\mathcal{A} = \mathcal{B}(\Omega)$. This allows him to consider the *w*-topology on $\mathcal{M}(\Omega \times S)$, but it is considerably more than we require here. On the other hand, Schäl (1975) uses a separable metric S , whereas we use a metrizable Suslin space S , so his result does not follow in its entirety from our result. Note also that Schäl's result contains a third equivalent property which we do not wish to consider here.

Corollary 2.2 *Suppose that Ω is a topological space, with $\mathcal{A} := \mathcal{B}(\Omega)$ countably generated, and suppose that S is metrizable. For every $\Pi \subset \mathcal{M}(\Omega \times S)$ the following are equivalent:*

- (a) Π is relatively *ws*-compact.
- (b) Π is relatively *w*-compact and Π^Ω is relatively *s*-compact.

PROOF. (a) \Rightarrow (b): Elementary; the *w*-topology on $\mathcal{M}(\Omega \times S)$ is coarser than the *ws*-topology and $\pi \mapsto \pi^\Omega$ is continuous.

(b) \Rightarrow (a): Continuity of $\pi \mapsto \pi^\Omega$ causes the marginal projection onto Ω of the *ws*-closure $\bar{\Pi}$ of Π to be contained in the *s*-closure of Π^Ω . Hence, that projection is relatively *s*-compact. It follows by Proposition 2.3 that $\bar{\Pi}$ is metrizable. Therefore, it is enough to prove relative *sequential ws*-compactness of Π . But (b) implies elementarily that Π^S is relatively *w*-compact; hence it is also relatively sequentially *w*-compact (recall that $\mathcal{M}(S)$ is metrizable – see the comments just after Theorem 2.5). So the desired relative sequential compactness of Π follows by Theorem 2.5(ii). QED

Lemma 2.1 If $\Pi \subset \mathcal{M}(\Omega \times S)$ is such that $\Pi^S \subset \mathcal{M}(S)$ is tight in the sense of Definition 2.1, then Π is tight in the sense of Definition 2.2(i).

PROOF. Since Π^S is tight, there exist compact sets K_ϵ , $\epsilon > 0$, as in Definition 2.1. Then part (b) of Proposition 2.1 applies, with Γ_ϵ the constant multifunction equal to K_ϵ . By Proposition 2.1, this shows that Π is tight in the sense of Definition 2.2(i). QED

Lemma 2.2 Suppose that S is metrizable.

- (i) Every w -convergent sequence in $\mathcal{M}(S)$ is tight in the classical sense of Definition 2.1.
- (ii) Every sequence (π_n) in $\mathcal{M}(\Omega \times S)$ such that (π_n^S) is w -convergent (and in particular every ws -convergent sequence (π_n)) is tight in the sense of Definition 2.2(i).

PROOF. (i) Let (ν_n) be w -convergent in $\mathcal{M}(S)$. Since S is Suslin, every single measure ν_n , $n \in \mathbb{N}$, is tight (alias Radon) by Theorem III.69 in Dellacherie and Meyer (1975). So, given the metrizability of S , it follows by Theorem 8 on p. 241 of Billingsley (1968) (see also LeCam (1957)) that the entire sequence of (ν_n) is tight.

(ii) For (π_n) in $\mathcal{M}(\Omega \times S)$ and $\nu_\infty \in \mathcal{M}(S)$, let $\pi_n^S \xrightarrow{w} \nu_\infty$. By part (i), (π_n^S) is tight in the classical sense. So by Lemma 2.1 (π_n) is tight in the sense of Definition 2.2(i). QED

PROOF OF THEOREM 2.5. (a) \Rightarrow (b): This follows directly from Lemma 2.1.

(b) \Rightarrow (c): *A fortiori*.

(c) \Rightarrow (d): Apply Theorem 2.4.

(d) \Rightarrow (e): Recall that the marginal projections $\pi \mapsto \pi^\Omega$ and $\pi \mapsto \pi^S$ are continuous.

(e) \Rightarrow (c) (if S is metrizable): Let (π_n) be an arbitrary sequence in $\mathcal{M}(\Omega \times S)$. Then (π_n^S) is relatively sequentially w -compact, so Lemma 2.2(ii) applies: (π_n) is tight. Since Π^Ω is relatively s -compact, (π_n) is also ws -tight.

(e) \Rightarrow (a) (if S is Polish): Since S is Polish, the converse Prohorov Theorem 6.2 in Billingsley (1968) applies. Hence, the relative (sequential) w -compactness of Π^S implies tightness of Π^S in the sense of Definition 2.1. QED

By following ideas of Balder (1995, 1998, 1999), we can completely characterize the ws -convergence of sequences in $\mathcal{M}(\Omega \times S)$. This is done by means of Theorem 2.2, provided that the Suslin space S is metrizable for its original topology. A similar characterization can also be given for non-metrizable S , but it would only hold for tight sequences; cf. Balder (1995, 1998, 1999). As applications in the next section will show, this characterization forms a powerful tool to study ws -convergence and ws -closure. It extends Corollary 3.16 of Balder (1995) and Theorem 4.8 of Balder (1999).

Theorem 2.6 Suppose that S is metrizable. For every (π_n) in $\mathcal{M}(\Omega \times S)$, marginally dominated by $\mu \in \mathcal{M}(\Omega)$, and every π_∞ in $\mathcal{M}(\Omega \times S)$ the following are equivalent:

- (a) $\pi_n \xrightarrow{ws} \pi_\infty$ in $\mathcal{M}(\Omega \times S)$,
- (b) $(\tilde{\phi}_{\pi_n})$ is uniformly μ -integrable, π_∞^Ω is absolutely continuous with respect to μ and every subsequence $(\pi_{n'})$ of (π_n) has a further subsequence $(\pi_{n''})$ such that $\tilde{\delta}_{\pi_{n''}} \xrightarrow{\mu, K} \tilde{\delta}_{\pi_\infty}$.

PROOF. (a) \Rightarrow (b): Uniform integrability of $(\tilde{\phi}_{\pi_n})$ holds by Proposition 2.2. Also, continuity of the marginal projection on S gives $\pi_n^S \xrightarrow{w} \pi_\infty^S$ in $\mathcal{M}(S)$. By Lemma 2.2(ii) it follows that (π_n) is tight. Since (π_n) is also evidently bounded, we may invoke Theorem 2.2. This gives that to every subsequence $(\pi_{n'})$ of (π_n) there correspond a further subsequence $(\pi_{n''})$ and a $\tilde{\delta}_* \in \mathcal{T}(\Omega; S)$ such that $\tilde{\delta}_{\pi_{n''}} \xrightarrow{\mu, K} \tilde{\delta}_*$. It remains to show that $\tilde{\delta}_* = \tilde{\delta}_{\pi_\infty}$ μ -a.e. (observe that π_∞^Ω is absolutely continuous with respect to μ by Definition 1.2). We already saw that $(\tilde{\phi}_{\pi_n})$ is uniformly μ -integrable, so it follows by Proposition 2.4 that $\pi_n \xrightarrow{ws} \mu \otimes \tilde{\delta}_*$ (see the proof of Corollary 2.1). Since the ws -topology is Hausdorff, (a) gives $\mu \otimes \tilde{\delta}_* = \pi_\infty = \mu \otimes \tilde{\delta}_{\pi_\infty}$, whence $\tilde{\delta}_*(\omega) = \tilde{\delta}_{\pi_\infty}(\omega)$ for μ -a.e. ω .

(b) \Rightarrow (a): Similar to the proof of Corollary 2.1, Proposition 2.4 implies that every subsequence $(\pi_{n'})$ of (π_n) has a further subsequence $(\pi_{n''})$ such that $\pi_{n''} \xrightarrow{ws} \pi_\infty$. By contraposition, this fact immediately implies (a). QED

3 Developments and applications

We begin this section by giving some applications of Theorem 2.6, the characterization result for ws -convergence of sequences in $\mathcal{M}(\Omega \times S)$. The following characterization of ws -convergence could be made part of a broader portmanteau-type theorem, quite similar to what was done in Balder (1995, 1998, 1999).

Theorem 3.1 Suppose that S is metrizable. For every (π_n) and π_∞ in $\mathcal{M}(\Omega \times S)$ the following are equivalent:

- (a) $\pi_n \xrightarrow{ws} \pi_\infty$ in $\mathcal{M}(\Omega \times S)$,
- (b) $\lim_n \int_{\Omega \times S} g \, d\pi_n = \int_{\Omega \times S} g \, d\pi_\infty$ for every bounded $\mathcal{A} \otimes \mathcal{B}(S)$ -measurable function $g : \Omega \times S \rightarrow \mathbb{R}$ such that $g(\omega, \cdot)$ is continuous on S for every $\omega \in \Omega$.
- (c) $\liminf_n \int_{\Omega \times S} g \, d\pi_n \geq \int_{\Omega \times S} g \, d\pi_\infty$ for every normal integrand g on $\Omega \times S$ such that

$$\lim_{\alpha \rightarrow \infty} \sup_n \int_{\{g \leq -\alpha\}} \max(-g, 0) \, d\pi_n = 0.$$

In (c) the following *integration convention* is made: we set $\int_{\Omega \times S} g \, d\pi_\infty := \int_{\Omega \times S} \max(g, 0) \, d\pi_\infty - \int_{\Omega \times S} \max(-g, 0) \, d\pi_\infty$, it being understood that $(+\infty) - (+\infty)$ is by definition equal to $+\infty$.

PROOF. (a) \Rightarrow (c): Fix any $\alpha \geq 0$ and let $\beta_\alpha := \liminf_n \int_{\Omega \times S} \max(g, -\alpha) \, d\pi_n$. There is a subsequence $(\pi_{n'})$ of (π_n) for which $\beta_\alpha = \lim_{n'} \int_{\Omega \times S} \max(g, -\alpha) \, d\pi_{n'}$. By Theorem 2.6, $(\pi_{n'})$ has a further subsequence $(\pi_{n''})$ with $\tilde{\delta}_{\pi_{n''}} \xrightarrow{\mu, K} \tilde{\delta}_{\pi_\infty}$ and $(\phi_{\pi_{n'}})$ is μ -uniformly integrable. Then Proposition 2.4(ii) implies $\beta_\alpha \geq \int_{\Omega \times S} \max(g, -\alpha) \, d\pi_\infty$. In turn, this gives $\beta_\alpha \geq \int_{\Omega \times S} g \, d\pi_\infty$. Letting α go to infinity gives the desired inequality, because

$$\int_{\Omega \times S} g \, d\pi_n \geq \int_{\Omega \times S} \max(g, -\alpha) \, d\pi_n - \int_{\{g \leq -\alpha\}} \max(-g, 0) \, d\pi_n.$$

for all $n \in \mathbb{N}$.

(c) \Rightarrow (b) \Rightarrow (a): Elementary: for the first implication, apply (c) to both g and $-g$, and for the second one apply (b) to $g(\omega, s) := 1_A(\omega)c(s)$. QED

Our next application of Theorem 2.6 is an upper semicontinuity result for the pointwise support sets of a ws -convergent sequence. Similar results were obtained for narrow convergence of transition probabilities in Balder (1995, 1998). Recall that the *support* of a measure ν in $\mathcal{M}(S)$ is defined as follows:

$$\text{supp } \nu := \cap_{F \subset S} \{F : F \text{ closed}, \nu(S \setminus F) = 0\}.$$

Recall also from Dal Maso (1993) that the Kuratowski upper limit set (alias limes superior) $\text{Ls}_n B_n$ of a sequence (B_n) of subsets of S is defined as the set of all $s \in S$ such that (s_{n_j}) converges to s for some subsequence (s_{n_j}) , $s_{n_j} \in S_{n_j}$. If S is metrizable, it is easy to see that the following identity holds:

$$\text{Ls}_n B_n = \cap_{p=1}^{\infty} \text{cl}(\cup_{n=p}^{\infty} B_n). \quad (3.1)$$

Theorem 3.2 Suppose that S is metrizable. If $\pi_n \xrightarrow{ws} \pi_\infty$ for (π_n) and π_∞ in $\mathcal{M}(\Omega \times S)$, then

$$\text{supp } \tilde{\delta}_{\pi_\infty}(\omega) \subset \text{Ls}_n \text{supp } \tilde{\delta}_{\pi_n}(\omega) \text{ for } \mu\text{-a.e. } \omega \text{ in } \Omega$$

for every marginally dominating measure $\mu \in \mathcal{M}(\Omega)$. Moreover,

$$\text{supp } \delta_{\pi_\infty}(\omega) \subset \text{Ls}_n \text{supp } \delta_{\pi_n}(\omega) \text{ for } \pi_\infty^\Omega\text{-a.e. } \omega \text{ in } \Omega,$$

whence

$$\pi_\infty(\{(\omega, s) \in \Omega \times S : s \notin \text{Ls}_n \text{supp } \delta_{\pi_n}(\omega)\}) = 0.$$

Lemma 3.1 Suppose that S is metrizable. If $\tilde{\delta}_n \xrightarrow{\mu,K} \tilde{\delta}_\infty$ for $(\tilde{\delta}_n)$ and $\tilde{\delta}_\infty$ in $\mathcal{T}(\Omega; S)$ and for $\mu \in \mathcal{M}(\Omega)$, then

$$\text{supp } \tilde{\delta}_\infty(\omega) \subset \text{Ls}_n \text{supp } \tilde{\delta}_n(\omega) \text{ for } \mu\text{-a.e. } \omega \text{ in } \Omega.$$

PROOF. Because of (2.3) there is a μ -null set N with $\nu_{m,\omega} := \frac{1}{m} \sum_{n=1}^m \tilde{\delta}_n(\omega) \xrightarrow{w} \tilde{\delta}_\infty(\omega)$ in $\mathcal{M}(S)$ for every $\omega \notin N$. Fix an arbitrary $\omega \notin N$. For every $p \in \mathbb{N}$ the portmanteau Theorem 2.1 of Billingsley (1968) gives $\tilde{\delta}_\infty(\omega)(G_p) \leq \liminf_m \nu_{m,\omega}(G_p) \leq \liminf_m \frac{1}{m} \sum_{n=1}^{p-1} \tilde{\delta}_n(\omega)(S) = 0$, where G_p denotes the open set $\Omega \setminus \text{cl} \cup_{n \geq p} \text{supp } \tilde{\delta}_n(\omega)$. It follows that $\tilde{\delta}_\infty(\omega)(\cup_p G_p) = 0$. By (3.1) this proves the result. QED

For nonmetrizable S this lemma continues to hold, but in a more complicated form. This can be gleaned from analogous results for narrow convergence in $\mathcal{R}(\Omega; S)$ given in Balder (1995, 1999).

PROOF OF THEOREM 3.2. By Theorem 2.6 there is a subsequence $(\pi_{n'})$ of (π_n) such that $\tilde{\delta}_{\pi_{n'}} \xrightarrow{\mu,K} \tilde{\delta}_{\pi_\infty}$. So the first result follows by Lemma 3.1, for the inclusion $\text{Ls}_{n'} \text{supp } \tilde{\delta}_{\pi_{n'}}(\omega) \subset \text{Ls}_n \text{supp } \tilde{\delta}_{\pi_n}(\omega)$ is evident.

The second result is an obvious consequence of the first one: For every $n \in \mathbb{N}$ and ω one has trivially $\text{supp } \tilde{\delta}_{\pi_n}(\omega) \subset \text{supp } \delta_{\pi_n}(\omega)$ by (2.2), with equality of these two sets whenever $\phi_{\pi_\infty}(\omega) > 0$. Observe here that (2.2) continues to hold for π_∞ because of Definition 1.2. The third result also follows from (2.2). QED

Remark 3.1 As follows from Lemma 3.1, $\tilde{\delta}_*$ in Theorem 2.2 has the following property:

$$\text{supp } \tilde{\delta}_*(\omega) \subset \text{Ls}_n \text{supp } \tilde{\delta}_{\pi_n}(\omega) \text{ for } \mu\text{-a.e. } \omega \text{ in } \Omega.$$

As a first application where Theorem 3.2 comes in handy, we generalize the main compactness result of Yushkevich (1997):

Proposition 3.1 Suppose that S is metrizable. Let $\Gamma : \Omega \rightarrow 2^S$ be a multifunction such that

$$\Gamma(\omega) \text{ is compact for every } \omega \in \Omega,$$

$$\text{gph } \Gamma := \{(\omega, s) \in \Omega \times S : s \in \Gamma(\omega)\} \text{ is } \mathcal{A} \otimes \mathcal{B}(S)\text{-measurable.}$$

Also, let $M \subset \mathcal{M}(\Omega)$ be an s -compact set. Then $\Pi_\Gamma := \{\pi \in \mathcal{M}(\Omega \times S) : \pi^\Omega \in M, \pi((\Omega \times S) \setminus \text{gph } \Gamma) = 0\}$ is sequentially ws -compact.

This extends Theorem 1 of Yushkevich (1997), where M is a singleton and (Ω, \mathcal{A}) is a measurable Lusin space (note that compactness is understood to be sequential compactness in that reference – see p. 459 of Yushkevich (1997)). Because Yushkevich (1997) works with a singleton M , his version of the above proposition could also be proven by means of standard Young measure theory (this fact was also observed in Yushkevich (1997)).

PROOF. Clearly, Π_Γ is ws -tight by Proposition 2.1. So by Theorem 2.4 Π_Γ is relatively sequentially ws -compact. Therefore, any sequence (π_n) in Π_Γ has a subsequence $(\pi_{n'})$ that ws -converges to some $\pi_* \in \mathcal{M}(\Omega \times S)$. Observe already that this implies $\pi_*^\Omega \in M$ by (ws,s) -continuity of $\pi \mapsto \pi^\Omega$. By (2.2) and by definition of Π_Γ we have $\text{supp } \tilde{\delta}_{\pi_{n'}}(\omega) \subset \Gamma(\omega)$ for all n' for μ -a.e. ω . Since $\Gamma(\omega)$ is certainly closed for every ω , it follows by Theorem 3.2 that $\text{supp } \tilde{\delta}_\infty(\omega)$ is also contained in $\Gamma(\omega)$ for μ -a.e. ω . Hence, $\pi_* \in \Pi_\Gamma$. QED

Next, following Balder (1984a, 1984b, 1995, 1998, 1999), we enrich S by considering $S \times \hat{\mathbb{N}}$. Here $\hat{\mathbb{N}} := \mathbb{N} \cup \{\infty\}$ is the Alexandrov compactification of \mathbb{N} (which is metrizable and compact), and $S \times \hat{\mathbb{N}}$ is equipped with the product topology. For $n \in \hat{\mathbb{N}}$ let $\epsilon_n \in \mathcal{P}(\hat{\mathbb{N}})$ be the Dirac probability measure concentrated at the point n . It turns out that such enrichment can be obtained entirely *for free*:

Lemma 3.2 Suppose that S is metrizable. For every (ν_n) and ν_∞ in $\mathcal{M}(S)$ the following are equivalent:

- (a) $\nu_n \xrightarrow{w} \nu_\infty$ in $\mathcal{M}(S)$,
- (b) $\nu_n \times \epsilon_n \xrightarrow{w} \nu_\infty \times \epsilon_\infty$ in $\mathcal{M}(S \times \hat{\mathbb{N}})$.

The nontrivial implication $(a) \Rightarrow (b)$ follows directly from Corollary 2.6 in Balder (1999). The following result, which generalizes Corollary 4.9 in Balder (1999), is now immediate by Theorem 2.6.

Theorem 3.3 *Suppose that S is metrizable. For every (π_n) and π_∞ in $\mathcal{M}(\Omega \times S)$ the following are equivalent:*

- (a) (π_n) converges in the ws -topology to $\pi_\infty \in \mathcal{M}(\Omega \times S)$,
- (b) $(\pi_n \times \epsilon_n)$ converges in the ws -topology to $\pi_\infty \times \epsilon_\infty \in \mathcal{M}(\Omega \times (S \times \hat{\mathbb{N}}))$.
- (c) $\liminf_n \int_{\Omega \times S} g(\omega, s, n) \pi_n(d\omega, s) \geq \int_{\Omega \times S} g(\omega, s, \infty) \pi_\infty(d\omega, s)$ for every normal integrand g on $\Omega \times (S \times \hat{\mathbb{N}})$ which is bounded from below.

The refined portmanteau-type theorems for ws -convergence, obtained by Galdéano (1997) and Galdéano and Truffert (1998), follow easily from Theorem 3.3 and the preceding results. This is quite similar to applications of Young measure theory to lower closure type results in Balder (1995, 1998, 1999). For instance, Theorem 2.1 of Galdéano and Truffert now follows by invoking Theorem 3.1 and the “free enrichment principle” explained above. As another example, we shall now essentially derive Theorem 1.2 of Galdéano and Truffert (who use a Polish space S):

Proposition 3.2 *Suppose that S is metrizable. For every (π_n) and π_∞ in $\mathcal{M}(\Omega \times S)$ the following are equivalent:*

- (a) $\pi_n \xrightarrow{w_s} \pi_\infty$ in $\mathcal{M}(\Omega \times S)$,
- (b) $\pi_n(\Omega \times S) \rightarrow \pi_\infty(\Omega \times S)$ and $\limsup_n \pi_n(\text{gph } \Gamma_n) \leq \pi_\infty(\text{gph } \Gamma_\infty)$ for every collection $\{\Gamma_n : n \in \mathbb{N} \cup \{\infty\}\}$ of multifunctions $\Gamma_n : \Omega \rightarrow 2^S$ such that

$$\text{gph } \Gamma_n \text{ is } \mathcal{A} \otimes \mathcal{B}(S)\text{-measurable for every } n \in \mathbb{N} \cup \{\infty\},$$

$$\Gamma_n(\omega) \text{ is closed for every } \omega \in \Omega \text{ and } n \in \mathbb{N} \cup \{\infty\},$$

$$\text{Ls}_n \Gamma_n(\omega) \subset \Gamma_\infty(\omega) \text{ for every } \omega \in \Omega.$$

PROOF. (a) \Rightarrow (b): The first statement in (b) is obvious. To prove the second one, we define $g : \Omega \times S \times \hat{\mathbb{N}} \rightarrow \{-1, 0\}$ by $g(\omega, s, n) := -1_{\text{gph } \Gamma_n}(\omega, s)$. Then it follows easily from the given properties of (Γ_n) that $g(\omega, \cdot, \cdot)$ is lower semicontinuous on $S \times \hat{\mathbb{N}}$ for every $\omega \in \Omega$. In view of (a), we can apply Theorem 3.3(c) to g , which easily yields the upper semicontinuity statement in (b).

(b) \Rightarrow (a): By Definition 1.1 it is clear that (a) holds if and only if $\pi_n^A \xrightarrow{w} \pi_\infty^A$ for an arbitrary $A \in \mathcal{A}$, where $\pi_n^A := \pi_n(A \times \cdot)$. Hence, by the portmanteau Theorem 2.1 of Billingsley (1968), here considered in $\mathcal{M}(S)$ instead of $\mathcal{P}(S)$, it is enough to prove that $\limsup_n \pi_n(A \times F) \leq \pi_\infty(A \times F)$ for every closed $F \subset S$. Define F_n as the set of all $s \in S$ with $\text{dist}(s, F) \leq 1/n$. Then the F_n are closed and $\text{Ls}_n F_n = F_\infty := F$. So we may apply (b) to $\Gamma_n(\omega) := F_n$ for $\omega \in A$ and $\Gamma_n(\omega) := \emptyset$ otherwise. This gives precisely $\limsup_n \pi_n(A \times F) \leq \pi_\infty(A \times F)$. QED

4 A new multidimensional Fatou lemma

A well-known area of applications of the Young measure apparatus is formed by lower closure results “without convexity”; see Balder (1984a,b,c, 1985, 1995, 1998, 1999). We illustrate the usefulness of the results developed thus far by giving a new type of Fatou’s lemma in several dimensions:

Theorem 4.1 *Given $\mu \in \mathcal{M}(\Omega)$ and $d \in \mathbb{N}$, let $(\tilde{\phi}_n)$ and $\tilde{\phi}_\infty$ be nonnegative functions in $\mathcal{L}_{\mathbb{R}}^1(\Omega, \mu)$ such that $(\tilde{\phi}_n)$ converges to $\tilde{\phi}_\infty \in \mathcal{L}_{\mathbb{R}}^1(\Omega, \mu)$ in the weak topology $\sigma(\mathcal{L}_{\mathbb{R}}^1(\Omega, \mu), \mathcal{L}_{\mathbb{R}}^\infty(\Omega))$. Let (f_n) be a sequence of \mathcal{A} -measurable functions from Ω into \mathbb{R}^d such that $\tilde{\phi}_n f_n^i$ is μ -integrable for every $n \in \mathbb{N}$ and such that*

$$a^i := \lim_n \int_{\Omega} \tilde{\phi}_n(\omega) f_n^i(\omega) \mu(d\omega) \text{ exists for } i = 1, \dots, d,$$

$$(\max(-f_n^i, 0) \tilde{\phi}_n) \text{ is uniformly } \mu\text{-integrable for } i = 1, \dots, d.$$

Then there exists a \mathcal{A} -measurable function f_ from Ω into \mathbb{R}^d such that $\int_{\Omega} \tilde{\phi}_\infty f_*^i d\mu \leq a^i$ for $i = 1, \dots, d$ and*

$$f_*(\omega) \in \text{Ls}_n \{f_n(\omega)\} \text{ for } \mu\text{-a.e. } \omega \text{ in } \Omega \text{ with } \tilde{\phi}_\infty(\omega) > 0. \quad (4.1)$$

PROOF. Take $S := \mathbb{R}^d$ and define $\pi_n \in \mathcal{M}(\Omega \times S)$ by

$$\pi_n(E) := \int_{\Omega} \tilde{\phi}_n(\omega) 1_{E_\omega}(f_n(\omega)) \mu(d\omega), \quad E \in \mathcal{A} \otimes \mathcal{B}(S),$$

Of course, $\sup_n \int_{\Omega} \max(-f_n^i, 0) \tilde{\phi}_n d\mu < +\infty$ holds for every i , by the uniform integrability hypothesis. Together with the existence of the limit a^i , this means that

$$\sup_n \int_{\Omega} \tilde{\phi}_n |f_n^i| d\mu = \sup_n \int_{\Omega \times S} |s^i| \pi_n(d(\omega, s)) < +\infty, \quad i = 1, \dots, d \quad (4.2)$$

Hence, for $h(\omega, s) := \sum_{i=1}^d |s^i|$ we obtain $\sup_n \int_{\Omega \times S} h d\pi_n < +\infty$. Also, it is obvious that the set $\{s \in \mathbb{R}_+^d : \sum_{i=1}^d |s^i| \leq \beta\}$ is compact for every $\beta \in \mathbb{R}_+$. Therefore, part (i) of Definition 2.2 is fulfilled. Part (ii) of that definition is also fulfilled, because

$$\pi_n^\Omega(A) = \int_A \tilde{\phi}_n d\mu \rightarrow \int_A \tilde{\phi}_\infty d\mu =: \lambda(A) \text{ for every } A \in \mathcal{A}.$$

Hence, (π_n) is ws -tight. By Theorem 2.4, there exist a subsequence $(\pi_{n'})$ of (π_n) and a measure π_* in $\mathcal{M}(\Omega \times S)$ such that $\pi_{n'} \xrightarrow{ws} \pi_*$. Then it follows by Theorem 3.1 that

$$a^i = \lim_{n'} \int_{\Omega \times S} s^i \pi_{n'}(d(\omega, s)) \geq \int_{\Omega \times S} s^i \pi_*(d(\omega, s)) \text{ for } i = 1, \dots, d. \quad (4.3)$$

The preceding gives $\lambda = \pi_*^\Omega$. Hence, by (2.1), π_* disintegrates as $\pi_* = \lambda \otimes \delta_*$ for some transition probability δ_* with respect to (Ω, \mathcal{A}) and $(\mathbb{R}^d, \mathcal{B}(\mathbb{R}^d))$. So the above can be rewritten as $a^i \geq \int_{\Omega} [\int_S s^i \delta_*(\omega)(ds)] \lambda(d\omega)$ for $i = 1, \dots, d$. Besides, Theorem 3.2 gives $\text{supp } \delta_*(\omega) \subset \text{Ls}_n\{f_n(\omega)\}$ for λ -a.e. ω .

The space Ω can now be partitioned into a nonatomic part Ω^{na} and a purely atomic part Ω^{pa} . First, we deal with Ω^{pa} which is, by its definition, the union of at most countably many μ -atoms A_j , with $\mu(A_j) > 0$. On each A_j the functions $\tilde{\phi}_n$ and f_n are a.e. constant, say with values $\beta_{n,j} \in \mathbb{R}$ and $s_{n,j} \in \mathbb{R}^d$. We now split Ω^{pa} further into \tilde{A} , the union of all A_j for which $\lambda(A_j) > 0$ and its complement $\Omega^{pa} \setminus \tilde{A}$. Then it is evident that $\Omega^{pa} \setminus \tilde{A}$ has λ -measure zero. On all A_j weak convergence of $(\tilde{\phi}_n)$ to $\tilde{\phi}_\infty$ comes down to $\lim_n \beta_{n,j} = \beta_{\infty,j}$. Also, (4.2) implies that $\sup_n \sum_j \beta_{n,j} |s_{n,j}^i| < +\infty$ for $i = 1, \dots, d$. Hence, it follows that $\sup_n |s_{n,j}^i| < +\infty$ for every j with $\beta_{\infty,j} > 0$ (that is, with $\lambda(A_j) > 0$). Hence, by a preliminary diagonal subsequence selection argument we can suppose without loss of generality that on \tilde{A} the sequence (f_n) converges pointwise λ -a.e. some limit function f_* . Since $\text{supp } \delta_*(\omega) \subset \text{Ls}_n\{f_n(\omega)\} = \{f_*(\omega)\}$ for λ -a.e. ω in \tilde{A} , we conclude that $\delta_*(\omega)$ is the point measure $\epsilon_{f_*(\omega)}$ for such ω . Clearly, this meets (4.1) on \tilde{A} .

Next, on Ω^{na} the measure μ is nonatomic, whence also λ , which is μ -absolutely continuous. Thus, an application of Lyapunov's theorem for Young measures (Theorem 5.3 in Balder (1999)) gives the existence of a measurable function f_* from Ω^{na} into S such that $f_*(\omega) \in \text{Ls}_n\{f_n(\omega)\}$ for λ -a.e. ω in Ω^{na} and $\int_{\Omega^{na}} f_*^i d\lambda = \int_{\Omega^{na}} [\int_S s^i \delta_*(\omega)(ds)] \lambda(d\omega) \leq a^i$ for all i .

Finally, substituting the effect of these decompositions into (4.3) gives

$$a^i \geq \int_{\Omega^{na}} \left[\int_S s^i \delta_*(\omega)(ds) \right] \lambda(d\omega) + \int_{\tilde{A}} f_*^i d\lambda = \int_{\Omega^{na} \cup \tilde{A}} f_*^i d\lambda \text{ for } i = 1, \dots, d.$$

By choosing $f_* \equiv 0$ on the λ -null set $\Omega^{pa} \setminus \tilde{A}$, it is easy to see that f_* is now as stated in the theorem. QED

Theorem 4.1 generalizes the multidimensional Fatou lemma of Balder (1984a), which subsumes both the original Fatou lemma of Schmeidler (1970) and the one of Artstein (1979). All those results work with $\phi_n \equiv 1$ for all n , and the above result does not seem to follow from any of them. The following example shows that the positivity condition $\tilde{\phi}_\infty(\omega) > 0$ in (4.1) is indispensable.

Example 4.1 Let $\Omega := [0, 1]$ be equipped with the Lebesgue σ -algebra \mathcal{A} and with the Lebesgue measure μ . Let $d := 1$, $\tilde{\phi}_n \equiv n^{-1}$, $\tilde{\phi}_\infty \equiv 0$ and $f_n \equiv n$. Then $\lim_n \int_\Omega \tilde{\phi}_n f_n d\mu = 1$, and $\text{Ls}_n f_n(\omega) = \emptyset$ for all ω . By $\tilde{\phi}_\infty \equiv 0$, this is still in agreement with (4.1).

We conclude with a new application of Theorem 4.1. Several applications of the multidimensional Fatou lemma of Balder (1984a), which is generalized by Theorem 4.1, were already given in Balder (1984c).

Example 4.2 A decision maker is uncertain about the state of nature in $\Omega := \mathbb{R}$ (equipped with the Lebesgue σ -algebra \mathcal{A} and the Lebesgue measure λ), which she believes to be distributed according to a normal distribution with variance 1 and unknown mean $\theta \in [-\theta_0, \theta_0]$. Her θ_0 is a given bound. Denote the corresponding normal densities by p_θ . A “most optimistic scenario” for the decision maker is defined to be an optimal solution of the minimization problem

$$(P) : \text{minimize } J^0(\theta, u) := \int_\Omega g^0(\omega, u(\omega)) p_\theta(\omega) \lambda(d\omega)$$

over all possible decision rules u and all θ , $|\theta| \leq \theta_0$, subject to certain constraints

$$J^i(\theta, u) := \int_A g^i(\omega, u(\omega)) p_\theta(\omega) \lambda(d\omega) \leq \alpha^i, \quad i = 1, \dots, m.$$

Here $\alpha^1, \dots, \alpha^m$ are given constants in \mathbb{R} . Also, a *decision rule* is defined to be a measurable function u from $\Omega := \mathbb{R}$ into $Z := \mathbb{R}^p$, such that $u(\omega) \in U(\omega)$ for all $\omega \in \Omega$, where $U : \Omega \rightarrow 2^Z$ is a compact nonempty-valued multifunction with $\mathcal{A} \times \mathcal{B}(Z)$ -measurable graph $\text{gph } U$. Further, the functions $g^i : \text{gph } U \rightarrow (-\infty, +\infty]$, $i = 0, \dots, m$, are $\mathcal{A} \otimes \mathcal{B}(Z)$ -measurable, and $g^i(\omega, \cdot)$ is lower semicontinuous on $U(\omega)$ for every $\omega \in \Omega$. Moreover, we suppose that $\gamma := \inf_{0 \leq i \leq m} \inf_{(\omega, z) \in \text{gph } U} g^i(\omega, z) > -\infty$. Hence, the above integrals exist.

We shall now prove the existence of a “most optimistic scenario” (θ_*, u_*) for problem (P) by means of Theorem 4.1, supposing that (P) has at least one feasible solution pair (u, θ) . Let $a^0 := \inf(P)$; then there exists a minimizing sequence (θ_n, u_n) for (P) . By compactness of $[-\theta_0, \theta_0]$ we may suppose, without loss of generality, that (θ_n) converges to some $\theta_* \in [-\theta_0, \theta_0]$. Also, by compactness of $[\gamma, \alpha^i]$ we may suppose without loss of generality that $(J^i(\theta_n, u_n))$ converges to some $a^i \in [\gamma, \alpha^i]$ for $i = 1, \dots, m$. By continuity of $\theta \mapsto p_\theta(\omega)$ for each $\omega \in \mathbb{R}$, it follows from Scheffé’s Theorem 16.11 in Billingsley (1986) that $\int_\Omega |p_{\theta_n} - p_{\theta_*}| d\lambda \rightarrow 0$. Hence, $\mu_n \xrightarrow{s} \mu_\infty$ for $\mu_n(A) := \int_A p_{\theta_n} d\lambda$ and $\mu_\infty(A) := \int_A p_{\theta_*} d\lambda$. Now define $f_n : \Omega \rightarrow \mathbb{R}^{m+1}$ by

$$f_n^i(\omega) := g^i(\omega, u_n(\omega)) p_{\theta_n}(\omega), \quad i = 0, \dots, m;$$

then it is evident that f_n^i is λ -integrable for every n and that $\lim_n \int_\Omega f_n^i d\lambda = a^i$. By Theorem 4.1 there exists a \mathcal{A} -measurable function f_* from Ω into \mathbb{R}^{m+1} such that

$$\int_\Omega f_*^i p_{\theta_*} d\lambda \leq a^i \text{ for } i = 0, \dots, m \tag{4.4}$$

and such that for λ -a.e. ω in Ω there exists a subsequence (f_{n_ω}) – possibly depending upon ω – with

$$f_{n_\omega}(\omega) \rightarrow f_*(\omega). \tag{4.5}$$

For all coordinates $i = 0, \dots, m$ we have here $f_{n_\omega}^i(\omega) := g^i(\omega, u_{n_\omega}(\omega))$, with $u_{n_\omega}(\omega)$ in the compact subset $U(\omega)$. By taking a convergent subsequence in (4.5) and by subsequently using the lower semicontinuity of $g^i(\omega, \cdot)$, it follows that for λ -a.e. ω there exists at least one point $z_\omega \in U(\omega)$ for which $f_*^i(\omega) = \lim_{n_\omega} f_{n_\omega}^i(\omega) \geq g^i(\omega, z_\omega)$, $i = 0, \dots, m$. By the implicit measurable selection Theorem III.38 in Castaing and Valadier (1977) it thus follows that there exists a measurable selection u_* of U with the same inequalities, i.e., $f_*^i(\omega) \geq g^i(\omega, u_*(\omega))$ for $i = 0, \dots, m$. If we substitute this in (4.4), we find

$$J^i(\theta_*, u_*) \leq a^i \leq \alpha^i, \quad i = 0, \dots, m.$$

So (θ_*, u_*) meets the constraints of (P) and $J^0(\theta_*, u_*) \leq a^0 := \inf(P)$. Hence, (θ_*, u_*) is an optimal solution of (P) .

5 A non-sequential Prohorov-type theorem

Here we extend the non-sequential (i.e., topological) part (ii) of Prohorov's Theorem 2.1 to the *ws*-topology. We show it to generalize the corresponding criterion for relative *ws*-compactness in Jacod and Mémin (1981). Our proof uses truncation of transition measures and reduces the situation to one where results from Young measure theory can be applied. We just mention that several other purely topological results from Young measure theory can be extended so as to yield counterparts for the *ws*-topology.

Theorem 5.1 *Suppose that $\Pi \subset \mathcal{M}(\Omega \times S)$ is *ws*-tight. Then Π is relatively *ws*-compact.*

Together with Theorem 2.4, this completely extends Prohorov's Theorem 2.1 to the *ws*-topology. Theorem 5.1 can be stated differently when S is a Polish space. We present the following counterpart to Theorem 2.5:

Theorem 5.2 *For $\Pi \subset \mathcal{M}(\Omega \times S)$ consider the following statements:*

- (a) $\Pi^\Omega \subset \mathcal{M}(\Omega)$ is relatively *s*-compact and $\Pi^S \subset \mathcal{M}(S)$ is tight.
- (b) Π is *ws*-tight.
- (c) Π is relatively *ws*-compact.
- (d) $\Pi^\Omega \subset \mathcal{M}(\Omega)$ is relatively *s*-compact and $\Pi^S \subset \mathcal{M}(S)$ is relatively *w*-compact.

The following hold:

- (i) In general (a) \Rightarrow (b) \Rightarrow (c) \Rightarrow (d).
- (ii) If S is Polish, then (a) \Leftrightarrow (b) \Leftrightarrow (c) \Leftrightarrow (d).

The proof is almost completely contained in that of Theorem 2.5 and will be omitted. Theorem 2.8 of Jacod and Mémin (1981), who use a Polish space S throughout, comes down to the equivalence (c) \Leftrightarrow (d) in the above theorem.

In the remainder of this section we prove Theorem 5.1 by means of an extension of Prohorov's theorem for the narrow topology for transition probabilities. This result was given in Theorem 2.3 of Balder (1988) for a metrizable Lusin space S . Subsequently, in Theorem 2.2 of Balder (1989a), it was extended to the situation where S is completely regular and Suslin, as used in this paper. For the reader's convenience we include its proof as given in Balder (1989a). Recall that the narrow topology was defined in Definition 1.3.

Theorem 5.3 (Theorem 2.2 of Balder (1989a)) *If for $\mu \in \mathcal{M}(\Omega)$ and $\Delta \subset \mathcal{R}(\Omega; S)$ the set $\{\mu \otimes \delta : \delta \in \Delta\}$ in $\mathcal{M}(\Omega \times S)$ is tight, then Δ is relatively narrowly compact.*

PROOF. Recall from section 1 that $\delta \mapsto \mu \otimes \delta$ is a homeomorphism from $\mathcal{R}(\Omega; S)$ into $\mathcal{M}(\Omega \times S)$. Therefore, it is enough to prove that Δ is relatively narrowly compact in $\mathcal{R}(\Omega; S)$.

Preliminary case: First, we suppose in addition that S is metrizable. To prove relative compactness of Δ for the narrow topology it is enough to demonstrate that Theorem 2.3 in Balder (1988) remains valid for a metrizable Suslin space S instead of the metrizable Lusin space used there. Observe first that everything said on pp. 266-270 of that same reference continues to hold for a metrizable Suslin space S , except for the line that immediately follows the definition of the function \hat{h} . Recall this definition from p. 270 of Balder (1988): $\hat{h} := h$ on $\Omega \times S$ and $\hat{h} := +\infty$ on $\Omega \times (\hat{S} \setminus S)$. Here h is as in Definition 2.2 and \hat{S} stands for the Hilbert cube compactification of S . To prove that \hat{h} is $\mathcal{A} \otimes \mathcal{B}(\hat{S})$ -measurable, the metrizable Lusin hypothesis of Balder (1988) is of immediate use, since it implies that S belongs to $\mathcal{B}(\hat{S})$ by Definition III.16 of Dellacherie and Meyer (1975). However, in case S is merely metrizable Suslin we can still prove that \hat{h} is $\mathcal{A}_\mu \otimes \mathcal{B}(\hat{S})$ -measurable and end up with a standard $\mathcal{A} \otimes \mathcal{B}(\hat{S})$ -measurable modification of \hat{h} . Here \mathcal{A}_μ stands for the μ -completion of the σ -algebra \mathcal{A} . This goes as follows. Let \hat{d} be a metric on the Hilbert cube and let $\beta \in \mathbb{R}$ be arbitrary. Observe that the set $\hat{h}^{-1}[0, \beta]$ in $\Omega \times S$ equals $C := h^{-1}[0, \beta]$. Define $u(\omega, \hat{s}) := \inf_{s \in C_\omega} \hat{d}(\hat{s}, s)$; if $C_\omega = \emptyset$ we set $u(\omega, \cdot)$ equal to $+\infty$. By the measurable projection Theorem III.23 in Castaing and Valadier (1977), $u(\cdot, \hat{s})$ is \mathcal{A}_μ -measurable for every $\hat{s} \in \hat{S}$. Also, $u(\omega, \cdot)$ is clearly continuous on \hat{S} .

for every $\omega \in \Omega$. By Lemma III.14 of Castaing and Valadier (1977) it follows that u is $\mathcal{A}_\mu \otimes \mathcal{B}(\hat{S})$ -measurable. Now by Definition 2.2 C_ω is compact in S , whence in \hat{S} ; of course, this also means that C_ω is closed in \hat{S} . Hence, C coincides with $u^{-1}(\{0\})$. We conclude therefore that that \tilde{h} is measurable with respect to $\mathcal{A}_\mu \otimes \mathcal{B}(\hat{S})$. At this point, the approximation argument involving the μ -completion of \mathcal{A} on p. 269 of Balder (1988) can be imitated (or, more directly, Lemma A.1 in Balder (1984b) can be applied). This gives a $\mathcal{A} \otimes \mathcal{B}(\hat{S})$ -measurable modification $\tilde{h} : \Omega \times \hat{S} \rightarrow [0, +\infty]$ of \tilde{h} , for which $\tilde{h}(\omega, \cdot) = h(\omega, \cdot)$ for μ -a.e. ω . After this, the proof on pp. 270-271 of Balder (1988) can be resumed to conclude that Δ is relatively compact for the narrow topology.

General case. Following Theorem 2.1 we already argued that S can be given a weak metric d , whose topology is not finer than the given topology on S . Moreover, we recorded there that the resulting metric space (S, d) is also Suslin and that its Borel σ -algebra coincides with the original σ -algebra $\mathcal{B}(S)$ on S . Now observe that h in Definition 2.2 is *a fortiori* such that for every $\omega \in \Omega$ the function $h(\omega, \cdot)$ is inf-compact for the d -topology on S . By the preliminary case above it follows that Δ is certainly relatively “new-narrowly” compact, where “new-narrowly” indicates that we have switched from the original topology to the d -topology on S . We now finish by demonstrating that, as a consequence of the given tightness, the new-narrow topology coincides on $\Sigma := \{\delta \in \mathcal{R}(\Omega; S) : I_h(\delta) \leq \sigma\}$ with the original narrow topology. Here $I_h(\delta) := \int_{\Omega \times S} h \, d(\mu \otimes \delta)$ and $\sigma := \sup_{\delta \in \Delta} I_h(\delta)$. Evidently, on all of $\mathcal{R}(\Omega; S)$ the new-narrow topology is certainly not finer than the original narrow topology. So it remains to prove the converse inclusion, relative to Σ . For this it is enough, by Theorem 2.2 in Balder (1988), to prove that $\delta \mapsto I_g(\delta)$ is new-narrowly continuous for any $\mathcal{A} \times \mathcal{B}(S)$ -measurable $g : \Omega \times S \rightarrow [0, +\infty]$ such that $g(\omega, \cdot)$ is lower semicontinuous for every $\omega \in \Omega$. Let $g_\epsilon := g + \epsilon h$, $\epsilon > 0$. Then every g_ϵ is $\mathcal{A} \times \mathcal{B}(S)$ -measurable and such that, for every $\omega \in \Omega$, the function $g_\epsilon(\omega, \cdot)$ is inf-compact; *a fortiori*, the latter makes $g_\epsilon(\omega, \cdot)$ also d -inf-compact, whence d -lower semicontinuous. So, again by Theorem 2.2 of Balder (1988), the functional I_{g_ϵ} is new-narrowly lower semicontinuous on all of $\mathcal{R}(\Omega; S)$. The identity $I_g(\delta) = \sup_{\epsilon > 0} (I_{g_\epsilon}(\delta) - \epsilon\sigma)$, which holds for every $\delta \in \Sigma$, then implies that I_g is new-narrowly lower semicontinuous on Σ . QED

PROOF OF THEOREM 5.1. By Proposition 2.2 there exists a dominating measure μ for Π^Ω such that the corresponding set of densities $\{\tilde{\phi}_\pi : \pi \in \Pi\}$ is uniformly integrable with respect to μ . For $p \in \mathbb{N}$ and $\tilde{\delta} \in \mathcal{T}(\Omega; S)$ we define $\tilde{\delta}^p \in \mathcal{T}(\Omega; S)$ by truncation:

$$\tilde{\delta}^p(\omega) := \begin{cases} \tilde{\delta}(\omega) & \text{if } \tilde{\delta}(\omega)(S) \leq p \\ \text{null measure on } S & \text{otherwise} \end{cases}$$

Fix p . Since $\tilde{\delta}_\pi^p \leq \tilde{\delta}_\pi$ for every $\pi \in \Pi$, tightness of Π as in Definition 2.2 implies

$$\sup_{\pi \in \Pi} \int_{\Omega} [\int_S h(\omega, s) \tilde{\delta}_\pi^p(\omega)(ds)] \mu(d\omega) \leq \sup_{\pi \in \Pi} \int_{\Omega \times S} h \, d\pi < +\infty,$$

in view of (2.2). This shows that for $\Delta^p := \{\frac{1}{p} \tilde{\delta}_\pi^p : \pi \in \Pi\}$ the tightness condition of Theorem 5.3 is met. The fact that Δ^p does not lie in $\mathcal{R}(\Omega; S)$, but in the set of all transition subprobabilities with respect to (Ω, \mathcal{A}) and $(S, \mathcal{B}(S))$ does not impede application of Theorem 5.3, since it is well-known that Young measure theory extends immediately to transition subprobabilities. Theorem 5.3 now implies that Δ^p is relatively compact for the narrow topology. Hence, $\Pi^p := \{\mu \otimes \tilde{\delta}_\pi^p : \pi \in \Pi\}$ is relatively *ws*-compact. We define $T^p : \Pi \rightarrow \Pi^p$ by $T^p(\pi) := \mu \otimes \tilde{\delta}_\pi^p$. Let \mathcal{U} be an arbitrary ultrafilter on Π . To prove relative *ws*-compactness of Π , it is enough to prove that \mathcal{U} *ws*-converges in $\mathcal{M}(\Omega \times S)$. By Proposition 4.12 of Choquet (1969) the collection $T^p(\mathcal{U})$ is an ultrafilter on Π^p . By relative *ws*-compactness of Π^p , demonstrated above, it follows that $T^p(\mathcal{U})$ *ws*-converges to some limit in the *ws*-closure of Π^p (apply Proposition 4.15 of Choquet (1969)). Clearly, this limit must be of the form $\mu \otimes \tilde{\eta}_p$, with $\tilde{\eta}_p \in \mathcal{T}(\Omega; S)$ such that $\tilde{\eta}_p(\omega)(S) \leq p$ for μ -a.e. ω (use Definition 1.1). Uniformly in p , the following bound obviously holds:

$$(\mu \otimes \tilde{\eta}_p)(\Omega \times S) \leq \sup_{\pi \in \Pi} \pi(\Omega \times S) < +\infty. \quad (5.1)$$

Further, by Definition 1.1 the inequality $\tilde{\delta}_\pi^{p+1}(\omega)(B) \geq \tilde{\delta}_\pi^p(\omega)(B)$ for all $\omega \in \Omega$ and all $B \in \mathcal{B}(S)$ leads to $(\mu \otimes \tilde{\eta}_{p+1})(A \times B) \geq (\mu \otimes \tilde{\eta}_p)(A \times B)$ for every $A \in \mathcal{A}$ and $B \in \mathcal{B}(S)$ (first, take B to be open for the weak metric d and use Theorem A6.6 of Ash (1972); thereafter, approximate as in Corollary 4.3.7 of that same reference). This implies $\mu \otimes \tilde{\eta}_{p+1} \geq \mu \otimes \tilde{\eta}_p$ on $\mathcal{A} \times \mathcal{B}(S)$. Because of this monotonicity, the limit $\pi_* := \lim_p \mu \otimes \tilde{\eta}_p$ forms a measure on $\mathcal{A} \otimes \mathcal{B}(S)$, which is bounded by (5.1); so π_* belongs to $\mathcal{M}(\Omega \times S)$. We claim that the ultrafilter \mathcal{U} *ws*-converges to π_* . To this end, fix any $A \in \mathcal{A}$ and $c \in \mathcal{C}_b(S)$. Then the above definition of truncation gives for every $p \in \mathbb{N}$ and $\pi \in \Pi$

$$\alpha_\pi^p := \left| \int_{A \times S} c \, d\pi - \int_A \left[\int_S c(s) \tilde{\delta}_\pi^p(\omega)(ds) \right] \mu(d\omega) \right| \leq \|c\|_\infty \int_{\{\omega \in \Omega : \tilde{\phi}_\pi(\omega) > p\}} \tilde{\phi}_\pi \, d\mu,$$

where we use (2.2) and the associated identity $\tilde{\delta}_\pi(\cdot)(S) = \tilde{\phi}_\pi$. By uniform μ -integrability of $\{\tilde{\phi}_\pi : \pi \in \Pi\}$, this implies $\lim_{p \rightarrow \infty} \sup_{\pi \in \Pi} \alpha_\pi^p = 0$. Now for any p

$$\left| \int_{A \times S} c \, d\pi - \int_{A \times S} c \, d\pi_* \right| \leq \alpha_\pi^p + \beta_\pi^p + \gamma^p,$$

with $\beta_\pi^p := |\int_{A \times S} c \, d(\mu \otimes \tilde{\delta}_\pi^p) - \int_{A \times S} c \, d(\mu \otimes \tilde{\eta}_p)|$ and $\gamma^p := |\int_{A \times S} c \, d(\mu \otimes \tilde{\eta}_p) - \int_{A \times S} c \, d\pi_*|$. For any fixed p the above *ws*-convergence of $T^p(\mathcal{U})$ to $\mu \otimes \tilde{\eta}_p$ implies that β_π^p converges to 0 along \mathcal{U} . Finally, $\lim_p \gamma^p = 0$ follows by an obvious application of the monotone convergence theorem for the positive and negative parts of the bounded function c , using (5.1). Together, this proves that $\int_{A \times S} c \, d\pi$ converges to $\int_{A \times S} c \, d\pi_*$ along \mathcal{U} . Since c and A were arbitrary, this proves that \mathcal{U} *ws*-converges to π_* in $\mathcal{M}(\Omega \times S)$. QED

A Second proof of Theorem 2.2

This appendix is devoted to a second proof of Theorem 2.2, which is similar to arguments given in Balder (1991, 1995, 1998, 1999) and is based on the direct use of Komlós' Theorem 2.3. We need the following lemma, which uses Prohorov's theorem:

Lemma A.1 *Let (ν_m) in $\mathcal{M}(S)$ be bounded and tight in the classical sense. Suppose that (c_i) is a countable subset of $\mathcal{C}_b(S)$ that separates the points of $\mathcal{M}(S)$. If $\lim_m \int_S c_i \, d\nu_m$ exists for every $i \in \mathbb{N}$, then there exists $\nu_* \in \mathcal{M}(S)$ such that $\nu_m \xrightarrow{w} \nu_*$ in $\mathcal{M}(S)$.*

PROOF. Denote $\alpha_i := \lim_m \int_S c_i \, d\nu_m$, $i \in \mathbb{N}$. By Prohorov's Theorem 2.1 there exist a subsequence of (ν_m) and $\nu_* \in \mathcal{M}(S)$ such that this subsequence *w*-converges to ν_* . This gives $\alpha_i = \int_S c_i \, d\nu_*$ for every $i \in \mathbb{N}$. If we did not have $\nu_m \xrightarrow{w} \nu_*$, there would exist a subsequence $(\nu_{m'})$ of (ν_m) and a $c \in \mathcal{C}_b(S)$ such that $\beta := \lim_{m'} \int_S c \, d\nu_{m'}$ exists and $\beta \neq \int_S c \, d\nu_*$. However, by another application of Theorem 2.1 there exist a further subsequence of $(\nu_{m'})$ and $\nu_{**} \in \mathcal{M}(\Omega \times S)$ such that this further subsequence *w*-converges to ν_{**} . Just as above, it follows that $\alpha_i = \int_S c_i \, d\nu_{**}$ for every $i \in \mathbb{N}$ and, moreover, $\beta = \int_S c \, d\nu_{**}$. But (c_i) separates the points of $\mathcal{M}(S)$, so it follows that $\nu_* = \nu_{**}$, which is impossible by $\beta > \int_S c \, d\nu_*$. This gives the desired contradiction. QED

SECOND PROOF OF THEOREM 2.2. By (2.2) $\tilde{\phi}_n := \tilde{\phi}_{\pi_n} = \tilde{\delta}_{\pi_n}(\cdot)(S)$, so boundedness of (π_n) amounts to

$$\sup_n \int_\Omega \tilde{\phi}_n \, d\mu < +\infty. \tag{A.1}$$

So by Theorem 2.3 there exist $\tilde{\phi}_* \in \mathcal{L}_\mathbb{R}^1(\Omega, \mu)$ and a preliminary subsequence (k) of (n) with the following property: For every subsequence $(\tilde{\phi}_{k_j})$ of $(\tilde{\phi}_k)$ there is a μ -null set $M \in \mathcal{A}$ such that

$$\lim_{m \rightarrow \infty} \frac{1}{m} \sum_{j=1}^m \tilde{\phi}_{k_j}(\omega) = \tilde{\phi}_*(\omega) \tag{A.2}$$

for every $\omega \in \Omega \setminus M$. By Fatou's lemma this yields

$$\int_{\Omega} \tilde{\phi}_* d\mu \leq \sup_n \int_{\Omega} \tilde{\phi}_n d\mu < +\infty. \quad (\text{A.3})$$

After this choice of the preliminary subsequence (k) , we can mimick the proof of the Prohorov-Komlós Theorem 3.8 for transition probabilities in Balder (1999). This goes as follows. By Lemma III.31 of Castaing and Valadier (1977) (already used before to obtain the weak metric on S) there is an at most countable subset (c_i) of $\mathcal{C}_b(S)$ that separates the points of $\mathcal{M}(S)$. In view of (A.1) and boundedness of each function c_i on S we have that for each $i \in \mathbb{N}$

$$\sup_n \int_{\Omega} |\phi_{n,i}| d\mu < +\infty, \quad (\text{A.4})$$

where $\phi_{n,i}(\omega) := \int_S c_i(s) \tilde{\delta}_{\pi_n}(\omega)(ds)$. Also, for h as in Definition 2.2(i), we define $(\phi_{n,0})$ by $\phi_{n,0}(\omega) := \int_S h(\omega, s) \tilde{\delta}_{\pi_n}(\omega)(ds)$. Then Definition 2.2(i) gives that (A.4) also holds for $i = 0$. By repeated application of Theorem 2.3 and a diagonal subsequence selection argument it follows by (A.4) that there exist a subsequence (n') of (k) and functions $\phi_{*,i} \in \mathcal{L}_{\mathbb{R}}^1(\Omega, \mu)$, $i \in \mathbb{N} \cup \{0\}$, such that for every further subsequence (n'_j) of (n') the following is true. There is a null set $N \in \mathcal{A}$ such that for every $\omega \in \Omega \setminus N$

$$\lim_{m \rightarrow \infty} \frac{1}{m} \sum_{j=1}^m \phi_{n'_j, i}(\omega) = \phi_{*,i}(\omega). \quad (\text{A.5})$$

First, we take (n'_j) to be (n') itself. We denote the corresponding exceptional μ -null-sets for (A.2) and (A.5) respectively by M' and N' . Let $\nu_{m,\omega} \in \mathcal{M}(S)$ be defined by $\nu_{m,\omega} := \frac{1}{m} \sum_{n'=1}^m \tilde{\delta}_{\pi_{n'}}(\omega)$. We prepare for applying Lemma A.1 to $(\nu_{m,\omega})$ for $\omega \notin N' \cup M'$. First, (A.5) implies that for every $\omega \notin N'$

$$\lim_{m \rightarrow \infty} \int_S c_i d\nu_{m,\omega} = \phi_{*,i}(\omega). \quad (\text{A.6})$$

Also, by (A.2) the preliminary selection guarantees that $\sup_m \frac{1}{m} \sum_{n'=1}^m \tilde{\delta}_{\pi_{n'}}(\omega)(S) < +\infty$ for every $\omega \notin M'$; thus, $(\nu_{m,\omega})$ is a bounded sequence of measures. Moreover, for every $\omega \notin N' \cup M'$ we also have $\sup_m \int_S h(\omega, s) \nu_{m,\omega}(ds) < +\infty$ by invoking (A.4) for $i = 0$. Thus, $(\nu_{m,\omega})$ is also tight in the classical sense of Definition 2.1. Together, this justifies the application of Lemma A.1 to the sequence $(\nu_{m,\omega})$ for any fixed $\omega \notin N' \cup M'$. This gives the existence of $\nu_{*,\omega} \in \mathcal{M}(S)$ such that $\nu_{m,\omega} \xrightarrow{w} \nu_{*,\omega}$ for every $\omega \notin N' \cup M'$. We define $\tilde{\delta}_*(\omega) := \nu_{*,\omega}$ for $\omega \notin N' \cup M'$; on $N' \cup M'$ we choose $\tilde{\delta}_*(\omega)$ identically equal to some fixed element of $\mathcal{M}(S)$. Now w -convergence as in (2.3) holds (for $\tilde{\delta}_*$ in the role of $\tilde{\delta}_\infty$) for every ω not in the null-set $N' \cup M'$. Also, the above definition shows that $\tilde{\delta}_*$ belongs to $\mathcal{T}(\Omega; S)$, since all $\omega \mapsto \nu_{m,\omega}$, $m \in \mathbb{N}$, are transition measures. Observe already that $\tilde{\delta}_*(\omega)(S) = \lim_m \nu_{m,\omega}(S) = \lim_m \frac{1}{m} \sum_{n'=1}^m \tilde{\phi}_{n'}(\omega)$ for $\omega \notin N' \cup M'$. Combined with (A.3) this shows $\int_{\Omega} \tilde{\delta}_*(\omega)(S) \mu(d\omega) < +\infty$. Observe that in the limit (A.6) gives

$$\int_S c_i(s) \tilde{\delta}_*(\omega)(ds) = \phi_{*,i}(\omega) \text{ for every } \omega \notin N' \cup M'. \quad (\text{A.7})$$

It remains to show that the *same* $\tilde{\delta}_*$ can also be used in (2.3) if we take any other subsequence (n'_j) (rather than (n') itself). We can denote the corresponding μ -null-sets for (A.2) and (A.5) simply by M and N respectively. Repetition of the above gives a transition measure, say $\tilde{\delta}_{**}$, for which w -convergence as in (2.3) holds, this time of course with $\tilde{\delta}_{**}$ in the role of $\tilde{\delta}_\infty$, for every $\omega \notin N \cup M$. Repeating (A.7), we get $\int_S c_i(s) \tilde{\delta}_{**}(\omega)(ds) = \phi_{*,i}(\omega)$ for every $\omega \notin N \cup M$. Since (c_i) separates the points of $\mathcal{M}(S)$, this shows that $\tilde{\delta}_{**}(\omega) = \tilde{\delta}_*(\omega)$ for all ω outside the null set $N \cup N' \cup M \cup M'$. So, indeed, $\tilde{\delta}_*$ can be used in (2.3) for any subsequence (n'_j) , and $\tilde{\delta}_{n'} \xrightarrow{\mu,K} \tilde{\delta}_*$ has been proven. QED

References

- Aldous, D.J. (1977). Limit theorems for subsequences of arbitrarily-dependent sequences of random variables, *Z. Wahrscheinlichkeitstheorie Verw. Geb.* **40** 59-82.
- Artstein, Z. (1979). A note on Fatou's lemma in several dimensions, *J. Math. Econom.* **6** 277-282.
- Ash, R.B. (1972). *Real Analysis and Probability*, Academic Press.
- Balder, E.J. (1979). On a useful compactification for optimal control problems, *J. Math. Anal. Appl.* **72** 391-398.
- (1984a). A unifying note on Fatou's lemma in several dimensions, *Math. Oper. Res.* **9** 267-275.
- (1984b). A general approach to lower semicontinuity and lower closure in optimal control theory, *SIAM J. Control Optim.* **22** 570-598.
- (1984c). Existence results without convexity conditions for general problems of optimal control with singular components, *J. Math. Anal. Appl.* **101** 527-539.
- (1985). An extension of Prohorov's theorem for transition probabilities with applications to infinite-dimensional lower closure problems, *Rend. Circ. Mat. Palermo* (II) **34** 427-447.
- (1988). Generalized equilibrium results for games with incomplete information, *Math. Oper. Res.* **13** 265-276.
- (1989a). On Prohorov's theorem for transition probabilities, *Sém. Anal. Convexe* **19** 9.1-9.11.
- (1989b). On compactness of the space of policies in stochastic dynamic programming, *Stoch. Processes Appl.* **32** 141-150.
- (1989c). Unusual applications of a.e. convergence. In: G.A. Edgar and L. Sucheston (Eds.), *Almost Everywhere Convergence*, Academic Press, New York, 31-53.
- (1990). New sequential compactness results for spaces of scalarly integrable functions, *J. Math. Anal. Appl.* **151** 1-16.
- (1991). On equivalence of strong and weak convergence in L_1 -spaces under extreme point conditions, *Israel J. Math.* **75** 21-47.
- (1992). Existence without explicit compactness in stochastic dynamic programming, *Math. Oper. Res.* **17** 572-580.
- (1995). Lectures on Young Measures, *Cahiers du Centre de Recherche de Mathématiques de la Décision (CEREMADE)* **9517**, Université Paris-Dauphine, Paris.
- and Hess, C. (1996). Two generalizations of Komlós' theorem with lower closure-type applications, *J. Convex Anal.* **3** 25-44.
- (1998). Lectures on Young measure theory and its applications in economics (Proceedings Grado School on Measure Theory and Real Analysis). *Rend. Sem. Matem. Trieste*, in press (<http://www.math.uu.nl/publications/preprints/1052.ps.gz>).
- (1999). New fundamentals of Young measure theory. In: A. Ioffe, S. Reich and I. Shafrir (Eds.), *Calculus of Variations and Optimal Control*. Chapman and Hall/CRC Research Notes in Mathematics **411**, CRC Press, Boca Raton, 24-48.
- and Yannelis, N.C. (1999). Upper semicontinuity of the Cournot-Nash equilibrium correspondence: a general approach, forthcoming.
- Berliocchi, H. and Lasry, J.M. (1973). Intégrandes normales et mesures paramétrées en calcul des variations, *Bull. Soc. Math. France* **101** 129-184.
- Bertsekas, D.P. and Shreve, S.E. (1978). *Stochastic Optimal Control: the Discrete Time Case*, Academic Press, New York.
- Billingsley, P. (1968). *Convergence of Probability Measures*, Wiley, New York.
- (1986). *Probability and Measure*, Wiley, New York.
- Bourbaki, N. (1974). *Topologie Générale*. Hermann, Paris.
- Brooks, J. and Chacon, R.V. (1980). Continuity and compactness of measures, *Adv. in Math.* **37** 16-26.

- Castaing, C. and Valadier, M. (1977). *Convex Analysis and Measurable Multifunctions*, Lecture Notes in Math. **580**, Springer-Verlag, Berlin.
- Chatterji, S.D. (1973). Les martingales et leurs applications analytiques. In: J.L. Bretagnolle et al. (Eds.), *Ecole d'Eté de Probabilités: Processus Stochastiques*, Lecture Notes in Math. **307**, Springer-Verlag, Berlin, pp. 27-164.
- (1985). A subsequence principle theory, *Jber. d. Dt. Math.-Verein.* **87** 91-107.
- Choquet, G. (1969). *Lectures on Analysis*, Benjamin, Reading, Mass.
- Dal Maso, G. (1993). *An Introduction to Γ -Convergence*, Birkhäuser, Boston.
- Dellacherie, C., and Meyer, P.-A. (1975). *Probabilités et Potentiel*, Hermann, Paris (English translation: North-Holland, Amsterdam).
- Galdéano, F. (1997). *Convergence Étroite de Mesures Définies sur un Espace Produit*, Thèse de Doctorat, Université de Perpignan, Académie de Montpellier.
- and Truffert, A. (1998). On the narrow convergence of measures defined on a product space. Preprint.
- Gänssler, P. (1971). Compactness and sequential compactness inspaces of measures, *Z. Wahrscheinlichkeitsth. Verw. Geb.* **17** 124-146.
- Gaposhkin, V.F. (1972). Convergence and limit theorems for sequences of random variables, *Theory Probab. Appl.* **17** No. 3 379-400.
- Jacod, J. and Mémin, J. (1981). Sur un type de convergence intermédiaire entre la convergence en loi et la convergence en probabilité. In: J. Azéma and M. Yor (Eds.), *Séminaire de Probabilités XV*, Lecture Notes in Math. **850**, Springer-Verlag, Berlin, 529-546.
- and — (1983). Rectification à: "Sur un type de convergence intermédiaire entre la convergence en loi et la convergence en probabilité". In: J. Azéma and M. Yor (Eds.), *Séminaire de Probabilités XVII*, Lecture Notes in Math. **986**, Springer-Verlag, Berlin, 509-511.
- Jawhar, A. (1984). Mesures de transition et applications, *Sém. Analyse Convexe Montpellier* **14** 13.1-13.62.
- Komlós, J. (1967). A generalisation of a problem of Steinhaus, *Acta Math. Acad. Sci. Hungar.* **18** 217-229.
- LeCam, L.M. (1957). Convergence in distribution of stochastic processes, *Univ. California Publ. Statist.* **2**, no. 11, 207-236.
- Neveu, J. (1965). *Mathematical Foundations of the Calculus of Probability*, Holden-Day, San Francisco.
- Nowak, A. (1988). On the weak topology on a space of probability measures induced by policies, *Bull. Polish Acad. Sci. Math.* **36** 181-186.
- Prohorov, Yu. V. (1956). Convergence of random processes and limit theorems in probability theory, *Theory Probab. Appl.* **1** 157-214.
- Schäl, M. (1975). On dynamic programming: compactness of the space of policies, *Stoch. Processes Appl.* **3** 345-364.
- Schmeidler, D. (1970). Fatou's lemma in several dimensions, *Proc. Amer. Math. Soc.* **24** 300-306.
- Schwartz, L. (1975). *Radon Measures*, Oxford University Press
- Stout, W.F. (1974). *Almost Sure Convergence*, Academic Press, New York.
- Valadier, M. (1973). Désintégration d'une mesure sur un produit, *C. R. Acad. Sc. Paris* **276** 33-35.
- (1991). La loi forte des grands nombres comme conséquence d'autres théorèmes, *Sém. Analyse Convexe Montpellier* **21** 15.1-15.6.
- Yushkevich, A. (1997). The compactness of a policy space in dynamic programming via an extension theorem for Carathéodory functions, *Math. Oper. Res.* **22** 458-467.