

Effect of supplementing coconut or krabok oil, rich in medium-chain fatty acids on ruminal fermentation, protozoa and archaeal population of bulls

P. Panyakaew^{1,2,3}, N. Boon⁴, G. Goel^{3,5}, C. Yuangklang², J. Th. Schonewille⁶,
W. H. Hendriks^{1,6} and V. Fievez^{3†}

¹Department of Animal Sciences, Wageningen University, Wageningen, The Netherlands; ²Department of Animal Science, Faculty of Natural Resources, Rajamangala University of Technology Isan, Sakon-Nakhon Campus, Sakon Nakhon 47160 Thailand; ³Department of Animal Production, Laboratory for Animal Nutrition and Animal Product Quality (Lanupro), Faculty of Bioscience Engineering, Ghent University, Proefhoevestraat 10, 9090 Melle, Belgium; ⁴Department of Biochemical and Microbial Technology, Laboratory of Microbial Ecology and Technology (LabMET), Faculty of Bioscience Engineering, Ghent University, Ghent, Belgium; ⁵Department of Biotechnology and Bioinformatics, Jaypee University of Information Technology, Solan, India; ⁶Department of Farm Animal Health, Animal Nutrition Division, Faculty of Veterinary Medicine, Utrecht University, Utrecht, The Netherlands

(Received 20 December 2012; Accepted 26 August 2013)

Medium-chain fatty acids (MCFA), for example, capric acid (C10:0), myristic (C14:0) and lauric (C12:0) acid, have been suggested to decrease rumen archaeal abundance and protozoal numbers. This study aimed to compare the effect of MCFA, either supplied through krabok (KO) or coconut (CO) oil, on rumen fermentation, protozoal counts and archaeal abundance, as well as their diversity and functional organization. KO contains similar amounts of C12:0 as CO (420 and 458 g/kg FA, respectively), but has a higher proportion of C14:0 (464 v. 205 g/kg FA, respectively). Treatments contained 35 g supplemental fat per kg DM: a control diet with tallow (T); a diet with supplemental CO; and a diet with supplemental KO. A 4th treatment consisted of a diet with similar amounts of MCFA (i.e. C10:0 + C12:0 + C14:0) from CO and KO. To ensure isolipidic diets, extra tallow was supplied in the latter treatment (KO + T). Eight fistulated bulls (two bulls per treatment), fed a total mixed ration predominantly based on cassava chips, rice straw, tomato pomace, rice bran and soybean meal (1.5% of BW), were used. Both KO and CO increased the rumen volatile fatty acids, in particular propionate and decreased acetate proportions. Protozoal numbers were reduced through the supplementation of an MCFA source (CO, KO and KO + T), with the strongest reduction by KO. Quantitative real-time polymerase chain reaction assays based on archaeal primers showed a decrease in abundance of Archaea when supplementing with KO and KO + T compared with T and CO. The denaturing gradient gel electrophoresis profiles of the rumen archaeal population did not result in a grouping of treatments. Richness indices were calculated from the number of DGGE bands, whereas community organization was assessed from the Pareto–Lorenz evenness curves on the basis of DGGE band intensities. KO supplementation (KO and KO + T treatments) increased richness and evenness within the archaeal community. Further research including methane measurements and productive animals should elucidate whether KO could be used as a dietary methane mitigation strategy.

Keywords: krabok oil, coconut oil, rumen protozoa, Archaea, microbial community organization

Implication

Krabok seeds are widely available in South-East Asian forests, contain large amounts of fat and as such are of interest to be used as a fat source in animal diets by local farmers. As krabok oil (KO) almost exclusively consists of lauric and myristic acid, it not only should be considered an energy supplier and its potential as modifier of the rumen microbial community was assessed here. KO showed potential to reduce rumen protozoal and archaeal numbers and shift

rumen fermentation towards more propionate at smaller quantities than coconut oil (CO), another more common source of lauric and myristic acid.

Introduction

Lauric (C12:0) and myristic (C14:0) acid show potential to reduce the number of rumen Archaea and protozoa (e.g. Dohme *et al.*, 1999; Machmüller *et al.*, 2003b; Soliva *et al.*, 2003; Klevenhusen *et al.*, 2011), which make these medium-chain fatty acids (MCFA) important candidates for dietary methane mitigation in ruminants. The reduction in archaeal

† E-mail: Veerle.Fievez@Ugent.be

numbers might be partially because of the defaunating effect of MCFA (Dohme *et al.*, 2001), as some methanogens are endosymbionts or ectosymbionts of protozoa (Finlay *et al.*, 1994), although independent action of MCFA sources on rumen *Archaea* has also been reported (Dohme *et al.*, 1999). From batch *in vitro* incubations, C12:0 has been identified as the most powerful of the two MCFAs (Soliva *et al.*, 2003 and 2004). In contrast, Rusitec and *in vivo* data (e.g. Dohme *et al.*, 2001; Machmüller *et al.*, 2003b; Klevenhusen *et al.*, 2011) indicated a similar effect of C12:0 and C14:0 on rumen *Archaea*, although protozoal numbers were suppressed less by C14:0 than by C12:0 (Dohme *et al.*, 2001). *In vivo* changes in archaeal numbers through the addition of coconut oil (CO), which is particularly rich in C12:0 (470 g/kg FA) and C14:0 (180 g/kg FA), are equivocal (Machmüller *et al.*, 2003a; Pilajun and Wanapat, 2011). Furthermore, not only the total number of *Archaea*, but also the composition of the methanogenic population might be altered by MCFA supplementation, for example, studies using order-specific probes revealed *Methanococcales* to be less sensitive to C14:0 (Machmüller *et al.*, 2003b), whereas C12:0 seemed to have a greater effect on the free-living *Methanomicrobiales* than on the *Methanobacteriales*, which are frequently associated with protozoa (Klevenhusen *et al.*, 2011). Similarly, CO changed the diversity, dominance and functional organization of *Archaea* during an *in vitro* batch study, as evidenced from DGGE profiles (Patra and Yu, 2013), although Hristov *et al.* (2009), using the same molecular approach, could not attribute specific changes in abundance of methanogens to CO supplementation to dairy cows.

Although former observations indicate a different mode of action of C12:0 and C14:0, most comparative studies were conducted *in vitro*, whereas *in vivo* studies mainly used either C12:0 (e.g. Klevenhusen *et al.*, 2011), C14:0 (e.g. Machmüller *et al.*, 2003b) or CO (e.g. Hristov *et al.*, 2009), with a fixed C12:0/C14:0 ratio. However, in practice, dietary fats with significant amounts of C14:0 in addition to C12:0 might be of major importance because of the lower palatability of C12:0 (Soliva *et al.*, 2004). Krabok oil (KO), an oil from krabok seeds (*Irvingia malayana* Oliv. ex A. w. Benn), is such a dietary oil source that contains similar amounts of C12:0, but is particularly richer in C14:0 (464 g/kg FA) as compared with CO. Hence, the

comparison of two natural oil sources with similar amounts of C12:0 but varying in C14:0 content is considered of practical relevance.

The objective of this *in vivo* study was to compare the effect of CO and KO on rumen fermentation pattern, number of protozoa and *Archaea*, as well as their diversity and functional organization. Unfortunately, the experimental infrastructure did not allow for concomitant monitoring of methane emissions.

Material and methods

Animals

Eight rumen-fistulated bulls were used in a 4 × 4 Latin square design (Table 1). The experimental diets were fed as a total mixed ration (TMR) and consisted of (g/kg TMR): cassava chips, 421; rice straw, 211; dry tomato pomace, 158; molasses, 73.7; rice bran, 52.6; soya bean meal, 31.6; urea, 21; salt, 10.5; di-calcium phosphate, 7.4; oyster meal, 5.3; mineral premix, 5.3; and sulfur, 3.2. Water was provided *ad libitum*. The animals were adjusted to the TMR for 14 days followed by 4 experimental periods of 21 days (Table 1), which were separated from each other by a washout period of 21 days during which the basal diet was fed without external fat. At the beginning of each experimental period, the bulls were weighed to adjust the feed supply to 1.5% of BW, which was kept constant during the experimental period and supplied in two daily meals. Rumen sampling took place between days 15 and 21 of the experimental period.

Treatments

The four isolipidic diets in this study were: (1) TMR with 35 g/kg DM of beef tallow (T); (2) TMR with 35 g/kg DM of coconut oil (CO); (3) TMR with 35 g/kg DM of krabok oil (KO); and (4) TMR with 29 g/kg DM of krabok oil + 6 g/kg DM tallow (KO + T). Treatment KO + T was designed in an attempt to provide similar amounts of MCFA as by CO (Table 1). Besides C12:0 and C14:0, C10:0 was also included in this calculation as the latter has been shown to effectively reduce methane production and/or the number of methanogens and protozoa (Dohme *et al.*, 2000 and 2001; Goel *et al.*, 2009). Dietary inclusion of oil supplements in the KO + T treatment

Table 1 Overview of the sequence of experimental treatments in a 4 × 4 Latin square design experiment with four treatments (supplementation (35 g/kg DM) with either beef T, CO, KO or KO + T) and eight bulls (B) in each of the four periods (P), the amounts of oil added to each treatment and the supply of medium-chain fatty acids (C10:0, C12:0 and C14:0) from CO or KO (average ± s.d. of four batches of CO and KO)

Period	T	CO	KO	KO + T
P1	B1 and B2	B3 and B4	B5 and B6	B7 and B8
P2	B5 and B6	B1 and B2	B7 and B8	B3 and B4
P3	B3 and B4	B7 and B8	B1 and B2	B5 and B6
P4	B7 and B8	B5 and B6	B3 and B4	B1 and B2
Tallow (g/kg DM)	35	0	0	6
Coconut oil (g/kg DM)	0	35	0	0
Krabok oil (g/kg DM)	0	0	35	29
Total C10:0 + C12:0 + C14:0 (g/kg DM)	0	23.9 ± 0.76	31.5 ± 0.67	26.1 ± 0.56

T = tallow; CO = coconut oil; KO = krabok oil.

Table 2 Fatty-acid composition (g/100 g fatty acids) of the CO, KO, T and TMR

Fatty acid	CO	KO	T	TMR
C8:0	0.21	0.00	0.01	0.00
C10:0	2.10	1.58	0.05	0.28
C12:0	45.8	42.0	0.12	0.12
C14:0	20.5	46.4	3.97	0.38
C16:0	11.1	4.49	26.8	15.7
C18:0	3.22	0.41	25.7	6.42
C18:1 c9	13.1	2.57	23.5	25.9
C18:1 c11	0.42	0.43	1.16	0.96
C18:2 n-6	2.63	0.32	0.53	42.2
C18:3 n-3	0.02	0.04	0.39	0.00
C18:2 c9t11	–	–	0.37	0.23

CO = coconut oil; KO = krabok oil; T = tallow; TMR = total mixed ration.

was based on fatty-acid compositions of the oils used in a concomitant *in vitro* experiment (Panyakaew *et al.*, 2013). Although oils were from a similar source for the current *in vivo* experiment, the composition of the four oil batches used in the current study deviated slightly (Table 2). Nevertheless, differences were small, allowing the original objectives to be assessed, as MCFA supplies of the treatments CO and KO + T did not differ significantly ($P > 0.05$) (Table 1).

Rumen sample collection

Rumen contents were collected on days 15, 18 and 21 of each period at 4 and 8 h after the morning feeding (0700 h) by sampling in different directions of the rumen. Samples were divided into three portions: the first portion of rumen contents (10 ml) that was stored at -80°C until DNA extraction; the second portion rumen aliquot (10 ml) was acidified with 0.2 ml of phosphoric: formic acid (10:1) and stored at -20°C until volatile fatty-acid (VFA) analysis; and the third portion of rumen contents (1 ml) was fixed with 10% formalin saline solution (37% to 40% formalin in 0.9% (w/v) normal saline solution, 1:9) and stored at 4°C for microscopical protozoal countings.

VFAs

Samples for VFA analysis were pooled per day and cow, and centrifuged at $15\,000 \times g$ (4°C) for 10 min (HKFI Ltd., Mainland, China). The supernatant was filtered through a nylon membrane (0.2 μm) and transferred to vials. The VFA analysis was performed using gas chromatography (Schimadzu GC-14A, 's-Hertogenbosch, Belgium) (Van Ranst *et al.*, 2010).

Protozoal counts

Microscopical counting was done using a haemocytometer (Boeco, Hamburg, Germany) of 0.1 mm depth. Twenty aliquots per sample were counted (Galyean, 1989).

Archaea analysis

DNA extraction

The rumen samples were pooled per day and bull before DNA extraction. Total DNA was extracted from 0.4 g rumen contents using the QIAamp Stool Kit DNA extraction method following

the protocol for 'Isolation of DNA from stool for Pathogen Detection' in the handbook supplied by Qiagen Ltd. (Crawley, UK), according to the manufacturer's recommendation.

Polymerase chain reaction (PCR) for denaturing gradient gel electrophoresis (DGGE) analysis

Amplification of a fragment of the 16S rDNA gene of *Archaea* for DGGE was performed using a single PCR with a forward GC-ARCH915 and reverse UNI-b-rev primer as described by Yu *et al.* (2008). The PCR mixture was based on the one described by Yu and Morrison (2004), with adjustments according to the manufacturer's instructions (Fermentas, St. Leon-Rot, Germany). PCR amplifications were performed in 25- μl volumes containing KCl (50 mM), MgCl_2 (1.5 mM), dNTP mix (200 μM), both primers (200 nM each), 0.6 U *Taq* DNA polymerase, 480 ng bovine serum albumin/ μl , 1 μl DNA extract and 2.5 μl 10X *Taq* buffer. Adjustments to 25 μl were made with DNase-, RNase-free filter-sterilized water (Sigma, Bornem, Belgium).

Amplification conditions were according to Yu and Morrison (2004) with small adjustments: initial denaturation at 94°C for 5 min; 10 cycles of touch-down PCR (denaturation at 94°C for 0.5 min, annealing for 0.5 min with an $0.5^{\circ}\text{C}/\text{cycle}$ decrement at a temperature of 5°C above the annealing temperature of 70°C and DNA synthesis at 72°C for 1 min); 25 cycles of regular PCR (denaturation at 94°C for 0.5 min, annealing at 65°C for 0.5 min, DNA-synthesis at 72°C for 1 min); and a final extension step at 72°C for 10 min.

DGGE analysis

Archaea-specific PCR fragments were loaded onto a 7.5% (w/v) polyacrylamide gel with denaturing gradients ranging from 45% to 60%. The electrophoresis was run for 16 h at 60°C and 50 V. DGGE patterns were visualized by staining with SYBR Green I nucleic acid gel stain (Molecular Probes, Eugene, OR, USA). Each gel included all treatments ($n = 4$) and sampling days ($n = 3$) per bull. To normalize the differences among gels, *Archaea*-specific fragments of one rumen reference sample were loaded in triplicate on each gel that was used as a standard for comparison of gels. The obtained DGGE patterns were analysed with the BioNumerics software

version 3.5 (Applied Maths, Kortrijk, Belgium). Similarities were calculated by the Pearson correlation. The clustering algorithm of Ward was used to calculate dendrograms (Boeckaert *et al.*, 2007). The DGGE profiles allow to gain insight into changes in diversity and functional organization of rumen microbial communities, which are assessed here through various indices and approaches:

(i) the range-weighted richness (R_r), calculated according to Boon *et al.* (2011) based on the total number of bands (N), and the denaturing gradient comprised between the first and the last band of the pattern (D_g) as: $R_r = N^2 \times D_g$; (ii) the Shannon (H), (iii) dominance (c) and (iv) evenness index (e), calculated as described by Patra *et al.* (2012) based on the peak surface of each band (n_i), the sum of all peak surfaces of all bands (N) and the total number of bands (S) as: $H = -\sum [(n_i/N) \times \ln(n_i/N)]$; $c = \sum(n_i/N)^2$; $e = H/\ln(S)$; (v) the Pareto–Lorenz evenness curves (PL curves), based on the DGGE profile using the Lorenz–Gini method (Mertens *et al.*, 2005; Marzorati *et al.*, 2008; Wittebolle *et al.*, 2009) and constructed by ranking the respective DGGE bands for each sample from high to low intensity and using the cumulative proportion of bands and their respective cumulative proportions of intensities as X- and Y-axis; and (vi) the community organization, which describes the evenness (equality) of the archaeal population and is the normalized area between a given PL-curve and the perfect evenness line. community organization ranges from 0 to 100 and represents a single value that describes a specific degree of evenness. The higher the community organization coefficient, the more uneven a community (Boon *et al.*, 2011).

Quantitative PCR (qPCR)

Archaea rRNA gene copies present in the DNA extract of ruminal digesta samples were quantified as described by Boeckaert *et al.* (2007) and Boon *et al.* (2003), using an ABI Prism SDS 7000 instrument (Applied Biosystems, Lennik, Belgium). Amplification reactions were carried out in 25 μ l volumes containing 12.5 μ l SYBR Green PCR Master Mix (Applied Biosystems, Warrington, UK), 6 μ l RNA-free water and 5 μ l diluted (1 : 20) DNA extract, as well as the ARCH915 and UNI-b-rev primers at a final concentration of 300 nM each. Measurements were taken in triplicate for each run. A standard curve for qPCR was constructed using six different

DNA concentrations ($n=3$), ranging from 1.13 copies to 1.13×10^8 copies of DNA per μ l. An *Archaea* 492 bp PCR fragment inserted in a TOPO vector was used as a template for the standard curve. Average amplification efficiency, slope and R^2 of qPCR assays were 102% (calculated as $E = 10^{(-1/\text{slope})} - 1$), -3.27 and 0.99 , respectively.

Statistical analysis

Before statistical analysis, protozoa numbers were averaged over a day (i.e. average of 3, 6, 9 and 12 h after feeding) and log transformed. Daily average protozoal numbers, VFA concentrations and proportions, *archaeal* log copies from qPCR and the indices as calculated from the DGGE profiles were subjected to a linear mixed model using SPSS 21 (SPSS Inc., IBM, Chicago, USA), according to the model $Y_{ijk} = \mu + A_i + B_j + C_k + \varepsilon_{ijk}$ where Y_{ijk} is the overall response; μ the overall of mean; A_i the mean effect of treatment ($i=1$ to 4) (fixed factor); B_j the mean effect of the period ($j=1$ to 4) (random factor); C_k the mean effect of animal ($k=1$ to 8) (random factor) and ε_{ijk} the residual error. Differences between means were considered to be significant at the $P < 0.05$ level, tendencies at $P < 0.10$. Treatment differences were distinguished using a two-sided Dunnett's test.

Results

Rumen fermentation characteristics

Treatments with MCFA resulted in a higher concentration of total VFA ($P < 0.05$), a lower proportion of acetate ($P < 0.01$) and higher proportion of propionate ($P < 0.01$) in comparison with the control treatment T (Table 3). Feeding CO and KO + T diets resulted in a similar proportion of acetate and propionate. The KO diet was associated with the lowest proportion of acetate ($P < 0.05$) and the highest proportions of propionate ($P < 0.001$) and minor VFA (iso-butyrate, valerate and iso-valerate) ($P < 0.05$). CO and KO + T also tended to increase proportions of minor VFA ($P < 0.10$) as compared with T.

Protozoa

All treatments supplemented with a MCFA source (CO, KO and KO + T) showed a lower number of protozoa than the control treatment (T). Treatments with the same level of MCFA reduced the protozoa numbers to a similar extent

Table 3 Total VFA concentrations and proportions of individual VFA in the rumen samples of bulls on a TMR supplemented (35 g/kg DM) with either beef T, CO, KO or a combination of KO + T

Fatty acid	T	CO	KO	KO + T	s.e.m.
Total VFA (mmol/l)	73.7 ^a	80.3 ^b	79.6 ^b	80.7 ^b	7.16
Acetate (mmol/mol total VFA)	650 ^a	617 ^b	597 ^b	609 ^b	17.8
Propionate (mmol/mol total VFA)	179 ^a	220 ^b	252 ^c	231 ^{bc}	24.6
Butyrate (mmol/mol total VFA)	140 ^a	131 ^{ab}	117 ^c	128 ^b	12.9
Other VFA (mmol/mol total VFA)*	30 ^a	32 ^{+b}	35 ^b	32 ^{+b}	3.3

VFA = volatile fatty acid; TMR = total mixed ration; T = tallow; CO = coconut oil; KO = krabok oil.

^{a-c}Means within a row with different superscripts differ ($P < 0.05$).

^{+b}Means within a row with different superscripts tend to differ ($P < 0.10$).

*Other VFA include isobutyrate, isovalerate and valerate.

Table 4 Protozoa counts, qPCR of Archaea and diversity and community structure indices of rumen Archaea, on the basis of archaeal DGGE profiles (range-weighted richness, Shannon H-index, evenness, dominance, and community organization) of the rumen samples of bulls on a TMR supplemented (35 g/kg DM) with either beef T, CO, KO or KO + T

	T	CO	KO	KO + T	s.e.m.
Protozoa ($\times 10^5$ /ml)	9.2 ^a	6.3 ^b	4.5 ^c	6.5 ^b	2.41
Archaea (log copy/g rumen sample)	5.31 ^a	5.33 ^a	4.75 ^b	5.09 ^{+b}	0.231
Range-weighted richness	4.96 ^a	4.92 ^a	5.69 ^b	5.75 ^b	0.270
Shannon H-index	1.32 ^{ab}	1.25 ^b	1.39 ^{+b}	1.45 ^a	0.635
Evenness	0.80	0.79	0.78	0.84	0.036
Dominance	0.32 ^{ab}	0.37 ^a	0.31 ^{ab}	0.29 ^b	0.028
Community organization	61.2 ^{ab}	63.6 ^a	57.5 ^{bc}	53.2 ^c	2.18

DGGE = denaturing gradient gel electrophoresis; TMR = total mixed ration; T = tallow; CO = coconut oil; KO = krabok oil.

^{a-c}Means within a row with different superscripts differ ($P < 0.05$).

^{+b}Means within a row with different superscripts tend to differ ($P < 0.10$).

(0.17 and 0.15 log units for CO and KO + T, respectively) (Table 4), whereas KO (with the highest levels of MCFA) had the lowest number of protozoa, which represented a decrease by 0.30 log unit compared with the control (T).

Molecular characterization of Archaea

Quantitative real-time PCR assays showed a decrease in the numbers of Archaea ($P < 0.05$ and $P = 0.06$) when supplementing krabok oil (KO and KO + T) compared with the CO and control treatment (T) (Table 4).

The DGGE profile of the archaeal population in the rumen did not result in a grouping of treatments, nor of animals (Figure 1). This could be caused by inter-animal variation in the methanogenic population. To check this, the DGGE profile of the Archaea present in the rumen when animals received the control treatment (T treatment) is presented (Figure 2). Three main groups of archaeal profiles with differences in predominant bands were observed. This clustering was not determined by the prior treatment (Table 1) and did not represent bull pairs receiving the same treatment sequence (Table 1), suggesting both the washout and the adaptation period, guaranteeing an interval of at least 35 days before rumen sampling, were enough to remove the effect of the prior treatment in the current study.

To evaluate whether stress, such as the supplementation of a MCFA source provoked shifts in diversity and organization of the archaeal population, irrespective of the inter-animal variation, several indices were calculated from the DGGE profiles (Table 4). Greater differences were observed between KO (KO and KO + T treatments) and CO supplementation than between any of the MCFA treatments and T. The rumen of bulls receiving KO (both KO and KO + T treatments) seemed colonized by a more diverse (based on the range-weighted richness, R_r and Shannon index, H) and less-dominated (based on the dominance index and the community organization) archaeal community (Table 4). The latter was also illustrated in the PL curves (Figure 3): in case of CO supplementation, 30% of all bands of the archaeal community accounted for 80% of their cumulative abundance, whereas 40% of all bands were required to cover this level of abundance when supplementing with KO + T.

Discussion

CO and KO markedly increased propionate proportions (23% to 41%) at the expense of acetate and decreased the numbers of protozoa in the rumen. This is in line with *in vitro* studies by Panyakaew *et al.* (2013), who reported that MCFA supply by CO and KO (0.8 or 1.2 mg/ml of incubation fluid) increased the proportion of propionate by 35% to 49%, which was accompanied by an inhibition in methane production of 13% to 38%. Propionate formation is an alternative to H_2 formation, which is typically favoured when concentrations of dissolved H_2 increase as hydrogen gas formation under these circumstances becomes thermodynamically unfavourable (Janssen, 2010). As the latter is the main energy source of methanogenic Archaea, greater propionate proportions might be indicative of less CH_4 formation per unit of rumen-fermented feed (Janssen, 2010).

The number of protozoa decreased by 0.15 to 0.31 log units in the CO and KO supplemented diets compared with the control diet. Defaunating effects of CO have been reported before: for example, a reduction in ciliate protozoa counts of 0.91 and 1.46 log units were reported in sheep when supplementing 35 and 70 g/kg DM of CO (Machmüller and Kreuzer, 1999). Another study from the same group (Machmüller *et al.*, 2000) reported a 0.59 log unit decrease in protozoal numbers when sheep were fed CO at a concentration of 56 g/kg DM, which is 1.5 times more than the amount used in our study. A similar supplementation rate (50 g/kg DM) to dairy cows, provoked a decrease of 1.05 log units in total protozoa, with a shift in the protozoal community towards *Epidinium* and *Isotricha*, whereas *Entodinium* and *Dasytricha* seemed more sensitive to CO (Reveneau *et al.*, 2012).

In their study on the effect of individual fatty acids, Dohme *et al.* (2001) suggested C10:0 and C12:0 to provoke an equally effective defaunating effect, whereas C14:0 did not reduce protozoal numbers. However, the supplementation rates of C10:0 + C12:0 in the CO treatment were highest (16.8 g/kg DM), whereas the KO and KO + T treatments supplied lower amounts of these MCFA (15.3 and 12.6 g/kg DM, respectively). Nevertheless, the KO treatment resulted in




Figure 1 Cluster analysis of the denaturing gradient gel electrophoresis (DGGE) profile of *Archaea* present in the rumen of bulls on a total mixed ration supplemented (35 g/kg DM) with either beef tallow (T), coconut oil (CO), krabok oil (KO) or KO + T. To guarantee readability, the reference numbers of 'bull' and 'treatment' codes were combined when replicates clustered, as indicated by vertical lines next to the profile.




Figure 2 Denaturing gradient gel electrophoresis (DGGE) profile indicates diversity in the rumen archaeal community, despite uniform diet fed to all bulls (total mixed ration supplemented with 35 g/kg DM of beef tallow). Three major clusters can be distinguished, indicated as 1 (grouping bull 7 and 8), 2 (grouping bulls 1, 4 and 5) and 3 (grouping bulls 2, 3 and 6).




Figure 3 Pareto-Lorenz distribution curves based on denaturing gradient gel electrophoresis (DGGE) profiles of archaeal communities in the rumen of bulls supplemented (35 g/kg DM) with either beef tallow (T), coconut oil (CO), krabok oil (K) or KO + T. Line A indicates the cumulative band intensity corresponding to a combination of 20% of all the bands for the KO + T treatment (52%). Lines B, C and D indicate the proportion of bands required to cover 80% of the cumulative band intensities for T and CO (30.2%), KO (33.3%) and KO + T (40.0%) treatments, respectively.

the strongest effect on protozoal numbers, whereas reductions by the CO and KO + T treatments were similar, which is in line with the total amount of C10:0 + C12:0 + C14:0 supplemented (Table 1). The latter might suggest C14:0 to be equally effective in reducing protozoal numbers as C10:0 and C12:0 or a supportive effect of C14:0 when combined with C12:0 and C10:0, as has been suggested for ruminal methanogens and methane production (Soliva *et al.*, 2004).

Cluster analysis on the DGGE profiles of rumen *Archaea* from bulls monitored in the current study (supplementation of maximum 35 g/kg CO or KO) showed no treatment grouping.

The result was similar to that of Kongmun *et al.* (2011), supplying a somewhat lower dose (26 g/kg of CO) to rumen-fistulated swamp buffalo bulls. The latter supplementation rate also did not change the archaeal numbers as assessed by real-time PCR (qPCR). Similarly, in the current study, no change in the archaeal numbers were induced by CO supplementation. Nevertheless, supplementing these levels of KO (35 g/kg DM) reduced the archaeal numbers (−0.56 log copies/g rumen contents). Even when reducing the level of krabok oil (KO + T) to supply similar amounts of MCFA as through CO, archaeal numbers tended to be lower (−0.22 log copies/g rumen contents). This would suggest C14:0 to be more powerful in reducing methanogenic numbers than C12:0, as the main difference between KO and CO is its greater C14:0 content. This is in contrast with what has been suggested by Soliva *et al.* (2003 and 2004), from their *in vitro* studies using the Hohenheim gas test. They reported a stronger decrease in the number of rumen *Archaea* with an increasing proportion of C12:0. A 2/1 proportion of C12:0/C14:0 (a ratio which is similar to what is observed for CO) decreased the total archaeal counts by 1.54 log units, whereas a 1/1 ratio (similar to that found in KO) decreased total *Archaea* by 0.44 log units (Soliva *et al.*, 2003). In a follow-up study (Soliva *et al.*, 2004), this was confirmed, although a synergistic effect of C12:0 in combination with C14:0 was suggested, with archaeal numbers being lowest for a C12:0/C14:0 ratio of 4/1 and 3/2. Differences between the two *in vitro* studies and our study might be related to the nature of the experiment (i.e. *in vitro* v. *in vivo*), where *in vitro* methodology eventually reflects a more acute action, whereas *in vivo* measurements were taken after at least 2 weeks of supplementation of the MCFA source. Lower (acute) toxicity of C14:0 has been proposed to potentially

induce a lag-time in response, which might result in an underestimation of its effectiveness when assessed through batch *in vitro* incubations (Soliva *et al.*, 2004).

In the current study, there was no noticeable relationship between the protozoal population, which was affected by both CO and KO, and the archaeal population. Similarly, in the Rusitec study on CO (54 g/kg DM) (Dohme *et al.*, 1999), the decrease in archaeal numbers occurred, irrespective of the faunating status (defaunated or faunated) of the rumen inoculum. This could suggest MCFA to act independently on protozoa and methanogens.

As the cluster analysis of DGGE data of the archaeal community did not allow distinction across treatments, we assessed whether treatment affected the diversity of the archaeal community on the basis of measures such as richness (R_r and H) and community organization (evenness, dominance and community organization) (Table 4). For all parameters, contrasts were largest between the CO and KO + T treatments, whereas the KO and KO + T treatments did not differ. Richness (R_r and H), as assessed from the number of DGGE bands, was increased by KO supplementation, whereas CO supplementation did not change diversity indices. This is in contrast with Pilajun and Wanapat (2011), who suggested an increased diversity of methanogens, from their *in vivo* study on CO (36 g/kg DM) supplemented to swamp buffaloes, although no indices were calculated from the DGGE profiles presented in their paper, or at least these were not shown. Furthermore, the archaeal community showed a more dominated structure (greater dominance and community organization) when supplementing CO as compared with KO or KO + T. Increases in dominance and community organization were also shown when supplementing CO *in vitro* (Patra and Yu, 2013). Nevertheless, a recent *in vivo* study on dairy cattle fed lauric acid, showed a decrease in the prevalence of *Methanobrevibacter*, the most predominant *Archaea*, whereas the prevalence of the less-abundant *Methanosphaera* increased (Hristov *et al.*, 2012). This reduction in dominance within the archaeal community was hypothesized to be linked to strong decreases (about 2 log-units) in protozoal numbers with which *Methanobrevibacter* might be associated. However, in our study, protozoal numbers were reduced both by CO, KO and KO + T treatments, whereas treatments containing krabok oil (KO and KO + T treatments) showed greater diversity and reduced dominance as compared with the CO treatment. Hence, differences in archaeal diversity and dominance seemed independent of changes in protozoal numbers in our study. Morgavi *et al.* (2012) showed that the DGGE profiles of the rumen methanogenic community only differed between faunated and defaunated animals when protozoa had been removed for more than 2 years, whereas short-term defaunation had not yet provoked changes in the DGGE profiles generated from rumen methanogens. In our study, changes in organization and diversity of the archaeal community seemed to be more linked to reductions in archaeal numbers (provoked only by KO and KO + T treatments) rather than reductions in protozoal numbers (provoked both by CO,

KO and KO + T treatments). Obviously, it would be of utmost interest to link changes in microbial community diversity and organization to function (i.e. methane production in this case) in the future, as reduced dominance and larger diversity were suggested to allow methanogens to adapt more easily to different niches (e.g. free-living, particulate associated, ectosymbionts or endosymbionts of protozoa) and fluctuating substrate concentrations (in particular H_2) (Attwood *et al.*, 2011), which might be induced by methane mitigation strategies such as oil supplementation (Janssen, 2010).

Although both *in vivo* and *in vitro* research suggested C14:0 to be less effective in modifying rumen metabolism (e.g. Hristov *et al.*, 2012), KO, which is particularly enriched in C14:0, seemed to induce shifts in rumen fermentation patterns, protozoal numbers and archaeal abundance, diversity and community structure at lower doses than CO. Further research should elucidate whether KO could be used as a methane mitigation strategy as rumen methanogen numbers do not always seem a key factor determining CH_4 production (e.g. Popova *et al.*, 2011).

Conclusions

Both KO and CO supplementation shifted rumen fermentation towards propionate at the expense of acetate and reduced protozoal numbers. However, decreases in archaeal numbers and changes in the diversity and the structure of the archaeal community were limited to the KO supplemented treatments.

Acknowledgements

This experiment was funded by the EU-community (project TH/Asia-Link/014-141-176 and Food-CT-23006-36241, Prosa-febeef), Paiwan Panyakaew further received grants from the Rajamangala University of Technology Isan, Thailand and Lotus-Erasmus Mundus Action 2 (EU-South-East Asian mobility project).

References

- Attwood GT, Altermann E, Kelly WJ, Leahy SC, Zhang L and Morrison M 2011. Exploring rumen methanogen genomes to identify targets for methane mitigation strategies. *Animal Feed Science and Technology* 166–167, 65–75.
- Boeckaert C, Fievez V, Van Hecke D, Verstraete W and Boon N 2007. Changes in rumen biohydrogenation intermediates and ciliate protozoa diversity after algae supplementation to dairy cattle. *European Journal of Lipid Science and Technology* 109, 767–777.
- Boon N, Top EM, Verstraete W and Siciliano SD 2003. Bioaugmentation as a tool to protect the structure and function of an activated-sludge microbial community against a 3-chloroaniline shock load. *Applied and Environmental Microbiology* 69, 1511–1520.
- Boon N, Pycke BFG, Marzorati M and Hammes F 2011. Nutrient gradients in a granular activated carbon biofilter drives bacterial community organization and dynamics. *Water Research* 45, 6355–6361.
- Dohme F, Machmüller A, Wasserfallen A and Kreuzer M 2000. Comparative efficiency of various fats rich in medium-chain fatty acids to suppress ruminal methanogenesis as measured with RUSITEC. *Canadian Journal of Animal Science* 80, 473–482.
- Dohme F, Machmüller A, Wasserfallen A and Kreuzer M 2001. Ruminal methanogenesis as influenced by individual acids supplemented to complete ruminant diets. *Letters in Applied Microbiology* 32, 47–51.

- Dohme F, Machmüller A, Estermann BL, Pfister P, Wasserfallen A and Kreuzer M 1999. The role of the rumen ciliate protozoa for methane suppression caused by coconut oil. *Letters in Applied Microbiology* 29, 187–192.
- Finlay BJ, Esteban G, Clarke KJ, Williams AG, Embley TM and Hirt RP 1994. Some rumen ciliates have endosymbiotic methanogens. *FEMS Microbiology Letters* 117, 157–162.
- Galyean M 1989. Laboratory procedure in Animal Nutrition Research. Department of Animal and Range Sciences, New Mexico State University, USA.
- Goel G, Arvidsson K, Vlaeminck B, Bruggeman G, Deschepper K and Fievez V 2009. Effects of capric acid on rumen methanogenesis and biohydrogenation of linoleic and α -linolenic acid. *Animal* 3, 810–816.
- Hristov AN, Callaway TR, Lee C and Dowd SE 2012. Rumen bacterial, archaeal, and fungal diversity of dairy cows in response to ingestion of lauric or myristic acid. *Journal of Animal Science* 90, 4449–4457.
- Hristov AN, Vander Pol M, Agle M, Zaman S, Schneider C, Ndegwa P, Vaddella VK, Johnson K, Shingfield KJ and Karnati SKR 2009. Effect of lauric acid and coconut oil on ruminal fermentation, digestion, ammonia losses from manure and milk fatty acid composition in lactating cows. *Journal of Dairy Science* 92, 5561–5582.
- Janssen PH 2010. Influence of hydrogen on rumen methane formation and fermentation balances through microbial growth kinetics and fermentation thermodynamics. *Animal Feed Science and Technology* 160, 1–22.
- Klevenhusen F, Meile L, Kreuzer M and Soliva CR 2011. Effects of monolaurin on ruminal methanogens and selected bacterial species from cattle, as determined with the rumen simulation technique. *Anaerobe* 17, 232–238.
- Kongmun P, Wanapat M, Pakdee P, Navanukraw C and Yu Z 2011. Manipulation of rumen fermentation and ecology of swamp buffalo by coconut oil and garlic powder supplementation. *Livestock Science* 135, 84–92.
- Machmüller A and Kreuzer M 1999. Methane suppression by coconut oil and associated effects on nutrient and energy balance in sheep. *Canadian Journal of Animal Science* 79, 65–72.
- Machmüller A, Ossowski DA and Kreuzer M 2000. Comparative evaluation of the effects of coconut oil, oilseeds and crystalline fat on methane release, digestion and energy balance in lambs. *Animal Feed Science and Technology* 85, 41–60.
- Machmüller A, Soliva CR and Kreuzer M 2003a. Effect of coconut oil and defaunation treatment on methanogenesis in sheep. *Reproduction, Nutrition, Development* 43, 41–55.
- Machmüller A, Soliva CR and Kreuzer M 2003b. Methane-suppressing effect of myristic acid in sheep as affected by dietary calcium and forage proportion. *British Journal of Nutrition* 90, 529–540.
- Marzorati M, Wittebolle L, Boon N, Daffonchio D and Verstraete W 2008. How to get more out of molecular fingerprints: practical tools for microbial ecology. *Environmental Microbiology* 10, 1571–1581.
- Mertens B, Boon N and Verstraete W 2005. Stereo specific effect of hexachlorocyclohexane on activity and structure of soil methanotrophic communities. *Environmental Microbiology* 7, 660–669.
- Morgavi DP, Martin C, Jouany JP and Ranilla MJ 2012. Rumen protozoa and methanogenesis: not a simple cause-effect relationship. *British Journal of Nutrition* 107, 388–397.
- Panyakaew P, Goel G, Lourenço M, Yuangklang C and Fievez V 2013. Medium-chain fatty acids from coconut or krabok oil inhibit *in vitro* rumen methanogenesis and conversion of non-conjugated dienoic biohydrogenation intermediates. *Animal Feed Science and Technology* 180, 18–25.
- Patra AK and Yu Z 2013. Effects of coconut and fish oils on ruminal methanogenesis, fermentation, and abundance and diversity of microbial populations *in vitro*. *Journal of Dairy Science* 96, 1–11.
- Patra AK, Stiverson J and Yu Z 2012. Effects of quillaja and yucca saponins on communities and select populations of rumen bacteria and archaea, and fermentation *in vitro*. *Journal of Applied Microbiology* 113, 1329–1340.
- Pilajun R and Wanapat M 2011. Methane production and methanogen population in rumen liquor of swamp buffalo as influenced by coconut oil and mangosteen peel powder supplementation. *Journal of Animal and Veterinary Advances* 10, 2523–2527.
- Popova M, Martin C, Eugene M, Mialon MM, Doreau M and Morgavi DP 2011. Effect of fibre- and starch-rich finishing diets on methanogenic Archaea diversity and activity in the rumen of feedlot bulls. *Animal Feed Science and Technology* 166–167, 113–121.
- Reveneau C, Karnati SKR, Oelker ER and Firkins JL 2012. Interaction of unsaturated fat or coconut oil with monensin in lactating dairy cows fed 12 times daily. I. Protozoal abundance, nutrient digestibility, and microbial protein flow to the omasum. *Journal of Dairy Science* 95, 2046–2060.
- Soliva CR, Hindrichsen IK, Meile L, Kreuzer M and Machmüller A 2003. Effects of mixtures of lauric and myristic acid on rumen methanogens and methanogenesis *in vitro*. *Letters Applied Microbiology* 37, 35–39.
- Soliva CR, Meile L, Hindrichsen IK, Kreuzer M and Machmüller A 2004. Myristic acid supports the immediate inhibitory effect of lauric acid on ruminal methanogens and methane release. *Anaerobe* 10, 269–276.
- Van Ranst G, Fievez V, Vandewalle M, Van Waes C, De Riek J and Van Bockstaele E 2010. Influence of damaging and wilting red clover on lipid metabolism during ensiling and *in vitro* rumen incubation. *Animal* 4, 1528–1540.
- Wittebolle L, Marzorati M, Clement L, Balloi A, Daffonchio D, Heylen K, De Vos P, Verstraete W and Boon N 2009. Initial community evenness favours functionality under selective stress. *Nature* 458, 623–626.
- Yu Z and Morrison M 2004. Comparisons of different hypervariable regions of *rrs* genes for use in fingerprinting of microbial communities by PCR-denaturing gradient gel electrophoresis. *Applied and Environmental Microbiology* 70, 4800–4806.
- Yu Z, García-González R, Schanbacher FL and Morrison M 2008. Evaluations of different hypervariable regions of archaeal 16S rRNA genes in profiling of methanogens by Archaea-specific PCR and denaturing gradient gel electrophoresis. *Applied and Environmental Microbiology* 74, 889–893.