



Depsidones inhibit aromatase activity and tumor cell proliferation in a co-culture of human primary breast adipose fibroblasts and T47D breast tumor cells

Suthat Chottanapund^{a,b,c,d,e,*}, M.B.M. Van Duursen^e, Anne Zwartsen^{b,e}, Supatchaya Timtavorn^b, Panida Navasumrit^{a,b,c}, Prasat Kittakoop^{a,b,c}, Sanya Sureram^b, Mathuros Ruchirawat^{a,b,c}, Martin Van den Berg^e

^a Chulabhorn Graduate Institute, Kamphaeng Phet 6 Road, Laksi, Bangkok 10210, Thailand

^b Chulabhorn Research Institute, Kamphaeng Phet 6 Road, Laksi, Bangkok 10210, Thailand

^c Center of Excellence on Environmental Health and Toxicology (EHT), CHE, Ministry of Education, Bangkok, Thailand

^d Bamrasnaradura Infectious Diseases Institute, Ministry of Public Health, Thailand

^e Institute for Risk Assessment Sciences, Utrecht University, Utrecht, The Netherlands

ARTICLE INFO

Keywords:

Depsidones
Aromatase
Breast cancer
Co-culture

ABSTRACT

Naturally occurring depsidones from the marine fungus *Aspergillus unguis* are known to have substantial anti-cancer activity, but their mechanism of action remains elusive. The purpose of this study was to examine the anti-aromatase activity of two common depsidones, unguinol and aspergillusidone A, in a co-culture system of human primary breast adipose fibroblasts and hormonal responsive T47D breast tumor cells. Using this *in vitro* model it was shown that these depsidones inhibit the growth of T47D tumor cells most likely via inhibition of aromatase (CYP19) activity. The IC₅₀ values of these depsidones were compared with the aromatase inhibitors letrozole and exemestane. Letrozole and exemestane had IC₅₀ values of respectively, 0.19 and 0.14 μM, while those for Unguinol and Aspergillusidone A were respectively, 9.7 and 7.3 μM. Our results indicate that among the depsidones there maybe aromatase inhibitors with possible pharmacotherapeutical relevance.

1. Introduction

Depsidones are secondary metabolites mostly found in lichens, but also in some higher plants, where they play a role in the protection against insects and microbes or sun light. Depsidones are esters which compose of polyphenolic depsides and cyclic ethers, but the chemical structure of the depsidones formed is highly dependent on the type of fungus and its environment [1]. Generally, the chemical structure of depsidones resembles prostaglandins and leukotrienes in humans, which makes these compounds may have beneficial health effects in humans. Indeed, several studies have already reported biological activities of these naturally occurring depsidones like anti-proliferative actions [2–4], antimalarial and cytotoxic properties [5], antibacterial activity against the multidrug-resistant *Staphylococcus aureus* [6], radical scavenging, antioxidant activities and antifungal activities [7–9]. On the other hand, because of their structural similarity, leukotriene depsidones may induce allergic reactions in humans [10]. Also, apoptosis or inflammatory processes that involve free oxygen

radicals by depsidones have been reported to cause cytotoxicity in rat hepatocytes [11] and rat thymocytes [12]. More recently, *in vitro* studies from our laboratory indicate that some depsidones are potent inhibitors of the aromatase enzyme (CYP 19) [7,9]. The aromatase enzyme is responsible for the conversion of androgens into estrogens and is used as a therapeutic target for breast cancer treatment [13,14]. In view of the above biological properties of these depsidones novel cancer treatments may arise. In this respect, the inhibition of aromatase activity may be especially relevant for treatment of estrogen-dependent tumors, such as breast tumors.

Aromatase inhibitors, such as exemestane and letrozole, are currently used as hormonal therapy in estrogen positive postmenopausal breast cancer patients [15]. To detect inhibitory properties of aromatase, studies are frequently performed using relatively simple *in vitro* models such as human placental tissue in which radiolabeled androgens are used as substrates [16]. Other methods such as HPLC separation with UV detection [17] and recombinant enzyme systems are often less sensitive and laborious [18]. Stresser et al. developed a high throughput

* Corresponding author at: Laboratory of Environmental Toxicology, Chulabhorn Research Institute, 54 Kamphaeng Phet 6 Talat Bang Khen, Lak Si, Bangkok 10210, Thailand.
E-mail address: suthat.c@bidi.mail.go.th (S. Chottanapund).

screening method using the fluorometric substrate O-benzyl fluorescein benzyl ester (DBF) as a substrate for microsomal aromatase activity [19]. This method was also found suitable for detecting aromatase inhibition in our present co-culture study. In our previous study, we used a more realistic *in vitro* breast cancer model that consists of a co-culture of primary human breast fibroblasts with hormonal positive T47D breast cancer cells to study effects on aromatase activity [20]. In this co-culture model, paracrine interactions between various cell types are implemented leading to a more physiologically relevant test model to assess local effects of compounds on aromatase activity and subsequent (anti)tumor effects.

The purpose of this co-culture study was to examine the inhibitory properties of the common depsidones unguinol and aspergillusidone A, which are isolated from the marine-derived fungus *A. unguis* CRI282-03, on aromatase activity in primary human breast fibroblasts and its subsequent effect on the proliferation of T47D breast tumor cells. The results were compared with two clinical relevant aromatase inhibitors, letrozole and exemestane.

2. Materials and methods

2.1. Depsidones

Depsidones were isolated from *Aspergillus unguis* CRI282-03, a fungus isolated from the marine environment by the laboratory of natural products from the Chulaborn Research Institute (Bangkok, Thailand) [2,7]. For our present study two depsidones, unguinol (UNG) and aspergillusidone A (ASP-A), were selected. This was based on a preliminary study using the microsomal placental assay, which indicated inhibitory properties for aromatase [7]. Structures of UNG and ASP-A are shown in Fig. 1A and B. Range finding studies in our laboratory indicated cytotoxicity of UNG and ASP-A in the mM range [2,7], which is significantly higher than concentrations used in our present co-culture study. For comparison the IC₅₀ values of letrozole and exemestane were also determined under similar experimental conditions (See Fig. 1C and D). Cytotoxic effects of letrozole and exemestane have been reported earlier to be in the μM range and therefore concentrations in our co-culture experiments did not exceed 0.25 μM.

2.2. Breast cancer cell culture and incubation

The T47D cell line was obtained from ATCC (Rockville, MD, USA). These cells were grown in culture medium comprising of RPMI 1640, supplemented with 10% FBS, 1% penicillin/streptomycin, 1% l-gluta-

mine, 1% glucose, 1% sodium pyruvate, and 0.08% insulin solution. Seeding concentration was 5×10^4 cells/ml and cells were cultured at 37 °C in a humidified atmosphere with 5% CO₂ until 80–90% confluent.

2.3. Primary breast adipose fibroblasts (BAFs)

After written informed consent, BAFs were obtained from three breast cancer patients, who attended the Bamrasnaradura Infectious Disease Institute (Nonthaburi, Thailand) for modified radical mastectomy. The research protocol was approved by the Medical Ethical Committee of the Bamrasnaradura Infectious Disease Institute. About 5–10 g of macroscopically normal breast tissue was collected, while the remaining tissue was sent for routine pathological examination. Directly after the tissue was obtained via surgery, it was stored in a 4 °C saline solution (0.9% NaCl) and immediately transported to our laboratory. BAFs were isolated from this breast tissue as described earlier [21,22]. These BAFs were cultured as adherent cultures in RPMI 1640-medium w/o phenol red (Gibco/Invitrogen 11835) supplemented with Pen/Strep (Invitrogen 15140), FCS (Invitrogen 10270) and Insulin 10^{-3} M (144 mg/25 ml) (sigma I-5500) at a seeding concentration of 5×10^4 cells/ml. The cells were maintained in a humidified atmosphere at 37 °C with 5% CO₂. BAFs were sub-cultured if cell cultures in the bottles were confluent, which was usually once a week.

2.4. Breast cancer cells in co-culture with BAFs

Approximately three weeks after isolation, fibroblasts were used to establish a co-culture together with the T47D tumor cells [20,22,23]. On day one, BAFs were plated at a density of approximately 4×10^3 cells/well in a 96-well plate for proliferation or 5×10^4 cells/ml in a 25 cm² flask for gene expression and grown in culture medium as described above. The day after, mono-cultures of breast cancer cells and plated primary BAFs were washed with phosphate buffered saline (PBS) and placed on the assay medium in which heat inactivated FCS was replaced with heat-inactivated, charcoal/dextran-treated FCS (Hyclone, SH30068.03). On day 4, BAFs were washed with PBS and the T47D cells were trypsinized and seeded on top of the BAFs at a density of 4×10^3 cells/well for proliferation or 2.5×10^5 cells/flask for gene expression. On day 5, fresh assay medium was added to the co-cultures after which the cells were exposed to the test compounds for 96 h [24]. Final solvent concentration was 0.1% v/v in the medium. Initial concentrations were 1, 5, and 10 nM for testosterone, 50 nM for exemestane, 30 nM for letrozole, and 2 μM for depsidones to screen for aromatase inhibition in the presence of testosterone. In this co-culture model testosterone is converted by aromatase in the BAFs to estradiol, which in turn induces cell proliferation in the estrogen positive T47D tumor cells.

2.5. Cell proliferation

Cell proliferation was determined after treatment with testosterone by performing an MTT assay as described earlier [21]. After 96 h of exposure to the test compounds, 1 ml of MTT (5 mg MTT/ml in PBS) was added for 4 h. Next, the medium was aspirated and 100 μl of DMSO was added to dissolve the accumulated formazan crystals. The absorbance was measured at 550 nm using 96-well plates in a Spectramax plus 384 microplate reader (Molecular Devices, California, USA).

2.6. Modified high-throughput screen to identify inhibitors of Aromatase (CYP19) [19]

Aromatase activity was measured by the conversion of the fluorometric substrate O-benzylfluorescein benzyl ester (DBP) as described earlier [19] with the following minor modifications. Assays were conducted in 96-well microtiter plates (Catalog No. 3915, Corning Costar, Cambridge, MA) using a 200 μl volume containing a serially




Fig. 1. Chemical structures of a) unguinol (UNG), b) aspergillusidone A (ASP-A), c) letrozole and d) exemestane.




Fig. 2. Effect of testosterone on the proliferation of T47D cells in a mono- or co-culture with human breast adipose fibroblasts (BAFs). Data are expressed as mean proliferation \pm SD (N = 3). Statistical significance compared with vehicle-treated cells in the same experimental model is indicated as follows: reduced proliferation in T47D mono-culture: * = $p < 0.01$; increased proliferation in co-culture T47D and BAFs: # = $p < 0.01$.

dilution of 1:2 of UNG, ASP-A, exemestane or letrozole. DBF was obtained from Gentest Corporation (Woburn, MA). In this method the metabolite production of DBP by aromatase is linear in time up to 30 min and 2 μ M DBP, which was used to determine IC_{50} values of the test compounds. The enzymatic reaction was terminated by adding 40 μ l 2 N NaOH. To develop adequate signal to background ratio, the 96-well plates (with lid) were incubated for 2 h at 37 $^{\circ}$ C. Fluorescence in each individual well was measured in top-read mode using a FLUOstar Model 403 fluorescence plate reader (BMG LabTechnologies, Inc., Durham, NC). DBF metabolite concentrations were measured using an excitation wavelength of 485 nm and an emission wavelength of 538 nm.

2.7. Analysis of gene expression

After the cells were treated total RNA was extracted. RNA was reverse-transcribed into cDNA using the Qiagen high capacity cDNA reverse-transcription kit (Qiagen, Texas, USA). The gene expression of *pS2* and *Ki-67* was performed using a Roche-Light-cycler-480 one step RT PCR kit (Roche, Indiana, USA). Relative quantification of gene expression was expressed as the ratio of the intensity of the target gene to the housekeeping gene *beta-actin*. The RT-PCR conditions for each gene are described below.

2.8. RNA isolation and gene expression

Total RNA was isolated using RNeasy[®] mini kit (Qiagen, Texas, USA) according to the manufacturer's instruction. Cells were centrifuged and lysed with RLT buffer and β -mercaptoethanol. One volume of 70% ethanol was added and this was vigorously mixed. The sample was transferred to a RNeasy spin column, centrifuged at 12,000g for 15 s, and then 350 μ l washing buffer RW1 was added on the column and again centrifuged at 12,000 g for 15 s. The 80 μ l of DNase I reaction mixture was directly added on the membrane of the column and incubated for 15 min. Thereafter, the column was washed with RW1 buffer and RPE buffer twice. Bound RNA was eluted with 40 μ l of RNase-free water and centrifuged at 12,000g for 1 min. The purification and concentration of total RNA were measured by the ratio of absorbance at 260/280 and 260/230 NM using Nanodrop (Nanodrop Technologies, Inc., Delaware, USA).

Obtained RNA was stored at -20° C in aliquots of 10 ng/ μ l. The reaction mixture contained 0.1 μ M of primers, 1X Quantitect SYBR Green RT-PCR Master mix, 0.2 μ l QuantiTect RT mix and 2 μ l template RNA (10 ng/ μ l) in a total volume of 20 μ l. The mixture was reverse transcribed to cDNA at 50 $^{\circ}$ C for 20 min. After reverse transcription, the PCR reaction was initiated by heating at 95 $^{\circ}$ C for 15 min, then followed by denaturation at 95 $^{\circ}$ C for 10 s, annealing at 57 $^{\circ}$ C for 25 s, an

extension at 68 $^{\circ}$ C for 30 s and acquisition at 82 $^{\circ}$ C for 5 s for 45 cycles. Primers coding for the estrogen-responsive *pS2* gene were used as described earlier by Lee et al. [25,26]. Primers for *Ki-67* amplification were designed in our laboratory as described previously [23]. Expression of *β -ACTIN* (forward: 5'-TCTACAATGAGCTGCG-3' and reverse: 5'-AGGTAGTCAGCTAGGT-3') was used as a reference housekeeping gene. All primers were run through the National Center for Biotechnology Information (NCBI) blast (nucleotide nonredundant database) to confirm specificity. The efficiency was determined for a dilution range of cDNA and primers were only used with an efficiency between 95% and 105%. After each RT-qPCR, a melt curve was run to ensure that primer-dimers and other non-specific products were omitted.

2.9. Data analysis

Each experiment and each concentration were performed in triplicate. Data were analyzed using Microsoft Excel and IC_{50} values calculated by interpolation. Means and standard deviations were calculated of all nine values ($n = 9$). The statistical significance of differences of the means were calculated using the Student's *t*-test or one-way ANOVA. Differences were considered statistically significant if $P < 0.05$.

3. Results

3.1. Exposure of mono and co-cultured T47D cells with BAFs to testosterone

Firstly, aromatase activity and cell proliferation were determined in a co-culture of BAFs and T47D cells to validate the performance of the co-culture model. BAFs were obtained from one patient, since it has been established that aromatase activity can vary largely between individuals [23]. As a result, this variation may influence the sensitivity and functionality of this co-culture system significantly [21,23]. Adding testosterone upto 10 nM increased proliferation of T47D cells in the co-culture up to 30%. In contrast, adding testosterone to a mono-culture of T47D cells caused a maximum reduction in cell growth of 50% at 10 nM ($p < 0.01$) (See Fig. 2). These observations confirm the aromatization of testosterone to estradiol in the co-culture of T47D and BAFs, which is followed by a subsequent increase in estrogen-dependent cell proliferation of the tumor cells. Furthermore, the reduction of cell proliferation in the mono culture of T47D cells by testosterone confirms our earlier observations that androgens can reduce the proliferation of androgen receptor positive breast cancer cell lines like that of the T47D cells [23].

The inhibition of cell proliferation of T47D cells in co-culture with BAFs by the different test compounds at different concentrations of testosterone was also studied. As could be expected the aromatase inhibitors letrozole and exemestane inhibited T47D cell proliferation at




Fig. 3. Inhibition of T47D cell proliferation by letrozole (a), exemestane (b), UNG (c) and ASP-A (d) when co-cultured with human breast adipose fibroblasts at different testosterone concentrations. In the presence of the compounds tested, testosterone-induced proliferation (see Fig. 2) could no longer be observed in the T47D-BAF co-culture. Data are expressed as mean proliferation ($N = 3$).

concentrations of respectively 30 and 50 nM (See Fig. 3A and B).

At concentrations of 2 μM UNG and ASP-A inhibition of T47D cell proliferation in these co-cultures was observed, albeit at approximately two orders of magnitude higher concentrations than those of letrozole and exemestane (see Fig. 3C and D). These results indicate that UNG and ASP-A may act as aromatase inhibitors in these T47D and BAF co-cultures. In the subsequent experiments the relative potency of these compounds as aromatase inhibitors were studied in further detail for confirmation.

3.2. Aromatase inhibition in co-cultures of T47D cells and BAFs

The inhibitory potency of letrozole, exemestane, UNG and ASP-A on aromatase activity was determined at a fixed DBP concentration of 2 μM . A concentration-dependent inhibition of aromatase was observed for all four compounds. IC_{50} values for letrozole, exemestane, UNG and ASP-A were 0.19 μM (95% CI; 0.13–0.28 μM), 0.14 μM (95% CI; 0.11–0.18 μM), 9.73 μM (95% CI; 7.31–12.96 μM) and 7.26 μM (95% CI; 4.99–10.57 μM), respectively. Concentration – response relationships for these compounds are shown in Fig. 4A–D.

3.3. *pS2* and *Ki67* gene expression in co-cultures of T47D cells and BAFs

When a co-culture of T47D cells and BAFs was exposed to testosterone only (1, 5, 10 nM), a concentration dependent increase in the expression of *pS2* and *Ki67* mRNA was observed ($p < 0.01$) (Fig. 5A and B). The induction of both cell proliferation markers was

considered to be induced by estradiol production via aromatase in the BAFs. To confirm this, the T47D and BAF co-cultures were exposed to 1, 5 or 10 nM testosterone in combination with the aromatase inhibitor exemestane (50 nM). Adding exemestane completely negated the induction of *pS2* and *Ki67* mRNA expression, which confirmed the functional role of aromatase in this estradiol production. Similarly, UNG and ASP-A (2 μM) fully negated the testosterone-induced expression of *pS2* and *Ki67* mRNA, which again confirms their role as aromatase inhibitors.

4. Discussion

In this study, we clearly show that the naturally occurring despidones unguinol and aspergillusidone A can effectively inhibit aromatase activity and subsequently reduce estrogen-dependent breast cancer cell proliferation in an *in vitro* co-culture model with T47D breast cancer cells and primary BAFs.

4.1. Physiological relevance of the co-culture system

This co-culture system consist of T47D breast tumor cells in combination with breast adipose tissue fibroblasts (BAFs). In our earlier studies, we established that this T47D cell line expresses both the estrogen α and androgen receptors and is very suitable for use in a co-culture with BAFs to mimic the *in vivo* breast cancer situation [23,27].

The role of steroid hormones in breast tumors is complex due to the various paracrine interactions between different cell types in the breast




Fig. 4. Inhibition of aromatase activity in co-cultures of T47D cells and human breast adipose fibroblasts by letrozole (a), exemestane (b), UNG (c) and ASP-A (d). Data are expressed as mean DBP intensity as marker for aromatase activity compared to vehicle-treated control cells \pm SD (N = 3).




Fig. 5. Effect of testosterone, exemestane, UNG and ASP-A on pS2 (A) and Ki67 (B) mRNA expression in co-cultures of T47D cells and human breast adipose fibroblasts. Data are expressed as means \pm SD (n = 9). * Statistically significantly different from vehicle-treated control co-cultures ($p < 0.001$).

tissue. We and others have previously shown that (part of) the intercellular communication between the tumor cells and surrounding BAFs remains operational in our co-culture model [21,28,29]. Therefore, the effects studied in the co-culture model used in present study have more physiological relevance and predictive value compared to effects a monoculture of breast tumor cells *in vitro*. For estrogen dependent breast tumors the presence of surrounding BAFs is especially relevant, as the local estrogen production is situated in BAFs due to the presence of the aromatase enzyme. In several earlier studies with these breast co-cultures we have shown their relevance for studying natural compounds like androgens, melatonin and resveratrol [20–23]. Most co-culture studies have focused on estrogen dependent tumors. However, a substantial number of breast tumors are not hormonal responsive and for these clinical treatment options are very limited [30–33]. At present, little information is available about the paracrine interactions and intercellular communications between non-hormonal responsive (triple negative) breast tumor cells and BAFs. In the quest for finding new compounds with anti-tumor properties for these non-hormonal responsive breast tumors these co-cultures may offer a more advanced *in vitro* testing system.

4.2. Depsidones as aromatase inhibitors

Our institute has already published several studies on depsidones that were isolated from *A. unguis* CRI282-03, which had anti-aromatase activity [2,9,34]. These previous studies with *A. unguis*-derived ASP-F, D, E and Unguinol showed that inhibition of aromatase activity by these depsidones was in the same order of magnitude, ranging between 0.5 and 1 μM [7]. This suggests that the differences in chemical structure with respect to halogen substitution and hydroxyl placement, have no apparent effect on the aromatase inhibitory potency of the depsidones tested. Although ASP-A was not tested in that study, the IC₅₀ for aromatase inhibition for unguinol was 0.6 μM compared with 9.7 μM in present study. This difference can most likely be attributed to the difference in experimental set-up; the study of Sureram et al. used recombinant human aromatase enzyme, while in this study primary human breast cells were used. In present study, the bioavailability of the tested depsidones might have been affected by metabolism or active excretion from the cells. Moreover, in present study, the aromatase inhibiting potencies of UNG and ASP-A were two orders of magnitude lower than the clinically used therapeutic exemestane. Yet, the aromatase inhibiting properties occurred at non-cytotoxic concentrations, which is relevant to assess with respect to breast cancer therapeutic. Other depsidones; e.g. fungal-derived norstictic acid and plant-derived phomopsidone A and atrovirisidone B have been shown to cause cytotoxicity in breast cancer cell lines. Norstictic acid inhibited MDA-MB-231 proliferation by inhibition of proto-oncogenic tyrosine kinase (c-Met) activity [35]. Phomopsidone A, which has the structure similar to breast cancer drug named Taxol, also inhibited the proliferation of breast cancer cell line *in vitro* [36]. Atrovirisidone B displayed cytotoxicity in MCF-7 cells with an IC₅₀ value of approximately 23 μM [37]. Clearly, there are more types of depsidones with bioactive properties that might be relevant for breast cancer research. To determine the clinical relevance of our findings with respect to aromatase inhibition, clearly a more toxicological and pharmacological comparison between these depsidone and standard anti-aromatase is needed. These studies should then also address possible adverse effects of depsidones; e.g. caused by cytotoxicity at the higher dose levels.

5. Conclusion

Our results demonstrate that in a co-culture system of estrogen-dependent T47D breast tumor cells and primary breast adipose fibroblasts two depsidones, unguinol and aspergillusidone A, can act as aromatase inhibitors leading to inhibition of tumor cell proliferation. The multiple anti-cancer activities of the depsidones warrant further

studies for their potential clinical applications.

Conflict of interest statement

All authors declared no financial and personal relationships with other people or organizations that could inappropriately influence (bias) this work.

Acknowledgements

This work was supported by the Chulabhorn Graduate Institute, Chulabhorn Research Institute (Thailand), and The Institute for Risk Assessment Sciences at the Utrecht University (The Netherlands).

References

- [1] M. Hauck, S.-R. Jürgens, C. Leuschner, Norstictic acid: correlations between its physico-chemical characteristics and ecological preferences of lichens producing this depsidone, *Environ. Exp. Bot.* 68 (3) (2010) 309–313.
- [2] S. Sureram, et al., Directed biosynthesis through biohalogenation of secondary metabolites of the marine-derived fungus *Aspergillus unguis*, *RSC Adv.* 3 (2013) 1781–1788.
- [3] S.U. Lee, et al., Protuboxepins A and B and protubonines A and B from the marine-derived fungus *Aspergillus* sp. SF-5044, *J. Nat. Prod.* 74 (5) (2011) 1284–1287.
- [4] T. Jiang, et al., Cerebroside analogues from marine-derived fungus *Aspergillus flavipes*, *J. Asian Nat. Prod. Res.* 6 (4) (2004) 249–257.
- [5] P. Khumkomkhet, et al., Antimalarial and cytotoxic depsidones from the fungus *Chaetomium brasiliense*, *J. Nat. Prod.* 72 (8) (2009) 1487–1491.
- [6] T. Kokubun, W.K. Shiu, S. Gibbons, Inhibitory activities of lichen-derived compounds against methicillin- and multidrug-resistant *Staphylococcus aureus*, *Planta Med.* 73 (2) (2007) 176–179.
- [7] S. Sureram, et al., Depsidones: aromatase inhibitors and radical scavenging agents from the marine-derived fungus *Aspergillus unguis* CRI282-03, *Planta Med.* 78 (6) (2012) 582–588.
- [8] R. Abdou, et al., Botryorhodines A–D: antifungal and cytotoxic depsidones from *Botryosphaeria rhodina*, an endophyte of the medicinal plant *Bidens pilosa*, *Phytochemistry* 71 (1) (2009) 110–116.
- [9] P. Chomcheon, et al., Aromatase inhibitory: radical scavenging, and antioxidant activities of depsidones and diaryl ethers from the endophytic fungus *Corynespora cassiicola* L36, *Phytochemistry* 70 (3) (2009) 407–413.
- [10] Z. Zheng, et al., Trivarinic acid: a potent depeptide human leukocyte elastase inhibitor, *Biol. Pharm. Bull.* 35 (12) (2012) 2247–2251.
- [11] E.R. Correche, et al., Cytotoxic and apoptotic effects on hepatocytes of secondary metabolites obtained from lichens, *Altern. Lab. Anim.* 32 (6) (2004) 605–615.
- [12] V. Pavlovic, et al., Effect of four lichen acids isolated from *Hypogymnia physodes* on viability of rat thymocytes, *Food Chem. Toxicol.* 51 (2012) 160–164.
- [13] E.A. Thompson Jr., P.K. Siiteri, The involvement of human placental microsomal cytochrome P-450 in aromatization, *J. Biol. Chem.* 249 (17) (1974) 5373–5378.
- [14] Y. Toma, et al., Diverse functions of aromatase: O-deethylation of 7-ethoxycoumarin, *Endocrinology* 137 (9) (1996) 3791–3796.
- [15] A. Brodie, Q. Lu, B. Long, Aromatase and its inhibitors, *J. Steroid Biochem. Mol. Biol.* 69 (1–6) (1999) 205–210.
- [16] P. Mak, F.D. Cruz, S. Chen, A yeast screen system for aromatase inhibitors and ligands for androgen receptor: yeast cells transformed with aromatase and androgen receptor, *Environ. Health Perspect.* 107 (11) (1999) 855–860.
- [17] J. Fauss, W. Pyerin, Rapid assay of aromatase activity by fast liquid chromatography: spectroscopic evaluation of metabolite profiles indicate unexpected pitfalls, *Anal. Biochem.* 210 (2) (1993) 421–423.
- [18] D.J. Zhou, D. Pompon, S.A. Chen, Stable expression of human aromatase complementary DNA in mammalian cells: a useful system for aromatase inhibitor screening, *Cancer Res.* 50 (21) (1990) 6949–6954.
- [19] D.M. Stresser, et al., A high-throughput screen to identify inhibitors of aromatase (CYP19), *Anal. Biochem.* 284 (2) (2000) 427–430.
- [20] S. Chottanapund, et al., Anti-aromatase effect of resveratrol and melatonin on hormonal positive breast cancer cells co-cultured with breast adipose fibroblasts, *Toxicol. In Vitro* 28 (7) (2014) 1215–1221.
- [21] M. Heneweer, et al., Co-culture of primary human mammary fibroblasts and MCF-7 cells as an *in vitro* breast cancer model, *Toxicol. Sci.* 83 (2) (2005) 257–263.
- [22] J.A. van Meeuwen, et al., (Anti)estrogenic effects of phytochemicals on human primary mammary fibroblasts: MCF-7 cells and their co-culture, *Toxicol. Appl. Pharmacol.* 221 (3) (2007) 372–383.
- [23] S. Chottanapund, et al., Effect of androgens on different breast cancer cells co-cultured with or without breast adipose fibroblasts, *J. Steroid Biochem. Mol. Biol.* 138 (2013) 54–62.
- [24] K. Sonne-Hansen, A.E. Lykkesfeldt, Endogenous aromatization of testosterone results in growth stimulation of the human MCF-7 breast cancer cell line, *J. Steroid Biochem. Mol. Biol.* 93 (1) (2005) 25–34.
- [25] Y. Lee, et al., A ginsenoside-Rh1: a component of ginseng saponin, activates estrogen receptor in human breast carcinoma MCF-7 cells, *J. Steroid Biochem. Mol. Biol.* 84 (4) (2003) 463–468.
- [26] D. Lu, et al., Transcriptional regulation of the estrogen-inducible pS2 breast cancer

- marker gene by the ERR family of orphan nuclear receptors, *Cancer Res.* 61 (18) (2001) 6755–6761.
- [27] J.A. Aka, S.X. Lin, Comparison of functional proteomic analyses of human breast cancer cell lines T47D and MCF7, *PLoS One* 7 (2) (2012) pe31532.
- [28] M. Heneweer. Co-culture of primary human mammary fibroblasts and MCF-7 cells as an in vitro breast cancer model Institute for Risk Assessment Sciences (IRAS). Vol. Ph. D., Utrecht: Utrecht University. 175, 2005.
- [29] Y. Miki, et al., The advantages of co-culture over mono cell culture in simulating in vivo environment, *J. Steroid Biochem. Mol. Biol.* 131 (3–5) (2012) 68–75.
- [30] Y. Ogawa, et al., Androgen receptor expression in breast cancer: relationship with clinicopathological factors and biomarkers, *Int. J. Clin. Oncol.* 13 (5) (2008) 431–435.
- [31] A. Bosch, et al., Triple-negative breast cancer: molecular features, pathogenesis, treatment and current lines of research, *Cancer Treat. Rev.* 36 (3) (2010) 206–215.
- [32] F. Akiyama, H. Iwase, Triple negative breast cancer: clinicopathological characteristics and treatment strategies, *Breast Cancer* 16 (4) (2009) 252–253.
- [33] E. Rodler, L. Korde, J. Gralow, Current treatment options in triple negative breast cancer, *Breast Dis.* 32 (1–2) (2016) 99–122.
- [34] N. Sultana, A.J. Afolayan, A new depsidone and antibacterial activities of compounds from *Usnea undulata* Stirton, *J. Asian Nat. Prod. Res.* 13 (12) (2011) 1158–1164.
- [35] H.Y. Ebrahim, et al., Norstictic acid inhibits breast cancer cell proliferation: migration, invasion, and In vivo invasive growth through targeting C-Met, *Phytother. Res.* 30 (4) (2016) 557–566.
- [36] W. Zhang, et al., Phomopsidone A, a novel depsidone metabolite from the mangrove endophytic fungus *Phomopsis* sp. A123, *Fitoterapia* 96 (2014) 146–151.
- [37] D. Permana, et al., Atrovirisidone B, a new prenylated depsidone with cytotoxic property from the roots of *Garcinia atroviridis*, *Z. Naturforsch. C* 60 (7–8) (2005) 523–526.