



Draft Genome Sequence of *Streptococcus suis* S10, a Virulent Strain Used in Experimental Pig Infections

Rogier A. Gaiser,^{a*} Aldert L. Zomer,^b Jerry M. Wells,^a Peter van Baarlen^a

^aHost-Microbe Interactomics, Wageningen University & Research, Wageningen, The Netherlands

^bVeterinary Medicine, Department of Infectious Diseases and Immunology, Utrecht University, Utrecht, The Netherlands

ABSTRACT Here, we report the draft whole-genome sequence of *Streptococcus suis* strain S10, isolated from the tonsils of a healthy pig. *S. suis* S10 belongs to the highly virulent serotype 2, which includes isolates that cause infectious diseases, including meningitis, in pigs and human. The genome contains a complete prophage that encodes a candidate virulence gene.

Streptococcus suis bacteria are commonly part of the porcine tonsillar microbiota (1), comprising carriage strains and strains that cause infectious disease in pigs and humans (2). Here, we report the genome sequence of *S. suis* strain S10, sampled in 1992 from porcine tonsils, which has been used for experimental infections of pigs (3).

S. suis S10 was cultured without agitation at 37°C with 5% CO₂ in Todd-Hewitt Broth (THB) (Oxoid, UK). Genomic DNA was extracted from 2 ml of exponentially growing bacteria that were lysed in Nuclei Lysis solution (Promega) with proteinase K and protein precipitation solution (Promega) following the manufacturer's protocol. DNA was precipitated in isopropanol and purified using phenol-chloroform-isoamyl alcohol following the protocols described by Barker (4). DNA purity and quality were assessed by gel electrophoresis and spectrometric analysis (ND-1000; NanoDrop Technologies).

DNA library preparation using the Illumina Nextera XT kit following the manufacturer's protocol and genomic DNA sequencing using an Illumina HiSeq 2500 instrument on a paired-end library (125 cycles) were carried out at BaseClear B.V. (Leiden, The Netherlands). FASTQ sequence files were generated using the Illumina CASAVA pipeline 1.8.3. A BaseClear in-house pipeline carried out quality assessment of paired-end reads by Illumina Chastity filtering, removal of reads containing adapters, PhiX control signal, and FASTQC Quality Control Tool 0.10.0. CLC Genomics Workbench 8 trimmed low-quality bases and assembled reads into contigs. KmerGenie (5) determined the optimal k-mer size. SSPACE Premium Scaffold 2.3 (6) linked contigs and placed them into scaffolds. GapFiller 1.10 (7) closed gapped regions within scaffolds. For all *in silico* analyses, default parameters were used.

Sequencing generated 4,268,836 paired-end reads (average length, 126 bases) from which 2,011,091 bases were aligned (266.38-fold coverage). Filtered reads were assembled in 32 gap-closed scaffolds. Quast (8) showed that the GC content was 41.25%, the N_{50} and L_{50} values were 170,355 and 5, and the N_{75} and L_{75} values were 86,491 and 9. The number of uncalled bases per assembled 100 kbp was 0.10.

The S10 genome was annotated via Prokka (9). ANItools Web (10) showed that *S. suis* strain P1/7 (11) was most identical to S10 (99.97%; not shown). CONTIGuator (12) aligned S10 genomic contigs to the P1/7 genome; 23 S10 contigs with a combined base content of 1,997,858 base pairs were mapped to the total of 2,007,491 base pairs of P1/7.

In contig_00009, Phage Search Tool Enhanced Release (PHASTER) (13) predicted a complete prophage containing 64 coding sequences (CDS) spanning a region of

Citation Gaiser RA, Zomer AL, Wells JM, van Baarlen P. 2019. Draft genome sequence of *Streptococcus suis* S10, a virulent strain used in experimental pig infections. *Microbiol Resour Announc* 8:e00227-19. <https://doi.org/10.1128/MRA.00227-19>.

Editor Julie C. Dunning Hotopp, University of Maryland School of Medicine

Copyright © 2019 Gaiser et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Peter van Baarlen, peter.vanbaarlen@wur.nl.

* Present address: Rogier A. Gaiser, Karolinska Institutet, Department of Dental Medicine, Huddinge, Stockholm, Sweden.

Received 3 April 2019

Accepted 9 May 2019

Published 6 June 2019




FIG 1 Phage Search Tool Enhanced Release (PHASTER) analysis output displaying the 64 coding sequences (CDS) of phage origin and their functional annotations; the annotated region shown is part of the largest S10 region that did not align to *S. suis* strain P1/7. Att, attachment site; Hyp, hypothetical protein; Int, integrase; Oth, other protein; PLP, phage-like protein.

62,145 base pairs (Fig. 1). The predicted S10 phage contains a candidate virulence gene with high identity (E value, $7E-164$) with the *S. mitis* (14) and *S. pneumoniae* (15) phage-encoded gene *pblB*, which encodes a platelet adhesin that plays a role during multiple steps in endocardial infection, including direct binding of bacteria to platelets.

The genome sequence of *S. suis* S10 will enable the research community to make targeted gene deletion mutants.

Data availability. The genome sequence of strain *S. suis* S10 has been deposited at the European Molecular Biology Laboratory, European Bioinformatics Institute (EMBL-EBI); the raw reads have been deposited at the European Nucleotide Archive (ENA) under the study identifier [PRJEB30600](https://www.ebi.ac.uk/ena/submit/studies/PRJEB30600).

ACKNOWLEDGMENTS

This work was funded as part of the ITN project TRAIN-ASAP, funded by the Marie Curie Actions under the Seventh Framework Programme for Research and Technological Development of the EU (grant agreement no. 289285). *S. suis* strain S10 was obtained from Hilde Smith, Wageningen Bioveterinary Research, Wageningen University & Research, The Netherlands.

REFERENCES

1. Arends JP, Hartwig N, Rudolph M, Zanen HC. 1984. Carrier rate of *Streptococcus suis* capsular type 2 in palatine tonsils of slaughtered pigs. *J Clin Microbiol* 20:945–947.
2. Goyette-Desjardins G, Auger JP, Xu J, Segura M, Gottschalk M. 2014. *Streptococcus suis*, an important pig pathogen and emerging zoonotic agent—an update on the worldwide distribution based on serotyping and sequence typing. *Emerg Microbes Infect* 3:e45. <https://doi.org/10.1038/emi.2014.45>.
3. Vecht U, Wisselink HJ, van Dijk JE, Smith HE. 1992. Virulence of *Streptococcus suis* type 2 strains in newborn germfree pigs depends on phenotype. *Infect Immun* 60:550–556.
4. Barker K. 2005. *At the bench: a laboratory navigator*, updated edition. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY.
5. Chikhi R, Medvedev P. 2014. Informed and automated k-mer size selection for genome assembly. *Bioinformatics* 30:31–37. <https://doi.org/10.1093/bioinformatics/btt310>.
6. Boetzer M, Henkel CV, Jansen HJ, Butler D, Pirovano W. 2011. Scaffolding pre-assembled contigs using SSPACE. *Bioinformatics* 27:578–579. <https://doi.org/10.1093/bioinformatics/btq683>.
7. Boetzer M, Pirovano W. 2012. Toward almost closed genomes with GapFiller. *Genome Biol* 13:R56. <https://doi.org/10.1186/gb-2012-13-6-r56>.
8. Gurevich A, Saveliev V, Vyahhi N, Tesler G. 2013. QUAST: quality assessment tool for genome assemblies. *Bioinformatics* 29:1072–1075. <https://doi.org/10.1093/bioinformatics/btt086>.
9. Seemann T. 2014. Prokka: rapid prokaryotic genome annotation. *Bioinformatics* 30:2068–2069. <https://doi.org/10.1093/bioinformatics/btu153>.
10. Han N, Qiang Y, Zhang W. 2016. ANItools Web: a Web tool for fast genome comparison within multiple bacterial strains. *Database (Oxford)* 2016:baw084. <https://doi.org/10.1093/database/baw084>.
11. Holden MT, Hauser H, Sanders M, Ngo TH, Cherevach I, Cronin A, Goodhead I, Mungall K, Quail MA, Price C, Rabbinowitsch E, Sharp S, Croucher NJ, Chieu TB, Mai NT, Diep TS, Chinh NT, Kehoe M, Leigh JA, Ward PN, Dowson CG, Whatmore AM, Chanter N, Iversen P, Gottschalk M, Slater JD, Smith HE, Spratt BG, Xu J, Ye C, Bentley S, Barrell BG, Schultsz C, Maskell DJ, Parkhill J. 2009. Rapid evolution of virulence and drug resistance in the emerging zoonotic pathogen *Streptococcus suis*. *PLoS One* 4:e6072. <https://doi.org/10.1371/journal.pone.0006072>.
12. Galardini M, Biondi EG, Bazzicalupo M, Mengoni A. 2011. CONTIGuator: a bacterial genomes finishing tool for structural insights on draft genomes. *Source Code Biol Med* 6:11. <https://doi.org/10.1186/1751-0473-6-11>.
13. Arndt D, Grant JR, Marcu A, Sajed T, Pon A, Liang Y, Wishart DS. 2016. PHASTER: a better, faster version of the PHAST phage search tool. *Nucleic Acids Res* 44:W16–W21. <https://doi.org/10.1093/nar/gkw387>.
14. Bensing BA, Siboo IR, Sullam PM. 2001. Proteins PblA and PblB of *Streptococcus mitis*, which promote binding to human platelets, are encoded within a lysogenic bacteriophage. *Infect Immun* 69:6186–6192. <https://doi.org/10.1128/IAI.69.10.6186-6192.2001>.
15. Tunjungputri RN, Mobegi FM, Cremers AJ, van der Gaast-de Jongh CE, Ferwerda G, Meis JF, Roeleveld N, Bentley SD, Pastura AS, van Hijum SA, van der Ven AJ, de Mast Q, Zomer A, de Jonge MI. 2017. Phage-derived protein induces increased platelet activation and is associated with mortality in patients with invasive pneumococcal disease. *mBio* 8:e01984-16. <https://doi.org/10.1128/mBio.01984-16>.