

REFERENCES

REFERENCES

1. Bossuyt PM, Reitsma JB. The STARD initiative. *Lancet* 2003;361:71.
2. Bossuyt PM, Reitsma JB, Bruns DE *et al*. The STARD statement for reporting studies of diagnostic accuracy: explanation and elaboration. *Clin Chem* 2003;49:7-18.
3. Moons KG, Grobbee DE. Diagnostic studies as multivariable, prediction research. *J Epidemiol Community Health* 2002;56:337-338.
4. Moons KG, van Es GA, Michel BC, Buller HR, Habbema JD, Grobbee DE. Redundancy of single diagnostic test evaluation. *Epidemiology* 1999;10:276-281.
5. Grobbee DE, Miettinen OS. Clinical epidemiology. Introduction to the discipline. *Neth J Med* 1995;47:2-5.
6. Rutten FH, Moons KG, Hoes AW. Improving the quality and clinical relevance of diagnostic studies. *BMJ* 2006;332:1129.
7. Miettinen OS. The modern scientific physician: 4. The useful property of a diagnostic. *CMAJ* 2001;165:910-911.
8. Moons KG, Grobbee DE. When should we remain blind and when should our eyes remain open in diagnostic studies? *J Clin Epidemiol* 2002;55:633-636.
9. Swets JA. Measuring the accuracy of diagnostic systems. *Science* 1988;240:1285-1293.
10. Rutten FH, Moons KG, Cramer MJ *et al*. Recognising heart failure in elderly patients with stable chronic obstructive pulmonary disease in primary care: cross sectional diagnostic study. *BMJ* 2005;331:1379.
11. Weller SC, Mann NC. Assessing rater performance without a "gold standard" using consensus theory. *Med Decis Making* 1997;17:71-79.
12. Beghi E, Tonini C. Surgery for epilepsy: assessing evidence from observational studies. *Epilepsy Res* 2006;70:97-102.
13. Engel J, Jr., Wiebe S, French J *et al*. Practice parameter: Temporal lobe and localized neocortical resections for epilepsy: Report of the Quality Standards Subcommittee of the American Academy of Neurology, in Association with the American Epilepsy Society and the American Association of Neurological Surgeons. *Neurology* 2003;60:538-547.

14. Engel J, Cascino GD, Shields WD. Surgically remediable syndromes. In: Engel J, Pedley TA, eds. *Epilepsy - a comprehensive textbook*. Philadelphia: Lippincott-Raven, 1998:1687.
15. Engel J, Jr. Finally, a randomized, controlled trial of epilepsy surgery. *N Engl J Med* 2001;345:365-367.
16. Lüders HO. Protocols and outcome statistics from epilepsy surgery centers. In: Lüders HO, Comair YG, eds. *Epilepsy Surgery*. Philadelphia: Lippincott Williams & Wilkins, 2001:973.
17. Engel J, Jr. *Dutch Collaborative Epilepsy Surgery Program*. 2 edition. New York: Raven Press, Ltd, 1993.
18. Altman DG. Inter-rater agreement. *Practical statistics for medical research*. London: Chapham and Hall, 1991:403.
19. Landis JR, Koch GG. An application of hierarchical kappa-type statistics in the assessment of majority agreement among multiple observers. *Biometrics* 1977;33:363-374.
20. Ransohoff DF, Feinstein AR. Problems of spectrum and bias in evaluating the efficacy of diagnostic tests. *N Engl J Med* 1978;299:926-930.
21. Lijmer JG, Mol BW, Heisterkamp S *et al*. Empirical evidence of design-related bias in studies of diagnostic tests. *JAMA* 1999;282:1061-1066.
22. Oostenbrink R, Moons KG, Bleeker SE, Moll HA, Grobbee DE. Diagnostic research on routine care data: prospects and problems. *J Clin Epidemiol* 2003;56:501-506.
23. Duncan JS. Imaging and epilepsy. *Brain* 1997;120 (Pt 2):339-377.
24. Sperling MR, O'Connor MJ, Saykin AJ *et al*. A noninvasive protocol for anterior temporal lobectomy. *Neurology* 1992;42:416-422.
25. Duchowny MS, Harvey AS, Sperling MR, Williamson PD. Indications and criteria for surgical intervention. In: Engel J, Jr., Pedley TA, eds. *Epilepsy: A Comprehensive Textbook*. Philadelphia: Lippincott-Raven Publishers, 1997:1677.
26. Rosenow F, Lüders H. Presurgical evaluation of epilepsy. *Brain* 2001;124:1683-1700.
27. Jaeschke R, Guyatt G, Sackett DL. Users' guides to the medical literature. III. How to use an article about a diagnostic test. A. Are the results of the study valid? Evidence-Based Medicine Working Group. *JAMA* 1994;271:389-391.

28. Sackett DL, Haynes RB, Tugwell P. *Clinical epidemiology; a basic science for clinical medicine*. Boston: Little, Brown & Co, 1985.
29. Feinstein AR. *Clinical Epidemiology: the architecture of clinical research*. Philadelphia: WB Saunders Company, 1985.
30. Weijnen CF, Numans ME, de Wit NJ *et al*. Testing for *Helicobacter pylori* in dyspeptic patients suspected of peptic ulcer disease in primary care: cross sectional study. *BMJ* 2001;323:71-75.
31. Oostenbrink R, Moons KG, Donders AR, Grobbee DE, Moll HA. Prediction of bacterial meningitis in children with meningeal signs: reduction of lumbar punctures. *Acta Paediatr* 2001;90:611-617.
32. Reid MC, Lachs MS, Feinstein AR. Use of methodological standards in diagnostic test research. Getting better but still not good. *JAMA* 1995;274:645-651.
33. Knottnerus JA. Application of logistic regression to the analysis of diagnostic data: exact modeling of a probability tree of multiple binary variables. *Med Decis Making* 1992;12:93-108.
34. Begg CB. Biases in the assessment of diagnostic tests. *Stat Med* 1987;6:411-423.
35. Bossuyt PM, Reitsma JB, Bruns DE *et al*. Towards complete and accurate reporting of studies of diagnostic accuracy: the STARD initiative. *BMJ* 2003;326:41-44.
36. DellaBadia Jr J, Bell WL, Keyes Jr JW, Mathews VP, Glazier SS. Assessment and cost comparison of sleep-deprived EEG, MRI and PET in the prediction of surgical treatment for epilepsy. *Seizure* 2002;11:303-309.
37. Kilpatrick C, Cook M, Kaye A, Murphy M, Matkovic Z. Non-invasive investigations successfully select patients for temporal lobe surgery. *J Neurol Neurosurg Psychiatry* 1997;63:327-333.
38. Henkel A, Noachtar S, Pfander M, Lüders HO. The localizing value of the abdominal aura and its evolution: a study in focal epilepsies. *Neurology* 2002;58:271-276.
39. Brekelmans GJ, van Emde BW, Velis DN, Lopes da Silva FH, van Rijen PC, van Veelen CW. Comparison of combined versus subdural or intracerebral electrodes alone in presurgical focus localization. *Epilepsia* 1998;39:1290-1301.
40. O'Brien TJ, So EL, Mullan BP *et al*. Subtraction SPECT co-registered to MRI improves postictal SPECT localization of seizure foci. *Neurology* 1999;52:137-146.

41. Oliveira AJ, da Costa JC, Hilario LN, Anselmi OE, Palmieri A. - Localization of the epileptogenic zone by ictal and interictal SPECT with ^{99m}Tc-ethyl cysteinate dimer in patients with medically refractory epilepsy. *Epilepsia* 1999;40:693-702.
42. Tatum WO, Sperling MR, O'Connor MJ, Jacobstein JG. Interictal single-photon emission computed tomography in partial epilepsy. Accuracy in localization and prediction of outcome. *J Neuroimaging* 1995;5:142-144.
43. Velasco TR, Wichert-Ana L, Leite JP *et al.* Accuracy of ictal SPECT in mesial temporal lobe epilepsy with bilateral interictal spikes. *Neurology* 2002;59:266-271.
44. Lee DS, Lee SK, Kim YK *et al.* The usefulness of repeated ictal SPET for the localization of epileptogenic zones in intractable epilepsy. *Eur J Nucl Med Mol Imaging* 2002;29:607-614.
45. Antel SB, Li LM, Cendes F *et al.* Predicting surgical outcome in temporal lobe epilepsy patients using MRI and MRSI. *Neurology* 2002;58:1505-12.
46. Armon C, Radtke RA, Friedman AH, Dawson DV. Predictors of outcome of epilepsy surgery: multivariate analysis with validation. *Epilepsia* 1996;37:814-21.
47. McIntosh AM, Wilson SJ, Berkovic SF. Seizure outcome after temporal lobectomy: current research practice and findings. *Epilepsia* 2001;42:1288-1307.
48. Berkovic SF, McIntosh AM, Kalnins RM *et al.* Preoperative MRI predicts outcome of temporal lobectomy: an actuarial analysis. *Neurology* 1995;45:1358-63.
49. Spencer SS, Theodore WH, Berkovic SF. Clinical applications: MRI, SPECT, and PET. *Magn Reson Imaging* 1995;13:1119-1124.
50. Cascino GD, Trenerry MR, So EL *et al.* Routine EEG and temporal lobe epilepsy: relation to long-term EEG monitoring, quantitative MRI, and operative outcome. *Epilepsia* 1996;37:651-656.
51. Berg AT, Vickrey BG, Langfitt JT *et al.* The Multicenter Study of Epilepsy Surgery: Recruitment and Selection for Surgery. *Epilepsia* 2003;44:1425-1433.
52. Campos MG, Godoy J, Mesa MT, Torrealba G, Gejman R, Huete I. Temporal lobe epilepsy surgery with limited resources: results and economic considerations. *Epilepsia* 2000;41:18-21.
53. Williamson PD, Jobst BC. Epilepsy surgery in developing countries. *Epilepsia* 2000;41 Suppl 4:S45-S50.

54. Ogden-Epker M, Cullum CM. Quantitative and qualitative interpretation of neuropsychological data in the assessment of temporal lobectomy candidates. *Clin Neuropsychol* 2001;15:183-195.
55. Spencer SS, Berg AT, Vickrey BG *et al.* Initial outcomes in the Multicenter Study of Epilepsy Surgery. *Neurology* 2003;61:1680-1685.
56. Uijl SG, Leijten FS, Parra J, Arends JB, van Huffelen AC, Moons KG. What is the current evidence on decision-making after referral for temporal lobe epilepsy surgery? A review of the literature. *Seizure* 2005;14:534-540.
57. Spencer SS. When should temporal-lobe epilepsy be treated surgically? *Lancet Neurol* 2002;1:375-382.
58. Janszky J, Schulz R, Ebner A. Clinical features and surgical outcome of medial temporal lobe epilepsy with a history of complex febrile convulsions. *Epilepsy Res* 2003;55:1-8.
59. Lüders H, Acharya J, Baumgartner C *et al.* Semiological seizure classification. *Epilepsia* 1998;39:1006-1013.
60. Marks WJ, Jr., Laxer KD. Semiology of temporal lobe seizures: value in lateralizing the seizure focus. *Epilepsia* 1998;39:721-726.
61. Kotagal P. Automotor seizures. In: Lüders HO, Noachtar S, eds. *Epileptic Seizures; Pathophysiology and Clinical Semiology*. Philadelphia: Churchill Livingstone, 2000:449.
62. Meiners LC, van Gils AD, De Kort G, van der Graaf Y, Jansen GH, van Veelen CW. Fast fluid-attenuated inversion recovery (FLAIR) compared with T2-weighted spin-echo in the magnetic resonance diagnosis of mesial temporal sclerosis. *Invest Radiol* 1999;34:134-142.
63. Risinger MW, Engel J, Jr., Van Ness PC, Henry TR, Crandall PH. Ictal localization of temporal lobe seizures with scalp/sphenoidal recordings. *Neurology* 1989;39:1288-1293.
64. DeLong ER, DeLong DM, Clarke-Pearson DL. Comparing the areas under two or more correlated receiver operating characteristic curves: a nonparametric approach. *Biometrics* 1988;44:837-845.
65. Little RA. Regression with missing X's: a review. *J Am Stat Assoc* 1992;87:1227-1237.
66. Greenland S, Finkle WD. A critical look at methods for handling missing covariates in epidemiologic regression analyses. *Am J Epidemiol* 1995;142:1255-1264.

67. Harrell FE, Jr., Lee KL, Mark DB. Multivariable prognostic models: issues in developing models, evaluating assumptions and adequacy, and measuring and reducing errors. *Stat Med* 1996;15:361-387.
68. So EL. Value and limitations of seizure semiology in localizing seizure onset. *J Clin Neurophysiol* 2006;23:353-357.
69. Serles W, Caramanos Z, Lindinger G, Pataraja E, Baumgartner C. Combining ictal surface-electroencephalography and seizure semiology improves patient lateralization in temporal lobe epilepsy. *Epilepsia* 2000;41:1567-1573.
70. Deacon C, Wiebe S, Blume WT, McLachlan RS, Young GB, Matijevic S. Seizure identification by clinical description in temporal lobe epilepsy: how accurate are we? *Neurology* 2003;61:1686-1689.
71. Diehl B, Lüders HO. Temporal lobe epilepsy: when are invasive recordings needed? *Epilepsia* 2000;41 Suppl 3:S61-S74.
72. Alpherts WC, Vermeulen J, Hendriks MP *et al.* Long-term effects of temporal lobectomy on intelligence. *Neurology* 2004;62:607-611.
73. Leijten FS, Alpherts WC, van Huffelen AC, Vermeulen J, van Rijen PC. The effects on cognitive performance of tailored resection in surgery for nonlesional mesiotemporal lobe epilepsy. *Epilepsia* 2005;46:431-439.
74. Dodrill CB. Overview: presurgical neuropsychological evaluation. In: Lüders HO, Comair YG, eds. *Epilepsy Surgery*. Philadelphia: Lippincott Williams & Wilkins, 2001:475.
75. Debets RM, Sadzot B, van Isselt JW *et al.* Is 11C-flumazenil PET superior to 18FDG PET and 123I-iomazenil SPECT in presurgical evaluation of temporal lobe epilepsy? *J Neurol Neurosurg Psychiatry* 1997;62:141-150.
76. Debets RM, van Veelen CW, Maquet P *et al.* Quantitative analysis of 18/FDG-PET in the presurgical evaluation of patients suffering from refractory partial epilepsy. Comparison with CT, MRI, and combined subdural and depth. EEG. *Acta Neurochir Suppl (Wien)* 1990;50:88-94.
77. Juhasz C, Chugani HT. Imaging the epileptic brain with positron emission tomography. *Neuroimaging Clin N Am* 2003;13:705-16, viii.
78. Carne RP, O'Brien TJ, Kilpatrick CJ *et al.* MRI-negative PET-positive temporal lobe epilepsy: a distinct surgically remediable syndrome. *Brain* 2004;127:2276-2285.

79. Henry TR. Positron Emission Tomography in Epilepsy Surgery Evaluation. In: Lüders HO, Comair YG, eds. *Epilepsy Surgery*. Philadelphia: Lippincott Williams & Wilkins, 2001:257.
80. O'Brien TJ, Hicks RJ, Ware R, Binns DS, Murphy M, Cook MJ. The utility of a 3-dimensional, large-field-of-view, sodium iodide crystal--based PET scanner in the presurgical evaluation of partial epilepsy. *J Nucl Med* 2001;42:1158-1165.
81. Baron JC, Bousser MG, Comar D, Soussaline F, Castaigne P. Noninvasive tomographic study of cerebral blood flow and oxygen metabolism in vivo. Potentials, limitations, and clinical applications in cerebral ischemic disorders. *Eur Neurol* 1981;20:273-284.
82. Pantano P, Baron JC, Samson Y, Bousser MG, Derouesne C, Comar D. Crossed cerebellar diaschisis. Further studies. *Brain* 1986;109 (Pt 4):677-694.
83. Donders AR, van der Heijden GJ, Stijnen T, Moons KG. Review: a gentle introduction to imputation of missing values. *J Clin Epidemiol* 2006;59:1087-1091.
84. Ollenberger GP, Byrne AJ, Berlangieri SU *et al*. Assessment of the role of FDG PET in the diagnosis and management of children with refractory epilepsy. *Eur J Nucl Med Mol Imaging* 2005;32:1311-1316.
85. Ryvlin P, Bouvard S, Le Bars D *et al*. Clinical utility of flumazenil-PET versus [18F] fluorodeoxyglucose-PET and MRI in refractory partial epilepsy. A prospective study in 100 patients. *Brain* 1998;121 (Pt 11):2067-2081.
86. Willinsky RA, Taylor SM, TerBrugge K, Farb RI, Tomlinson G, Montanera W. Neurologic complications of cerebral angiography: prospective analysis of 2,899 procedures and review of the literature. *Radiology* 2003;227:522-528.
87. Loddenkemper T, Morris HH, III, Perl J. Carotid artery dissection after the intracarotid amobarbital test. *Neurology* 2002;59:1797-1798.
88. Macken, M. P. and Morris, H. H. Complications of the Wada test. A review of the Cleveland Clinic experience. *Epilepsia* 40[suppl 7], 84. 1999.
Ref Type: Abstract
89. Martin RC, Grote CL. Does the Wada test predict memory decline following epilepsy surgery? *Epilepsy & Behavior* 2002;3:4-15.
90. Baxendale S, Thompson P, Duncan J, Richardson M. Is it time to replace the Wada test? *Neurology* 2003;60:354-355.

91. Kirsch HE, Walker JA, Winstanley FS *et al.* Limitations of Wada memory asymmetry as a predictor of outcomes after temporal lobectomy. *Neurology* 2005;65:676-680.
92. Baxendale S, Thompson P, Harkness W, DUNCAN J. Predicting memory decline following epilepsy surgery: a multivariate approach. *Epilepsia* 2006;47:1887-1894.
93. Lineweaver TT, Morris HH, Naugle RI, Najm IM, Diehl B, Bingaman W. Evaluating the contributions of state-of-the-art assessment techniques to predicting memory outcome after unilateral anterior temporal lobectomy. *Epilepsia* 2006;47:1895-1903.
94. Alpherts WC, Vermeulen J, van Veelen CW. The wada test: prediction of focus lateralization by asymmetric and symmetric recall. *Epilepsy Res* 2000;39:239-249.
95. De Renzi E, Vignolo LA. The token test: A sensitive test to detect receptive disturbances in aphasics. *Brain* 1962;85:665-678.
96. Leijten FS, Alpherts WC, van Huffelen AC, Vermeulen J, van Rijen PC. The effects on cognitive performance of tailored resection in surgery for nonlesional mesiotemporal lobe epilepsy. *Epilepsia* 2005;46:431-439.
97. Cohen J. The factorial structure of the WAIS between early adulthood and old age. *J Consult Psychol* 1957;21:283-290.
98. Engel J, Jr., Van Ness PC, Rasmussen TB. Outcome with respect to epileptic seizures. In: Engel J, Jr., ed. *Surgical Treatment of the Epilepsies*. New York: Raven Press Ltd, 1993:609.
99. Loring DW, Meador KJ, Lee GP, Smith JR. Structural versus functional prediction of memory change following anterior temporal lobectomy. *Epilepsy Behav* 2004;5:264-268.
100. Wada JA. Youthful season revisited. *Brain Cogn* 1997;33:7-10.
101. Dinner DS. Use of the intracarotid amobarbital procedure in the lateralization of the epileptogenic zone. In: Lüders HO, Comair YG, eds. *Epilepsy Surgery*. Philadelphia: Lippincott Williams & Wilkins, 2001:537.
102. Sabsevitz DS, Swanson SJ, Morris GL, Mueller WM, Seidenberg M. Memory outcome after left anterior temporal lobectomy in patients with expected and reversed Wada memory asymmetry scores. *Epilepsia* 2001;42:1408-1415.
103. Stroup E, Langfitt J, Berg M, McDermott M, Pilcher W, Como P. Predicting verbal memory decline following anterior temporal lobectomy (ATL). *Neurology* 2003;60:1266-1273.

104. Loring DW, Meador KJ, Lee GP *et al.* Wada memory asymmetries predict verbal memory decline after anterior temporal lobectomy. *Neurology* 1995;45:1329-1333.
105. Spencer DC, Morrell MJ, Risinger MW. The role of the intracarotid amobarbital procedure in evaluation of patients for epilepsy surgery. *Epilepsia* 2000;41:320-325.
106. Lee GP, Westerveld M, Blackburn LB, Park YD, Loring DW. Prediction of verbal memory decline after epilepsy surgery in children: effectiveness of Wada memory asymmetries. *Epilepsia* 2005;46:97-103.
107. Lacruz ME, Alarcon G, Akanuma N *et al.* Neuropsychological effects associated with temporal lobectomy and amygdalohippocampectomy depending on Wada test failure. *J Neurol Neurosurg Psychiatry* 2004;75:600-607.
108. Chelune GJ. Hippocampal adequacy versus functional reserve: predicting memory functions following temporal lobectomy. *Arch Clin Neuropsychol* 1995;10:413-432.
109. Loring DW, Meador KJ. Wada testing to define hippocampal function. In: Lüders HO, Comair YG, eds. *Epilepsy Surgery*. Philadelphia: Lippincott Williams & Wilkins, 2001:531.
110. Grote CL, Wierenga C, Smith MC *et al.* Wada difference a day makes: interpretive cautions regarding same-day injections. *Neurology* 1999;52:1577-1582.
111. Loddenkemper T, Morris HH, Lineweaver TT, Kellinghaus C. Repeated intracarotid amobarbital tests. *Epilepsia* 2007;48:553-558.
112. Wellmer J, Fernandez G, Linke DB, Urbach H, Elger CE, Kurthen M. Unilateral intracarotid amobarbital procedure for language lateralization. *Epilepsia* 2005;46:1764-1772.
113. Schulze-Bonhage A, Quiske A. Validity of language lateralisation by unilateral intracarotid Wada test. *J Neurol Neurosurg Psychiatry* 2004;75:1367.
114. Jeha LE, Najm IM, Bingaman WE *et al.* Predictors of outcome after temporal lobectomy for the treatment of intractable epilepsy. *Neurology* 2006;66:1938-1940.
115. Kelemen A, Barsi P, Eross L *et al.* Long-term outcome after temporal lobe surgery--prediction of late worsening of seizure control. *Seizure* 2006;15:49-55.
116. Yun CH, Lee SK, Lee SY, Kim KK, Jeong SW, Chung CK. Prognostic factors in neocortical epilepsy surgery: multivariate analysis. *Epilepsia* 2006;47:574-579.
117. Janszky J, Pannek HW, Fogarasi A *et al.* Prognostic factors for surgery of neocortical temporal lobe epilepsy. *Seizure* 2006.

118. Janszky J, Janszky I, Schulz R *et al.* Temporal lobe epilepsy with hippocampal sclerosis: predictors for long-term surgical outcome. *Brain* 2005.
119. Jeong SW, Lee SK, Hong KS, Kim KK, Chung CK, Kim H. Prognostic factors for the surgery for mesial temporal lobe epilepsy: longitudinal analysis. *Epilepsia* 2005;46:1273-1279.
120. Spencer SS, Berg AT, Vickrey BG *et al.* Predicting long-term seizure outcome after resective epilepsy surgery: the multicenter study. *Neurology* 2005;65:912-918.
121. Clusmann H, Kral T, Fackeldey E *et al.* Lesional mesial temporal lobe epilepsy and limited resections: prognostic factors and outcome. *J Neurol Neurosurg Psychiatry* 2004;75:1589-1596.
122. Clusmann H, Schramm J, Kral T *et al.* Prognostic factors and outcome after different types of resection for temporal lobe epilepsy. *J Neurosurg* 2002;97:1131-1141.
123. Hennessy MJ, Elwes RD, Rabe-Hesketh S, Binnie CD, Polkey CE. Prognostic factors in the surgical treatment of medically intractable epilepsy associated with mesial temporal sclerosis. *Acta Neurol Scand* 2001;103:344-350.
124. Hennessy MJ, Elwes RD, Honavar M, Rabe-Hesketh S, Binnie CD, Polkey CE. Predictors of outcome and pathological considerations in the surgical treatment of intractable epilepsy associated with temporal lobe lesions. *J Neurol Neurosurg Psychiatry* 2001;70:450-458.
125. Jeong SW, Lee SK, Kim KK, Kim H, Kim JY, Chung CK. Prognostic factors in anterior temporal lobe resections for mesial temporal lobe epilepsy: multivariate analysis. *Epilepsia* 1999;40:1735-1739.
126. Berg AT, Walczak T, Hirsch LJ, Spencer SS. Multivariable prediction of seizure outcome one year after resective epilepsy surgery: development of a model with independent validation. *Epilepsy Res* 1998;29:185-194.
127. Radhakrishnan K, So EL, Silbert PL *et al.* Predictors of outcome of anterior temporal lobectomy for intractable epilepsy: a multivariate study. *Neurology* 1998;51:465-471.
128. Cohen-Gadol AA, Wilhelmi BG, Collignon F *et al.* Long-term outcome of epilepsy surgery among 399 patients with nonlesional seizure foci including mesial temporal lobe sclerosis. *J Neurosurg* 2006;104:513-524.
129. Tonini C, Beghi E, Berg AT *et al.* Predictors of epilepsy surgery outcome: a meta-analysis. *Epilepsy Res* 2004;62:75-87.

130. Uijl SG, Leijten FS, Arends JB, Parra J, van Huffelen AC, Moons KG. The added value of [18F]-Fluoro-D-deoxyglucose Positron Emission Tomography in screening for temporal lobe epilepsy surgery. *Epilepsia* 2007.
131. Steyerberg EW, Eijkemans MJ, Harrell FE, Jr., Habbema JD. Prognostic modelling with logistic regression analysis: a comparison of selection and estimation methods in small data sets. *Stat Med* 2000;19:1059-1079.
132. Moons KG, Bots ML, Salonen JT *et al.* Prediction of stroke in the general population in Europe (EUROSTROKE): Is there a role for fibrinogen and electrocardiography? *J Epidemiol Community Health* 2002;56 Suppl 1:i30-i36.
133. Sperling MR, O'Connor MJ, Saykin AJ, Plummer C. Temporal lobectomy for refractory epilepsy. *JAMA* 1996;276:470-475.
134. van Donselaar CA, Carpay JA. [Epilepsy; Guidelines for diagnosis and treatment: Dutch Neurological Society (NVN)]. 2 edition. Utrecht: Roto Smeets Grafiservices, 2006.
135. Karceski S, Morrell MJ, Carpenter D. Treatment of epilepsy in adults: expert opinion, 2005. *Epilepsy Behav* 2005;7 Suppl 1:S1-64.
136. Lhatoo SD, Solomon JK, McEvoy AW, Kitchen ND, Shorvon SD, Sander JW. A prospective study of the requirement for and the provision of epilepsy surgery in the United Kingdom. *Epilepsia* 2003;44:673-676.
137. Uijl SG, Leijten FS, Arends J, Parra J, van Huffelen AC, Moons KG. Decision-making in temporal lobe epilepsy surgery: The contribution of basic non-invasive tests. *submitted* 2007.
138. Uijl SG, Leijten FS, Arends JB *et al.* The intracarotid amobarbital or Wada test: unilateral or bilateral? *submitted* 2007.
139. Drane DL, Goldstein J, Gilliam F. When is the Wada test necessary for temporal lobectomy. In: Miller JW, Silbergeld DL, eds. *Epilepsy Surgery. Principles and Controversies*. New York: Taylor & Francis Group, 2006:239.
140. Ojemann JG, Binder JR, Trenerry MR. Will fMRI replace the Wada test? In: Miller JW, Silbergeld DL, eds. *Epilepsy Surgery. Principles and Controversies*. New York: Taylor & Francis Group, 2006:336.
141. Uijl SG, Leijten FS, Arends J, Parra J, van Huffelen AC, Moons KG. Prediction of seizure freedom after temporal lobe epilepsy surgery: The value of combining predictors. *submitted* 2007.

142. Vickrey BG, Hays RD, Engel J, Jr. *et al.* Outcome assessment for epilepsy surgery: the impact of measuring health-related quality of life. *Ann Neurol* 1995;37:158-166.
143. Spencer SS, Berg AT, Vickrey BG *et al.* Health-related quality of life over time since resective epilepsy surgery. *Ann Neurol* 2007.
144. Langfitt JT, Westerveld M, Hamberger MJ *et al.* Worsening of quality of life after epilepsy surgery: Effect of seizures and memory decline. *Neurology* 2007;68:1988-1994.
145. Tellez-Zenteno JF, Dhar R, Hernandez-Ronquillo L, Wiebe S. Long-term outcomes in epilepsy surgery: antiepileptic drugs, mortality, cognitive and psychosocial aspects. *Brain* 2006.
146. Uijl SG, Moons KG, Leijten FS, Veltman PH, Budde A, van Donselaar CA. How underutilized is epilepsy surgery? *submitted* 2007.
147. Schiller Y, Cascino GD, Sharbrough FW. Chronic intracranial EEG monitoring for localizing the epileptogenic zone: an electroclinical correlation. *Epilepsia* 1998;39:1302-1308.
148. Isnard J. [Drug-resistant partial epilepsy. Invasive electrophysiological explorations]. *Rev Neurol (Paris)* 2004;160 Spec No 1:5S138-5S143.

