

# References

## References

---

1. **Abbas, A.K., K.M. Murphy, and A. Sher, (1996).** Functional diversity of helper T lymphocytes. *Nature*, 383: p. 787-793.
2. **Agneessens, J., et al., (1997).** Epidemiological observations on gastrointestinal nematode infections in grazing cow-calf pairs in Belgium. *Vet Parasitol*, 69: p. 65-75.
3. **Albers, G., (1981).** Genetic resistance to experimental *Cooperia oncophora* infections in calves. *PhD thesis. Agricultural university, Wageningen*.
4. **Almeria, S., et al., (1997).** Isolation and phenotypic characterization of abomasal mucosal lymphocytes in the course of a primary *Ostertagia ostertagi* infection in calves. *Vet Immunol Immunopathol*, 57: p. 87-98.
5. **Almeria, S., et al., (1997).** Quantification of cytokine gene expression in lamina propria lymphocytes of cattle following infection with *Ostertagia ostertagi*. *J Parasitol*, 83: p. 1051-1055.
6. **Anderson, M.C. and J.F. Michel, (1977).** Density-dependent survival in populations of *Ostertagia ostertagi*. *Int J Parasitol*, 7: p. 321-329.
7. **Anderson, R.M. and R.M. May, (1985).** Herd immunity to helminth infection and implications for parasite control. *Nature*, 315: p. 493-496.
8. **Armour, J., (1989).** The influence of host immunity on the epidemiology of trichostrongyle infections in cattle. *Vet Parasitol*, 32: p. 5-19.
9. **Armour, J., et al., (1987).** Pathophysiological and parasitological studies on *Cooperia oncophora* infections in calves. *Res Vet Sci*, 42: p. 373-381.
10. **Axelsen, J., et al., (1986).** Grazing management and nematode parasite control in cattle in the temperate climatic zone of Australia. *Aust J Exp Agric*, 26: p. 267-273.
11. **Baker, D.G. and L.J. Gershwin, (1992).** Seasonal patterns of total and *Ostertagia*-specific IgE in grazing cattle. *Vet Parasitol*, 44: p. 211-221.
12. **Baldwin, C.L., I.W. Morrison, and J. Naessens, (1988).** Differentiation antigens and functional characteristics of bovine leucocytes. In Z. Trnka and M. Miyasaka (Eds.) *Comparative aspects of Differentiation antigens in lympho-haematopoietic tissues.*, Marcel Dekker Inc., New York, N. Y.: p. 455-465.
13. **Baldwin, C.L., et al., (2002).** Activation of bovine peripheral blood gd T cells for cell division and IFN-g production. *Vet Immunol Immunopathol*, 87: p. 251-259.
14. **Balic, A., V.M. Bowles, and E.N. Meeusen, (2000).** Cellular profiles in the abomasal mucosa and lymph node during primary infection with *Haemonchus contortus* in sheep. *Vet Immunol Immunopathol*, 75: p. 109-120.
15. **Balic, A., V.M. Bowles, and E.N. Meeusen, (2000).** The immunobiology of gastrointestinal nematode infections in ruminants. *Adv Parasitol*, 45: p. 181-241.
16. **Balic, A., V.M. Bowles, and E.N. Meeusen, (2002).** Mechanisms of immunity to *Haemonchus contortus* infection in sheep. *Parasite Immunol*, 24: p. 39-46.
17. **Banchereau, J. and R.M. Steinman, (1998).** Dendritic cells and the control of immunity. *Nature*, 392: p. 245-252.
18. **Bao, S., et al., (1996).** Interleukin-5 mRNA expressed by eosinophils and gamma/delta T cells in parasite-immune sheep. *Eur J Immunol*, 26: p. 552-556.
19. **Barger, I.A., et al., (1985).** Regulation of *Haemonchus contortus* populations in sheep exposed to continuous infection. *Int J Parasitol*, 15: p. 529-533.

20. Barnes, E.H. and R.J. Dobson, (1990). Population dynamics of *Trichostrongylus colubriformis* in sheep: model to predict the worm population over time as a function of infection rate and host age. *Int J Parasitol*, 20: p. 365-373.
21. Barnes, E.H., R.J. Dobson, and I.A. Barnes, (1995). Worm control and anthelmintic resistance: adventures with a model. *Parasitology Today*, 11: p. 56-63.
22. Barth, E.E., W.F. Jarrett, and G.M. Urquhart, (1966). Studies on the mechanism of self-cure reaction in rats injected with *Nippostrongylus brasiliensis*. *Immunology*, 10: p. 459-464.
23. Baxter, E.H. and G.G. Rousseau, (1979). Glucocorticoid hormone action: an overview. In *J.D. Baxter and G.G. Rousseau (Eds.). Glucocorticoid Hormone Action.*, Springer-Verlag, Heidelberg: p. 1-24.
24. Behm, C.A. and K.S. Ovington, (2000). The role of eosinophils in Parasitic Helminth infections: Insights from genetically Modified Mice. *Parasitology Today*, 16: p. 202-209.
25. Ben Smith, A., et al., (1999). The relationship between circulating and intestinal *Heligmosomoides polygyrus* specific IgG1 and IgA and resistance to primary infections. *Parasite Immunol*, 21: p. 587-593.
26. Bird, A.F. and J. Bird, (1991). The structure of nematodes. 2nd Ed, Academic Press limited, London.
27. Bischof, R.J. and E.N. Meeusen, (2002). Cellular kinetics of an allergic-type response in a sheep mammary gland model of inflammation. *Clin Exp Allergy*, 32: p. 619-626.
28. Blackwell, N.M. and K.J. Else, (2001). B cells and antibodies are required for resistance to the parasitic gastrointestinal nematode *Trichuris muris*. *Infect Immun*, 69: p. 3860-3868.
29. Bland, P.W. and L.G. Warren, (1986). Antigen presentation by epithelial cells of the rat small intestine. II. Selective induction of suppressor T cells. *Immunology*, 58: p. 9-14.
30. Boonstra, A., et al., (2003). Flexibility of mouse classical and plasmacytoid-derived dendritic cells in directing T helper 1 and 2 cells development: dependency on antigen dose and differential toll-like receptor ligation. *Exp Med*, 197: p. 101-109.
31. Borgsteede, F.H., (1977). The epidemiology of gastro-intestinal helminth infections in young cattle in the Netherlands. *PhD thesis, Utrecht, The Netherlands*.
32. Borgsteede, F.H. and J. Hendriks, (1979). Experimental infections with *Cooperia oncophora* (Railliet, 1918) in calves. Results of single infections with two graded dose levels of larvae. *Parasitology*, 78: p. 331-342.
33. Borgsteede, F.H., et al., (2000). Nematode parasies of adult dairy cattle in the Netherlands. *Vet Parasitol*, 89: p. 287-296.
34. Borgsteede, F.H. and W.P. van der Burg, (1982). Worm burdens in cows. II. An analysis of the population of nematodes in the abomasum of adult dairy cows. *Vet Parasitol*, 10: p. 323-330.
35. Brown, W.C., et al., (1994). CD4+ T-cell clones obtained from cattle chronically infected with *Fasciola hepatica* and specific for adult worm antigen express both unrestricted and Th2 cytokine profiles. *Infect Immun*, 62: p. 818-827.
36. Brown, W.C., A.C. Rice-Ficht, and D.M. Estes, (1998). Bovine type 1 and type 2 responses. *Vet Immunol Immunopathol*, 63: p. 45-55.
37. Brown, W.C., et al., (1993). Heterogeneity in cytokine profiles of *Babesia bovis*-specific bovine CD4+ T cells clones activated in vitro. *Infect Immun*, 61: p. 3273-3281.
38. Buddle, B.M., et al., (1992). Association of blood eosinophilia with the expression of resistance in Romney lambs to nematodes. *Int J Parasitol*, 22: p. 955-960.

## References

---

39. **Burrows, R.O., P.J. Best, and J.M. Preston, (1980).** Tryichostrongylid egg output of dairy cows. *Vet Rec*, 107: p. 399-401.
40. **Canals, A. and L.C. Gasbarre, (1990).** *Ostertagia ostertagi*: isolation and partial characterization of somatic and metabolic antigens. *Int J Parasitol*, 20: p. 1047-1054.
41. **Canals, A., et al., (1997).** Cytokine profile induced by a primary infection with *Ostertagia ostertagi* in cattle. *Vet Immunol Immunopathol*, 58: p. 63-75.
42. **Cassel, D.J. and R. H. Schwartz, (1994).** A quantitative analysis of antigen presenting cell function: activated B cells stimulate naive CD4+ cells but are inferior to dendritic cells in providing costimulation. *J Exp Med*, 180: p. 1829-1840.
43. **Cella, M., F. Sallusto, and A. Lanzavecchia, (1997).** Origin, maturation and antigen presenting function of dendritic cells. *Curr Opin Immunol*, 9: p. 10-16.
44. **Charleston, W.A.G., (1965).** Pathogenesis of experimental haemonchosis in sheep with special reference to the development of resistance. *J Comp Pathol*, 75: p. 55-67.
45. **Chiodini, R.J. and W.C. Davis, (1992).** The cellular immunology of bovine paratuberculosis: the predominant response is mediated by cytotoxic gamma/delta T cells which prevent CD4+ activity. *Microb Pathogenesis*, 13: p. 447-463.
46. **Claerebout, E., et al., (1999).** Larval migration inhibition activity in abomasal mucus and serum from calves infected with *Ostertagia ostertagi*. *Res Vet Sci*, 66: p. 253-257.
47. **Coop, R., A. Sykes, and K. Angus, (1979).** The pathogenicity of daily intakes of *Cooperia oncophora* in growing calves. *Vet Parasitol*, 5: p. 261-269.
48. **Coop, R.L. and I. Kyriazakis, (1999).** Nutrition-parasite interaction. *Vet Parasitol*, 84: p. 187-204.
49. **Corry, D.B. and F. Kheradmand, (1999).** Induction and regulation of the IgE response. *Nature*, 402: p. 18-23.
50. **Cox, D.D. and A.C. Todd, (1962).** Survey of gastointestinal parasitism in Wisconsin dairy cattle. *J Am Vet Med Assoc*, 141: p. 706-709.
51. **Crofton, H. and J. Withlock, (1968).** Changes in sex ratio in *Haemonchus contortus cayugensis*. *Cornell Veterinarian*, 58: p. 388-392.
52. **Crowe, S.E. and M.H. Perdue, (1992).** Gastrointestinal tract hypersensitivity: Basic mechanisms of pathophysiology. *Gastroenterology*, 103: p. 1075-1095.
53. **Dainichi, T., et al., (2001).** Nippocystatin, a cysteine protease inhibitor from *Nippostrongylus brasiliensis*, inhibits antigen processing and modulates antigen-specific immune response. *Infect Immun*, 69: p. 7380-7386.
54. **Dawkins, H.J., R.G. Windon, and G.K. Eagleson, (1989).** Eosinophil responses in sheep selected for high and low responsiveness to *Trichostrongylus colubriformis*. *Int J Parasitol*, 19: p. 199-205.
55. **De Graaf, D., et al., (1992).** Antigenic differences between the life cycle stages of *Cooperia oncophora*. *Res Vet Sci*, 53: p. 390-392.
56. **De Marez, T., et al., (1997).** Induction and suppression of lymphocyte proliferation by antigen extracts of *Ostertagia ostertagi*. *Vet Imm Immunopath*, 57: p. 69-77.
57. **Dewhirst, L.W. and M.F. Hansen, (1961).** Methods to differentiate and estimate worm burdens in cattle. *Vet Med*, 56: p. 84-89.
58. **Dorny, P., et al., (1997).** The influence of a *Cooperia oncophora* priming on a concurrent challenge with *Ostertagia ostertagi* and *C. oncophora* in calves. *Vet Parasitol*, 70: p. 143-151.

59. **Douch, P.G., et al., (1983).** In vitro bioassay of sheep gastrointestinal mucus for nematode paralysing activity mediated by substances with some properties characteristic of SRS-A. *Int J Parasitol*, 13: p. 207-212.
60. **Dubucquoi, S., et al., (1994).** Interleukin 5 synthesis by eosinophils: association with granules and immunoglobulin-dependent secretion. *J Exp Med*, 179: p. 703-708.
61. **Ehigiator, H.N., A.W. Stadnyk, and T.D. Lee, (2000).** Extract of *Nippostrongylus brasiliensis* stimulates polyclonal type-2 immunoglobulin response by inducing de novo class switch. *Infect Immun*, 68: p. 4913-4922.
62. **Ekkens, M.J., et al., (2002).** Memory Th2 effector cells can develop in the absence of B7-1/B7-2 interactions and effector Th cells after priming with an intestinal nematode parasite. *J Immunol*, 168: p. 6344-6351.
63. **Else, K.J. and F.D. Finkelman, (1998).** Intestinal nematode parasites, cytokines and effector mechanisms. *Int J Parasitol*, 28: p. 1145-1158.
64. **Else, K.J., et al., (1994).** Cytokine mediated regulation of chronic intestinal helminth infection. *J Exp Med*, 179: p. 347-351.
65. **Else, K.J. and R.K. Grencis, (1991).** Cellular immune responses to the murine nematode parasite *Trichuris muris*. I. Differential cytokine production during acute or chronic infection. *Immunology*, 72: p. 508-513.
66. **Else, K.J. and R.K. Grencis, (1996).** Antibody-independent effector mechanisms in resistance to the intestinal nematode parasite *Trichuris muris*. *Infect Immun*, 64: p. 2950-2954.
67. **Else, K.J., L. Hultner, and R.K. Grencis, (1992).** Modulation of cytokine production and response phenotypes in murine trichuriasis. *Parasite Immunol*, 14: p. 441-449.
68. **Else, K.J. and D. Wakelin, (1988).** The effects of H-2 and non H-2 genes on the expulsion of the nematode *Trichuris muris* from inbred and congenic mice. *Parasitology*, 96: p. 543-550.
69. **Emery, D.L., (1996).** Vaccination against worm parasites of animals. *Vet Parasitol*, 64: p. 31-45.
70. **Emery, D.L., et al., (1999).** Induction of protective immunity to *Trichostrongylus colubriformis* in neonatal Merino lambs. *Int J Parasitol*, 29: p. 1037-1046.
71. **Emery, D.L., et al., (1992).** Studies of stage-specific immunity against *Trichostrongylus colubriformis* in sheep: immunization by normal and truncated infections. *Int J Parasitol*, 22: p. 215-220.
72. **Emery, D.L., B.M. Wagland, and S.J. McClure, (1993).** Rejection of heterologous nematodes by sheep immunized with larval or adult *Trichostrongylus colubriformis*. *Int J Parasitol*, 23: p. 841-846.
73. **Estes, D.M., N.M. Closser, and G.K. Allen, (1994).** IFN-gamma stimulates IgG2 production from bovine B cells costimulated with anti-mu and mitogen. *Cell Immunol*, 154: p. 287-295.
74. **Eysker, M. and F.N. Kooyman, (1993).** Notes on necropsy and herbage processing techniques for gastrointestinal nematodes of ruminants. *Vet Parasitol*, 46: p. 205-213.
75. **Eysker, M. and H.W. Ploeger, (2000).** Value of present diagnostic methods for gastrointestinal nematode infections in ruminants. *Parasitology*, 120: p. S109-S119.
76. **Eysker M., et al., (1998).** The efficacy of Michel's dose and move system on gastrointestinal nematode infections in dairy calves. *Vet Parasitol*, 75: p. 99-114.
77. **Eysker M., et al., (1998).** The effect of repeated moves to clean pasture on the build up of gastrointestinal nematodes in calves. *Vet Parasitol*, 76: p. 81-94.

## References

---

78. **Ferlin, W.G., et al., (1996).** CD40 signaling induces interleukin-4 independent IgE switching in vivo. *Eur J Immunol*, 26: p. 2911-2915.
79. **Ferrick, D.A., et al., (1995).** Differential production of interferon-gamma and interleukin-4 in response to Th1- and Th2-stimulating pathogens by gamma delta T cells in vivo. *Nature*, 373: p. 255-257.
80. **Fiel, C.A., et al., (2001).** Resistance of *Cooperia* to ivermectin treatments in grazing cattle of the Humid Pampa, Argentina. *Vet Parasitol*, 97: p. 211-217.
81. **Finkelman, F.D., et al., (1991).** Regulation and biological function of helminth-induced cytokine responses. *Immunology Today*, 12: p. 62-67.
82. **Finkelman, F.D., et al., (1997).** Cytokine regulation of host defense against parasitic gastrointestinal nematodes: lessons from studies with rodent models. *Annu Rev Immunol*, 15: p. 505-533.
83. **Finkelman, F.D., et al., (1997).** Cytokine regulation of host defense against parasitic gastrointestinal nematodes. *Annu Rev Immunol*, 15: p. 505-533.
84. **Frankena, K., (1987).** The interaction between *Cooperia* spp. and *Ostertagia* spp. (nematoda: trichostrongylidae) in cattle. *PhD thesis. Agricultural University, Wageningen*.
85. **Galli, S.J., K.M. Zsebo, and E.N. Geissler, (1994).** The kit ligand, stem cell factor. *Adv Immunol*, 55: p. 1-96.
86. **Garside, P., et al., (2000).** Immunopathology of intestinal helminth infection. *Parasite Immunol*, 22: p. 605-612.
87. **Gasbarre, L., (1997).** Effects of gastrointestinal nematode infection on the ruminant immune system. *Vet Parasitol*, 72: p. 327-343.
88. **Gasbarre, L., E.A. Leighton, and C.J. Davies, (1993).** Influence of host genetics upon antibody responses against gastrointestinal nematode infections in cattle. *Vet Parasitol*, 46: p. 81-91.
89. **Gasbarre, L., E.A. Leighton, and T. Sonstegard, (2001).** Role of the bovine immune system and genome in resistance to gastrointestinal nematode. *Vet Parasitol*, 98: p. 51-64.
90. **Gasbarre, L.C., (1994).** *Ostertagia ostertagi*: changes in lymphoid populations in the local lymphoid tissues after primary or secondary infection. *Vet Parasitol*, 55: p. 105-114.
91. **Gasbarre, L.C., E.A. Leighton, and C.J. Davies, (1990).** Genetic control of immunity to gastrointestinal nematodes of cattle. *Vet Parasitol*, 37: p. 257-272.
92. **Gause, W.C., et al., (1996).** *H. polygyrus*: B7-independence of the secondary type 2 response. *Exp Parasitol*, 84: p. 264-273.
93. **Gause, W.C., J.F. Urban Jr, and M.J. Stadecker, (2003).** The immune response to parasitic helminths: insights from murine models. *Trends Immunol*, 24: p. 269-277.
94. **Genchi, C., M. Madonna, and G. Traldi, (1989).** Epidemiology of *Ostertagia ostertagi* in dairy cow from different breeding systems. *Parasitologia*, 31: p. 123-132.
95. **Gill, H.S., (1991).** Genetic control of acquired resistance to haemonchosis in Merino lambs. *Parasite Immunol*, 13: p. 617-628.
96. **Gill, H.S., et al., (2000).** Induction of T-helper-1 and T-helper-2 immune responses during *Haemonchus contortus* infection in sheep. *Immunology*, 99: p. 458-463.
97. **Gill, H.S., et al., (1993).** Isotype-specific antibody responses to *Haemonchus contortus* in genetically resistant sheep. *Parasite Immunol*, 15: p. 61-67.
98. **Gill, H.S., et al., (1994).** Antibody-containing cells in the abomasal mucosa of sheep with genetic resistance to *Haemonchus contortus*. *Res Vet Sci*, 56: p. 41-47.

99. **Githiori, J.B., et al., (2000).** Use of a 14.2 kDa recombinant *Cooperia oncophora* protein in an ELISA for herd health monitoring of nematode infections in first grazing season calves. *Vet Parasitol*, 91: p. 63-78.
100. **Gordon, H.M. and H.V. Whitlock, (1939).** A new technique for counting nematode eggs in sheep faeces. *J Counc Sci Ind Res*, 12: p. 50-52.
101. **Gorrell, M.D., et al., (1988).** Lymphocyte phenotypes in the intestinal mucosa of sheep infected with *Trichostrongylus colubriformis*. *Clin Exp Immunol*, 72: p. 274-279.
102. **Greenwald, R.J., et al., (1997).** Effects of blocking B7-1 and B7-2 interactions during a type 2 in vivo immune response. *J Immunol*, 158: p. 4088-4096.
103. **Greenwald, R.J., et al., (1999).** B7-2 is required for the progression but not the type 2 immune response to a gatrointestinal nematode parasite. *J Immunol*, 162: p. 4133-4139.
104. **Grencis, R.K., et al., (1993).** The in vivo role of stem cell factor (c-kit ligand) on mastocytosis and host protective immunity to the intestinal nematode *Trichinella spiralis* in mice. *Parasite Immunol*, 15: p. 55-59.
105. **Grencis, R.K. and G.M. Entwistle, (1997).** Production of an interferon-gamma homologue by an intestinal nematode: functionally significant or interesting artefact? *Parasitology*, 115: p. S101-S105.
106. **Grencis, R.K., J. Riedlinger, and D. Wakelin, (1985).** L3T4-positive T lymphoblasts are responsible for transfer of immunity to *Trichinella spiralis* in mice. *Immunology*, 56: p. 213-218.
107. **Hanrahan, L.A., G.W. Benz, and R.D. Schultz, (1984).** Experimentally induced *Cooperia oncophora* infection in calves: lymphocyte blastogenic and delayed hypersensitivity responses. *Am J Vet Res*, 45: p. 855-862.
108. **Harari, Y. and G.A. Castro, (1991).** Reconstruction of immune-mediated ion secretion in gut mucosa of the athymic rat. *Parasite Immunol*, 13: p. 313-328.
109. **Hayday, A., (2000).** gd Cells: a right time and a right way for a conserved third way of protection. *Ann Rev Immunol*, 18: p. 975-1026.
110. **Hein, W., L. Dudler, and C.R. Mackay, (1989).** Surface expression of differentiation antigens on lymphocytes in the ileal and jejunal Peyer's patches of lambs. *Immunology*, 68: p. 365-370.
111. **Hein, W.R. and C.R. Mackay, (1991).** Prominence of gd T cells in the ruminant immune system. *Immunology Today*, 12: p. 30-34.
112. **Hilderson, H., et al., (1993).** Characterisation of *Ostertagia ostertagi* antigens by the different bovine immunoglobulin isotypes. *Res Vet Sci*, 55: p. 203-208.
113. **Howard, C.J., et al., (1992).** Immunity to bovine virus diarrhoea virus in calves: the role of different T-cell subpopulations analysed by specific depletion in vivo with monoclonal antibodies. *Vet Immunol Immunopathol*, 32: p. 303-314.
114. **Howard, C.J., et al., (1997).** Identification of two distinct populations of dendritic cells in afferent lymph that vary in their ability to stimulate T cells. *J Immunol*, 159: p. 5372-5282.
115. **Howard, C.J., P. Sopp, and K.R. Parsons, (1992).** L-selectin expression differentiates T cells isolated from different lymphoid organs in cattle but does not correlate with memory. *Immunology*, 77: p. 228-234.
116. **Huntley, J.F., et al., (1987).** Systemic release of a mast cell proteinase following nematode infections in sheep. *Parasite Immunol*, 9: p. 603-614.

## References

---

117. **Huntley, J.F., G. Newlands, and H.R. Miller, (1984).** The isolation and characterization of globule leucocytes: their derivation from mucosal mast cells in parasitized sheep. *Parasite Immunol*, 6: p. 371-390.
118. **Huntley, J.F., et al., (1992).** The influence of challenge dose, duration of immunity, or steroid treatment on mucosal mast cells and on the distribution of sheep mast cell proteinase in *Haemonchus*-infected sheep. *Parasite Immunol*, 14: p. 429-440.
119. **Huntley, J.F., et al., (1998).** IgE antibody during infection with the ovine abomasal nematode, *Teladorsagia circumcincta*: primary and secondary responses in serum and gastric lymph of sheep. *Parasite Immunol*, 20: p. 565-571.
120. **Hurst, S.D. and T.A. Barret, (1996).** T-cell responses to enteric antigens. *Semin Gastrointest Dis*, 7: p. 118-123.
121. **Isenstein, R.S., (1963).** The life history of *Cooperia oncophora* (Railliet, 1898) Ransom, 1907, a nematode parasite of cattle. *Parasitology*, 49: p. 235-240.
122. **James, S.P., (1993).** The gastrointestinal mucosal immune system. *Dig Dis*, 11: p. 146-156.
123. **Jarrett, E.E. and H.R. Miller, (1982).** Production and activities of IgE in helminth infection. *Progress in Allergy*, 31: p. 178-233.
124. **Jarrett, W.F., et al., (1960).** Immunological studies on *Dictyocaulus viviparus* infection. Immunity produced by the administration of irradiated larvae. *Immunology*, 3: p. 145-151.
125. **Jones, D.G., (1993).** The eosinophil. *J Comp Pathol*, 108: p. 317-335.
126. **Kanobana, K., H.W. Ploeger, and L. Vervelde, (2002).** Immune expulsion of the trichostrongylid *Cooperia oncophora* is associated with increased eosinophilia and mucosal IgA. *Int J Parasitol*, 32: p. 1389-1398.
127. **Kanobana, K., et al., (2001).** Characterization of host responder types after a single *Cooperia oncophora* infection: kinetics of the systemic immune response. *Parasite Immunol*, 23: p. 641-653.
128. **Kantele, A., et al., (1997).** Homing potentials of circulating lymphocytes in humans depend on the site of activation: oral, but not parenteral, typhoid vaccination induces circulating antibody-secreting cells that all bear homing receptors directing them to the gut. *J Immunol*, 158: p. 574-579.
129. **Katona, I.M., J.F. Urban, Jr., and F.D. Finkelman, (1988).** The role of L3T4+ and Lyt-2+ T cells in the IgE response and immunity to *Nippostrongylus brasiliensis*. *J Immunol*, 140: p. 3206-3211.
130. **Klesius, P.H., (1988).** Immunity to *Ostertagia ostertagi*. *Vet Parasitol*, 27: p. 159-167.
131. **Kloosterman, A., (1971).** Observations on the epidemiology of trichostrongylosis of calves. *PhD thesis. Agricultural university, Wageningen*.
132. **Kloosterman, A., G. Albers, and R. van den Brink, (1978).** Genetic variation among calves in resistance to nematode parasites. *Vet Parasitol*, 4: p. 353-368.
133. **Kloosterman, A. and K. Franken, (1988).** Interactions between lungworms and gastrointestinal worms in calves. *Vet Parasitol*, 26: p. 305-320.
134. **Kloosterman, A., H.K. Parmentier, and H.W. Ploeger, (1992).** Breeding cattle and sheep for resistance to gastrointestinal nematodes. *Parasitology Today*, 8: p. 330-335.
135. **Kloosterman, A., H.W. Ploeger, and K. Franken, (1991).** Age resistance in calves to *Ostertagia ostertagi* and *Cooperia oncophora*. *Vet Parasitol*, 39: p. 101-113.

136. **Kooyman, F.N., et al., (1997).** Production of a monoclonal antibody specific for ovine immunoglobulin E and its application to monitor serum IgE responses to *Haemonchus contortus* infection. *Parasitology*, 114: p. 395-406.
137. **Kooyman, F.N., et al., (2002).** Serum immunoglobulin E response in calves infected with the lungworm *Dictyocaulus viviparus* and its correlation with protection. *Parasite Immunol*, 24: p. 47-56.
138. **Korenaga, M., et al., (1994).** Regulatory effect of anti-interleukin-5 monoclonal antibody on intestinal worm burden in a primary infection with *Strongyloides venezuelensis* in mice. *Int J Parasitol*, 24: p. 951-957.
139. **Koyama, K., H. Tamauchi, and Y. Ito, (1995).** The role of CD4+ and CD8+ cells in the protective immunity to the murine nematode parasite *Trichuris muris*. *Parasite Immunol*, 17: p. 161-165.
140. **Kraehenbuhl, J.-P. and M.R. Neutra, (1992).** Molecular and cellular basis of immune protection of mucosal surfaces. *Physiol Rev*, 72: p. 853-879.
141. **Kusama, Y., et al., (1995).** Mechanisms of eosinophilia in Balb/c-nu/+ and congenitally athymic Balb/C-nu/nu mice infected with *Toxocara canis*. *Immunology*, 84: p. 461-468.
142. **Laemmli, U.K., (1970).** Cleavage of structural proteins during the assembly of the head of bacteriophage. *Nature*, 227: p. 680-687.
143. **Lamkhioued, B., et al., (1995).** Human eosinophils express a receptor for secretory component. Role in secretory IgA-dependent activation. *Eur J Immunol*, 25: p. 117-125.
144. **Larsen, M., et al., (1997).** Biological control of gastro-intestinal nematodes - facts, future, or fiction? *Vet Parasitol*, 72: p. 479-492.
145. **Lascelles, A. K., et al., (1986).** The mucosal immune system with particular reference to ruminant animals. In: *The ruminant immune system in health and disease*. I.W. Morrison (Ed.). Cambridge University Press, Cambridge. p 429-457.
146. **Leighton, E.A., K.D. Murrell and L.C. Gasbarre, (1989).** Evidence for genetic control of nematode egg-shedding rates in calves. *J Parasitol*, 75: p. 498-504.
147. **Lenshow, D.J., T.L. Walunas, and J.A. Bluestone, (1996).** CD28/B7 system of costimulation. *Annu Rev Immunol*, 14: p. 233-258.
148. **Lightowers, M.W. and M.D. Rickard, (1988).** Excretory-secretory products of helminth parasites: effects on host immune responses. *Parasitology*, 96: p. S123-166.
149. **Madden, K.B., et al., (1991).** Antibodies to IL-3 and IL-4 suppress helminth-induced intestinal mastocytosis. *J Immunol*, 147: p. 1387-1391.
150. **Mansour, M.M., et al., (1990).** Bovine immune recognition of *Ostertagia ostertagi* larval antigens. *Vet Immunol Immunopathol*, 24: p. 361-371.
151. **McClure, S.J. and D.L. Emery, (1994).** Cell-mediated responses against gastrointestinal nematode parasites of ruminants. In: *Cell mediated immunity in Ruminants*. B.M. Goddeeris and I.W. Morrison (Eds.), Boca Ranton, CRC Press, p 250
152. **McClure, S.J., et al., (1998).** Attempts to generate immunity against *Trichostrongylus colubriformis* and *Haemonchus contortus* in young lambs by vaccination with viable parasites. *Int J Parasitol*, 28: p. 739-746.
153. **McClure, S.J., et al., (1995).** Depletion of IFN-gamma, CD8+ or TCR gamma delta+ cells in vivo during primary infection with an enteric parasite (*Trichostrongylus colubriformis*) enhances protective immunity. *Immunol Cell Biol*, 73: p. 552-555.

## References

---

154. **McClure, S.J., et al., (1992).** A serial study of rejection of *Trichostrongylus colubriformis* by immune sheep. *Int J Parasitol*, 22: p. 227-234.
155. **McDonald, A.S. and E.J. Pearce, (2002).** Cutting edge: polarized Th induction by transferred antigen pulsed dendritic cells is dependent upon IL-4 or IL-12 production by recipient cells. *J Immunol*, 168: p. 3127-3130.
156. **Meeusen, E.N., (1999).** Immunology of helminth infection with special references to immunopathology. *Vet Parasitol*, 84: p. 259-273.
157. **Meeusen, E.N. and A. Balic, (2000).** Do Eosinophils have a Role in the Killing of Helminth Parasites? *Parasitology Today*, 16: p. 95-101.
158. **Mertens, B., E. Gobright, and H.F. Seow, (1996).** The nucleotide sequence of the bovine interleukin-5-encoding cDNA. *Gene*, 176: p. 273-274.
159. **Mes, T.H., et al., (2001).** A novel method for the isolation of gastro-intestinal nematode eggs that allows automated analysis of digital images of eg preparations and high throughput screening. *Parasitology*, 123.
160. **Michel, J.F., (1963).** The phenomena of host resistance and the course of infection of *Ostertagia ostertagi* in calves. *Parasitology*, 53: p. 63-84.
161. **Miller, H.R., (1984).** The protective mucosal response against gastrointestinal nematodes in ruminants and laboratory animals. *Vet Immunol Immunopathol*, 6: p. 167-259.
162. **Miller, H.R., (1996).** Mucosal mast cells and the allergic response against nematode parasites. *Vet Immunol Immunopathol*, 54: p. 331-336.
163. **Miller, H.R., (1996).** Prospects for the immunological control of ruminant gastrointestinal nematodes: natural immunity, can it be harnessed? *Int J Parasitol*, 26: p. 801-811.
164. **Miller, T.A., (1971).** Vaccination against the canine hookworm disease. *Adv Parasitol*, 9: p. 153-183.
165. **Mosmann, T.R., et al., (1986).** Two types of murine helper T cell clone. I. Definition according to profiles of lymphokine activities and secreted proteins. *J Immunol*, 136: p. 2605-2616.
166. **Motegi, Y., et al., (2000).** Role of secretory IgA, secretory component, and eosinophils in mucosal inflammation. *Int Arch Allergy Immunol*, 122 Suppl 1: p. 25-27.
167. **Munn, E.A., (1997).** Rational design of nematode vaccines: hidden antigens. *Int J Parasitol*, 27: p. 359-366.
168. **Nawa, Y., et al., (1994).** Selective effector mechanisms for the expulsion of intestinal helminths. *Parasite Immunol*, 16: p. 333-338
169. **Nawa, Y. and H.R.P. Miller, (1979).** Adoptive transfer of the intestinal mast cell response in rats infected with *Nippostrongylus brasiliensis*. *Cell Immunol*, 42: p. 225-239.
170. **Newton, S.E. and E.A. Munn, (1999).** The development of vaccines against gastrointestinal nematode parasites, particularly *Haemonchus contortus*. *Parasitology Today*, 15: p. 116-122.
171. **Nieuwland, M.G., et al., (1995).** Systemic antibody responses of calves to low molecular weight *Cooperia oncophora* antigens. *Vet Parasitol*, 59: p. 231-239.
172. **Owen, R.L., (1994).** M cells-entryways of opportunity for enteropathogens. *J Exp Med*, 180: p. 7-9.
173. **Paciorkowski, N., L.D. Shultz, and T.V. Rajan, (2003).** Primed peritoneal B lymphocytes are sufficient to transfer protection against *Brugia pahangi* infection in mice. *Infect Immun*, 71: p. 1370-1378.

174. **Parciorkowsky, N., et al., (2000).** B1 B lymphocytes play an critical role in host protection against lymphatic filarial parasites. *J Exp Med*, 191: p. 731-736.
175. **Park, Y.H., et al., (1994).** A subset of mammary gland GD T lymphocytes downregulates BoCD4 T lymphocyte response to *Staphylococcus aureus* in cattle with intramammary infection. *Korean J Immunol*, 16: p. 19-27.
176. **Parmentier, H.K., et al., (1995).** Low molecular weight *Cooperia oncophora* antigens: characterization and humoral immune responses in calves mono-infected with 100,000 infective larvae. *Vet Parasitol*, 59: p. 219-230.
177. **Paterson, S. and M.E. Viney, (2002).** Host immune responses are necessary for density dependence in nematode infections. *Parasitology*, 125: p. 283-292.
178. **Ploeger, H., et al., (2000).** Cross-sectional serological survey on gastrointestinal and lung nematode infections in first and second-year replacement stock in the Netherlands: relation with management practices and use of anthelmintics. *Vet Parasitol*, 90: p. 285-304.
179. **Ploeger, H. and A. Kloosterman, (1993).** Gastrointestinal infections and weight gain in dairy replacement stock: first-year calves. *Vet Parasitol*, 46: p. 223-241.
180. **Ploeger, H.W., et al., (1990).** Effect of nematode infections on growth performance of calves after stabling on commercial dairy farms. *Vet Parasitol*, 36: p. 71-81.
181. **Ploeger, H.W., et al., (1990).** Effect of naturally occuring nematode infections in the first and the second grazing season on the growth performance of second year cattle. *Vet Parasitol*, 36: p. 57-70.
182. **Ploeger, H.W., et al., (1994).** Quantitative estimation of the level of exposure to gastrointestinal nematode infection in first-year calves. *Vet Parasitol*, 55: p. 287-315
183. **Ploeger, H.W., A. Kloosterman, and F.W. Rietveld, (1995).** Acquired immunity against *Cooperia* spp. and *Ostertagia* spp. in calves: effect of level exposure and timing of the midsummer increase. *Vet Parasitol*, 58: p. 61-74.
184. **Ploeger, H.W., et al., (1996).** Production of dairy replacement stock in relation to level of exposure to gastrointestinal nematode infection in the first grazing season: second year calves and heifers. *Vet Parasitol*, 22.
185. **Poot, J., et al., (1997).** Use of cloned excretory/secretory low-molecular-weight proteins of *Cooperia oncophora* in a serological assay. *J Clin Microbiol*, 35: p. 1728-1733.
186. **Press, C., S. McClure, and T. Landsverk, (1991).** Computer-assisted morphometric analysis of absorptive and follicle- associated epithelia of Peyer's patches in sheep foetuses and lambs indicates the presence of distinct T- and B-cell components. *Immunology*, 72: p. 386-392.
187. **Pritchard, D.I., (1993).** Immunity to helminths: is too much IgE parasite- -rather than host-protective? *Parasite Immunol*, 15: p. 5-9.
188. **Pritchard, D.I., R.J. Quinell, and E.A. Walsh, (1995).** Immunity in humans to *Necator americanus*: IgE, parasite weighth and fecundity. *Parasite Immunol*, 17: p. 71-75.
189. **Romagnani, S., (1992).** Human TH1 and TH2 subsets: regulation of differentiation and role in protection and immunopathology. *Int Arch Allergy Immunol*, 98: p. 279-285.
190. **Ron, Y. and J. Sprent, (1987).** T cell priming in vivo: a major role for B cells in presenting antigen to T cells in lymph nodes. *J Immunol*, 138: p. 2848.
191. **Rothkotter, H., R. Pabst, and M. Bailey, (1999).** Lymphocyte migration in the intestinal mucosa: entry, transit and emigration of lymphoid cells and the influence of antigen. *Vet Immunol Immunopathol*, 72: p. 157-165.

## References

---

192. **Rothwell, T.L., (1989).** Immune expulsion of parasitic nematodes from the alimentary tract. *Int J Parasitol*, 19: p. 139-168.
193. **Sanderson, C.J., (1992).** Interleukin-5, eosinophils, and disease. *Blood*, 79: p. 3101-3109.
194. **Sasaki, O., et al., (1993).** Ablation of eosinophils with anti-IL-5 antibody enhances the survival of intracranial worms of *Angiostrongylus cantonensis* in the mouse. *Parasite Immunol*, 15: p. 349-354.
195. **Schallig, H.D., et al., (1994).** Serum antibody responses of Texel sheep experimentally infected with *Haemonchus contortus*. *Res Vet Sci*, 57: p. 63-68.
196. **Schallig, H.D., M.A. van Leeuwen, and A.W. Cornelissen, (1997).** Protective immunity induced by vaccination with two *Haemonchus contortus* excretory secretory proteins in sheep. *Parasite Immunol*, 118: p. 297-304.
197. **Schallig, H.D., M.A. van Leeuwen, and W.M. Hendrikx, (1994).** Immune responses of Texel sheep to excretory/secretory products of adult *Haemonchus contortus*. *Parasitology*, 108: p. 351-357.
198. **Schierack, P., et al., (2003).** Parasite-specific immunomodulatory functions of filarial cystatin. *Infect Immun*, 71: p. 2422-2429.
199. **Seaton, D.S., et al., (1989a).** Development of immunity to incoming radiolabelled larvae in lambs continuously infected with *Ostertagia circumcincta*. *Res Vet Sci*, 46: p. 241-246.
200. **Shaw, R.J., T.K. Gatehouse, and M.M. McNeill, (1998).** Serum IgE responses during primary and challenge infections of sheep with *Trichostrongylus colubriformis*. *Int J Parasitol*, 28: p. 293-302.
201. **Shaw, R.J., et al., (1997).** Quantification of total sheep IgE concentration using anti-ovine IgE monoclonal antibodies in an enzyme immunoassay. *Vet Immunol Immunopathol*, 57: p. 253-265.
202. **Simon, H.U., (1997).** Apoptosis and *in vivo* distribution and clearance of eosinophils in normal and *Trichinella spiralis*-infected rats. *J Leukocyte Biol*, 62: p. 309-317.
203. **Slifka, M.K., et al., (1998).** Humoral immunity due to long-lived plasma cells. *Immunity*, 8: p. 363-372.
204. **Smith, G. and D.T. Galligan, (1988).** Mathematical models of the population biology of *Ostertagia ostertagi* and *Teladorsagia circumcincta*, and the economic evaluation of disease control strategies. *Vet Parasitol*, 27: p. 73-83.
205. **Smith, W.D. and K.W. Angus, (1980).** *Haemonchus contortus*: attempts to immunise lambs with irradiated larvae. *Res Vet Sci*, 29: p. 45-50.
206. **Smith, W.D., et al., (1986).** Transfer of immunity to *Ostertagia circumcincta* and IgA memory between identical sheep by lymphocytes collected from gastric lymph. *Res Vet Sci*, 41: p. 300-306.
207. **Smith, W.D., et al., (1984).** Resistance to *Haemonchus contortus* transferred between genetically histocompatible sheep by immune lymphocytes. *Res Vet Sci*, 37: p. 199-204.
208. **Sopp, P. and C.J. Howard, (2001).** INF- $\gamma$  and IL-4 production by CD4, CD8 and WC1  $\gamma\delta$  TCR+ T cells from cattle lymph nodes and blood. *Vet Immunol Immunopathol*, 81: p. 85-96.
209. **Stear, M.J., et al., (1997a).** The genetic basis of resistance to *Ostertagia circumcincta* in lambs. *Vet J*, 154: p. 111-119.
210. **Stear, M.J., et al., (1997b).** How hosts control worms. *Nature*, 389: p. 27.
211. **Stear, M.J. and S.C. Bishop, (1999a).** The curvilinear relationship between worm length and fecundity of *Teladorsagia circumcincta*. *Int J Parasitol*, 29: p. 777-780.

212. **Stear, M.J., S. Strain, and S.C. Bishop, (1999).** Mechanisms underlying resistance to nematode infection. *Int J Parasitol*, 29: p. 51-56; discussion 73-55.
213. **Stevenson, L.M., et al., (1994).** Local eosinophil- and mast cell-related responses in abomasal nematode infections of lambs. *FEMS Immunol Med Microbiol*, 8: p. 167-173.
214. **Stevenson, L.M. and D.G. Jones, (1994).** Cross-reactivity amongst recombinant haematopoietic cytokines from different species for sheep bone-marrow eosinophils. *J Comp Pathol*, 111: p. 99-106.
215. **Stevenson, L.M., et al., (1998).** Eosinophil-specific biological activity of recombinant ovine interleukin-5. *Vet Immunol Immunopathol*, 66: p. 359-365.
216. **Sutton, B.J. and H.J. Gould, (1993).** The human IgE network. *Nature*, 366: p. 421-428.
217. **Tezuka, H., et al., (2002).** Recombinant *Dirofilaria immitis* polyprotein that stimulates murine B cells to produce nonspecific polyclonal immunoglobulin E antibody. *Infect Immun*, 70: p. 1235-1244.
218. **Thamsborg, S.M., A. Roepstorff, and M. Larsen, (1999).** Integrated and biological control of parasites in organic and conventional production systems. *Vet Parasitol*, 84: p. 169-186.
219. **Thatcher, E.F., L.J. Gershwin, and N.F. Baker, (1989).** Levels of serum IgE in response to gastrointestinal nematodes in cattle. *Vet Parasitol*, 32: p. 153-161.
220. **Tivol, E.A., et al., (1995).** Loss of CTLA-4 leads to massive lymphoproliferations and fetal multiorgan tissue destruction, revealing a critical negative regulatory role of CTLA-4. *Immunity*, 3: p. 542.
221. **Urban, J.F. Jr., et al., (1992).** The importance of Th2 cytokines in protective immunity to nematodes. *Immunol rev*, 127: p. 205.
222. **Urban, J.F. Jr., I.M. Katona, and F.D. Finkelman, (1991).** *Heligmosomoides polygyrus*: CD4+ but not CD8+ T cells regulate the IgE response and protective immunity in mice. *Exp Parasitol*, 73: p. 500-511.
223. **Vallance, B.A., et al., (1999).** IL-5 contributes to worm expulsion and muscle hypercontractility in a primary *T. spiralis* infection. *Am J Physiol*, 277: p. G400-408.
224. **Vallance, B.A., et al., (2001).** Mast cell-independent impairment of host defense and muscle contraction in *T. spiralis*-infected W/W(V) mice. *Am J Physiol Gastrointest Liver Physiol*, 280: p. G640-648.
225. **Vallance, B.A. and S.M. Collins, (1998).** The effect of nematode infection upon intestinal smooth muscle function. *Parasite Immunol*, 20: p. 249-253.
226. **Vallance, B.A., et al., (1999).** CD4 T cells and major histocompatibility complex class II expression influence worm expulsion and increased intestinal muscle contraction during *Trichinella spiralis* infection. *Infect Immun*, 67: p. 6090-6097.
227. **Vallance, B.A., et al., (2000).** Interleukin-5 deficient mice exhibit impaired host defence against challenge *Trichinella spiralis* infections. *Parasite Immunol*, 22: p. 487-492.
228. **Van den Broeck, W., E. Cox, and B.M. Goddeeris, (1999).** Induction of immune responses in pigs following oral administration of purified F4 fimbriae. *Vaccine*, 17: p. 2020-2029.
229. **Van der Kleij, D., et al., (2002).** Triggering of innate immune responses by schistosome egg glycolipids and their carbohydrate epitope GalNAc beta 1-4 (Fuc alpha 1-2Fuc alpha 1-3) GlcNAc. *J Infect Dis*, 185: p. 531-539.
230. **Van Die, I., et al., (1999).** Core alpha 1-3-fucose is a common modification of N-glycans in parasitic helminths and constitutes an important epitope of IgE from *Haemonchus* infected sheep. *Febs Lett*, 463: p. 189-193.

## References

---

231. **Van Diemen, P.M., et al., (1997).** Low molecular weight *Cooperia oncophora* antigens. Potential to discriminate between susceptible and resistant calves after infection. *Int J Parasitol*, 27: p. 587-593.
232. **Van Diemen, P.M., et al., (1996).** Recognition of low molecular weight *Cooperia oncophora* antigens after primary and trickle infection of calves with third-stage infective larvae. *Int J Parasitol*, 26: p. 1305-1310.
233. **Vercruyse, J. and E. Claerebout, (2001).** Treatment vs non-treatment of helminth infections in cattle: defining the threshold. *Vet Parasitol*, 98: p. 195-214.
234. **Vermillion, D.L., P.B. Ernst, and S.M. Collins, (1991).** T-lymphocyte modulation of intestinal muscle function in the *Trichinella*-infected rat. *Gastroenterology*, 101: p. 31-38.
235. **Vervelde, L., et al., (2001).** Protection studies with recombinant excretory/secretory proteins of *Haemonchus contortus*. *Parasite Immunol*, 24: p. 189-201.
236. **Viney, M., (2002).** How do host immune responses affect nematode infections? *Trends Parasitol*, 18: p. 63-66.
237. **Walcheck, B. and M.A. Jutila, (1994).** Bovine gamma delta T cells express high levels of functional peripheral lymph node homing receptor (L-selectin). *Int Immunol*, 6: p. 81-91.
238. **Waller, P., (1997).** Anthelmintic resistance. *Vet Parasitol*, 72: p. 391-412.
239. **Waller, P.J., (1978).** Host-induced effects on the morphology of *Ostertagia circumcincta* in grazing sheep. *Int J Parasitol*, 8: p. 365-370.
240. **Waterhouse, P., et al., (1995).** Lymphoproliferative disorders with early lethality in mice deficient in CTLA-4. *Science*, 270: p. 932.
241. **Whelan, M., et al., (2000).** A filarial nematode-secreted product signals dendritic cells to acquire a phenotype that drives development of Th2 cells. *J Immunol*, 164: p. 6453-6460.
242. **Williams, D.J., J. Newson and J. Naessens, (1990).** Quantitation of bovine immunoglobulin isotypes and allotypes using monoclonal antibodies. *Vet Immunol Immunopathol*, 24: p. 267-283.
243. **Wilson, R.A., et al., (1996).** T-cell subsets in blood and lymphoid tissues obtained from fetal calves, mature calves and adult bovine. *Vet Immunol Immunopathol*, 53: p. 49-60.
244. **Windon, R.G., J.K. Dineen, and J.D. Kelly, (1980).** The segregation of lambs into 'responders' and 'non responders': response to vaccination with irradiated *Trichostrongylus colubriformis* larvae before weaning. *Int J Parasitol*, 10: p. 65-73.
245. **Winter, M.D., C. Wright, and D.L. Lee, (1997).** The mast cell and eosinophil response of young lambs to a primary infection with *Nematodirus battus*. *Parasitology*, 114: p. 189-193.
246. **Wyatt, C.R., et al., (1994).** Differential distribution of gdT-cell receptor lymphocyte subpopulation in blood and spleen of young and adult cattle. *Vet Immunol Immunopathol*, 40: p. 187-199.
247. **Yatsuda, A.P., et al., (2002).** *Cooperia punctata* trickle infections: parasitological parameters and evaluation of a *Cooperia* recombinant 14.2 kDa protein ELISA for estimating cumulative exposure of calves. *Vet Parasitol*, 105: p. 131-138.
248. **Zarlenga, D.S., et al., (1998).** Identification and semi-quantitation of *Ostertagia ostertagi* eggs by enzymatic amplification of ITS-1 sequences. *Vet Parasitol*, 77: p. 245-257.
249. **Dalton, J.P. and G. Mulcahy, (2001).** Parasite vaccines--a reality? *Vet Parasitol*, 98: p. 149-167.