

## Role of the Growth-Associated Protein B-50/GAP-43 in Neuronal Plasticity

*W. H. Gispen, H. B. Nielander, P. N. E. De Graan, A. B. Oestreicher,  
L. H. Schrama, and P. Schotman\**

*Rudolf Magnus Institute, Institute of Molecular Biology and Medical Biotechnology,  
and Laboratory for Physiological Chemistry, University of Utrecht, Padualaan 8,  
3584 CH Utrecht, The Netherlands*

### Contents

Abstract	Functional Aspects of B-50
Introduction	Role in Neuritogenesis
Expression	Role in NGF-Induced Neuronal Differentiation of PC12 cells
Neuron Specificity?	Translocation of B-50 with NGF-Induced Neuritogenesis
GAP-43, pp46, Fl, P-57, Neuromodulin	Localization in Neuronal Growth Cones
Primary Structure of B-50	Changes in B-50 Distribution During Neuronal Polarization
Anomalous Behavior in SDS-PAGE	Role of B-50 Phosphorylation in Signal Trans- duction in the Growth Cone
Primary Structure	B-50 Phosphorylation and LTP
Secondary Structure	Role of B-50 in Presynaptic Signal Transduction and Neurotransmitter Release
Homology to Other Proteins	Acknowledgments
Purified Protein and Primary Translation Product	References
Isolation Procedures	
Posttranslational Modifications	
Microheterogeneity	
Domains in B50	
Gene Structure	
Membrane Association	
Interaction with G <sub>0</sub> Protein	
Phosphorylation by PKC	
Calmodulin Binding	
Interaction with Cytoskeletal Elements	

\*Author to whom all correspondence and reprint requests should be addressed.

## Abstract

The neuronal phosphoprotein B-50/GAP-43 has been implicated in neuritogenesis during developmental stages of the nervous system and in regenerative processes and neuronal plasticity in the adult. The protein appears to be a member of a family of acidic substrates of protein kinase C (PKC) that bind calmodulin at low calcium concentrations. Two of these substrates, B-50 and neurogranin, share the primary sequence coding for the phospho- and calmodulin-binding sites and might exert similar functions in axonal and dendritic processes, respectively. In the adult brain, B-50 is exclusively located at the presynaptic membrane. During neuritogenesis in cell culture, the protein is translocated to the growth cones, i.e., into the filopodia. In view of many positive correlations between B-50 expression and neurite outgrowth and the specific localization of B-50, a role in growth cone function has been proposed. Its phosphorylation state may regulate the local intracellular free calmodulin and calcium concentrations or vice versa. Both views link the B-50 protein to processes of signal transduction and transmitter release.

**Index Entries:** B-50; GAP-43; neuromodulin; growth-associated proteins; cDNA cloning; protein kinase C; calmodulin binding; immune localization; pheochromocytoma (PC12) cells; NGF induction; growth cones; transmitter release.

## Introduction

### Expression

The protein B-50 was originally described as a phosphoprotein in adult rat brain (Zwiers et al., 1976), where it is located at the presynaptic membrane (Gispen et al., 1985a). The identical protein GAP-43 was described as a prominent member of the family of growth-associated proteins (GAPs) in the nervous system. These represent a relatively small subset of proteins synthesized at strikingly higher levels during neurite outgrowth (Skene and Willard, 1981). It has been proposed that the induction of members of this family of GAPs is a prerequisite for axonal growth (Levine et al., 1981). The best characterized member of the family is GAP-43/B-50, in goldfish named GAP-48 (Benowitz and Lewis, 1983). It is consistently expressed during developmental and regenerative axonal growth and is delivered by fast axonal transport to the extending neurite, especially to the growth cones (Skene and Willard, 1981). The presumed mediating role of B-50 in neurite growth is documented by a large number of studies in different neuronal systems during the last decade (Benowitz and Routtenberg,

1987; Allsopp and Moss, 1989; Benowitz et al., 1989,1990; De la Monte et al., 1989; Erzurumlu et al., 1989,1990; Gorgels et al., 1989; Hoffman, 1989; O'Hara et al., 1989; Skene, 1989; Tetzlaff et al., 1989; Van der Neut et al., 1990; Van der Zee et al., 1989; Verhaagen et al., 1989,1990; Baizer et al., 1990; Biffo et al., 1990; DiFiglia et al., 1990; Goslin and Banker, 1990; McIntosh et al., 1990; Moya et al., 1990; Sheu et al., 1990; Woolf et al., 1990; Yamamoto and Kondo, 1990; Yankner et al., 1990).

### Neuron Specificity?

Although B-50 is usually referred to as a neuron-specific protein, a number of recent reports show that under certain circumstances, B-50 is present in other cell types of the nervous system (Vitkovic et al., 1988; Tetzlaff et al., 1989; Da Cunha and Vitkovic, 1990; Deloulme et al., 1990; Hammang et al., 1990). The general opinion is that B-50 is restricted to the nervous system and is expressed in most if not all neuronal cells at some stage. During axonal growth or regeneration, this expression is enhanced. In mature undamaged neurons, expression is mostly low, whereas neurons that are believed to be involved in ongoing synaptic remodeling, e.g., in human associative

brain areas (Neve et al., 1987, 1988; Ng et al., 1988; Benowitz et al., 1989) and rat hippocampal and olfactory areas (Oestreicher et al., 1986; Oestreicher and Gispen, 1986; Benowitz et al., 1988; De la Monte et al., 1989; Verhaagen et al., 1989) continue to express high levels of B-50.

### **GAP-43, pp46, F1, P-57, and Neuromodulin**

The protein B-50/GAP-43 is also known under several other names, reflecting independent characterization by various approaches: pp46, a major growth cone protein from fetal rat brain (Katz et al., 1985); F1, a rat hippocampal protein implicated in a model of information storage known as long-term potentiation (LTP; Lovinger et al., 1986); or P-57, now called neuromodulin, a calmodulin (CaM)-binding protein from bovine brain (Andreasen et al., 1983). Although the vast amount of data obtained from different fields of neuroscience has greatly expanded knowledge of the protein and its characteristics, the physiological role of B-50 remains to be solved.

## **Primary Structure of B-50**

### **Anomalous Behavior in SDS-PAGE**

Originally characterized as an acidic protein (pI 4.3–4.7), B-50 showed anomalous behavior in SDS-PAGE; the apparent mol wt increases with lowered percentage acrylamide (Schrama et al., 1987; Benowitz et al., 1987). In 11% gels, the rat protein (23.6 kDa) has an apparent mol wt of 48,000. The unusual behavior of B-50 in SDS-PAGE has been used to identify the protein in different neuronal systems (Benowitz and Routtenberg, 1987). It has, however, delayed the recognition that proteins studied under different names by several laboratories were in fact identical. The most notable example is the bovine protein P-57, or neuromodulin, discovered by its CaM-binding properties (Andreasen et al., 1983); its identity with B-50 was only recently recognized (Cimler et al., 1987).

## **Primary Structure**

The primary structure of B-50 has been determined in a growing number of species. In bovine, this was achieved by direct sequencing (Wakim et al., 1987) and in rat (Basi et al., 1987; Karns et al., 1987; Nielander et al., 1987; Rosenthal et al., 1987; Changelian et al., 1990), human (Ng et al., 1988; Kosik et al., 1988), mouse (Cimler et al., 1987), goldfish (Labate and Skene, 1989), and chicken (Baizer et al., 1990) by molecular cloning. So far, the cDNAs reported for one species are identical within the coding region.

The B-50 open reading frame contains two potential initiation codons. In all species studied, the first codon conforms better to the rules formulated by Kozak (1989) for eukaryotic translational start sites than the second (at position 5). In addition, evidence obtained *in vitro* (Nielander et al., 1990) and *in vivo* (Basi et al., 1987) suggests that in rat brain, translation starts at the first of the two methionines. Thus, it seems unlikely that the purified B-50 forms that lack the first four residues (Basi et al., 1987; Rosenthal et al., 1987; McMaster et al., 1988; Baudier et al., 1989) represent primary translation products.

It turned out that B-50 is an extremely hydrophilic protein of 226–247 amino acids, dependent on the species. There are no potential membrane-spanning domains in the molecule; a small amino-terminal region forms the only hydrophobic part of B-50. The primary sequence is well-conserved between species; it shows an abundance of charged, mainly acidic residues, a high percentage of alanine and proline, and an almost complete absence of residues with large nonpolar or aromatic side chains.

## **Secondary Structure**

Circular dichroism studies of the bovine protein have indicated that the secondary structure of the protein in the presence of 0.05% Lubrol PX is mainly random coil, with some  $\beta$ -sheet and almost no  $\alpha$ -helix (Masure et al., 1986). Based on sucrose density gradient sedimentation and fluorescence polarization measurements, an

axial ratio of approx 16:1 was reported (Masure et al., 1986). These properties are in accordance with the primary structure; for instance, the relative abundance of proline residues would disrupt potential  $\alpha$ -helices. The high density of charged residues, the hydrophylicity, and the lack of large, nonpolar side groups promoting internal folding predicts a highly extended structure (Labate and Skene, 1989). Proton nuclear magnetic resonance (NMR) studies gave no indications of ordered secondary structure in B-50 (Coggins et al., 1989).

Protein B-50 shows anomalous behavior in SDS-PAGE. Comparison with proteins showing a similar anomaly (Kleinschmidt et al., 1986; Takano et al., 1988; Graff et al., 1989a) reveals as a common feature the highly charged character, which could reduce SDS binding to the protein. Also, B-50 has been found to bind SDS poorly (Benowitz et al., 1987). The finding that different B-50 peptides show a similar anomaly in SDS-PAGE (Nielander et al., 1991) is in accordance with this explanation, since charged residues are spread over the whole molecule.

### Homology to Other Proteins

Recently, a 17-kDa neuron-specific protein designated as neurogranin, p17, or RC3 has been characterized as containing some strikingly conserved amino acid sequences with respect to B-50 (residues 1-4 and 34-52) (Represa et al., 1990; Watson et al., 1990; Baudier et al., 1991; Coggins et al., 1991). This might suggest an evolutionary link between neurogranin and B-50, particularly because both are neuron-specific PKC substrates expressed in dendritic and neurite extensions, respectively.

A similarity in amino acid composition but not so much in sequence is found with the particle-bound cytosolic protein CAP-23. This protein was recently discovered by virtue of the similarity of its physical properties (anomalous SDS-PAGE behavior) and subcellular distribution (membrane cytoskeleton-associated) to that of B-50 (Widmer and Caroni, 1990).

## Purified Protein and Primary Translation Product

### Isolation Procedures

Although it is mainly membrane-bound, B-50 is also found in soluble fractions during purification (Cimler et al., 1985; Skene and Virág, 1989; Changelian et al., 1990; Moss et al., 1990; Costello et al., 1991). Several purification procedures for B-50 make use of the membrane association of the protein, starting with a synaptosomal membrane preparation. The protein is easily extracted from the membrane fraction at high pH (Oestreicher et al., 1983) but not with high salt or ethylenediaminetetraacetic acid (EDTA; Skene and Willard, 1981), agents that usually remove nonintegral membrane proteins, although a combination of salt and detergent can release B-50 to some extent (Rodnight, 1982; Zwiers et al., 1979, 1980). Moss et al. (1990) and Meiri and Gordon-Weeks (1990) reported that a considerable part of B-50 from chick and rat brain, respectively, is contained in membrane skeleton preparations, insoluble even at very high concentrations of nonionic detergents.

For further purification of B-50, use has been made of several of its remarkable properties: its extreme acidity for purification from isoelectric focusing (IEF) gels, its resistance to heat and alkali (probably owing to its hydrophylicity), its solubility in 2.5% perchloric acid (a property shared with several other PKC substrates; Baudier et al., 1989), and the specific binding of CaM at low  $Ca^{2+}$ , for purification with CaM-Sepharose columns (Andreasen et al., 1983).

### Posttranslational Modifications

To date, the only known *in situ* posttranslational modifications of B-50 concern the addition of fatty acid chains to Cys residues (Skene and Virág, 1989), the phosphorylation by PKC (Aloyo et al., 1983) of Ser residues (Chan et al., 1986; Alexander et al., 1987). Posttranslational modifications seem to have little effect on electrophoretic behavior of purified B-50 in SDS-PAGE: The pri-



mary translation product of B-50, obtained by cell-free translation (Nielander et al., 197, 1990), and B-50 obtained by means of an *Escherichia coli* expression system (Au et al., 1989) comigrate in SDS-PAGE with the purified protein. Recently, however, a separation of phospho- and dephospho-B-50 by one-dimensional SDS-PAGE has been reported (Meiri et al., 1991).

### Microheterogeneity

Isoelectric focusing in a narrow pH range reveals 3–4 spots for B-50 (Zwiers et al., 1985; Oestreicher et al., 1989; Skene and Virág, 1989; Nelson et al., 1989; Deloulme et al., 1990). A remarkable heterogeneity in purified protein preparations is also encountered in other chromatographic separations (McMaster et al., 1988; Baudier et al., 1989; Schotman et al., 1989). This points to co- or posttranslational modifications, although some of the heterogeneity might be explained by interactions of the highly charged B-50 with chromatographic matrices (Schotman et al., 1989) or by the amphiphatic character of the protein showing an ion dependency for its division into aqueous and detergent phases (Basi et al., 1987; Dosemeci and Rodnight, 1987; Perrone-Bizzozero et al., 1988; Schotman et al., 1989; Skene and Virág, 1989).

The very large isoforms found on gel filtration might be caused by multimer formation (Chan et al., 1986; Masure et al., 1986; Benowitz et al., 1987; Schotman et al., 1989). There are also indications that B-50 is complexed to other proteins in the cell. After detergent extraction from the membrane, during a number of chromatographic purification steps, B-50, PKC, and phosphatidylinositol 4-phosphate (PIP) kinase remained in one complex (Zwiers et al., 1980; Aloyo et al., 1983; Van Dongen et al., 1985). Indeed, B-50 appears to be part of a large hydrophilic complex (Chan et al., 1986; Benowitz et al., 1987).

With respect to the observed heterogeneity of phosphorylated B-50, it has been argued (Schotman et al., 1989) that this cannot be fully

explained by different phosphorylation states mediated by PKC and phosphatases, in conjunction with fatty acylation of Cys3 and -4, because only one phosphorylation site for PKC has been found. This suggests the existence of additional modifications or interactions of B-50. No evidence could be obtained for any *N*-linked glycosylation of B-50 (Skene and Willard, 1981; Zwiers et al., 1985; Masure et al., 1986; Benowitz et al., 1987). Autophosphorylation of B-50 is unlikely since the primary structure of B-50 contains no potential ATP-binding sites. Obviously, membrane-associated and soluble B-50 might represent different isoforms of the protein. If not artifacts of the isolation procedure, B-50 isoforms lacking the first four residues (the sole hydrophobic part of the molecule) are good candidates for a soluble protein fraction (Skene and Virág, 1989). Products of B-50, resulting from specific proteolysis by brain enzymes, are B-60 (B-50 41-226) and a phosphorylation-inhibiting peptide (B-50 1-40; Zwiers et al., 1982; Zwiers and Coggins, 1990), and a 40-kDa protein (B-50 1-203; Oestreicher et al., 1989).

In metabolic labeling studies, the onset of charge heterogeneity was observed only after several hours, suggesting that the heterogeneity is caused by a modification that does not take place in the cell body where B-50 is translated (Skene and Virág, 1989). Interestingly, B-50 from cultured oligodendrocytes migrates as a single spot in IEF gels (Deloulme et al., 1990). This raises the question of whether there is a neuron-specific mechanism behind the heterogeneity observed in neuronal cells.

### Domains in B-50

Current knowledge of the protein structure and organization of the B-50 gene enables the assignment of certain functions to specific domains of the molecule. The main functional domains are contained by the amino terminal part, residues 1–56; those include sites for membrane association, phosphorylation by PKC, and CaM binding (Fig. 1).




Fig. 1. Functional domains in rat B-50 alias GAP-43 alias F1 alias neuromodulin. PKC, protein kinase C; CaM, calmodulin; P, phosphate; and PS, phosphatidylserine, indicate proposed sequence for CaM or neurofilament interaction. Numbers refer to rat B-50 amino acid sequence.

### Gene Structure

The protein is encoded by three exons in the rat (Grabczyk et al., 1990) and human gene (Schrama et al., 1990). Often, exons correspond to functional domains in a protein (Gilbert, 1986). The first exon encodes the amino-terminal 10 residues of B-50, responsible for the membrane association of the protein (Zuber et al., 1989a). The major part of B-50, including CaM-binding and phosphorylation sites, originates from a 568 or 595bp exon (in rat and human, respectively); the last 27 or 28 amino acids are encoded by a third exon.

Based on comparison of primary sequences of a number of mammalian species and goldfish, a division was made by Labate and Skene (1989) in a highly conserved *N*-terminal sequence of 57 residues and a large *C*-terminal part with a marked conservation in amino acid composition. Phylogenetic comparison has been used to locate sequences of possible interest, especially in the *C*-terminal part (Labate and Skene, 1989; Moss et al., 1990).

The distribution of prolines and cysteines in a protein is of special importance. Disulphide bridges between cysteines are main determinants

of tertiary structure, and clusters of prolines, for instance, can delineate functional domains. The *N*-terminal 59 residues of B-50 contain only one proline at position 27, whereas the remainder of B-50 is proline-rich, with an average distance of 10 residues and one adjacent pair of prolines at positions 184 and 185. (Residue numbering is based on the rat amino acid sequence; in other species, the distribution is very similar.)

### Membrane Association

The small amino-terminal hydrophobic part of B-50 (Met-Leu-Cys-Cys-Met) seems insufficient to explain the association of B-50 with the membrane. Many eukaryotic proteins contain covalently attached lipid, e.g., fatty acid, bound by amide, thioester, or *O*-acyl bonds (Schmidt, 1983; Low, 1987; Magee and Hanley, 1988). A remarkable feature of the B-50 amino-terminus is the presence of two adjacent cysteines (residues 3 and 4), the sole ones in the protein. Skene and Virág (1989) presented evidence for palmitoylation of B-50, presumably through a thioester bond at Cys3 and/or -4. Soon after synthesis in the cytosol, the protein becomes membrane-associated (Skene and Virág, 1989). The group of Benowitz could not detect acylation of B-50 in cultured embryonic neurons (Perrone-Bizzozero et al., 1989). The data from Skene and Virág (1989) suggest that acylation of B-50 is a dynamic process; in isolated growth cones, labeled palmitic acid was incorporated into B-50. For other proteins known to contain covalently bound fatty acids, often the attached lipid does not seem to be responsible for direct anchoring of the protein to the membrane (Schmidt, 1983; Low, 1987). The punctuate staining for B-50 observed in growing axons (Meiri et al., 1988) seems inconsistent with a direct interaction of B-50 with the lipid bilayer. Like B-50, the neural-specific protein SCG10 is tightly membrane-associated but lacks a membrane-spanning hydrophobic segment; it also contains two cysteines at positions 22 and 24. The intracellular localization of SCG10 resembles that of B-50, including the punctuate staining in growth cones (Stein et al., 1988).

The B-50 amino terminus was found to be blocked for direct amino acid sequencing (Rosenthal et al., 1987; McMaster et al., 1988; P. Schotman and L. H. Schrama, unpublished results), even after alkaline extraction, which largely uncouples acyl chains linked by a thioester bond. This indicates that still other posttranslational modifications may appear at the *N*-terminus. Proton NMR studies indicated the possible attachment of a polyisoprenoid to B-50 (Coggins et al., 1989; Coggins and Zwiers, 1991).

Whatever the exact nature of the modifications at the *N*-terminus, there is convincing evidence that the first 10 amino acids of B-50 contain all the determinants for membrane association. Zuber et al. (1989a) constructed recombinant proteins of the first 10 residues of B-50 with chloramphenicol acetyl transferase, a cytosolic protein. This appeared sufficient to direct the fusion protein to the membrane in transfected "neuronal" (PC12) and nonneuronal (COS) cell lines. Moreover, the fusion protein had the same distribution in PC12 (pheochromocytoma) cells as normal B-50. Similar results in PC12 cells with a fusion of B-50 1-10 to another cytosolic protein,  $\beta$ -galactosidase, were reported by Neve et al. (1990). Mutation of Cys3 and/or -4 in B-50 caused the protein to remain mainly cytosolic (Zuber et al., 1989a), confirming the importance of both residues for membrane targeting.

### Interaction With $G_0$ Protein

Trimeric GTP-binding (G) proteins transduce information from transmembrane receptors to intracellular systems (Gilman, 1987). The most abundant G protein in brain is  $G_0$ , a major growth cone protein. Its exact role is not clear, but it has been implicated in regulation of  $Ca^{2+}$  channels and phospholipase C activity (Banno et al., 1987; Brown and Birnbaumer, 1990).

Recently, it was shown that B-50 can regulate the binding of GTP- $\gamma$ -S to  $G_0$  (Strittmatter et al., 1990). The first 24 residues of B-50 stimulated GTP- $\gamma$ -S binding to  $G_0$  to the same level as the whole protein (although much higher concentrations of the peptide were needed), whereas the

first 10 amino acids did not stimulate this binding but competed with B-50 for binding to  $G_0$ . Apparently, both exon 1 (residues 1-10) and exon 2 encode information important for interaction with  $G_0$ . Sequence comparison of B-50 with domains of G-linked transmembrane receptors proposed to be involved in  $G_0$  activation reveals a homologous sequence with the consensus amino acid sequence hydrophobic-Leu-Cys-Cys-X-basic-basic (residues 1-7 in B-50; Strittmatter et al., 1990). In the three cases reviewed by Strittmatter et al. (1990), Cys residues in the transmembrane receptor proteins were subject to fatty acid acylation. Since the experiments of Strittmatter et al. were performed with alkali-extracted, presumably, nonpalmitoylated B-50, fatty acid acylation of B-50 may not be essential to the interaction of B-50 with  $G_0$ .

The nature of the proposed interaction and its physiological relevance remain to be established. Strittmatter et al. did not succeed in isolation of a  $G_0$ /B-50 complex, and modulation of GTPase activity by B-50, the physiological parameter for  $G_0$  function, has to our knowledge not been shown yet.

### Phosphorylation by PKC

In general, PKC plays an important role in cellular responses to extracellular stimuli (Miller, 1986; Nishizuka, 1988,1989). In the nervous system, B-50 is a major substrate of PKC. Initially, it was found that in vitro phosphorylation of B-50 was inhibited by adrenocorticotrophic hormone ACTH1-24 (Zwiers et al., 1976, 1978). Subsequent studies showed a correlation of the phosphorylation state of B-50 with the activity of a PIP kinase (Van Dongen et al., 1985; Gispén et al., 1985b, 1986; Van Hooff et al., 1988), LTP (Lovinger et al., 1986; Benowitz and Routtenberg, 1987; De Graan et al., 1988a), and the modulation of neurotransmitter release (Dekker et al., 1989a,b, 1990a,b; Heemskerk et al., 1990; De Graan et al., 1991).

Evidence from several laboratories suggested that Ser<sup>41</sup> might be the unique phospho site for PKC (Coggins and Zwiers, 1989; Nielander et al.,

1989; Apel et al., 1990). The fact that mutation of Ser<sup>41</sup> into Thr<sup>41</sup>, Ala<sup>41</sup>, or Gly<sup>41</sup> prohibited phosphorylation by PKC *in vitro* seems to prove this (Nielander et al., 1990; Chapman et al., 1991). Proteolytic digestion of *in situ* phosphorylated B-50 resulted into the same phospho-fragments obtained following *in vitro* phosphorylation by PKC, indicating that Ser<sup>41</sup> is of physiological significance (Van Hooff et al., 1988; Heemskerk et al., 1989; Oestreicher et al., 1989).

Phosphate is incorporated into the more acidic forms obtained by IEF and preferentially into the most acidic isoform of B-50 (Zwiers et al., 1985; Skene and Virág, 1989; Deloulme et al., 1990). Since there is a single phosphate acceptor site (Ser<sup>41</sup>), this preference for one of the isoforms suggests that posttranslational modifications of B-50 (*see* Membrane Association, p6) can regulate phosphorylation by PKC (Deloulme et al., 1990). Dephosphorylation of B-50 can be mediated by several phosphatases *in vitro* (Liu and Storm, 1989; Schrama et al., 1989; Dokas et al., 1990). The physiological relevance of that is as yet unclear.

Phosphorylation of B-50 by protein kinases other than PKC is still under study. Aloyo et al. (1983), Akers and Routtenberg (1985), and Chan et al. (1986) found no significant *in vitro* phosphorylation of B-50 from adult rat brain with cAMP-dependent kinase and Ca<sup>2+</sup>/CaM kinase I and II. Nevertheless, Katz et al. (1985) reported phosphorylation by a Ca<sup>2+</sup>/CaM-dependent kinase in growth cones. More recently, it has been shown that casein kinase II can phosphorylate B-50 *in vitro*, presumably at Ser<sup>192</sup> (rat: Pisano et al., 1988; mouse Ser<sup>193</sup>: Apel et al., 1991). To date, however, PKC is the only kinase documented to play a role in phosphorylation of B-50 *in situ*.

It remains to be shown to what extent the population of B-50 proteins is phosphorylated during different stages of neuronal maturation or in different subcellular compartments. Phosphorylation by PKC, as judged from the onset of appearance of the more acidic forms of B-50 on IEF in metabolic labeling studies, does not take place until several hours after synthesis in the cell body (Skene and Virág, 1989). From a study per-

formed with monoclonal antibodies (MAb) recognizing specifically the phospho-form, it was concluded that not only is the phosphorylated form spatially restricted to the distal axon and growing tip but that phosphorylation might occur subsequent to the onset of axonogenesis (Meiri et al., 1991).

### CaM Binding

The protein B-50 interacts with CaM in an unusual way. It binds to CaM-Sepharose at low Ca<sup>2+</sup> conditions and can be released by raising the Ca<sup>2+</sup> concentration (Andreasen et al., 1983), whereas most other CaM-binding proteins show a Ca<sup>2+</sup> dependency for CaM binding. The interaction of neuromodulin (B-50) with CaM has been studied in detail by the group of Storm (Liu and Storm, 1990; Estep et al., 1990; Chapman et al., 1991). Also, B-50 crosslinks to a CaM derivative in a 1:1 molar ratio (Andreasen et al., 1983), indicating that it has a single CaM-binding site. In accordance with its unusual CaM-binding characteristics, the structural properties of B-50 distinguish it from CaM-binding regions in other proteins (mainly basic amphiphilic  $\alpha$ -helical peptides (O'Neil and Degrado, 1990). Using proteolytic fragments and synthetic peptides of B-50, it was shown that the CaM-binding site encompasses residues 43–51 (Alexander et al., 1988). Interestingly, this domain weakly satisfies the proposed criteria for a "normal" CaM-binding site; it is the only region carrying a net positive charge, and it contains the only aromatic residue in B-50.

The interference of phosphorylation by PKC with CaM binding (Alexander et al., 1987; Graff et al., 1989a) suggests that Ser<sup>41</sup> forms part of the binding site. Interference by a phosphate group may be caused by steric hindrance, the introduction of extra negative charges (mimicked by mutating Ser<sup>41</sup> into Asp<sup>41</sup> (Chapman et al. 1991), an effect on the adjacent Phe residue, or an alteration of secondary structure, leading to a decreased affinity for CaM.

A comparable interaction between CaM binding and PKC-mediated phosphorylation is



observed for the myristoylated alanine-rich C kinase substrate (MARCKS) or 87-kDa protein, a protein of unknown function (Graff et al., 1989b). This protein shares other characteristics with B-50, such as enrichment in growth cones, acidity, and anomalous electrophoretic behavior in SDS-PAGE. Its interaction with CaM, however, is  $\text{Ca}^{2+}$ -dependent, and MARCKS is not restricted to the nervous system (Albert et al., 1987; Graff et al., 1989b).

The reduced CaM binding reported for B-60 (amino acids 41–226) suggests a possible involvement of sequences N-terminal to Ser<sup>41</sup> in the interaction of B-50 and CaM (Coggins and Zwiers, 1990)—more explicitly residues 38–40, since a synthetic peptide 38–55 bound CaM with an affinity comparable to native B-50 (Alexander et al., 1988).

Two other proteins have been described to bind to CaM-Sepharose with characteristics similar to B-50. The rat adipocyte protein pp170 (McDonald and Lawrence, 1989) and a protein of 16 kDa found in rat brain and SPM that is alkaline-resistant and heat-stable as well (Schrama et al., 1989), recently characterized as p17/neurogranin (Baudier et al., 1991), BICKS (B-50-immunoreactive C-kinase substrate, Coggins et al., 1991), or RC3 (Watson et al., 1990). Both proteins are also subject to phosphorylation by PKC.

Alexander et al. (1987) observed in vitro a reduced affinity of B-50 for CaM with increasing ionic strength. This and the structure of the proposed CaM-binding domain suggest that the B-50–CaM interaction is primarily ionic in nature (Alexander et al., 1988). The dependency on low calcium concentrations and the effects of ionic strength in vitro have raised questions on the physiological significance of the CaM–B-50 interaction (Skene 1989; Coggins and Zwiers, 1991). However, in native synaptosomal and growth cone membranes, complex formation between CaM and endogenous B-50 was shown with the crosslinker disuccinimidyl suberate (DSS) under physiological  $\text{Ca}^{2+}$  concentrations (De Graan et al., 1990). Moreover, the  $\text{Ca}^{2+}$  sensitivity of com-

plex formation was not decreased by raising the ionic strength (P. N. E. De Graan, unpublished results) as described for the in vitro system (Alexander et al., 1987). Recently, Houbre et al. (1991) demonstrated that the conserved amino acid sequence in B-50 and neurogranin that corresponds to the PKC phosphorylation site and CaM-binding domain is essential for the binding to phosphatidylserine, which mediates the phosphorylation by PKC.

### Interaction with Cytoskeletal Elements

The region C-terminal to residue 57 shows a sharp divergence in primary sequence between rat and fish (Labate and Skene, 1989). Several short regions of apparent sequence conservation between fish and mammalian species, potentially involved in interactions with other proteins, have been noted in this part of B-50 (motifs A–F; Labate and Skene, 1989). In motif B, repeated twice in the rat and human, the sequence (SEEK) is found noted for its similarity to the PKC phosphorylation site of the  $\alpha$ -subunit of the nicotinic acetylcholine receptor (Nielander et al., 1987). In rat B-50, however, there seems to be no significant phosphorylation by PKC of these sites (Nielander et al., 1990).

Sequence motif C (ETXESSQ) contains the postulated site for phosphorylation by casein kinase II (rat residue 192, mouse residue 193; Pisano et al., 1988; Apel et al., 1991). Indeed, the region containing motif C and part of motif F shows some homology to a casein kinase consensus sequence in the *c-myc* proto-oncogene product *myc* (Coggins and Zwiers, 1991). Ng et al. (1988) noted several internally homologous direct repeat regions in the human B-50 cDNA, forming a contiguous domain near the COOH-terminal end. In exon 3, coding for the 28 amino acids at the C-terminus, some structural similarity was found for goldfish B-50 and a mouse neurofilament (NF-L).

This homology might be responsible for the observed immunological crossreactivity between some B-50 antisera and NF-L proteins (G. J. A.



Ramakers, personal communication). The homologous sequence is located in a part of NF-L thought to be involved in interaction with other cytoskeletal elements. This structural similarity has led to the proposition that B-50 might interact with skeletal elements through its C-terminal domain (Labate and Skene, 1989).

Although there is little sequence homology, the amino acid composition of the large C-terminal part (residues 57–226 in rat) is strongly conserved, and such predicted physical characteristics as hydrophobicity and charge are very similar between species (Labate and Skene, 1989). The functionally important feature, which apparently can be accommodated by quite-relaxed sequence requirements, might be the extended, negatively charged rod-like structure (Labate and Skene, 1989). Recently, B-50 has been proposed to extend from the cytoplasmic side of the membrane inward (Gispen et al., 1985a; Meiri et al., 1988). With the amino terminus involved in membrane association and an extended structure, this would bring the carboxyl-terminus away from the membrane in the cytosolic space. The recent characterization of B-50 as a component of the neuronal membrane skeleton in chicken and rat (Allsopp and Moss, 1989; Moss et al., 1990; Meiri and Gordon-Weeks, 1990) is consistent with such a view.

## Functional Aspects of B-50

Posttranslationally, B-50 undergoes several modifications. Some of these, i.e., phosphorylation and palmitoylation, are reversible, involving fast-working enzymes. Specific proteolytic cleavage, e.g., C-terminally to residues 4 and 41, might provide alternative, irreversible ways to modify the protein. The general picture arises of a series of B-50 isoforms, with various types of modifications, perhaps related to differential localization within the cell. These different isoforms are capable of interacting with a number of other functional proteins, some of which are thought to be part of a multifunctional B-50-containing protein complex at the inner surface of the plasma membrane.

## Role in Neuritogenesis

A multitude of functions has been proposed for B-50 in neuronal growth, receptor-mediated processes, and transmitter release. So far, a unifying concept is missing. The best documented role is in neuronal growth (for refs., see Introduction), albeit that a causal relationship between growth and B-50 expression is still under debate (Gordon-Weeks, 1989; Megerian and Klein, 1990; Yankner et al., 1990; Baetge and Hammang, 1991; Costello et al., 1991). Important arguments in favor of such causal relationships are the following:

1. Transfection of COS cells with an expression vector containing this B-50 gene leads to formation of growth cone-like filopodia (Zuber et al., 1989b);
2. Transfected PC12 cells that overexpress B-50 display enhanced responsiveness to nerve growth factor (NGF) and regeneration of damaged neurite-like extensions (Yankner et al., 1990); and
3. In rat, PC12 cells antisense oligomers to B-50 blocked the NGF-induced increase in B-50 expression and at the same time, the enhanced neurite outgrowth (Schotman et al., 1990).

Future research should shed light on the precise role of B-50 in neuritogenesis.

## Role in NGF-Induced Neuronal Differentiation of PC12 Cells

Pheochromocytoma cells have been intensively studied as a model for the mechanism of neuronal differentiation. When rat PC12 cells are cultured in the presence of NGF, the cells stop proliferating and differentiate to a phenotype resembling cholinergic sympathetic neurons, elaborating neuritic extensions (Greene and Shooter, 1980). Also, B-50 has been identified in undifferentiated and NGF-treated PC12 cells (Van Hooff et al., 1989b). Treatment of PC12 cells

with NGF for 2 d results in a 2.5-fold increase in the B-50 level as measured by radioimmunoassay (RIA). Immunofluorescence microscopy shows that B-50 is detectable in fixed permeabilized PC12 cells, as intensified membrane-associated staining in the neurites, especially in the growth cones, of the NGF-differentiated cells and as diffuse staining in the cell bodies of the controls and the NGF-differentiated cells. In contrast, when living, nonpermeabilized PC12 cells are incubated with B-50 antibodies, then fixed and probed with a fluorescein-conjugated secondary antibody, no immunofluorescence is detectable, suggesting that B-50 is inside the cells.

### ***Translocation of B-50 with NGF-Induced Neuritogenesis***

With the aim to learn more about the putative growth-associated function of B-50, we have compared its ultrastructural localization in control proliferating PC12 cells, with its distribution in PC12 cells induced to differentiate by various agents (50 ng/mL NGF, 1 mM dibutyryl cAMP and substratum). The studies were performed on cells fixed in 2% paraformaldehyde/0.1% acrolein in PBS, embedded in gelatin, cryoprotected, and sectioned on an ultracycromicrotome at  $-90^{\circ}\text{C}$ . Grids holding 70–100 nm thick sections were incubated with B-50 antibodies and the antigen-antibody complexes were detected by protein-A gold (approx 10 nm) conjugates (Van Hooff et al., 1989a).

To our surprise, we found that in the cryosections of untreated control PC12 cells, the B-50 immunoreactivity (BIR) is mainly associated with intracellular membrane-rich and vesicular structures, including organelles of the lysosome family and Golgi apparatus. The plasma membrane is virtually devoid of label. In contrast, after 48 h of treatment with NGF or cAMP, BIR is most pronounced on the plasma membrane. The highest BIR is observed on plasma membrane surrounding sprouting microvilli, lamellipodia, and filopodia.

In addition, extending neurites and growth cones contain cytosolic BIR, which is partially

associated with chromaffin granules. In the NGF-differentiated PC12 cells, the cytosolic location of B-50 is similar to untreated cells. Although the induction of neurite formation in PC12 cells by NGF and cAMP is reported to occur by different pathways, B-50 distribution in dibutyryl cAMP-differentiated cells closely resembles that observed in the NGF-treated cells. Thus, in PC12 cells, various differentiating agents induce a change in the ultrastructural localization of B-50, facilitating association of the B-50 protein with the plasma membrane (Van Hooff et al., 1989b).

Recently, further ultrastructural studies of the time course of the NGF effect on B-50 distribution indicate that after approx 4 h, the beginning of the shift of B-50 to the plasma membrane can be observed, and after 8 h, the translocation appears to be fully established. When PC12 cells are grown for the same time (48 h) in the presence of epidermal growth factor (EGF, 50 ng/mL), a growth factor inducing proliferation in PC12 cells, the cell number increases, the B-50 levels increase by 50%, but few neuritic protrusions are formed. The ultrastructural distribution of B-50 in the case of EGF is rather similar to that in the untreated control cells (A. B. Oestreicher, unpublished results). This indicates that the translocation of B-50 appears to be linked to initiation and formation of neuritic protrusions. Below we discuss that after translocation of B-50 to the plasma membrane as an initial step in the neuronal differentiation, there follow stages of development in which the neuron undergoes polarization, and B-50 becomes restricted to the axonal domain.

### ***Localization in Growth Cones***

The subcellular distribution of B-50 in growth cones was studied by immunogold labeling of B-50 antibodies in ultrathin cryosections of isolated neuronal growth cones (Pfenninger, 1986). Ultrastructure of isolated neuronal growth cones clearly shows a dense matrix of smooth endoplasmic reticulum, round and oval vesicles of various sizes, and small mitochondria. Eighty percent of the B-50 immunoreactivity is located at the plasma membrane, whereas 20% is present

in the cytoplasm of the growth cone, occasionally in the proximity of vesicles (Van Lookeren Campagne et al., 1990). In other electron microscopic immunocytochemical studies on hippocampus (Van Lookeren Campagne et al., 1990) and pyramidal tract axons (Gorgels et al., 1989) during postnatal development of rat brain, we found that B-50 is predominantly located at the plasma membrane of growth cones *in situ*.

### Changes in B-50 Distribution During Neuronal Polarization

During development of the nervous system, neurons go through a developmental stage (polarization) in which neurons form two distinct classes of morphological and functional different processes: the axon and the dendrite. In studies of neuronal cell cultures dissociated from embryonic brain (Meiri et al., 1986,1988; Goslin et al., 1988; Burry et al., 1991; Ramakers et al., 1991), it has been observed that during the first days *in vitro*, the neurons produce short neurites with growth cones, both detectable by dense B-50 immunostaining. As the neurons develop, mature, and make connections, B-50 immunostaining is reduced in the processes and in some cases disappears completely.

For the development of embryonic hippocampal neurons in cell culture, Dotti et al. (1988) have documented the sequence of changes in neurite outgrowth and morphology, leading to polarization and maturation of the neurons. Goslin et al. (1988,1990; Goslin and Banker, 1990) showed by immunofluorescence that after morphological polarization of the pyramidal neurons, B-50 becomes selectively located in the axonal growth cone. In addition, B-50 is demonstrated to be absent from dendrites and their growth cones, identified by microtubule-associated protein-2 (MAP-2), a cytoskeletal marker for dendrites (Dotti et al., 1987).

Our immuno-electron microscopic studies of hippocampal neurons in cell culture at consecutive developmental stages show that B-50 immunoreactivity becomes restricted to the axonal domain, being located at the plasma membrane

of varicosities, some parts of the shaft and terminals. A condition for this restriction appears to be that the neurons make contacts and form synapses (Van Lookeren et al., 1991b; Van der Neut et al., 1990).

Evidence for this suggestion is the finding that polarized neurons with a long, thin MAP-2 negative axon grown solitary for more than 1 wk *in culture* display B-50 immunoreactivity at the plasma membrane of the MAP-2 positive dendrites and growth cones. Since the anti-B-50 antibodies used do not crossreact with neurogranin (A. B. Oestreicher, unpublished results), the immunoreactivity represents only B-50 protein. Hippocampal neurons cultured for 2–3 d *in vitro* initiate polarization by rapid outgrowth of one longer process in comparison to the other shorter, slower growing neurites. At this stage of development, we have detected B-50 immunoreactivity in all processes by immunofluorescence and ultrastructurally to be homogeneously distributed at the plasma membrane of growth cones, the axon, the dendrites, and the cell body. These results suggest that B-50 is not involved in functions exclusively specific for the growth cone but rather is functionally important for the neuronal membrane in general (*see below*).

In connection with the suggestion that B-50 is not specific for the growth cone, it is relevant to note that in the adult animal, dense B-50 immunolabeling is also detected at the plasma membrane of axon terminals and at that of unmyelinated thin axons in various regions (hippocampus, periaqueductal gray, olfactory system) of the central nervous system (CNS) (Fig. 2; Van Lookeren Campagne et al., 1990,1991a,b) and only little B-50 immunoreactivity in myelinated axons (Gorgels et al., 1989). Of interest is that in the mature CNS, B-50 immunoreactivity has been rarely detected in dendrites, although DiFiglia et al. (1990) have detected B-50 immunoreactivity not just in unmyelinated fibers and axon terminals but also in dendritic spines of the neuropil of rat neostriatum.

A recent study by Meiri et al. (1991) using two MAb to B-50 with distinct different specificity, one recognizing selectively the PKC-phosphory-




Fig. 2. Electron micrograph of apical dendrites, proximal to the pyramidal cells of the CA1 stratum radiatum in a P90 rat. The sections were immunostained for B-50 by preembedding immunoperoxidase labeling (A) or by postembedding immunogold labeling (B). Note the absence of immunoreactivity in the longitudinally-sectioned apical dendrites (Den<sub>1</sub>), clearly showing microtubules (arrow heads A, B). In the preembedding immunostained section, axon terminals (At) are most prominently stained, whereas in the postembedding immunogold (10 nm) labeled section, plasma membranes of all axon shafts (Ax) are labeled for B-50, in addition to those of the axon terminals. Sp, dendritic spine; mit, mitochondrion. Asterisks in A show unlabeled axon terminals. Scale bar: 0.5  $\mu$ m.

lated isoform of B-50 and the other reacting with both the PKC-phosphorylated and nonphosphorylated B-50, shows that during development of nervous tissue, the ratio of these forms is changed, and PKC-phosphorylated B-50 is restricted temporally and spatially to the distal

part of the outgrowing axons. Taken with recent ultrastructural data, one of the implications is that the functional state of B-50 is determined locally by the steady-state level of calcium and the dynamic balance between the activities of PKC, protein phosphatases, and other enzymes.



### **Role of B-50 Phosphorylation in Signal Transduction in the Growth Cone**

Nerve growth cone membranes isolated from fetal and neonatal rat brain are enriched in B-50 (De Graan et al., 1985; Katz et al., 1985). Phosphorylation of B-50 in these growth cone membranes *in vitro* is inhibited by ACTH<sub>1-24</sub> and stimulated by phorbol diesters, indicating that PKC is the active protein kinase. In addition, ACTH<sub>1-24</sub> stimulated the phosphatidylinositol 4,5-bisphosphate (PIP<sub>2</sub>) formation in growth cone membranes, suggesting the possibility of an inverse relationship between the degree of B-50 phosphorylation and PIP<sub>2</sub> labeling as shown for adult presynaptic membranes (Van Hooff et al., 1988).

An important question is whether extracellular signals can influence B-50 phosphorylation in intact nerve growth cones. To explore this, intact growth cones were isolated from the brain of 5-d-old rats and studied after prelabeling with [<sup>32</sup>P]-orthophosphate, using an immunoprecipitation method to selectively monitor the degree of B-50 phosphorylation. Phorbol 12,13-dibutyrate and dioctanoyl glycerol, activators of PKC, stimulate B-50 phosphorylation in the intact growth cones, confirming that PKC is involved in B-50 phosphorylation (Van Hooff et al., 1989a).

Depolarization induced by 30 mM K<sup>+</sup> produces a transient rise in B-50 phosphorylation. This response to K<sup>+</sup> depolarization is also demonstrated in adult rat brain preparations, such as the hippocampal slice (Dekker et al., 1989b) and isolated synaptosomes. The effect of K<sup>+</sup> depolarization on B-50 phosphorylation in growth cones can be blocked partially by atropine (0.1–1 mM). The suggestion of involvement of muscarinic receptors is supported by the finding that carbachol, a cholinergic receptor agonist, enhances B-50 phosphorylation in a concentration-dependent manner (50% at 1 mM). The carbachol (1 mM) stimulation can be blocked by atropine (0.1 μM). The carbachol stimulation can be further increased by concurrent K<sup>+</sup> depolarization (Van Hooff et al., 1989a,b). It is not yet clear how the receptor-

mediated changes in B-50 phosphorylation in isolated growth cones relate to modulation of GTP-γ-S binding to G<sub>0</sub> (Strittmatter et al., 1990).

### **B-50 Phosphorylation and LTP**

To investigate the role of B-50 phosphorylation in long-term potentiation (LTP), Routtenberg and co-workers (Linden and Routtenberg, 1989) elicited LTP by *in vivo* tetanization and subsequently measured B-50 phosphorylation using [γ-<sup>32</sup>P]-ATP. They found that tetanization is associated with an increase in post hoc B-50 phosphorylation and membrane-bound PKC activity. Using [<sup>32</sup>P]-orthophosphate labeling combined with immunoprecipitation of B-50, we measured B-50 phosphorylation in intact hippocampal slices after short-term phorbol ester treatment (De Graan et al., 1988b). The phorbol ester induced a long-lasting enhancement of synaptic efficacy resembling in many ways tetanus-induced LTP, which was accompanied by a long-lasting increase in B-50 phosphorylation (Fig. 3).

Recently, Gianotti et al. (1991) showed that in hippocampal slices (using our immunoprecipitation procedure), LTP is accompanied by an increase in B-50 phosphorylation. It is not yet known how long the increase in B-50 phosphorylation persists after tetanization. It is not yet clear whether B-50 phosphorylation is involved in the presynaptic mechanism underlying LTP. An intriguing possibility is that a long-term increase in B-50 phosphorylation is related to the prolonged increase in neurotransmitter release that has been reported to occur after tetanization (Dolphin et al., 1982).

### **Role of B-50**

#### **in Presynaptic Signal Transduction and Neurotransmitter Release**

In view of the possible involvement of PKC in transmitter release and of the presynaptic localization of the PKC substrate B-50, we started a series of studies to investigate the relationship between B-50 phosphorylation and transmitter release.






Fig. 3. Long-lasting effects of a 20-min phorbol dibutyrate (PDB) treatment on population spike (PS) amplitude B-50 phosphorylation. After treatment with  $10^{-6}$ M PDB (●) or  $4\alpha$ -phorbol didecanoate ( $4\alpha$ -PDD)(○), slices were incubated in or perfused with KRB. Field potentials were recorded in the stratum pyramidale of CA1. The average PS amplitude (solid lines, mean  $\pm$  SEM; and  $n = 4$ ) is expressed as a percentage of the PS during the 10-min control period (prior to the phorbol treatment). B-50 phosphorylation (broken lines, mean  $\pm$  SEM; and  $n = 4$ ) is expressed as a percentage of the value obtained for untreated control slices at each time point.

Stimulation of transmitter release from rat brain synaptosomes by high  $K^+$ , 4-amino-pyridine, or veratridine induces a concomitant phosphorylation of B-50 (Dekker et al., 1990a; Heemskerk et al., 1990). We more closely investigated  $K^+$ -induced phosphorylation of B-50. Like  $K^+$ -evoked release,  $K^+$ -evoked B-50 phosphorylation is dependent on extracellular  $Ca^{2+}$  (Dekker et al., 1989a, 1990a). Most likely, both processes are activated by an influx of extracellular  $Ca^{2+}$  into the synaptic terminal.  $K^+$ -evoked B-50 phosphorylation depends on PKC activity because it is inhibited by polymyxin B, H-7, and staurosporine, three inhibitors of PKC (Dekker et al., 1989a, 1990).

In synaptosomes,  $K^+$  and veratridine-induced phosphorylation of B-50 is transient; it is rapidly induced and returns to basal levels within 5 min (Dekker et al., 1990b). Such a time-dependency profile has also been reported for the amount of transmitter released/unit of time from synaptosomes after depolarization and for the amount of  $Ca^{2+}$  entering the synaptosomes for each unit of time (De Graan et al., 1991; Dekker et al., 1991), indicating that  $Ca^{2+}$  influx, PKC-mediated B-50 phosphorylation, and transmitter release may be closely related.

The clear correlation that has been found between B-50 phosphorylation and transmitter release did not answer the question of to what extent B-50 phosphorylation is essential to this process. To address this issue, specific B-50 antibodies that inhibit the phosphorylation of B-50 were tested on transmitter release. Insofar as B-50 is localized at the inner leaflet of the synaptic plasma membrane, exposed toward the cytosol (Gispén et al., 1985a; Van Hooff et al., 1989b), the antibodies needed to be introduced into the synaptosomes without disrupting the release machinery. Anti-B-50 antibodies were introduced using a streptolysin-O (SL-O) permeation technique (Dekker et al., 1989a). The effect of  $K^+$  depolarization on noradrenaline (NA) release can be mimicked in SL-O-permeated synaptosomes by increasing the  $Ca^{2+}$  concentration in the incubation buffer from  $10^{-8}$  to  $10^{-5}$  (Dekker et al., 1989a). The effect of  $Ca^{2+}$  cannot be attributable to an efflux of cytosolic NA, since  $Ca^{2+}$  did not stimulate the SL-O-induced efflux of the cytosolic marker protein lactate dehydrogenase (LDH). This confirms that the  $Ca^{2+}$ -dependent part of SL-O-induced NA efflux is vesicular NA release.

Next, permeated synaptosomes were incubated with anti-B-50 IgGs concomitantly with the  $\text{Ca}^{2+}$  trigger, and NA release and B-50 phosphorylation were measured (Fig. 4).  $\text{Ca}^{2+}$ -induced release of NA was inhibited by the anti-B-50 IgGs in a concentration-dependent way (Dekker et al., 1989a), as was B-50 phosphorylation. Control IgGs and heat-inactivated anti-B-50 IgGs were ineffective. Neither control nor anti-B-50 IgGs affected NA efflux at  $10^{-8}$  and  $10^{-7}$  M  $\text{Ca}^{2+}$ , showing that the antibodies did not influence the  $\text{Ca}^{2+}$ -independent efflux. Surprisingly, we found that in contrast to  $\text{Ca}^{2+}$ -induced NA release, phorbol ester-induced release was not inhibited by the anti-B-50 antibodies. Although we do not currently have an explanation for this, it may indicate that phorbol esters and  $\text{Ca}^{2+}$  have a different effect on transmitter release.

From these studies, we conclude that B-50 plays a crucial role in the molecular mechanism of NA release from rat cerebral cortex synaptosomes. At this moment, it is not clear whether B-50 is involved in the mechanism of release of transmitters other than NA.

Although it has been established that inhibition of B-50 phosphorylation by specific antibodies is accompanied by inhibition of calcium-induced NA release from permeated isolated nerve terminals, this does not imply that only the degree of B-50 phosphorylation is important for NA release. It is also possible that B-50 antibodies block properties of B-50, other than its phosphorylation or in conjunction with its phosphorylation, which are important to transmitter release. In favor of a direct involvement of PKC-mediated phosphorylation of B-50 in transmitter release is the following evidence:

1. Phorbol esters that directly activate PKC and induce B-50 phosphorylation enhance the release of a variety of neurotransmitters, and this enhancement can be antagonized by the kinase inhibitors polymyxin B, staurosporine, and H-7;
2. Both B-50 phosphorylation and neurotransmitter release can be enhanced by high  $\text{K}^+$ , veratridine, and 4-aminopyridine;




Fig. 4. Effect of anti-B-50 IgG on  $\text{Ca}^{2+}$ -induced release of NA from SL-O-permeabilized synaptosomes. Synaptosomes were permeabilized with 0.8 IU/mL SL-O and incubated without IgG (-), with different concentrations of anti-B-50 IgG (anti-B-50), or with control IgG (control) (open bars,  $\square$   $10^{-8}$ ; dotted bars,  $\square$   $10^{-7}$  M  $\text{Ca}^{2+}$ ; filled bars,  $\blacksquare$   $10^{-5}$  M  $\text{Ca}^{2+}$ ). Neither anti-B-50 nor control IgG (20  $\mu\text{g}$ /assay) affected NA release in nonpermeabilized synaptosomes. Data are expressed as SEM (three independent experiments). Identical results were obtained in a separate experiment using a different batch of anti-B-50 IgG. Data were analyzed by one-way analysis of variance (ANOVA) with a supplemental Newman-Keuls test. Asterisk denotes group significantly different ( $p < 0.05$ ) from all groups treated with  $10^{-7}$  or  $10^{-8}$  M  $\text{Ca}^{2+}$ .

3. The protein kinase inhibitor polymyxin B inhibits  $\text{K}^+$ -induced B-50 phosphorylation and  $\text{K}^+$ -induced neurotransmitter release; and
4. Anti-B-50 IgGs inhibit B-50 phosphorylation as well as  $\text{Ca}^{2+}$ -dependent transmitter release.

However, in disagreement with a direct role for PKC-mediated B-50 phosphorylation in neurotransmitter release are the observations that:

1. Phorbol esters enhance B-50 phosphorylation but not transmitter release in polarized tissue;
2. Antibodies to B-50 only partially interfere with transmitter release induced by phorbol esters, which stimulate PKC-mediated phosphorylation of B-50; and
3. The protein kinase inhibitors H-7 and staurosporine, both inhibiting B-50 phosphorylation, do not inhibit  $\text{Ca}^{2+}$ -dependent transmitter release.

In evaluating these data, it should be emphasized that the precise mechanisms of  $\text{Ca}^{2+}$  and phorbol ester-induced transmitter release remain to be elucidated. It may very well be that phorbol esters and  $\text{Ca}^{2+}$  affect neurotransmitter release in a different way, thus giving rise to the differential effects of inhibitors and antibodies on  $\text{Ca}^{2+}$  and phorbol ester-induced release. Alternatively, as suggested, the present results leave open the possibility that properties of B-50 other than its PKC-mediated phosphorylation are contributing to the transmitter release process and that these are affected by antibodies and inhibitors.

Such properties could involve the CaM-binding capacity of B-50. It has been shown that B-50 binds calmodulin *in vitro* under low  $\text{Ca}^{2+}$  conditions and releases CaM when  $\text{Ca}^{2+}$  is raised (Andreasen et al., 1983; Cimler et al., 1985; De Graan et al., 1990; Liu and Storm, 1990). The phosphorylation site and the CaM-binding site of B-50 are very close to each other (Alexander et al., 1988), and prephosphorylation of B-50 by PKC inhibits its property to bind CaM (Alexander et al., 1987). Data suggest that B-50 acts as a membrane-bound CaM store, releasing CaM after elevation of the  $\text{Ca}^{2+}$  levels or upon phosphorylation. A depolarization-induced rise of the intracellular concentration of  $\text{Ca}^{2+}$  *in vivo* would dissociate CaM and B-50. Once liberated from CaM, B-50 can be phosphorylated by PKC. PKC may be activated by  $\text{Ca}^{2+}$  entering the synaptic terminal after the depolarization. It is not clear whether such activation should involve the translocation of PKC from the cytosol to the membrane or whether  $\text{Ca}^{2+}$  directly activates membrane-bound PKC. Alternatively,  $\text{Ca}^{2+}$  may activate phospholipases generating diacylglycerol which could induce activation of PKC. PKC-mediated phosphorylation could prevent CaM from reassociating with B-50.

Phosphorylation of B-50 is also thought to be a modulatory event in the activity of PIP kinase, the enzyme that converts the membrane lipid PIP to  $\text{PIP}_2$  (Gispén et al., 1985b). In fact, it has been shown that an inverse relationship exists between B-50 phosphorylation and  $\text{PIP}_2$  labeling (Gispén,

1986). For instance, inhibition of B-50 phosphorylation in synaptosomal plasma membranes by  $\text{ACTH}_{1-24}$  or by B-50 antibodies induced a stimulation of  $\text{PIP}_2$  production (Oestreicher et al., 1983). Enhancement of B-50 phosphorylation by addition of purified PKC to the synaptosomal plasma membranes resulted in an inhibition of PIP kinase activity (De Graan et al., 1988a).

These data are consistent with the hypothesis that B-50 phosphorylation exerts a negative feedback control on PIP kinase activity. Increased levels of diacylglycerol, formed by receptor-mediated hydrolysis of  $\text{PIP}_2$  or phosphatidylcholine, resulting in PKC-mediated B-50 phosphorylation, would reduce replenishment of  $\text{PIP}_2$  as a mechanism to reduce further transmembrane signal transduction (Gispén et al., 1985b). It is interesting to note that recently, Houbre et al. (1991) showed that B-50 binds to phosphatidylserine and phosphatidic acid and pointed to a phosphatidylserine affinity site overlapping the phosphorylation site and the CaM-binding domain. Since phosphatidylserine is known as an activator of PIP kinase as well as of PKC, these data support the idea that B-50, PIP kinase, and PKC are part of a multifunctional complex associated with the membrane.

The postulated regulatory effect of B-50 on  $G_o$  protein (Strittmatter et al., 1990) could effect  $\text{PIP}_2$  hydrolysis in another way, since  $G_o$  might be involved in coupling some receptors to phospholipase C (Banno et al., 1987). Furthermore, this proposed interaction of B-50 with  $G_o$  forms a link to another important second messenger system of adenylate cyclase and cAMP. It is not yet evident whether phosphorylation of B-50 by PKC is of any relevance to its  $G_o$ -modulating effects. Ample evidence exists suggesting the involvement of G proteins in secretory processes (Howell et al., 1987), and it would be very interesting to investigate whether the apparent necessity of B-50 to secretion in the nerve terminal is related to modulation of  $G_o$  activity.

An intriguing question remains if there is a general main feature of B-50 responsible for its seemingly diverse roles in developing and mature neurons. A regulatory effect on vesicle fusion pro-

cesses, taking place during axonal growth, synaptic remodeling, and neurotransmitter release seems to be an attractive candidate for such a unifying feature.

## Acknowledgments

The authors are grateful to Menno van Lookeren Campagne for providing the electron micrographs and to Paul van der Most and Gerard Peek for artwork and photography, respectively.

## References

- Akers R. F. and Routtenberg A. (1985) Protein kinase C phosphorylates a 47  $M_r$  protein (F1) directly related to synaptic plasticity. *Brain Res.* **334**, 147–151.
- Albert K. A., Nairn A. C., and Greengard P. (1987) The 87-kDa protein, a major specific substrate for protein kinase C: purification from bovine brain and characterization. *Proc. Natl. Acad. Sci. USA* **84**, 7046–7050.
- Alexander K. A., Cimler B. M., Meier K. E., and Storm D. R. (1987) Regulation of calmodulin binding to P-57. *J. Biol. Chem.* **262**, 6108–6113.
- Alexander K. A., Wakim B. T., Doyle G. S., Walsh K. A., and Storm D. R. (1988) Identification and characterization of the calmodulin-binding domain of neuromodulin, a neurospecific calmodulin-binding protein. *J. Biol. Chem.* **263**, 7544–7549.
- Allsopp T. E. and Moss D. J. (1989) A developmentally regulated chicken neuronal protein associated with the cortical cytoskeleton. *J. Neurosci.* **9**, 13–24.
- Aloyo V. J., Zwiers H., and Gispen W. H. (1983) Phosphorylation of B-50 protein by calcium-activated, phospholipid-dependent protein kinase and B-50 protein kinase. *J. Neurochem.* **41**, 649–653.
- Andreasen T. K., Luetje C. W., Heideman W., and Storm D. R. (1983) Purification of a novel calmodulin binding protein from bovine cerebral cortex membranes. *Biochemistry* **22**, 4615–4618.
- Apel E. D., Byford M. F., Douglas A., Walsh K. A., and Storm D. R. (1990) Identification of the protein kinase C phosphorylation site in neuromodulin. *Biochemistry* **29**, 2330–2335.
- Apel E. D., Litchfield D. W., Clark R. H., Krebs E. G., and Storm D. R. (1991) Phosphorylation of neuromodulin (GAP-43) by casein kinase II: identification of phosphorylation sites and regulation by calmodulin. *J. Biol. Chem.* **266**, 10,544–10,551.
- Au D. C., Apel E. D., Chapman E. R., Estep R. P., Nicolson T. A., and Storm D. R. (1989) Expression of cDNAs encoding wild-type and mutant neuromodulins in *Escherichia coli*: comparison with the native protein from bovine brain. *Biochemistry* **28**, 8142–8148.
- Baetge E. E. and Hammang J. P. (1991) Neurite outgrowth in PC12 cells deficient in GAP-43. *Neuron* **6**, 21–30.
- Baizer L., Alkan S., Stocker K., and Ciment G. (1990) Chicken growth-associated protein (GAP)-43: primary structure and regulated expression of mRNA during embryogenesis. *Mol. Brain Res.* **7**, 61–68.
- Banno Y., Nagao S., Katada T., Nagata K., Ui M., and Nozawa Y. (1987) Stimulation by GTP-binding proteins ( $G_i$ ,  $G_o$ ) of partially purified phospholipase C activity from human platelet membranes. *Biochem. Biophys. Res. Comm.* **146**, 861–869.
- Basi G. S., Jacobson R. D., Virág I., Schilling J., and Skene J. H. P. (1987) Primary structure and transcriptional regulation of GAP-43, a protein associated with nerve growth. *Cell* **49**, 785–791.
- Baudier J., Bronner C., Kligman D., and Cole R. D. (1989) Protein kinase C substrates from bovine brain. Purification and characterization of neuromodulin, a neuron-specific calmodulin-binding protein. *J. Biol. Chem.* **264**, 1824–1828.
- Baudier J., Deloulme J. C., Van Dorsselaer A., Black D., and Matthes H. W. D. (1991) Purification and characterization of a brain-specific protein kinase C substrate, neurogranin (p17). *J. Biol. Chem.* **266**, 229–237.
- Benowitz L. I. and Lewis E. R. (1983) Increased transport of 44,000 to 49,000-dalton acidic proteins during regeneration of the goldfish optic nerve. *J. Neurosci.* **3**, 2153–2163.
- Benowitz L. I., Perrone-Bizzozero N. I., and Finklestein S. P. (1987) Molecular properties of the growth-associated protein GAP-43 (B-50). *J. Neurochem.* **48**, 1640–1647.
- Benowitz L. I. and Routtenberg A. (1987) A membrane phosphoprotein associated with neural development, axonal regeneration, phospholipid metabolism, and synaptic plasticity. *Trends Neurosci.* **10**, 527–532.
- Benowitz L. I., Apostolides P. J., Perrone-Bizzozero N. I., Finklestein S. P., and Zwiers H. (1988) Anatomical distribution of the growth-associated

- proteins GAP-43/B-50 in the adult rat brain. *J. Neurosci.* **8**, 339–352.
- Benowitz L. I., Perrone-Bizzozero N. I., Finklestein S. P., and Bird E. D. (1989) Localization of the growth-associated phosphoprotein GAP-43 (B-50, F1) in the human cerebral cortex. *J. Neurosci.* **9**, 990–995.
- Benowitz L. I., Rodriguez W. R., and Neve R. I. (1990) The pattern of GAP-43 immunostaining changes in the rat hippocampal formation during reactive synaptogenesis. *Mol. Brain Res.* **8**, 7–23.
- Biffo S., Verhaagen J., Schrama L. H., Schotman P., Danho W., and Margolis F. L. (1990) B-50/GAP43 expression correlates with process outgrowth in the embryonic mouse nervous system. *Eur. J. Neurosci.* **2**, 487–499.
- Brown A. M. and Birnbaumer L. (1990) Ionic channels and their regulation by G protein subunits. *Ann. Rev. Physiol.* **52**, 197–213.
- Burry R. W., Lah J. J., and Hayes D. M. (1991) Redistribution of GAP-43 during growth cone development in vitro: immunocytochemical studies. *J. Neurocytol.* **20**, 133–144.
- Chan S. Y., Murakami K., and Routtenberg A. (1986) Phosphoprotein F1: purification and characterization of a brain kinase C substrate related to plasticity. *J. Neurosci.* **6**, 3618–3627.
- Changelian P. S., Meiri K. F., Soppet D., Valenza H., Loewy A., and Willard M. B. (1990) Purification of the growth-associated protein GAP-43 by reversed phase chromatography: amino acid sequence analysis and cDNA identification. *Brain Res.* **510**, 259–268.
- Chapman E. R., Au D., Alexander A., Nicolson T. A., and Storm D. R. (1991) Characterization of the calmodulin binding domain of neuromodulin. *J. Biol. Chem.* **266**, 207–213.
- Cimler B. M., Andreasen T. J., Andreasen K. I., and Storm D. R. (1985) P-57 is a neural specific calmodulin-binding protein. *J. Biol. Chem.* **260**, 10,784–10,788.
- Cimler B. M., Giebelhaus D. H., Wakim B. T., Storm D. R., and Moon R. T. (1987) Characterization of murine cDNAs encoding P-57, a neural-specific calmodulin-binding protein. *J. Biol. Chem.* **262**, 12,158–12,163.
- Coggins P. J. and Zwiers H. (1989) Evidence for a single protein kinase C-mediated phosphorylation site in rat brain protein B-50. *J. Neurochem.* **53**, 1895–1901.
- Coggins P. J., McIntyre D. D., Vogel A. J., and Zwiers H. (1989) Neuronal protein B-50: a proton nuclear-magnetic-resonance study. *Biochem. Soc. Trans.* **17**, 785–786.
- Coggins P. J. and Zwiers H. (1990) Binding of the neuronal protein B-50, but not the metabolite B-60, to calmodulin. *J. Neurochem.* **54**, 274–277.
- Coggins P. J. and Zwiers H. (1991) B-50 (GAP-43): biochemistry and functional neurochemistry of a neuron-specific phosphoprotein. *J. Neurochem.* **56**, 1095–1106.
- Coggins P. J., Stanisiz J., Nagy A., and Zwiers H. (1991) Identification of a calmodulin-binding, B-50-immunoreactive C-kinase substrate (BICKS) in bovine brain. *Neurosci. Res. Commun.* **8**, 49–56.
- Colbran R. J. and Soderling T. R. (1990) Calcium/calmodulin dependent protein kinase II. *Curr. Topics Cell Reg.* **31**, 181–217.
- Costello B., Lin L-H., Meymandi A., Bock S., Norden J. J., and Freeman J. A. (1991) Expression of the growth and plasticity-associated neuronal protein, GAP-43, in PC12 pheochromocytoma cells. *Prog. Brain Res.* **89**, 47–67.
- Da Cunha A. and Vitkovic L. (1990) Regulation of immunoreactive GAP-43 expression in rat cortical macroglia is cell type specific. *J. Cell. Biol.* **111**, 209–215.
- De Graan P. N. E., Van Hooff C. O. M., Tilly B. C., Oestreicher A. B., Schotman P., and Gispen W. H. (1985) Phosphoprotein B-50 in nerve growth cones from fetal rat brain. *Neurosci. Lett.* **61**, 235–241.
- De Graan P. N. E., Dekker L. V., De Witt M., Schrama L. H., and Gispen W. H. (1988a) Modulation of B-50 phosphorylation and phosphoinositide metabolism in synaptic plasma membranes by protein kinase C, phorbol esters and ACTH. *J. Recept. Res.* **8**, 345–361.
- De Graan P. N. E., Heemskerk F. M. J., Dekker L. V., Melchers B. P. C., Gianotti C., and Schrama L. H. (1988b) Phorbol esters induce long and short-term enhancement of B-50/GAP-43 phosphorylation in rat hippocampal slices. *Neurosci. Res. Commun.* **3**, 175–182.
- De Graan P. N. E., Oestreicher A. B., De Wit M., Kroef M., Schrama L. H., and Gispen W. H. (1990) Evidence for the binding of calmodulin to endogenous B-50 (GAP-43) in native synaptosomal plasma membranes. *J. Neurochem.* **55**, 2139–2141.
- De Graan P. N. E., Oestreicher A. B., Schotman P., and Schrama L. H. (1991) Protein kinase C substrate B-50 (GAP-43) and neurotransmitter release. *Prog. Brain Res.* **89**, 187–207.



- Dekker L. V., De Graan P. N. E., Oestreicher A. B., Versteeg D. H. G., and Gispen W. H. (1989a) Inhibition of noradrenaline release by antibodies to B-50 (GAP-43). *Nature* **342**, 74–76.
- Dekker L. V., De Graan P. N. E., Versteeg D. H. G., Oestreicher A. B., and Gispen W. H. (1989b) Phosphorylation of B-50 (GAP43) is correlated with neurotransmitter release in rat hippocampal slices. *J. Neurochem.* **52**, 24–30.
- Dekker L. V., De Graan P. N. E., De Wit M., Hens J. J. H., and Gispen W. H. (1990a) Depolarization-induced phosphorylation of the protein kinase C substrate B-50 (GAP-43) in rat cortical synaptosomes. *J. Neurochem.* **54**, 1645–1652.
- Dekker L. V., De Graan P. N. E., Spierenburg H., De Wit M., Versteeg D. H. G., and Gispen W. H. (1990b) Evidence for a relationship between B-50 (GAP-43) and [<sup>3</sup>H]noradrenaline release in rat brain synaptosomes. *Eur. J. Pharmacol.* **188**, 113–122.
- Dekker L. V., De Graan P. N. E., and Gispen W. H. (1991) Transmitter release: target of regulation by protein kinase C? *Prog. Brain Res.* **89**, 209–233.
- De la Monte S., Federoff H. J., Ng S.-C., Grabczyk E., and Fishman M. C. (1989) GAP-43 gene expression during development: persistence in a distinctive set of neurons in the mature central nervous system. *Dev. Brain Res.* **46**, 161–168.
- Deloulme J. C., Janet T., Au D., Storm D. R., Sensenbrenner M., and Baudier J. (1990) Neuro-modulin (GAP43): a neuronal protein kinase C substrate is also present in 0-2A glial cell lineage. Characterization of neuro-modulin in secondary cultures of oligodendrocytes and comparison with the neuronal antigen. *J. Cell Biol.* **111**, 1559–1569.
- DiFiglia M., Roberts R. C., and Benowitz L. L. (1990) Immunoreactive GAP43 in the neuropil of adult rat neostriatum: localization in unmyelinated fibres, axon terminals, and dendritic spines. *J. Comp. Neurol.* **302**, 992–1001.
- Dokas L. A., Pisano M. R., Schrama L. H., Zwiers H., and Gispen W. (1990) Dephosphorylation of B-50 in synaptic plasma membranes. *Brain Res. Bull.* **24**, 321–329.
- Dolphin A. C., Errington M. L., and Bliss T. V. P. (1982) Long-term potentiation of the perforant path *in vivo* is associated with increased glutamate release. *Nature* **297**, 496–498.
- Dosemeci A. and Rodnight R. (1987) Demonstration by phase-partitioning in Triton X-114 solutions that phosphoprotein B-50 (F-1) from rat brain is an integral membrane protein. *Neurosci. Lett.* **74**, 325–330.
- Dotti C. G., Banker G. A., and Binder L. I. (1987) The expression and distribution of the microtubule-associated proteins tau and microtubule-associated protein 2 in hippocampal neurons in the rat *in situ* and in cell culture. *Neuroscience* **23**, 121–130.
- Dotti C. G., Sullivan C. A., and Banker G. A. (1988) The establishment of polarity by hippocampal neurons in culture. *J. Neurosci.* **8**, 1454–1468.
- Erzurumlu R. S., Jhaveri S., Moya K. L., and Benowitz L. I. (1989) Peripheral nerve regeneration induces elevated expression of GAP-43 in the brainstem trigeminal complex of adult hamsters. *Brain. Res.* **498**, 135–139.
- Erzurumlu R. S., Jhaveri S., and Benowitz L. I. (1990) Transient patterns of GAP-43 expression during the formation of barrels in the rat somatosensory cortex. *J. Comp. Neurol.* **292**, 443–456.
- Estep R. P., Alexander K. A., and Storm D. R. (1990) Regulation of free calmodulin levels in neurons by neuromodulin: relationship to neuronal growth and regeneration. *Curr. Topics Cell Reg.* **31**, 161–180.
- Gianotti C., Nunzi M. G., Gispen W. H., and Corradetti R. (1991) B-50 phosphorylation following different patterns of electrical stimulation in rat hippocampal slices. *Synaptic Plasticity Meeting*, Mainz, Germany.
- Gilbert W. (1986) Genes-in-pieces revisited. *Science* **228**, 823–824.
- Gilman A. G. (1987) G proteins: transducers of receptor-generated signals. *Ann. Rev. Biochem.* **56**, 615–649.
- Gispen W. H. (1986) Phosphoprotein B-50 and phosphoinositide in brain synaptic plasma membranes: a possible feedback relationship. *Biochem. Soc. Trans.* **14**, 163–165.
- Gispen W. H., Leunissen J. L. M., Oestreicher A. B., Verkleij A. J., and Zwiers H. (1985a) Presynaptic localization of B-50 phosphoprotein: the ACTH-sensitive protein kinase C substrate involved in rat brain phosphoinositide metabolism. *Brain Res.* **328**, 381–385.
- Gispen W. H., Van Dongen C. J., De Graan P. N. E., Oestreicher A. B., and Zwiers H. (1985b) The role of phosphoprotein B-50 in phosphoinositide metabolism in brain synaptic plasma membranes, in *Inositol and Phosphoinositides*, Bleasdale J. E., Hauser G., and Eichberg J., eds., Humana, Clifton, NJ, pp. 399–413.
- Gispen W. H., De Graan P. N. E., Chan S. Y., and Routtenberg A. (1986) Comparison between the neural acidic proteins B-50 and F1. *Prog. Brain Res.* **69**, 383–386.

- Gordon-Weeks P. R. (1989) GAP-43—What does it do in the growth cone. *Trends Neurosci.* **12**, 363–365.
- Gorgels T. G. M. F., Van Lookeren Campagne M., Oestreicher A. B., Gribnau A. A. M., and Gispen W. H. (1989) B-50/GAP-43 is localized at the cytoplasmic side of the plasma membrane in developing and adult rat pyramidal tract. *J. Neurosci.* **9**, 3861–3869.
- Goslin K., Schreyer D. J., Skene J. H. P., and Banker G. (1988) Development of neuronal polarity: GAP-43 distinguishes axonal from dendritic growth cones. *Nature* **366**, 672–680.
- Goslin K. and Banker G. (1990) Rapid changes in the distribution of GAP-43 correlate with the expression of neuronal polarity during normal development and under experimental conditions. *J. Cell Biol.* **110**, 1319–1331.
- Goslin K., Schreyer D. J., Skene J. H. P., and Banker G. (1990) Changes in the distribution of GAP-43 during development of neuronal polarity. *J. Neurosci.* **10**, 588–602.
- Grabczyk E., Zuber M. X., Federoff H. J., Ng S.-C., Pack A., and Fishman M. C. (1990) Cloning and characterization of the rat gene encoding GAP-43. *Eur. J. Neurosci.* **2**, 822–827.
- Graff J. M., Stumpo D. J., and Blackshear P. J. (1989a) Characterization of the phosphorylation sites in the chicken and bovine myristoylated alanine-rich C kinase substrate protein, a prominent cellular substrate for protein kinase C. *J. Biol. Chem.* **264**, 11,912–11,919.
- Graff J. M., Young T. N., Johnson J. D., and Blackshear P. J. (1989b) Phosphorylation-regulated calmodulin binding to a prominent cellular substrate for protein kinase C. *J. Biol. Chem.* **264**, 21,818–21,823.
- Greene L. A. and Shooter E. M. (1980) The nerve growth factor: biochemistry, synthesis and mechanism of action. *Ann. Rev. Neurosci.* **3**, 353–402.
- Hammang J. P., Messing A., and Baetge E. E. (1990) The C6 glioma cell line expresses the growth-associated protein GAP-43 in a developmentally regulated fashion. *Soc. Neurosci. Abstr.* **16**, 339.8.
- Heemskerk F. M. J., Schrama L. H., and Gispen W. H. (1989) Activation of protein kinase C by 4-aminopyridine dependent on Na<sup>+</sup> channel activity in rat hippocampal slices. *Neurosci. Lett.* **106**, 315–321.
- Heemskerk F. M. J., Schrama L. H., De Graan P. N. E., and Gispen W. H. (1990) 4-Aminopyridine stimulates B-50 (GAP-43) phosphorylation in rat synaptosomes. *J. Mol. Neurosci.* **2**, 11–17.
- Hoffman P. N. (1989) Expression of GAP-43, a rapidly transported growth-associated protein, and class II  $\beta$ -tubulin, a slowly transported cytoskeletal protein, are coordinated in regenerating neurons. *J. Neurosci.* **9**, 893–897.
- Howell T. W., Cockcroft S., and Gomperts B. D. (1987) Essential synergy between Ca<sup>2+</sup> and guanine nucleotides in exocytotic secretion from permeabilized mast cells. *J. Cell Biol.* **105**, 191–197.
- Houbre D., Duportail G., Deloulme J.-C., and Baudier J. (1991) The interactions of the brain-specific calmodulin-binding protein kinase C substrate, neuromodulin (GAP-43), with membrane phospholipids. *J. Biol. Chem.* **266**, 7121–7131.
- Karns L. R., Ng S.-C., Freeman J. A., and Fishman M. C. (1987) Cloning of complementary DNA for GAP-43, a neuronal growth-related protein. *Science* **236**, 597–600.
- Katz F., Ellis L., and Pfenninger K. H. (1985) Nerve growth cones isolated from fetal rat brain III: calcium-dependent protein phosphorylation. *J. Neurosci.* **5**, 1402–1411.
- Kleinschmidt J. A., Dingwall C., Maier G., and Franke W. W. (1986) Molecular characterization of a karyophilic, histone-binding protein: cDNA cloning, amino acid sequence and expression of nuclear protein N1/N2 of *Xenopus laevis*. *EMBO J.* **5**, 3547–3552.
- Kosik K. S., Orecchio L. D., Bruns G. A. P., Benowitz L. I., MacDonald G. P., Cox D. R., and Neve R. L. (1988) Human GAP-43: its deduced amino acid sequence and chromosomal localization in mouse and human. *Neuron* **1**, 127–132.
- Kozak M. (1989) The scanning model for translation: an update. *J. Cell Biol.* **108**, 229–241.
- Labate M. E. and Skene J. H. P. (1989) Selective conservation of GAP-43 structure in vertebrate evolution. *Neuron* **3**, 299–310.
- Levine J., Skene J. H. P., and Willard M. B. (1981) GAPs and fodrin. Novel axonally transported proteins. *Trends Neurosci.* **4**, 273–277.
- Linden D. J. and Routtenberg A. (1989) The role of protein kinase C in long-term potentiation: a testable model. *Brain Res. Rev.* **14**, 279–296.
- Liu Y. and Storm D. R. (1989) Dephosphorylation of neuromodulin by calcineurin. *J. Biol. Chem.* **264**, 12,800–12,804.
- Liu Y. and Storm D. R. (1990) Regulation of free calmodulin levels by neuromodulin: neuron growth and regeneration. *Trends Pharmacol. Sci.* **11**, 107–111.

- Lovinger D. M., Colley P. A., Akers R. F., Nelson R. B., and Routtenberg A. (1986) Direct relation of long-term synaptic potentiation to phosphorylation of membrane protein F1, a substrate for membrane protein kinase C. *Brain Res.* **399**, 205–211.
- Low M. G. (1987) Biochemistry of the glycosylphosphatidylinositol membrane protein anchors. *Biochem. J.* **22**, 1–13.
- Magee T. and Hanley M. (1988) Sticky fingers and CAAX boxes. *Nature* **335**, 114–115.
- Masure H. R., Alexander K. A., Wakim B. T., and Storm D. R. (1986) Physicochemical and hydrodynamic characterization of P-57, a neurospecific calmodulin binding protein. *Biochemistry* **25**, 7553–7560.
- McDonald J. R. and Lawrence J. C. (1989) Identification of an adipocyte protein that binds to calmodulin in the absence of  $Ca^{2+}$  and is phosphorylated in response to insulin and tumor-promoting phorbol esters. *J. Biol. Chem.* **262**, 9611–9618.
- McIntosh H., Daw N., and Parkinson D. (1990) GAP-43 in the cat visual cortex during postnatal development. *Visual Neurosci.* **4**, 585–593.
- McMaster D., Zwiers H., and Lederis K. (1988) The growth associated neuronal phosphoprotein B-50: improved purification, partial primary structure and characterization and localization of proteolysis products. *Brain Res. Bull.* **21**, 265–276.
- Megerian J. T. and Klein W. L. (1990) Evidence that neuromodulin (GAP-43, B-50, pp46, F1) is not essential for early neuritogenesis. *Soc. Neurosci. Abstr.* **16**, 339.11.
- Meiri K. F., Pfenninger K. H., and Willard M. B. (1986) Growth-associated protein, GAP-43, a polypeptide that is induced when neurons extend axons, is a component of growth cones and corresponds to pp46, a major polypeptide of a subcellular fraction enriched in growth cones. *Proc. Natl. Acad. Sci. USA* **83**, 3537–3541.
- Meiri K. F., Willard M. B., and Johnson M. I. (1988) Distribution and phosphorylation of the growth-associated protein GAP-43 in regenerating sympathetic neurons in culture. *J. Neurosci.* **8**, 2571–2581.
- Meiri K. F. and Gordon-Weeks P. R. (1990) GAP-43 in growth cones is associated with areas of membrane that are tightly bound to substrate and is a component of a membrane skeleton subcellular fraction. *J. Neurosci.* **10**, 256–266.
- Meiri K. F., Bickerstaff L. E., and Schwob J. E. (1991) Monoclonal antibodies show that kinase C phosphorylation of GAP-43 during axonogenesis is both spatially and temporally restricted in vivo. *J. Cell Biol.* **112**, 991–1005.
- Miller R. J. (1986) Protein kinase C: a key regulator of neuronal excitability? *Trends Neurosci.* **9**, 538–541.
- Moss D. J., Fernyhough P., Chapman K., Baizer L., Bray D., and Allsopp T. (1990) Chicken growth-associated protein GAP-43 is tightly bound to the actin-rich neuronal membrane skeleton. *J. Neurochem.* **54**, 729–736.
- Moya K. L., Benowitz L. I., and Schneider G. E. (1990) Abnormal retinal projections alter GAP-43 patterns in the diencephalon. *Brain Res.* **527**, 259–265.
- Nelson R. B., Linden D. J., Hyman C., Pfenninger K. H., and Routtenberg A. (1989) The two major phosphoproteins in growth cones are probably identical to two protein kinase C substrates correlated with persistence of long term potentiation. *J. Neurosci.* **9**, 381–389.
- Neve R. L., Perrone-Bizzozero N. I., Finklestein S., Zwiers H., Bird E., Kurnit D. M., and Benowitz L. I. (1987) The neuronal growth-associated protein GAP-43 (B-50, F1): neuronal specificity, developmental and regulation and regional distribution of the human and rat mRNAs. *Mol. Brain Res.* **2**, 177–183.
- Neve R. L., Finch E. A., Bird E. D., and Benowitz L. I. (1988) Growth-associated protein GAP-43 is expressed selectively in associative regions of the adult human brain. *Proc. Natl. Acad. Sci. USA* **85**, 3638–3642.
- Neve R. L., Dawes L. R., and Geller A. I. (1990) Fusion of the amino-terminal 10 amino acids of GAP-43 to  $\beta$ -galactosidase targets the chimeric protein to neuronal processes. *Soc. Neurosci. Abstr.* **16**, 27.6.
- Ng S.-C., De la Monte S. M., Conboy G. L., Kams L. R., and Fishman M. C. (1988) Cloning of human GAP-43: growth association and ischemic resurgence. *Neuron* **1**, 133–139.
- Nielander H. B., Schrama L. H., Van Rozen A. J., Kasperaitis M., Oestreicher A. B., De Graan P. N. E., Gispen W. H., and Schotman P. (1987) Primary structure of the neuron-specific phosphoprotein B-50 is identical to growth-associated protein GAP-43. *Neurosci. Res. Commun.* **1**, 163–172.
- Nielander H. B., Schrama L. H., Van Rozen A. J., Oestreicher A. B., Gispen W. H., and Schotman P. (1989) The calmodulin-binding protein B-50/GAP-43: localization of the proteolytic sites for *S. aureus*

- protease and the phosphorylation sites for protein kinase C. *J. Neurochem.* **S52**, S106A.
- Nielander H. B., Schrama L. H., Van Rozen A. J., Kasperaitis M., Oestreicher A. B., Gispen W. H., and Schotman P. (1990) Mutation of serine 41 in the neuron-specific protein B-50 (GAP-43) prohibits phosphorylation by protein kinase C. *J. Neurochem.* **55**, 1442-1445.
- Nishizuka Y. (1988) The molecular heterogeneity of protein kinase C and its implications for cellular regulation. *Nature* **334**, 661-665.
- Nishizuka Y. (1989) The family of protein kinase C for signal transduction. *JAMA* **262**, 1826-1833.
- Oestreicher A. B., Van Dongen C. J., Zwiers H., and Gispen W. H. (1983) Affinity-purified anti-B-50 protein antibody: interference with the function of the phosphoprotein B-50 in synaptic plasma membranes. *J. Neurochem.* **41**, 331-340.
- Oestreicher A. B. and Gispen W. H. (1986) Comparison of the immunocytochemical distribution of the phosphoprotein B-50 in the cerebellum and hippocampus of immature and adult rat brain. *Brain Res.* **375**, 267-279.
- Oestreicher A. B., Dekker L. V., and Gispen W. H. (1986) A radioimmunoassay for the phosphoprotein B-50: distribution in rat brain. *J. Neurochem.* **46**, 1366-1369.
- Oestreicher A. B., De Graan P. N., Schrama L. H., Lamme V. A. F., Bloemen R. J., Schotman P., and Gispen W. H. (1989) The protein kinase C phosphosite(s) in B-50 (GAP-43) are confined to 15K phosphofragments produced by *Staphylococcus aureus* V8 protease. *Neurochem. Int.* **14**, 361-372.
- O'Hara B. F., Fisher S., Oster-Granite M. L., Gearhart J. D., and Reeves R. H. (1989) Developmental expression of the amyloid precursor protein, growth-associated protein 43, and somatostatin in normal and trisomy 16 mice. *Dev. Brain Res.* **49**, 300-304.
- O'Neil K. T. and Degrado W. F. (1990) How calmodulin binds its targets: sequence independent recognition of amphiphilic  $\alpha$ -helices. *Trends Biochem. Sci.* **15**, 59-64.
- Perrone-Bizzozero N. I., Weiner D., Hauser G., and Benowitz L. I. (1988) Extraction of major acidic  $Ca^{2+}$  dependent phosphoproteins from synaptic membranes. *J. Neurosci. Res.* **20**, 346-350.
- Perrone-Bizzozero N. I., Benowitz L. I., Apostolides P. J., Franck E. R., Finklestein S. P., and Bizzozero O. A. (1989) Protein fatty acid acylation in developing cortical neurons. *J. Neurochem.* **52**, 1149-1155.
- Pfenninger K. H. (1986) Of nerve growth cones, leukocytes and memory: second messenger systems and growth-regulated proteins. *Trends Neurosci.* **9**, 562-565.
- Pisano M. R., Hegazy M. G., Reimann E. W., and Dokas L. A. (1988) Phosphorylation of protein B-50 (GAP-43) from adult rat brain cortex by casein kinase II. *Biochem. Biophys. Res. Commun.* **155**, 1207-1212.
- Ramakers G. J. A., Oestreicher A. B., Van Leeuwen F. W., De Graan P. N. E., and Gispen W. H. (1991) Developmental changes in B-50 (GAP-43) in primary cultures of cerebral cortex: B-50 immunolocalization, axonal elongation rate and growth cone morphology. *Int. J. Dev. Neurosci.* **9**, 231-241.
- Represa A., Deloulme J. C., Sensenbrenner M., Ben-Ari Y., and Baudier J. (1990) Neurogranin: immunocytochemical localization of a brain-specific protein kinase C substrate. *J. Neurosci.* **10**, 3782-3792.
- Rodnight R. (1982) Aspects of protein phosphorylation in the nervous system with particular reference to synaptic transmission. *Prog. Brain Res.* **56**, 1-25.
- Rosenthal A., Chan S. Y., Henzel W., Haskell C., Kuang W.-J., Chen E., Wilcox J. N., Ullrich A., Goeddel D. V., and Routtenberg A. (1987) Primary structure and mRNA localization of protein F1, a growth-related protein kinase C substrate associated with synaptic plasticity. *EMBO J.* **6**, 3641-3646.
- Schmidt M. F. G. (1983) Fatty acid binding: a new kind of posttranslational modification of membrane proteins. *Curr. Topics Microbiol. Immunol.* **102**, 101-112.
- Schotman P., Nielander H. B., Van Rozen A. J., Frankena H., Schrama L. H., and Gispen W. H. (1989) Microheterogeneity of the growth-associated neuronal protein B-50 (GAP-43). Contribution of phosphorylation by protein kinase C. *J. Chromatogr.* **483**, 301-309.
- Schotman P., Jap Tjoen San E. R. A., Schmidt-Michels M., and Gispen W. H. (1990) Antisense B-50 (GAP-43) prevents neurite outgrowth in PC12 cells. *Soc. Neurosci. Abstr.* **16**, 339.5.
- Schrama L. H., De Graan P. N. E., Oestreicher A. B., and Gispen W. H. (1987) B-50 phosphorylation, protein kinase C and the induction of excessive grooming behavior in the rat. *Adv. Exp. Med. Biol.* **221**, 393-409.
- Schrama L. H., Heemskerk F. M. J., and De Graan P. N. E. (1989) Dephosphorylation of protein kinase



- C phosphorylated B-50/GAP-43 by the calmodulin-dependent phosphatase calcineurin. *Neurosci. Res. Commun.* **5**, 141–147.
- Schrama L. H., Nielander H. B., Van Rozen A. J., Schotman P., and Gispen W. H. (1990) Organization of the rat and the human B-50 (GAP-43) gene. *Soc. Neurosci. Abstr.* **16**, 154.11.
- Sheu F. S., Kasamatsu T., and Routtenberg A. (1990) Protein kinase-C activity and substrate (F1/GAP-43) phosphorylation in developing cat visual cortex. *Brain Res.* **24**, 144–148.
- Skene J. H. P. and Willard M. B. (1981) Axonally transported proteins associated with axon growth in rabbit central and peripheral nervous systems. *J. Cell Biol.* **89**, 96–103.
- Skene J. H. P. (1989) Axonal growth-associated proteins. *Ann. Rev. Neurosci.* **12**, 127–156.
- Skene J. H. P. and Virág I. (1989) Posttranslational membrane attachment and dynamic fatty acylation of a neuronal growth cone protein, GAP-43. *J. Cell Biol.* **108**, 613–624.
- Stein R., Mori N., Matthews K., Lo L.-C., and Anderson D. J. (1988) The NGF-inducible SCG10 mRNA encodes a novel membrane-bound protein present in growth cones and abundant in developing neurons. *Neuron* **1**, 463–476.
- Strittmatter S. M., Valenzuela D., Kennedy T. E., Neer E. J., and Fishman M. C. (1990) G<sub>o</sub> is a major growth cone protein subject to regulation by GAP-43. *Nature* **344**, 836–841.
- Takano E., Maki M., Mori H., Hatanaka M., Marti T., Titani K., Kannagi R., Ooi T., and Murachi T. (1988) Pig heart calpastatin: identification of repetitive domain structures and anomalous behavior in polyacrylamide gel electrophoresis. *Biochemistry* **27**, 1964–1972.
- Tetzlaff W., Zwiers H., Lederis K., Cassar L., and Bisby M. A. (1989) Axonal transport and localization of B-50/GAP-43-like immunoreactivity in regenerating sciatic and facial nerves of the rat. *J. Neurosci.* **9**, 1303–1313.
- Van der Neut R., Oestreicher A. B., Gispen W. H., and Bär P. R. (1990) The expression of B-50/GAP-43 during development of rat spinal neurons in culture is regulated by interneuronal contact. *Neurosci. Lett.* **109**, 36–41.
- Van der Zee C. E. E. M., Nielander H. B., Vos J. P., Lopes da Silva S., Verhaagen J., Oestreicher A. B., Schrama L. H., Schotman P., and Gispen W. H. (1989) Expression of growth-associated protein B-50 (GAP43) in dorsal root ganglia and sciatic nerve during regenerative sprouting. *J. Neurosci.* **9**, 3505–3512.
- Van Dongen C. J., Zwiers H., Oestreicher A. B., and Gispen W. H. (1985) ACTH, phosphoprotein B-50, and polyphosphoinositide metabolism in rat brain membranes, in *Phospholipids in the Nervous System*, vol. 2, Horrocks L. A., ed., Raven, New York, pp. 49–59.
- Van Hooff C. O. M., De Graan P. N. E., Oestreicher A. B., and Gispen W. H. (1988) B-50 phosphorylation and polyphosphoinositide metabolism in nerve growth cone membranes. *J. Neurosci.* **8**, 1789–1795.
- Van Hooff C. O. M., Oestreicher A. B., De Graan P. N. E., and Gispen W. H. (1989a) Role of the growth cone in neuronal differentiation. *Mol. Neurobiol.* **3**, 101–133.
- Van Hooff C. O. M., Holthuis J., Oestreicher A. B., Boonstra J., De Graan P. N. E., and Gispen W. H. (1989b) Nerve growth factor-induced changes in the intracellular localization of the protein kinase C substrate B-50 in pheochromocytoma cells. *J. Cell Biol.* **108**, 1115–1125.
- Van Lookeren Campagne M., Oestreicher A. B., Van Bergen en Henegouwen P. M. P., and Gispen W. H. (1990) Ultrastructural immunocytochemical localization of B-50/GAP-43, a protein kinase C substrate, in isolated presynaptic nerve terminals and neuronal growth cones. *J. Neurocytol.* **18**, 479–489.
- Van Lookeren Campagne M., Oestreicher A. B., Buma P., Verkleij A. J., and Gispen W. H. (1991a) Ultrastructural localization of adrenocorticotrophic hormone and the phosphoprotein B-50/growth-associated protein 43 in freeze-substituted, lowicryl HM20-embedded mesencephalic central gray substance of the rat. *Neuroscience* **42**, 517–529.
- Van Lookeren Campagne M., Dotti C. G., Jap Tjoen San E. R. A., Gispen W. H., and Oestreicher A. B. (1991b) Ultrastructural distribution of B-50/GAP-43 in cultured hippocampal neurons during various developmental stages. *Soc. Neurosci. Abstr.* **17**, 525.17.
- Verhaagen J., Oestreicher A. B., Gispen W. H., and Margolis F. L. (1989) The expression of the growth associated protein B-50/GAP43 in the olfactory system of neonatal and adult rats. *J. Neurosci.* **9**, 683–691.
- Verhaagen J., Oestreicher A. B., Grillo M., Khew-Goodall Y.-S., Gispen W. H., and Margolis F. L. (1990) Neuroplasticity in the olfactory system: differential effects of central and peripheral lesions of the primary olfactory pathway on the expression



- of B-50/GAP43 and the olfactory marker protein. *J. Neurosci. Res.* **26**, 31–44.
- Vitkovic L., Stelsslinger H. W., Aloyo V. J., and Mersel M. (1988) The 43-kDa neuronal growth-associated protein (GAP-43) is present in plasma membranes of rat astrocytes. *Proc. Natl. Acad. Sci. USA* **85**, 8296–8300.
- Wakim B. T., Alexander K. A., Masure H. R., Cimler B. M., Storm D. R., and Walsh K. A. (1987) Amino acid sequence of P-57, a neurospecific calmodulin-binding protein. *Biochemistry* **26**, 7466–7470.
- Watson J. B., Battenberg E. F., Wong K. K., Bloom F. E., and Sutcliffe J. G. (1990) Subtractive cDNA cloning of RC3, a rodent cortex-enriched mRNA encoding a novel 78 residue protein. *J. Neurosci. Res.* **26**, 397–408.
- Widmer F. and Caroni P. (1990) Identification, localization, and primary structure of CAP-23, a particle-bound cytosolic protein of early development. *J. Cell Biol.* **11**, 3035–3047.
- Woolf C. J., Reynolds M. L., Molander C., O'Brien C., Lindsay R. M., and Benowitz L. I. (1990) The growth-associated protein GAP-43 appears in dorsal root ganglion cells and in the dorsal horn of the rat spinal cord following peripheral nerve injury. *Neuroscience* **34**, 465–478.
- Yamamoto M. and Kondo H. (1990) Gene expression of a neuronal growth-associated protein, GAP-43, in the paraganglionic carotid body as well as in the autonomic ganglia of normal adult rats. *Neurosci. Lett.* **117**, 275–279.
- Yankner B. A., Benowitz L. I., Villa-Komaroff L., and Neve R. L. (1990) Transfection of PC12 cells with the human GAP-43 gene: effects on neurite outgrowth and regeneration. *Mol. Brain Res.* **7**, 39–44.
- Zuber M. X., Strittmatter S. M., and Fishman M. C. (1989a) A membrane-targeting signal in the amino terminus of the neuronal protein GAP-43. *Nature* **341**, 345–348.
- Zuber M. X., Goodman D. W., Karns L. R., and Fishman M. C. (1989b) The neuronal growth-associated protein GAP-43 induces filopodia in nonneuronal cells. *Science* **224**, 1193–1195.
- Zwiers H., Veldhuis H. D., Schotman P., and Gispen W. H. (1976) ACTH, cyclic nucleotides, and brain protein phosphorylation *in vitro*. *Neurochem. Res.* **1**, 669–677.
- Zwiers H., Wiegant V. M., Schotman P., and Gispen W. H. (1978) ACTH-induced inhibition of endogenous rat brain protein phosphorylation *in vitro*: structure activity. *Neurochem. Res.* **3**, 455–463.
- Zwiers H., Tonnaer J., Wiegant V. M., Schotman P., and Gispen W. H. (1979) ACTH-sensitive protein kinase from rat brain membranes. *J. Neurochem.* **33**, 247–256.
- Zwiers H., Schotman P., and Gispen W. H. (1980) Purification and some characteristics of an ACTH-sensitive protein kinase and its substrate protein in rat brain membranes. *J. Neurochem.* **34**, 1689–1699.
- Zwiers H., Gispen W.H., Kleine L., and Mahler H.R. (1982) Specific proteolysis of a brain membrane phosphoprotein (B-50): effects of calcium and calmodulin. *Neurochem. Res.* **7**, 127–137.
- Zwiers H., Verhaagen J., Van Dongen C. J., De Graan P. N. E., and Gispen W. H. (1985) Resolution of rat brain synaptic phosphoprotein B-50 into multiple forms by twodimensional electrophoresis: evidence for multisite phosphorylation. *J. Neurochem.* **44**, 1083–1090.
- Zwiers H. and Coggins P. J. (1990) Corticotropin (ACTH) inhibits the specific proteolysis of the neuronal phosphoprotein B-50/GAP-43. *Peptides* **11**, 951–954.