

A Pan-African core complex in the Sinai, Egypt

Published in “Geologie en Mijnbouw: B. Blasband, P. Brooijmans, P. Dirks, W. Visser and White, S., 1997. A Pan-African core complex in the Sinai Egypt. *Geologie en Mijnbouw*, 73, 247-266

After the publication of this paper, additional research was performed in the Wadi Kid Complex. This research, presented in Appendix 1, confirmed the core complex model for this area and showed that the low-angle shear zone was active at ~595 Ma

Abstract

In the late Precambrian history of the Wadi Kid area in the Sinai, Egypt, two deformation phases are clearly recognized. The first phase, D1 (pre-620 Ma), produced a steep regional foliation, axial planar to upright F1 folds, in rocks of a lower-greenschist grade. This compressional phase of deformation is interpreted in terms of subduction in an island-arc setting.

The second phase, D2 (post-620 Ma), is mainly expressed by the widespread development of sub-horizontal mylonitic zones with a total thickness of 1.5 km. Shear sense indicators give a consistent regional transport direction to the northwest, with local indications of reversal to the southeast. This event is associated with regional LP/HT metamorphism, indicative of high thermal gradients. Because of the LP/HT metamorphism, the change in geochemical nature of the granitoids, and the orientation of the dykes, we interpret the mylonitic zones as low-angle normal shear zones related to core-complex development during an extensional event with the transport reversal being induced by doming. We postulate that orogenic collapse was responsible for the transition from the D1 compressional phase to the D2 extensional phase.

5.1 Introduction

The nature of the Pan-African orogeny in the Arabo-Nubian Shield has been subject of debate over the last years. Many geologists believe that a compressional regime in an island-arc setting was responsible for the main tectonic features (e.g. Shackleton et al. 1980; Bentor 1985; Vail

1985; Ragab 1993). However, in some recent studies large strike-slip zones are regarded as being the important tectonic feature; they are associated with extensional structures such as sedimentary basins and dykes (Stern 1985; Hussein 1988).

The Wadi Kid area in the Sinai, Egypt, was chosen as a key area to study the structural geology and tectonics of the Precambrian rocks of the Arabo-Nubian Shield. Just as two models exist for the overall evolution of this shield, there are also two different models for the evolution of the Precambrian in the Sinai. Shimron (1980, 1983) interprets the tectonic development in terms of a compressional island-arc setting with two deformation phases: a syn-sedimentary D1 phase, responsible for the folding of the bedding (S_0), and a D2 thrust-phase, mainly represented by the sub-horizontal foliation found throughout most of the Wadi Kid area. On the other hand, Reymer (1983, 1984) postulates that there was a D1 compressional phase which took place in an island-arc setting similar to Shimron's D1 phase, but he relates the D2 structures to the diapiric emplacement of granites.

The aim of the present study is to investigate in particular the D2 phase of deformation in the Wadi Kid area. Detailed field mapping, followed by a microstructural analysis of carefully selected samples, was undertaken. The combined results indicate that the sub-horizontal foliations, described by Shimron (1980, 1983) and Reymer (1983, 1984), are mylonitic in origin and related to extension. Consequently, we propose a radically different tectonic interpretation for the Wadi Kid area: that of a widespread core complex, resulting from orogenic collapse. The data supporting this conclusion are presented and discussed below, along with their implications for the tectonic evolution of the Wadi Kid area.

5.2 Geological background

5.2.1 Introduction

The Sinai peninsula consists at the surface mainly of sedimentary rocks (80%), ranging from Cambrian to Quaternary age, overlying the Precambrian crystalline basement. This basement forms a part of the Arabo-Nubian Shield, which extends over most of NE Africa and Arabia.

The recent geology of the Sinai is strongly influenced by the rifting of the Red Sea. Two orientations of faults are active since the Miocene: a NNE-SSW trend and a NW-SE trend. The metamorphic assemblages indicate that the total uplift did not exceed 15 km. The main phase of uplift took place during the Cambrian and was responsible for an uplift of at least 12 km. The post-Cretaceous tectonics are responsible for block rotation and another phase of uplift of 3 km (Kohn & Eyal 1981).

5.2.2 Lithology

The metamorphic and magmatic rocks making up the basement in the Wadi Kid area have been described by Shimron (1980, 1983, 1984), Furnes et al. (1985) and Reymer (1983, 1984). Their distribution is shown in Figure 5-1. The metamorphic and magmatic rock suites are referred to as 'Formations' in the literature and we follow this terminology. The Umm Zariq Fm., Malhaq Fm., Heib Fm. and Tarr Fm. contain metasedimentary and metavolcanic rocks. The Quneia Fm. and the Sharira Gabbro/Diorite Fm. contain plutonic and meta-plutonic rocks. All these




Figure 5-1 Generalized geological sketch-map of the Wadi Kid area, modified after Shimron (1987).

formations have been intruded by granites. The area is cross-cut by swarms of mafic and felsic dykes. The total thickness of the Precambrian metamorphic formations in the Sinai is at least 1.5 km.

The *Umm Zariq Fm.* is a metasedimentary sequence, mainly formed by metapelites; it contains also some metapsammities. The schistose metapelites contain biotite, muscovite,

retrograde chlorite and feldspar along with porphyroblasts of garnet, andalusite and cordierite, indicating amphibolite facies of the low-pressure/high-temperature (LP/HT) type. Relict sedimentary structures, such as bedding, cross-bedding, slump structures and fining upward sequences, are found in the less schistose lenses. The rocks of the Umm Zariq Fm. are interpreted as metagreywackes.

The *Malhaq Fm.* is a metavolcanic sequence (Furnes et al. 1985) containing mainly schists and minor non-schistose massive equivalents. The schistose metavolcanics consist of feldspars, quartz, biotite and hornblende, indicating upper-amphibolite facies. The massive equivalents occur as layers and blocks within the schists and contain biotite, muscovite, chlorite, some quartz and feldspar.

The *Heib Fm.* is a mixed sequence of metasediments and metavolcanics. Rhyolitic and andesitic volcanics can be recognized. The metasedimentary sequence contains conglomerates and turbidites (the Beda Turbidites). The rocks display a lower metamorphic grade than the Umm Zariq and Malhaq Fms and are less deformed. In the more schistose parts, biotite and muscovite are observed, indicating a higher metamorphic grade. The lower-grade parts overlie the higher-grade parts of the Heib Fm. and of the Malhaq Fm. The rocks are thought to be of island-arc volcanic origin, deposited in a marine environment (Furnes et al. 1985).

The *Tarr Fm.*, found in the southern parts of the Wadi Kid, is dominantly a metasedimentary sequence consisting of metaconglomerates, metapelites and metapsammities, in which melanges and felsic flows can be recognized. The formation displays greenschist-grade metamorphism which is lower than that of the Malhaq and Umm Zariq Fms.

The *Quneia Fm.* consists of foliated and lineated diorites and tonalites with occasional biotite-rich xenoliths. Identical intrusives occur in the Nuweiba area, 100 km N of the Wadi Kid Complex, and these were interpreted, from geochemical data, as subduction-related I-type intrusives (Ahmed et al. 1993). Similar intrusives in the Timna area, 150 km N of the Wadi Kid area, were interpreted as I-type granitoids related to crustal thickening in an island-arc environment (Beyth et al. 1994).

Furnes et al. (1985) concluded on basis of major and trace-element studies that the rocks of the metamorphic formations, described above, are subduction-related.

The *Sharira Gabbro/Diorite Fm.* occurs as a single large intrusive body in the northern part of the Wadi Kid Complex. It was described by Furnes et al. (1985) as a layered gabbroic and dioritic intrusion.

The Wadi Kid metamorphic complex is bordered to the north, west and south by *Younger Granites*. Three phases of granitoid intrusions were described (Bentor 1984; Ahmed et al. 1993). The oldest plutonic rocks are the deformed tonalitic to granodioritic rocks of the Quneia Fm. The second phase consists of muscovite granites, which only occur in the most northern part of the Wadi Kid area. The youngest are the biotite-rich granites. Geochemical data indicate that the oldest granitoids are I-type, the muscovite-rich granites S-type, and the youngest biotite granites A-type granites (Ahmed et al. 1993). The S- and A-type intrusives are younger than any of the metamorphic rocks and mostly show intrusive contacts with these rocks. Trace-element studies of late Precambrian A-type granites in southern Israel indicate that these were formed in a thinned crust due to extension, where mantle melts were able to rise near

to the surface (Beyth et al. 1994).

5.2.3 Metamorphism

Metamorphic grades of the Wadi Kid area range from lower-greenschist to amphibolite facies with a general trend of increase towards the central and northern parts of the area. Lower-greenschist-facies rocks dominate the southern areas. Sillimanite, garnet and andalusite porphyroblasts appear in the central Wadi Kid Complex where the metamorphic grade reaches upper-amphibolite facies. Here, staurolite is overgrown by andalusite and biotite, which indicates that a medium-pressure phase (> 5 kbar) preceded a LP/HT metamorphic phase. In the northern part of the area, where metavolcanic rocks are found, garnet and hornblende porphyroblasts occur. In the north-eastern part, migmatites are present.

Reymer et al. (1984) calculated a temperature of 700° C and a pressure of 3 to 4 kbar for the Malhaq and Umm Zariq Fms in the central Wadi Kid. Temperatures of 300° C and pressures of 2 kbar have been deduced for the lower-greenschist-facies rocks in the southeastern part of the complex. The metamorphic grade in the Heib Fm. increases from lower greenschist in the Bada Turbidites in the southeast, to higher greenschist in the west and north where biotite-schists appear. Reymer et al. (1984) estimated that the thermal gradient was higher than 50°C/km in the rocks showing amphibolite-grade metamorphism. Shimron (1987) constructed a clockwise P-T-t path for the metamorphic rocks in the Wadi Kid (Figure 5-2).




Figure 5-2 P-T-t path for aluminosilicate metamorphic rocks in the Wadi Kid area after Shimron (1987). The plot shows a clockwise path, starting with LP/LT (M1) and moving to medium P and T. After upper greenschist facies was reached (M1a), P remained constant at first and then started to go down but T continued to rise. Eventually, P and T both decreased during retrograde metamorphism related to uplift (M2).

5.2.4 Geochronology

The limited number of isotopic ages available for the wider Wadi Kid area are summarized in Table 5-1. From these data it can be concluded that deposition of sediments and volcanics took place after 770 Ma and until 650 Ma ago (Priem et al. 1984), when the metamorphism started. The main amphibolite phase of metamorphism started at about 620 Ma ago. Pulses of metamorphism and the intrusion of granites continued till the early Cambrian and were accompanied by the intrusion of dykes (Ayalon et al. 1987).

Table 5-1 Summary of isotopic ages of Precambrian rocks in the wider Wadi Kid area. Bio, biotite; WR, whole rock; Zr, zircon.

| Formation | Metamorphic grade | Age in Ma | Technique | Reference |
|--|-------------------|------------------------|------------------------|----------------------------------|
| Heib Fm. (rhyolitic flow) | - | 770 ± 150 | U-Pb (Zr) | H.N.A. Priem (unpubl. data) |
| Heib Fm. (granitic pebbles) | - | 848 ± 61 735 ± 25 | U-Pb (Zr) U-Pb (Zr) | H.N.A. Priem (unpubl. data) ” |
| Tarr Fm. | upper greenschist | 616 ± 30 | Rb - Sr (WR) | Halpern & Tristan (1981) |
| Umm Zariq Fm. | amphibolite | 609 ± 12 | Rb - Sr (WR) | Bielski (1982) |
| Malhaq Fm. | amphibolite | 609 ± 12 | Rb -Sr (WR) | Bielski (1982) |
| Metavolcanic Ataqa schists (10 km N of Wadi Kid area) | lower amphibolite | 600 ± 10 566 -- 529 | Rb - Sr (WR) K - Ar | Ayalon et al. (1987) ” |
| Def. diorite/tonalite | - | 608 ± 7 | Rb - Sr (WR) | Siender et al. (1974) |
| Monzodiorite (150 km N of Wadi Kid area) | - | 625 ± 5 | U-Pb (Zr) | Beyth et al. (1994) |
| Dykes | - | 591 ± 9 | Rb - Sr (WR) | Stern & Manton (1987) |
| Younger Granite | - | 580 -- 530 | K - Ar (Bio) | Bielski (1982) |

5.3 The structural geology

5.3.1 Introduction

The late Precambrian tectonic history of the Sinai has been divided by most authors (e.g. Shimron 1980, 1983; Reymer 1983, 1984) into two deformation phases, namely D1 and D2. The results of our structural study are presented in Figure 5-3. Our study focuses on the D2 event because of its controversial nature in the tectonic models for the Sinai.

5.3.2 D1 structures

The S1 foliation is well developed in the lower-grade parts of the Heib Fm., such as the Beda Turbidites in the southern Wadi Kid area. The S1 slaty cleavage, formed by white micas, is easily recognizable in the shaly layers within these meta-turbidites. It is oblique to the bedding. Inversion of the bedding can be recognized through bedding and cleavage relationships and through sedimentary structures such as cross bedding and coarsening upward sequences. Upright isoclinal F1 folds are thus associated with the formation of the axial planar S1 foliation. Shimron (1983) concluded that D1 has taken place at shallow depth, soon after deposition. We agree with this, since the Heib Fm. was interpreted to have been deposited in an island-arc




Figure 5-3 Map showing structures related to Precambrian tectonics in the Wadi Kid area.

environment (Furnes et al. 1985), since these rocks display a low metamorphic grade where sedimentary structures are recognizable, and since accretion is the oldest tectonic process recognized in other parts of the Arabo-Nubian Shield (Bentor 1985). The D1 structures were thus formed in a compressional island-arc setting. F1 folds and the associated cleavage are occasionally recognized in the Umm Zariq Fm., with a steep foliation formed by muscovite and biotite. The D1 structures in this formation were overprinted by a later foliation.

5.3.3 D2 structures

5.3.3.1 S2 foliation

The S2 foliation is widely developed and dominates the studied area. It is flat-lying to gently dipping ($< 30^\circ$, Figure 5-4) and in most areas it totally masks the earlier foliation. It is a disjunctive and sometimes anastomosing foliation, formed by biotite, muscovite and chlorite. The grains in the microlithons show a rather strong preferred orientation.

The general trend in the Wadi Kid Complex is that S2 is better developed to the west

and north. S2 is best developed in the Malhaq, Umm Zariq and Quneia Fms. It is continuous through the higher-grade parts of the Heib Fm. into the Quneia Fm., indicating that the deformed tonalites and diorites of the Quneia Fm. were intruded before the development of S2. Rocks containing D1 structures overlie rocks displaying D2 structures. Slightly foliated granites were observed in the western part of the Wadi Kid Complex. These intrusives were dated at 580-530 Ma as the youngest rocks of the Sinai basement (Bielski 1982) and are thus critical for the timing of the deformation.




Figure 5-4 Contoured stereographic plot (equal area) of the poles to the S2 foliation in the Wadi Kid area. Contour interval = 2.0%, significance level = 3.0.

5.3.3.2 Lineations associated with S2

A lineation is associated with the S2 foliation and arises from an alignment of minerals such as andalusite and hornblende or from elliptical objects such as elongate xenoliths and pebbles. The lineation is sub-horizontal ($< 30^\circ$), plunging either NW or SE (Figure 5-5), that is in opposite directions due to later folding. The two trends reflect one strain regime.

The mineral lineation is best developed on the S2 plane in the schists. It is mainly defined by individual minerals that grew in one uniform direction: amphiboles, sillimanite, andalusite, quartz and feldspar. Some mineral lineations are formed by elongated pods of biotite, muscovite and by quartzo-feldspathic segregations. At some places the mica pods, andalusites, quartz and feldspars are mechanically elongated and define a stretching lineation (5-6a). It is concluded that the lineation was formed partly by stretching, partly by orientated growth. The stretching lineation indicates an extensional strain in NW-SE direction; this will be discussed later. In a number of cases the elongated minerals, mainly andalusites and hornblendes, grew in random directions; this was observed in the Umm Zariq and Malhaq Fms. Their significance will likewise be discussed later.

Elongated pebbles were found in the conglomerates of the high-grade parts of the Heib

Fm., and elongated xenoliths in the deformed intrusives of the Quneia Fm. They are parallel to the general trend of the mineral lineation.




Figure 5-5 Contoured stereographic plot (equal area) of the lineations in the Wadi Kid area, showing maxima at NW and SE. Contour interval = 2.0%, significance level = 3.0.

5.3.3.3 Dykes

The area is cross-cut by vertical felsic and mafic dyke swarms of Precambrian age. They are sub-parallel and strike 35 to 50°, i.e. NE-SW (Figure 5-7). Both types of dykes intruded at 590-580 Ma (Halpern 1980; Stern & Manton 1987). This age is close to the D2 ages. Along with their orientation, the contemporaneity suggests that the intrusion was related to the D2 phase.

Dyke swarms require lateral extension in the crust and the intrusion plane will be perpendicular to σ_3 , the direction of greatest extension (Price & Cosgrove 1990). The dykes thus indicate extension in the NW-SE direction at the time of D2, the same direction as can be deduced from the stretching lineation.

5.3.4 Evidence of non-coaxial strains during D2

5.3.4.1 Introduction

Previously, D2-structures have been interpreted as due to an irrotational and a coaxial strain regime (e.g. Reymer 1983, 1984, Shimron 1983, 1984). However, in the course of this study we consistently found that rotational strains were present. Several possible indicators of non-coaxial strain were observed: extensional crenulation cleavage, S-C shearbands, rotated clasts with pressure shadows, asymmetric folds, deformed elliptical objects and quartz c-axis fabrics. The importance and reliability of these indicators will be discussed below.

Figure 5-3 shows the distribution of shear indicators with the sense of shear and the recorded stretching and mineral lineations. The strain was evaluated for elliptical markers.




Figure 5-6 Photographs. a) Lineated hornblende, Malhaq Fm., northern Wadi Kid area. Area shown is 4.9mm x 3.4mm. b) Deformed xenoliths, Quneia Fm., NE part of Wadi Kid area. c) Rotated clast with pressure shadows indicating sinistral sense of shear Malhaq Fm., central Wadi Kid area. Area shown is 4.9mm x 3.4mm. d) Macroscopic extensional crenulation cleavage, indicating sinistral sense of shear, Umm Zariq Fm, central Wadi Kid area. Area shown is 4.9mm x 3.4mm.

5.3.4.2 Finite Strain Markers

Throughout the Wadi Kid area, elliptical markers, namely xenoliths, conglomerate pebbles and boudins, were found. They can be used for a qualitative and quantitative strain analysis. These markers are regarded as good indicators of the total finite strain (Lisle 1985), which is very useful in areas with a complicated strain history. Different methods were developed to determine the strain of deformed elliptical objects, taking into account that the initial shape was not spherical and that the markers had a non-random initial orientation.

The tests for non-coaxiality were performed on deformed xenoliths, conglomerate pebbles and boudins. The xenoliths were used to perform a strain analysis with the Rf/Φ method (Dunnet 1969; Lisle 1985). The deformed conglomerate pebbles and boudins were used for qualitative analysis and in this framework their relationships with other structural features were studied.

In the northeastern part of the Wadi Kid area, elliptical mafic xenoliths were found in a foliated and lineated syn-kinematic diorite which intruded during D2 (Figure 5-6b). The foliation in the schistose xenoliths continues into the diorite. The stretching lineation of hornblendes in the diorite is sub-parallel with the longest axes of the xenoliths.





Figure 5-7 Rose diagram of the orientation of the mafic and felsic dykes in the Wadi Kid area.

The basic method, used to compute the strain, is that of Ramsay-Dunnet modified by Lisle (1985) for elliptical markers with an initial ellipticity (R_i). This is a graphical method to compute the ellipticity of the strain ellipse (R_s) from R_f (the final ellipticity) and the angle between the long axis of the elliptical marker and the orientation plane, in the deformed state, Φ_s . The results of the R_f/Φ method were compared to the harmonic mean, H , which gives a good approximation for R_s (Lisle 1977).


The xenoliths in the Wadi Kid area have an elliptical form and show a uniform direction of their principal axes, with the longest axis (X-axis) sub-parallel with the mineral lineation. The Z-axis is the axis of greatest shortening, so that $X > Y > Z$. The foliation was chosen as the reference plane. The X : Z ratio was measured on a plane parallel with the lineation and perpendicular to the foliation. Φ_s is the angle between the X-axis and the foliation in the case of the X-Z section, and that between the Y-axis and the foliation in the case of the Y-Z section. Φ_s was chosen positive when the orientation of the X-axis was dipping steeper to the southeast than the reference plane. The Y : Z ratio was measured on a plane perpendicular to the lineation and perpendicular to the foliation.

Eighty-five measurements were made on the X-Z plane. The final ellipticity, R_f for X-Z ranges from 1.58 to 13.5. Figure 5-8a shows the R_f/Φ plot for X-Z. The plot shows an asymmetric distribution of points around the mean vector $\Phi = 3.41^\circ$. From the R_f/Φ diagram, values for $R_s = 4.6$ and $R_i = 2-2.5$ are deduced. The harmonic mean, H , is 4.71. From H and the R_f/Φ plot it can be concluded that $R_s = 4.6-4.7$. The mean vector, H , is 3.41° and so the distribution of points is asymmetric with respect to $\Phi = 0$. The longest axis of the plot is not parallel to the vertical base axis, revealing an internal asymmetry.

Forty measurements were made on the Y-Z plane. The R_f for Y-Z ranges from 2.167 to 11. Figure 5-8b shows the R_f/Φ diagram for Y-Z. The distribution of the points is symmetrical around the $\Phi = 0$ axis. From the diagram, values for $R_s = 4.2$ and $R_i = 1.5-2.0$ are deduced. The harmonic mean, H , gives a value of 3.95. Since H is expected to give lower values than R_s



a



b

Figure 5-8 R_f/Φ plots of deformed xenoliths, Quneia Fm., NE part of Wadi Kid area. a) The X-Z section reveals the asymmetry with respect to $\Phi = 0$ and the internal asymmetry. b) In the Y-Z section no asymmetry is visible.

(Lisle 1977), 3.95 should be a good indication for the maximum of R_s in the Y-Z plane.

Other deformed elliptical objects, such as elongated conglomerate pebbles and asymmetric elongated boudins, occur throughout the Wadi Kid area. The former were found in the relatively high-grade parts of the Heib Fm. in the central Wadi Kid area and the latter where S2 was developed. These elliptical objects have their longest axis sub-parallel with the mineral lineation and plunge slightly steeper to the southeast than the foliation, thus showing the same

structural relationships with the lineation and the foliation as the xenoliths. The deformed conglomerates and boudins show, like the deformed xenoliths, a movement of the top block to the northwest. The identical morphology and structural relations of all the deformed elliptical markers indicate a similar deformation history for all the total finite strain markers.

Conclusions on finite strain markers: The Rf/Φ method is a suitable method to analyse the strain of the xenoliths, but additional data from statistic and geometrical methods are required to obtain a precise indication of the amount of strain and the initial form. Lisle (1985) showed that H gives a good approximation for R_s and used this method in cases of asymmetric distribution in Rf/Φ plots. Best values for the $X : Y : Z$ relationship were calculated to be 4.6 : 3.95 : 1. From this relationship it is clear that a large component of flattening is present.

The mean vector of the X-Z plane is 3.41° and thus asymmetric with respect to $\Phi = 0$. The Rf/Φ plot for this plane also shows an internal asymmetric distribution. The mean vector Φ for the Y-Z plane is close to 0° and the Rf/Φ plot shows a symmetric distribution, and thus no asymmetry with respect to the reference plane exists. Internal asymmetries in Rf/Φ plots are ascribed to a preferred initial orientation of the elliptical objects (Lisle 1985), which can be explained by the fact that xenoliths in syn-kinematic intrusions will have a preferred initial orientation. The asymmetry with respect to $\Phi = 0$ is best explained as due to non-coaxial deformation (Le Theoff 1979, Choukroune et al. 1987). The X-axis is parallel to the stretching lineation but plunges steeper to the southeast than the foliation and lineation, and thus the finite strain markers indicate a NW-vergent movement of the top block during a regional shear event.

The deformed pebbles, xenoliths and boudins show the same relationships with other structural features: lineation and foliation. Since they were deposited or initially formed under different conditions, their final stage of deformation must have been similar. The asymmetry is found in the X-Z plane, parallel with the lineation and the movement direction. No asymmetry was observed in the Y-Z direction. This indicates non-coaxial deformation with the axis of rotation perpendicular to the X-Z section and movement of the top block to the northwest.

5.3.4.3 Kinematic indicators

Rotated clasts with asymmetric pressure shadows were observed in a number of thin sections from the Wadi Kid area (5-6c). They are regarded as good indicators of the sense of displacement (Simpson & Schmid 1983) if applied as indicated by Ten Brink (1996). The rotated pressure shadows are of the displacement-controlled type with deformable fibers. These fibers indicate relatively high temperatures, allowing dynamic recrystallization. Table 5-2 and Figure 5-3 show that shear took place in two opposite directions, with the top-to-the-NW direction dominating. Some clasts are accompanied by symmetric pressure shadows. This can indicate a component of coaxial deformation, or a reversal of the shear direction, re-deforming the fibers.

Throughout the area, *extensional crenulation cleavages* (e.c.c.) were observed. They develop in the late stages of shear zone activity (Platt & Vissers 1980) and are regarded as a good indication of local shear movement.

The e.c.c.'s in the Wadi Kid are found both on meso-scale (Figure 5-6d) and on micro-

scale. They contain biotites, white micas, hornblendes and andalusites, depending on the metamorphic assemblage in the country rock. Table 5-2 and Figure 5-3 show that the e.c.c.'s, like the rotated clasts, display a top-to-the-NW as well as a top-to-the-SE shear, with the NW transport dominating. Hornblende occurs mostly in the e.c.c.'s with a sense of shear to the northwest. White mica appears in e.c.c.'s with a NW sense of shear as well as in e.c.c.'s with a SE sense of shear, but is relatively abundant in the latter e.c.c.'s. We relate the formation of the white mica with a phase of retrograde metamorphism and so the SE sense of shear postdated the NW sense of shear.

Although *asymmetric folds* can be used as shear indicators, they are not very reliable (Bell & Hammond 1984). Their reliability improves when they are viewed perpendicular to the stretching lineation, as was done in this study. During progressive deformation, fold axes tend to rotate parallel to the movement direction and are thus of little value in determining the sense of movement. The few asymmetric folds that were found in the Wadi Kid area indicate a NW sense of shear.

5.3.4.4 Quartz c-axes as indicators for non-coaxial deformation

Quartz fabrics record deformation caused by plastic deformation mechanisms and can give information about the type and conditions of deformation (Price 1985). Coaxial deformation produces symmetrical patterns with respect to the principal axes of finite strain and thus with respect to the foliation, whereas non-coaxial deformation produces an asymmetric fabric. The latter points to a mylonitic origin of a schistose rock. The type of slip system in the quartz grains can be deduced from the quartz c-axis fabric, thus providing an indication of the relative temperature at which the deformation took place.

The samples used in the quartz c-axis studies were cut parallel to the lineation and perpendicular to the foliation. Twelve samples were chosen from the Wadi Kid Complex. A minimum of 125 grain measurements were made on each sample. Figure 5-9 shows the location and quartz fabrics of the studied samples. Figure 5-10 shows the position of the different maxima in a quartz c-axis plot (after Fueten et al. 1991); this plot was used for the interpretation of the fabrics. Table 5-2 summarizes the results of the quartz c-axis studies. On the basis of their patterns, the quartz c-axis fabrics were divided into three groups. These groups indicate different slip systems and consequently different conditions of deformation.

Samples WJ 9, WJ 22 and WJ 42 belong to the *first group*. Their fabrics display single girdles and maxima are best developed at I or II. The maxima are at angles of 30° with the lineation. This fabric indicates slip along the basal plane in the <a> direction plus minor activity on the rhomb and prism in an <a> direction and is indicative of relatively low temperatures of deformation (Jessel & Lister 1990).

WJ 42 comes from the transitional area between low-grade and high-grade Heib Fm. and formed at relatively low temperature. The fabric asymmetry indicates top-to-the-NW movement. Rotated clasts in this sample show a reverse shear direction. Quartz c-axis fabrics will, in most cases, record the last stages of deformation (Lister & Hobbs 1980) and it is also likely that the last increments of deformation will be concentrated in the quartz-rich bands. The

rotated clasts are thus indicative of the earlier stages of deformation and the quartz fabric of WJ 42 is reflecting a later overprint. Sample WJ 9 shows the same characteristics of deformation as WJ 42, but the mineralogy of this rock indicates a higher metamorphic grade. The fabric shows movement of the topblock to the northwest indicating deformation after peak metamorphism.

Sample WJ 22 did not show any macroscopic or microscopic shear indicators, but a clear quartz fabric asymmetry was observed, indicating movement of the top block to the northwest. The c-axis fabric indicates basal $\langle a \rangle$ slip with a significant rhombic component, which takes place at low-to-medium temperatures. From WJ 22 it can be concluded that deformation continued to take place until the later stages of cooling and thus relatively late in the development of the Sinai basement.

The *second group* of quartz c-axis fabrics is distinguished from other groups by the fact that the fabrics contain a maximum at III. This maximum is connected with the maxima at I and/or II via a girdle, depending on the shear-direction. The maximum at III indicates that the prism $\langle a \rangle$ slip system was active, which indicates that deformation took place at medium temperatures (Tullis et al. 1973; Schmid & Casey 1986). The basal $\langle a \rangle$ and prism $\langle a \rangle$ slip systems are thus active in this group.

Samples B 20, B 29, P 37 and WJ 29 belong to this group. In sample B 29, c-axes appear between III and I, indicating that also the rhomb $\langle a \rangle$ slip system was active and that deformation took place at intermediate-high temperatures. This sample indicates a movement of the top block to the southeast. Sample WJ 29 is a biotite schist from the high-grade part of the Heib Fm. near the contact with granitoid rocks. It displays a NW-vergent sense of shear. Sample B 20 is from the relatively high-grade rocks of the Malhaq Fm. and indicates a top-to-the-SE movement. Sample P 37 comes from the slightly foliated tonalites in the Wadi Quneia/Wadi Ratma area and displays a movement of the top-block to the northwest. It shows less evidence for rhomb slip, as the prism and basal sections are more isolated.

The samples of the *third group* have strong maxima at I and at II, but in all cases one of the two maxima was better developed than the other and no maxima are found in III. The angles between the I and II maxima are 50° . They resemble small-circle girdle fabrics and are a special case of type-I crossed girdles (Schmid & Casey 1986). This fabric shows strong flattening. The active slip systems were basal $\langle a \rangle$ with rhomb $\langle c+a \rangle$, indicative of deformation at medium to higher temperatures.

Samples B 18, B 61, P 16 and P 39 belong to this third group. Samples B 18 and B 61 are from the high-grade biotite schists of the Malhaq Fm. Both show a NW sense of shear with a rather strong flattening. Sample P 16 is from the high-grade metapelites of the Umm Zariq Fm. The fabric indicates movement to the southeast but other shear indicators in this sample show a NW movement. At other outcrops in this area SE-vergent shear indicators were observed. Sample P 39 is from the gneissic tonalites of the Quneia Fm., and displays a NW sense of shear.

All studied samples show a well-developed quartz c-axis fabric. The individual fabrics display an asymmetry, indicating non-coaxial deformation. The three groups were distinguished on the basis of the fabric, which is related to the temperature of deformation. Deformation temperatures vary from relatively low to intermediate-high metamorphic conditions. The

Table 5-2 Results of quartz c-axis studies from the Wadi Kid area. For locations see Figure 5-9.

| Sample | Rock type | Metamorphic grade | Fabric group (see text) | Likely slip system | Movement deduced from quartz c-axis fabrics | Movement deduced from other features | Comments |
|--------|---------------------|--|-------------------------|---|---|--------------------------------------|--------------------------------------|
| B 18 | hbl-bio schist | amphibolite | 3 | basal <a> and \pm rhomb <c+a> | NW | NW (e.c.c. and rotated clasts) | A component of flattening |
| B 20 | hbl-bio schist | amphibolite | 2 | prism <a> with basal <a> | SE | NW (e.c.c.) | - |
| B 29 | hbl-bio schist | upper greenschist-- lower amphibolite | 2 | prism <a> with basal <a> and rhomb <a> | SE | SE (rotated clasts) | A component of flattening |
| B 61 | hbl-bio schist | upper amphibolite | 3 | basal <a> and \pm rhomb <c+a> | NW | NW (e.c.c.) | A small component of flattening |
| P 16 | and-gnt-bio schist | amphibolite | 3 | basal <a> and \pm rhomb <c+a> | SE | NW (e.c.c.) | Outcrops in this area show SE shear. |
| P 17 | sill-gnt-bio schist | amphibolite | 2 | prism <a> with basal <a> | SE | SE (e.c.c.) | A component of flattening |
| P 37 | deformed tonalite | - | 2 | prism <a> with basal <a> | NW | NW (deformed xenoliths) | A small component of flattening |
| P 39 | deformed tonalite | - | 3 | basal <a> and \pm rhomb <c+a> | NW | - | A component of flattening |
| WJ 9 | and-gnt-bio gneiss | amphibolite | 1 | basal <a> and minor prism and rhomb <a> | NW | NW (e.c.c.) | - |
| WJ 22 | deformed granite | - | 1 | basal <a> with significant rhomb <a> | NW | - | Deformed younger granite |
| WJ 29 | bio schist | greenschist | 2 | prism <a> with basal <a> | NW | NW (e.c.c.) | A component of flattening |
| WJ 42 | bio schist | greenschist | 1 | basal <a> | SE | NW (rotated clasts) | - |




Figure 5-9 Locations and quartz c-axis fabrics of the studied samples listed in Table 2. Contour interval = 2.0%, significance level = 3.0.

single-girdle fabrics (group 1) come from the lower-grade rocks, which are found at the upper structural levels. Higher-grade rocks (groups 2 and 3) were deformed at higher temperatures and are from a structurally lower level throughout the complex. These high-grade samples show a more coaxial type of deformation. Movement reversals appear also in the quartz c-axis fabrics and will be discussed later.

Sample WJ 22 shows non-coaxial deformation in a relatively late granite (530 Ma,




Figure 5-10 Positions of the different maxima in the quartz c-axis plot.

Bielski 1982) implying that the deformation continued after the intrusion of this body.

5.4 Discussion

5.4.1 Interpretation of regional structural data

Our study indicates that in the Wadi Kid area a clear distinction can be made between D1 and D2 structures on the basis of the types of structures and their metamorphic grades. The D1 structures consist of an axial planar slaty cleavage, namely S1 foliations related to F1 isoclinal folds. The F1 folds in lower-grade rocks were formed in a compressional regime. No indicators of non-coaxial deformation were observed in these relatively low-grade rocks.

The second deformation phase, D2, formed a flat-lying foliation (S2) which is the most important structure in the area. The abundant evidence of asymmetry indicates that the areas of intense foliation-development mark D2 shear zones.

The elongated minerals, formed on S2 surfaces, are mostly orientated in NW-SE direction. Stretching of the elongated minerals indicates that peak-metamorphism started before or during D2. The growth in a preferred parallel orientation of hornblende, andalusite and sillimanite indicates that peak-metamorphism continued during D2. Randomly orientated hornblendes and andalusites were also found and could indicate either that peak-metamorphism outlasted D2 or that the randomly orientated minerals grew in low-strain pockets, implying that the strain regime was not active through the whole shear zone during the entire deformation phase. The latter interpretation seems to fit best the observations in the Wadi Kid area.

The Wadi Kid area is cross-cut by dykes in NE-SW direction, perpendicular to the NW-SE trending lineation. This indicates that extension took place in NW-SE direction.

The indicators of non-coaxial strain were only found in the areas where S2 was developed. Table 5-3 summarizes the different types of shear indicators with the sense of movement. Deformed pebbles, boudins and xenoliths are indicators of total finite strain and all show a non-coaxial deformation with movement of the top block to the northwest. Other non-coaxial shear indicators, such as rotated clasts with pressure shadows, extensional crenulation cleavages and quartz c-axis fabrics, predominantly produce a NW sense of shear but some

Table 5-3 The numbers and senses of movement of the different shear indicators, observed in the Wadi Kid area.

| Type of shear sense indicator | Macro. | | Micro. | |
|----------------------------------|--------|----|--------|----|
| | NW | SE | NW | SE |
| Deformed elliptical objects | 10 | - | - | - |
| Rotated clasts | 5 | 4 | 12 | 5 |
| Extensional crenulation cleavage | 20 | 11 | 21 | 8 |
| S-C fabrics | 1 | - | 1 | - |
| Asymmetric folds | 6 | 1 | - | - |
| Quartz c-axis fabrics | - | - | 7 | 5 |

indicate the reverse direction. From the structural map (5- 3) it is difficult to indicate certain areas with one sense of shear and other areas with the reverse sense, but it was rare that within one outcrop (i.e. on a metre scale) shear indicators with opposite sense were found. The distribution of the sense of shear is thus heterogeneous.

The duality in shear sense may be ascribed to coaxial deformation when flexural slip took place in different limbs of a fold. However, this does not agree with the trend of the S2 foliation and the direction of extension, nor with many of the asymmetric features. Because these latter features are abundant and form synthetically a well-developed stretching lineation, the foliation can be interpreted as a mylonitic foliation resulting from flat-lying shear zones. The main movement of the top block in the Wadi Kid area was interpreted to be to the northwest, since finite strain markers and some 70% of the shear indicators were found to indicate a NW sense of shear.

5.4.2 A core-complex model for the Wadi Kid area

A cross section of the Wadi Kid area according to our interpretation is shown in Figure 5-11. The relationship between the sub-horizontal mylonitic sequences formed in an extensional regime, the upper-crustal low-grade rocks related to an early compressive tectonic phase, the high metamorphic gradients, and the syntectonic granitoids as observed in the area, indicate together a core-complex development. This is investigated below. Much has been published about core complexes in a variety of localities such as the Basin and Range province, U.S.A. (Davis 1980; Davis & Lister 1988), the Canadian Cordillera (Malavieille 1987; Liu & Furlong 1993), the Aegean Sea, Greece (Lister et al 1984), and Variscan core complexes in the Massif Central, France (Malavieille 1993). Below, some of the most important characteristics found in these core complexes are reviewed and compared to those of the Wadi Kid Complex.

The ideal core complex consists of a core of granitoids, gneisses and other lower crustal rocks, overlain by a major mylonite zone. Relatively low-grade rocks, preserving relicts of an earlier deformation phase, overlie this zone. The mylonite zone tends to be sub-horizontal with a near-constant direction of the stretching lineation through the whole area (Davis 1980). The mylonites form at depths of at least 10 km and at temperatures of at least 500° C, with a thermal




Figure 5-11 Schematic interpreted cross-section through the Wadi Kid Complex. High-angle normal faults were formed during the opening of the Gulf of Aqaba in the Miocene. The sedimentary cover is found to the northwest of the Wadi Kid area mapped in Figure 1. Length of section is approximately 50 km, height is approximately 10 km.

gradient of $\geq 50^\circ/\text{km}$, and at pressures of 3.5 to 4.5 kbar (Davis & Lister 1988). The mylonitic zones normally have a thickness of about 1 km but may reach a thickness of 3.5 km (Davis & Lister 1988). The mylonitic rocks are generally interpreted as gently dipping low-angle normal shear zones of the type envisaged by Wernicke (1985), and are related to continental extension. The mylonites contain indicators of non-coaxial deformation which show the general direction of movement. However, in many cases a mixed mode of deformation, i.e. non-coaxial and coaxial deformation, is found and reversal of the sense of shear is observed (Malavieille 1993). The complicated deformation history makes kinematic interpretation difficult but this is characteristic of strain development in an extensional core-complex setting (Malavieille 1993). Undeformed younger intrusives cross-cut all other sequences of the complex.

The core in the Wadi Kid metamorphic complex consists of deformed tonalites and diorites of the Quneia Fm., which have a gneissic appearance. Gneissic rocks were also described in other parts of the Sinai (Shimron 1980; Stern & Manton 1987). The mylonitic Malhaq and Umm Zariq Fms. and the northern and western high-grade parts of the Heib Fm. form the mylonitic drape over the core. This drape is approximately 1.5 km thick. These formations contain a sub-horizontal foliation with a constant NW-SE trending stretching lineation. The low-grade rocks of the Heib Fm., overlying the mylonitic formations, form the upper part of the core complex, with the D1 compressive phase as a relict of an earlier

deformation event. Metamorphism in the mylonitic formations is of a LP/HT type (Shimron 1987). The thermal gradient was $\geq 50^\circ/\text{km}$ and the foliated formations were formed at a depth of at least 10 km (Reymer et al. 1984).

In the Wadi Kid area a large variety and large number of shear sense indicators were observed. The axis of rotation of the sense of asymmetry from the kinematic indicators is always perpendicular to the stretching lineation. About 70% of the shear sense indicators show movement of the top block to the northwest; the rest shows movement to the southeast (Table 5-3). Part of the non-coaxial strain markers show a coaxial component. All the finite strain markers, indicators of the total strain, point to a sense of shear to the northwest. For this reason the mylonitic formations are interpreted as low-angle shear zones with a movement of the top block to the northwest.

The formation of core complexes is accompanied by extensive magmatism. Basaltic intrusives and syn-kinematic dykes intruded perpendicular to the direction of extension (Anderson & Cullers 1990; Hazlett 1990). Diorites and granodiorites, often syn-kinematic, are the oldest intrusives in core complexes (Barton 1990; Anderson & Cullers 1990). Younger granitoids, intruded during the extensional phase, have a more alkaline granitoid composition (Barton 1990; Anderson & Cullers 1990). In the Wadi Kid area the dykes strike NE-SW, perpendicular to the lineation and thus indicating extension in the NW-SE direction during D2. The older intrusives are I-type granitoids, and younger intrusives are A-type alkaline granites (Ahmed et al. 1993). In other parts of the Arabo-Nubian Shield the older I-type granites are interpreted to have been related to a phase of crustal thickening during orogenic compression, while the A-type granites are interpreted to have been related to extension and crustal thinning when melts were arising from the mantle (Beyth et al. 1994; Greiling et al. 1994).

The features observed in core complexes such as that in the Wadi Kid area, are best explained as developing during orogenic collapse (Dewey 1988). The orogenic history starts with a period of compression and crustal thickening. When convergence slows down, thermal re-equilibration will start. Erosion and convective thinning of the root will cause extension and associated with this, there will be an increase in magmatic activity. The thermal gradient will increase rapidly. The extension may be accommodated by low-angle shear zones (Wernicke 1985). The metamorphism will show a clockwise P-T-t path, characterizing uplift in an extensional setting after a compressional phase, as is the case in the Wadi Kid area.

5.4.3 Reversal in the sense of shear and coaxial overprint in core complexes

The presence of opposing senses of shear and of components of coaxial deformation posed problems in the kinematic interpretation of the Wadi Kid area, where some 70% of the shear indicators show a top-to-the-NW movement and the rest shows movement to the southeast. Since all finite strain markers show the top-to-the-NW movement, the main deformational event is interpreted to have this sense of shear. Reversal of shear sense and minor components of coaxial deformation are regarded as characteristic features of core complexes (Malavieille 1987, 1993; Davis & Lister 1988; Spencer 1984; Reynolds & Lister 1990; Gautier & Brun 1994). Reynolds & Lister (1990), Malavieille (1993) and Gautier & Brun (1994) suggested that upward bending of the lower plate as an isostatic response to tectonic denudation and doming




Figure 5-12 Model for the different stages of the formation of the Wadi Kid Core Complex. At stage A, shortening took place with subduction in an island-arc setting, and extensive crustal thickening (D1). At stage B, convection at the root initiated extension. At C the actual collapse started with the formation of low-angle normal faults, including the detachment fault (D2a). At stage D granitoids intruded, causing local reversal of the sense of shear due to upwarping of the detachment fault (D2b). Stage E shows the situation as observed today (Figure 11) with the late Precambrian shear directions. Ages are interpreted from literature data (Table 4).

Table 5-4 Timing of the different phases of metamorphism and deformation in the Wadi Kid area as deduced from the literature (see 'Geological background').

* In the literature (e.g. Bielski 1982, Halpern & Tristan 1981) some D2 rocks were dated pre-620 Ma, whereas D1 was also dated at pre-620 Ma. To the north, older ages were recorded for D1. No dates are available for the low-grade rocks in the Wadi Kid area (e.g. the Beda Turbidites). These dates, interpreted by Bielski (1982) and Halpern & Tristan (1981) as D1, may very well reflect D2 ages. Mafic xenoliths resembling rocks of the Malhaq and Heib Fms were found in deformed diorites dated at 620 Ma and indicate that volcanism took place well before 620 Ma ago.

| Age | Deformation | Deformation features | Metamorphism (facies) |
|----------------|---------------------------|--|---------------------------------------|
| pre-620 Ma* | D1 | S1, F1 folds. | M1 (lower greenschist) |
| 620* --560 Ma* | D2 (Main phase) | S2, lineation, deformed elliptical objects, rotated clasts, e.c.c.'s, asymmetric folds, quartz c-axis fabrics. | M2 (upper greenschist to amphibolite) |
| post-560 Ma | D2 (Shear sense reversal) | ” | ” |

above plutons can cause shear reversal. In such cases the shear zone will undergo uplift and flexure, and parts of the shear zone will eventually dip to the opposite direction. Movement in the initial direction will stop in these parts of the shear zone, and reverse movement will start. The reverse sense of shear is thus a later overprint feature.

Gravimetric and aeromagnetic studies indicate the presence of elongate NE-SW trending intrusive bodies in the Sinai (Folkman & Assael 1980a, b; Ghazala 1995). The reversal of shear sense took place in the later stages of the development of the Wadi Kid area and is thus related to the intrusion of the younger NE-SW trending granitic plutons near this area, when uplift at LP/HT conditions occurred. These intrusions caused the uplift and flexure of the shear zone, and normal movement to the southeast started at their southeasterly sides. The SE movement, as recorded in some 30% of the shear indicators, is thus an overprint. This is in accordance with our metamorphic studies, which show that white micas are predominantly present in the SE-vergent shear-indicators. The SE movement was due to normal movement in a shear zone along the SE side of an intruding granite, northwest of the Wadi Kid area, at the later stages of the formation of the core complex. Ayalon et al. (1987) found evidence in amphibolite schists for a second thermal pulse at 560-530 Ma, after the main phase at 600 Ma. This phase can be related to the intrusion of younger plutons and so the shear reversal event can be dated at 560-530 Ma.

The general trend of the quartz c-axis fabrics shows a transition from a stronger coaxial component at the lower structural levels to a stronger non-coaxial component at higher structural levels. This is in agreement with findings of Malavieille (1993) for quartz c-axis studies of the core complexes in the Basin and Range province, U.S.A. He concluded that low-angle normal shear zones will show a strain distribution through such a shear zone as described above for the quartz c-axis fabrics, contrary to compressional settings, where a homogeneous distribution of strain is to be expected (Malavieille 1993). He attributes this variation in the style of deformation through extensional shear zones to the different amount of rotational strain between the highly deformed upper crustal rocks of the hanging wall and the less deformed basement in the footwall, below the shear zone. This also explains the coaxial component of

strain, as observed in the deformed xenoliths within the gneissic lower-crustal diorites. Malavieille (1993) also demonstrated that deformation, in the shearzone, will be increasingly non-coaxial away from the axis of the dome. This all explains the complex strain patterns in the Wadi Kid area.

5.4.4 A model for the tectonic history of the Wadi Kid area

Our model for the tectonic development of the Wadi Kid area in the late Precambrian is summarized in Figure 5-12 and Table 5-4. In an island-arc setting, volcanics of the Heib and Tarr Fms. were deposited prior to 620 Ma. In this period, compression caused by the subduction was responsible for the D1 deformation phase. At the same time, or slightly afterwards, I-type granitoids intruded. After a phase of crustal thickening related to the subduction of an oceanic plate, orogenic collapse started with extension in NW-SE direction, starting at 620 Ma. The schistose units of the Malhaq, the Umm Zariq, and the northern and western Heib Fms, and the deformed tonalites and diorites testify to the low-angle shear zone. During the extensional phase, dyke swarms intruded with a NE-SW strike. All the units were intruded by granites. The main movement along the shear zone was to the northwest, as deduced from the indicators of finite strain. Doming of younger NE-SW trending granitoid bodies near the Wadi Kid area caused upwarping of the main shear zone and reversal of the sense of shear. The heterogeneous strain distribution through the shear zone was responsible for the coaxial component as observed in the deformed xenoliths and the c-axis fabrics. Deformation continued until after intrusion of the late granites.

The core-complex model explains why the D1 shortening phase is only found at the upper crustal levels, while the horizontal cleavage is observed at lower crustal levels.

5.5 Conclusions

The Precambrian rocks in the Wadi Kid area display two main phases of deformation: D₁ and D₂. The present study offers new data and interpretations and leads to the following conclusions:

- 1) D₁ is a compressional phase related to island-arc subduction as recorded from F₁ folds and the S₁ foliation. It took place before 620 Ma.
- 2) A D₂ deformation phase, dated at 620-580 Ma, was responsible for the main structural features in the area, i.e. the sub-horizontal foliation and the well-developed stretching lineation. The foliation was formed due to mylonitization in a low-angle normal shear zone. The mineralogy indicates that the deformation took place at LP/HT conditions.
- 3) Shear indicators and quartz c-axis fabrics confirm that non-coaxial deformation took place. Finite strain markers such as deformed xenoliths and pebbles indicate a movement of the top block to the northwest in the shear zones. Most of the shear indicators, such as rotated clasts with pressure shadows, extensional crenulation cleavage, and quartz c-axis fabrics also indicate NW-vergent shear, but some indicate movement to the southeast.
- 4) The reversal of movement along mylonites was caused by upwarping due to the intrusion of granitoid bodies near the Wadi Kid area at the latest stages of the Pan-African orogenesis, 560-

530 Ma ago.

5) Dyke swarms, dated at 590-580 Ma, striking NE-SW and perpendicular to the lineation indicate extension in NW-SE direction.

6) The structures and metamorphism observed, support a core complex model for the Wadi Kid area.

7) The transition from the D1 compressional regime to the D2 extensional regime was due to orogenic collapse.

