

References

- Akhmetiev, M.A. and Beniamovski, V.A., 2004. Palaeocene and Eocene of Western Eurasia (Russian sector) - stratigraphy, palaeogeography, climate. *Neues Jahrbuch für Geologie und Paläontologie - Abhandlungen*, 234: 137-181.
- Ali, J.R., Kent, D.V. and Hailwood, E.A., 2000. Magnetostratigraphic reinvestigation of the Palaeocene/Eocene boundary interval in Hole 690B, Maud Rise, Antarctica. *Geophysical Journal International*, 141: 639-646.
- Allison, P.A., Wignall, P.B. and Brett, C.T., 1995. Palaeo-oxygenation: Effects and recognition. In: D.W.J. Bosence and P.A. Allison (Editors), *Marine Palaeoenvironmental Analysis from Fossils*. Geological Society Special Publication, 83, pp. 97-112.
- Alroy, J., Koch, P.L. and Zachos, J.C., 2000. Global climate change and North American mammalian evolution. *Paleobiology*, 26(4 SUPPS): 259-288.
- Anderson, D.M., Lively, J.J., Reardon, E.M. and Price, C.A., 1985. Sinking characteristics of dinoflagellate cysts. *Limnology and Oceanography*, 30: 1000-1009.
- Anderson, D.M., Taylor, C.D. and Armbrust, E.V., 1987. The effects of darkness and anaerobiosis on dinoflagellate cyst germination. *Limnology and Oceanography*, 32: 340-351.
- Archer, D., in press. Methane hydrates and anthropogenic climate change. *Geophysical Review Letters*.
- Aubry, M.-P., 2001. Provincialism in the photic zone during the LPTM. In: A. Ash and S. Wing (Editors), *Climate and Biota of the Early Paleogene*, Powell, Wyoming.
- Aubry, M.-P., Berggren, W.A., Stott, L. and Sinha, A., 1996. The upper-Paleocene-lower Eocene stratigraphic record and the Paleocene-Eocene boundary carbon isotope excursion: implications for geochronology. In: R.W.O.B. Knox, R.M. Corfield and R.E. Dunay (Editors), *Correlation of the Early Paleogene in Northwest Europe*. Geological Society of London Special Publication, Volume 101. Geological Society of London London, pp. 353-380.
- Aubry, M.P., Lucas, S.G. and Berggren, W.A. (Editors), 1998. *Late Paleocene - Early Eocene Climatic and Biotic Events in the Marine and Terrestrial Records*. Columbia University Press, New York.
- Aubry, M.P. and Ouda, K., 2003. Introduction. In: K. Ouda and M.P. Aubry (Editors), *The Upper Paleocene-Lower Eocene of the Upper Nile Valley, Part 1, Stratigraphy*. Micropaleontology Press, New York, pp. ii-iv.
- Ayres, M.G., Bilal, M., Jones, R.W., Slentz, L.W., Tartir, M. and Wilson, A.O., 1982. Hydrocarbon habitat in main producing areas, Saudi Arabia. *American Association of Petroleum Geologists Bulletin*, 66: 1-9.
- Backer, L.C., Fleming, L.E., Rowan, A., Cheng, Y.-S., Benson, J., Pierce, R.H., Zaias, J., Bean, J., Bossart, G.D., Johnson, D., Quimbo, R. and Baden, D.G., 2003. Recreational exposure to aerosolized brevetoxins during Florida red tide events. *Harmful Algae*, 2: 19-28.
- Backhouse, J., 1988. Late Jurassic and Early Cretaceous palynology of the Perth Basin, western Australia. *Geological Survey of Western Australia Bulletin*, 135: 1-233.
- Backman, J., 1986. late Paleocene to middle Eocene calcareous nannofossil biochronology from Shatsky Rise, Walvis Ridge and Italy. *Palaeogeography, Palaeoclimatology, Palaeoecology*, 57(43-59).
- Backman, J., Moran, K., McInroy, D.B., Mayer, L.A. and Expedition 302 Scientists, 2006. *Proceedings of the Integrated Ocean Drilling Program, 302*. Integrated Ocean Drilling Program Management International, Inc., College Station TX.
- Bains, S., Corfield, R.M. and Norris, G., 1999. Mechanisms of climate warming at the end of the Paleocene. *Science*, 285: 724-727.
- Bains, S., Norris, R.D., Corfield, R.M. and Faul, K.L., 2000. Termination of global warmth at the Paleocene/Eocene boundary through productivity feedbacks. *Nature*, 407: 171-174.
- Barber, D.C., Dyke, A., Hillaire-Marcel, C., Jennings, A.E., Andrews, J.T., Kerwin, M.W., Bilodeau, G., McNeely, R., Southon, J., Morehead, M.D. and Gagnon, J.-M., 1999. Forcing of the cold event of 8,200 years ago by catastrophic drainage of Laurentide lakes. *Nature*, 400: 344-348.
- Batten, D.J., Gray, J. and Harland, R., 1999. Palaeoenvironmental significance of a monospecific assemblage of dinoflagellate cysts from the Miocene Clarkia Beds, Idaho, USA. *Palaeogeography, Palaeoclimatology, Palaeoecology*, 153: 161-177.
- Batten, D.J. and Lister, J.K., 1988. Early Cretaceous dinoflagellate cysts and chlorococcalean

References

- algae from freshwater and low salinity palynofacies in the English Wealden. *Cretaceous Research*, 9: 337-367.
- Beard, K.C., 1998. East of Eden; Asia as an important center of taxonomic origination in mammalian evolution. In: K.C. Beard and M.R. Dawson (Editors), Dawn of the age of mammals in Asia; Bulletin of the Carnegie Museum of Natural History No. 34, pp. 5-39.
- Beard, K.C. and Dawson, M.R., 1999. Intercontinental dispersal of Holarctic land mammals near the Paleocene/Eocene boundary; paleogeographic, paleoclimatic and biostratigraphic implications. *Bulletin de la Société Géologique de France*, 170(5): 697-706.
- Beerling, D.J., 2000. Increased terrestrial carbon storage across the Palaeocene-Eocene boundary. *Palaeogeography, Palaeoclimatology, Palaeoecology*, 161: 395-405.
- Benedek, P.N.v. and Gocht, H., 1981. *Thalassiphora pelagica* (Dinoflagellata, Tertiär): Elektronenmikroskopische Untersuchung und Gedanken zur Paläobiologie. *Palaeontographica (B)*, 180: 39-64.
- Berger, W.H., Smetacek, V.S. and Wefer, G., 1989. Ocean productivity and paleoproductivity - an overview. In: W.H. Berger, V.S. Smetacek and G. Wefer (Editors), Productivity of the Ocean: Present and Past, Life Sciences Research Report 44, pp. 1-34.
- Berggren, W.A., Kent, D.V., Obradovich, J.D. and Swisher, C.C., III, 1992. Towards a revised Paleogene geochronology. In: D.R. Prothero and W.A. Berggren (Editors), Eocene-Oligocene Climatic and Biotic Evolution. Princeton University Press, Princeton, pp. 29-45.
- Berggren, W.A., Kent, D.V., Swisher, C.C., III., and Aubry, M.-P., 1995. A revised Cenozoic geochronology and chronostratigraphy. In: W.A. Berggren, D.V. Kent and J. Hardenbol (Editors), Geochronology, Time Scales and Global Stratigraphic Correlation. SEPM (Society for Sedimentary Geology) Special Publication 54, Tulsa, Oklahoma, USA, pp. 129-212.
- Bertrand, P., Shimmield, G., Martinez, P., Grousset, F., Jorissen, F., Paterne, M., Pujol, C., Bouloubassi, I., Buat Menard, P., Peypuquet, J.-P., Beaufort, L., Sicre, M.-A., Lallier-Verges, E., Foster, J.M. and Ternois, Y., 1996. The glacial ocean productivity hypothesis: The importance of regional temporal and spatial studies. *Marine Geology*, 130: 1-9.
- Bice, K.L. and Marotzke, J., 2001. Numerical evidence against reversed thermohaline circulation in the warm Paleocene/Eocene ocean. *Journal of Geophysical Research*, 106(C6): 11,529-11,542.
- Blackman, R.B. and Tukey, J.W., 1958. The measurement of Power Spectra From the Point of View of Communication Engineering. Dover Publications, New York.
- Boersma, A., Premoli Silva, I. and Hallock, P., 1998. Trophic models for the well-mixed and poorly mixed warm oceans across the Paleocene/Eocene epoch boundary. In: M.-P. Aubry, S.G. Lucas and W.A. Berggren (Editors), Late Paleocene - early Eocene climatic and biotic events in the marine and terrestrial records. Columbia University Press, New York, pp. 204-213.
- Boessenkool, K.P., Van Gelder, M.-J., Brinkhuis, H. and Troelstra, S.R., 2001. Distribution of organic-walled dinoflagellate cysts in surface sediments from transects across the Polar Front offshore southeast Greenland. *Journal of Quaternary Science*, 16 (7): 661-666.
- Bohaty, S. and Zachos, J.C., 2003. Significant Southern Ocean warming event in the late middle Eocene. *Geology*, 31(11): 1017-1020.
- Bold, H.C., 1973. Morphology of Plants. Harper, New York, 668 pp.
- Bolle, M.-P., Pardo, A., Hinrichs, K.-U., Adatte, T., von Salis, K., Burns, S., Keller, G. and Muzylev, N., 2000. The Paleocene-Eocene transition in the marginal northeastern Tethys (Kazakhstan and Uzbekistan). *International Journal of Earth Sciences*, 89: 390-414.
- Boulter, M.C. and Manum, S.B., 1989. The Brito-Arctic igneous province flora around the Paleocene/Eocene boundary. In: O. Eldholm, J. Thiede and E. Taylor (Editors), Proceedings of the Ocean Drilling Program, Scientific Results. Ocean Drilling Program, College Station, TX, pp. 663-680.
- Bowen, G.J., Beerling, D.J., Koch, P.L., Zachos, J.C. and Quattlebaum, T., 2004. A humid climate state during the Palaeocene/Eocene thermal maximum. *Nature*, 432(7016): 495-499.
- Bowen, G.J., Bralower, T.J., Delaney, M.L., Dickens, G.R., Kelly, D.C., Koch, P.L., Kump, L.R., Meng, J., Sloan, L.C., Thomas, E., Wing, S.L. and Zachos, J.C., 2006. Eocene Hyperthermal Event Offers Insight Into Greenhouse Warming. *EOS, Transactions, American Geophysical Union*, 87(17): 165, 169.
- Bowen, G.J., Clyde, W.C., Koch, P.L., Ting, S.Y., Alroy, J., Tsubamoto, T., Wang, Y.Q. and Wang, Y., 2002. Mammalian dispersal at the Paleocene/Eocene boundary. *Science*, 295: 2062-2065.
- Bowen, G.J., Koch, P.K., Gingerich, P.D., Norris, R.D., Bains, S. and Corfield, R.M., 2001. Refined isotope stratigraphy across the continental Paleocene-Eocene boundary on Polecat Bench in the Northern Bighorn Basin. In: P.D. Gingerich (Editor), Paleocene-Eocene Stratigraphy and Biotic Change in the Bighorn and Clarks Fork Basins, Wyoming. University of Michigan Papers on Paleontology 33.
- Bowen, G.J., Koch, P.L., Meng, J., Ye, J. and Ting, S., 2005. Age and correlation of fossiliferous Late Paleocene-Early Eocene strata of the Erlan Basin, Inner Mongolia, China. *American*

References

- Museum Novitates, 3474: 1-26.
- Bradford, M.R. and Wall, D.A., 1984. The Distribution of Recent Organic-Walled Dinoflagellate Cysts in the Persian Gulf, Gulf of Oman, and northwestern Arabian Sea. *Palaeontographica*, 192: 16-84.
- Bralower, T.J., 2002. Evidence of surface water oligotrophy during the Paleocene-Eocene thermal maximum: nanofossil assemblage data from Ocean Drilling Program Site 690, Maud Rise, Weddell Sea. *Paleoceanography*, 17(2): 10.1029/2001PA000662.
- Bralower, T.J., Kelly, D.C. and Thomas, E., 2004. Comment on "Coccolith Sr/Ca records of productivity during the Paleocene-Eocene thermal maximum from the Weddell Sea" by Heather M. Stoll and Santo Bains. *Paleoceanography*, 17(PA1014): doi:10.1029/2003PA000953.
- Bralower, T.J., Thomas, D.J., Zachos, J.C., Hirschmann, M.M., Röhl, U., Sigurdsson, H., Thomas, E. and Whitney, D.L., 1997. High-resolution records of the late Paleocene thermal maximum and circum-Caribbean volcanism: Is there a causal link? *Geology*, 25(11): 963-966.
- Bralower, T.J., Zachos, J.C., Thomas, E., Parrow, M., Paull, C.K., Kelly, D.C., Premoli Silva, I., Sliter, W.V. and Lohmann, K.C., 1995. Late Paleocene to Eocene paleoceanography of the equatorial Pacific Ocean: stable isotopes recorded at Ocean Drilling Program Site 865, Allison Guyot. *Paleoceanography*, 10: 841-865.
- Brasier, M.D., 1985. Fossil indicators of nutrient levels. 1: Eutrophication and climate change. In: D.W.J. Bosence and P.A. Allison (Editors), *Marine Palaeoenvironmental Analysis from Fossils*, Geological Society Special Publication 83, pp. 113-132.
- Brenner, W., 2001. Organic-walled microfossils from the central Baltic Sea, indicators of environmental change and base for ecostratigraphic correlation. *Baltica*, 14: 40-51.
- Brinkhuis, H., 1994. Late Eocene to Early Oligocene dinoflagellate cysts from the Priabonian type-area (Northeast Italy); biostratigraphy and palaeoenvironmental interpretation. *Palaeogeography, Palaeoclimatology, Palaeoecology*, 107: 121-163.
- Brinkhuis, H. and Biffi, U., 1993. Dinoflagellate cyst stratigraphy of the Eocene/Oligocene transition in Central Italy. *Marine Micropaleontology*, 22: 131-183.
- Brinkhuis, H., Bujak, J.P., Smit, J., Versteegh, G.J.M. and Visscher, H., 1998. Dinoflagellate-based sea surface temperature reconstructions across the Cretaceous-Tertiary boundary. *Palaeogeography, Palaeoclimatology, Palaeoecology*, 141: 67-83.
- Brinkhuis, H., Huber, M., Schellenberg, S.A., Stickley, C.E., Sluijs, A., Warnaar, J., Williams, G.L., Exon, N.F. and Kennett, J.P., 2002a. Was Antarctica kept warm by subtropical waters in the Eocene? Part 1: Evidence from biotic endemism, American Geophysical Union Fall Conference, San Francisco, pp. Abstract volume, pp. F925.
- Brinkhuis, H., Huber, M., Schellenberg, S.A., Stickley, C.E., Sluijs, A., Warnaar, J., Williams, G.L., Exon, N.F. and Kennett, J.P., 2002b. Was Antarctica kept warm by subtropical waters in the Eocene? Part 2: climate model results, American Geophysical Union Fall Conference, San Francisco, pp. Abstract volume, pp. F926.
- Brinkhuis, H., Huber, M., Sluijs, A., Zachos, J.C. and Warnaar, J., 2003a. The end of the Early Eocene Climatic Optimum: Evidence for Concomitant Cooling of Southern Ocean Surface Waters and Global Deep Waters From Dinoflagellate Endemism, AGU Fall Conference, San Francisco. Abstract volume., San Francisco, pp. F887.
- Brinkhuis, H., Munsterman, D.K., Sengers, S., Sluijs, A., Warnaar, J. and Williams, G.L., 2003b. Late Eocene to Quaternary dinoflagellate cysts from ODP Site 1168, Off western Tasmania. In: N.F. Exon, J.P. Kennett and M. Malone (Editors), *Proceedings Ocean Drilling Program, Scientific Results*. Available from World Wide Web: http://www-odp.tamu.edu/publications/189_SR/105/105.htm. College Station, Texas, pp. 1-36.
- Brinkhuis, H., Powell, A.J. and Zevenboom, D., 1992. High-resolution dinoflagellate cyst stratigraphy of the Oligocene/Miocene transition interval in north-west and central Italy. In: M.J. Head and J.H. Wrenn (Editors), *Neogene and Quaternary Dinoflagellate Cysts and Acritarchs*. American Association of Stratigraphic Palynologists Foundation, Dallas, pp. 219-258.
- Brinkhuis, H., Romein, A.J.T., Smit, J. and Zachariasse, W.J., 1994. Danian-Selandian dinoflagellate cysts from lower latitudes with special reference to the El Kef section, NW Tunisia. *Geologiska Föreningens i Stockholm Förhandlingar (GFF; Transactions of the Geological Society in Stockholm)*, 116(46-48).
- Brinkhuis, H. and Schiøler, P., 1996. Palynology of the Geulhemmerberg Cretaceous/Tertiary boundary section (Limburg, SE Netherlands). *Geologie en Mijnbouw*, 75(2/3): 193-213.
- Brinkhuis, H., Schouten, S., Collinson, M.E., Sluijs, A., Damsté, J.S.S., Dickens, G.R., Huber, M., Cronin, T.M., Onodera, J., Takahashi, K., Bujak, J.P., Stein, R., van der Burgh, J., Eldrett, J.S., Harding, I.C., Lotter, A.F., Sangiorgi, F., Cittert, H.v.K.-v., de Leeuw, J.W., Matthiessen, J., Backman, J., Moran, K. and the Expedition, S., 2006. Episodic fresh surface waters in the Eocene Arctic Ocean. *Nature*, 441(7093): 606-609.
- Brinkhuis, H., Sengers, S., Sluijs, A., Warnaar, J. and Williams, G.L., 2003c. Latest Cretaceous to earliest Oligocene, and Quaternary dinoflagellate cysts from ODP Site 1172, East Tasman Plateau. In: N.F. Exon, J.P. Kennett and M. Malone (Editors), *Proceedings Ocean*

References

- Drilling Program, Scientific Results. Available from World Wide Web: http://www-odp.tamu.edu/publications/189_SR/106/106.htm. College Station, Texas, pp. 1-48.
- Bucefalo-Palliani, R., Mattioli, E. and Riding, J.B., 2002. The response of marine phytoplankton and sedimentary organic matter to the early Toarcian (Lower Jurassic) oceanic anoxic event in northern England. *Marine Micropaleontology*, 46(3-4): 223-245.
- Bujak, J.P. and Brinkhuis, H., 1998. Global warming and dinocyst changes across the Paleocene/Eocene Epoch boundary. In: M.-P. Aubry, S.G. Lucas and W.A. Berggren (Editors), *Late Paleocene - early Eocene biotic and climatic events in the marine and terrestrial records*. Columbia University Press, New York, pp. 277-295.
- Bujak, J.P. and Williams, G.L., 1979. Dinoflagellate diversity through time. *Marine Micropaleontology*, 4: 1-12.
- Butler, R.F., Gingerich, P.D. and Lindsay, E.H., 1981. Magnetic polarity stratigraphy and biostratigraphy of Paleocene and lower Eocene continental deposits, Clarks Fork Basin, Wyoming. *Journal of Geology*, 89: 299-316.
- Bybell, L.M., Gibson, T.G., Heilmann-Clausen, C. and Zachos, J.C., 2001. Events associated with the Late Paleocene Thermal Maximum in the Northeastern U.S., *Climate and Biota of the Early Paleogene 2001*, Powell, Wyoming, USA.
- Caballero, R. and Langen, P.L., 2005. The dynamic range of poleward energy transport in an atmospheric general circulation model. *Geophysical Research Letters*, 32(L02705): doi:10.1029/2004GL021581.
- Caldeira, K. and Wickett, M.E., 2003. Anthropogenic carbon and ocean pH. *Nature*, 425(6956): 365.
- Cande, S. and Kent, D.V., 1995. Revised calibration of the geomagnetic polarity timescale for the Late Cretaceous and Cenozoic. *Journal of Geophysical Research*, 111: 6093-6095.
- Cembella, A.D., Quilliam, M.A., Lewis, N.I., Bauder, A.G., Dell'Aversano, C., Thomas, K., Jellett, J. and Cusack, R.R., 2002. The toxigenic marine dinoflagellate *Alexandrium tamarense* as the probable cause of mortality of caged salmon in Nova Scotia. *Harmful Algal Blooms*, 1: 313-325.
- Charisi, S.D. and Schmitz, B., 1998. Paleocene to early eocene paleoceanography of the Middle East: the $\delta^{13}\text{C}$ and $\delta^{18}\text{O}$ isotopes from foraminiferal calcite. *Paleoceanography*, 13(1): 106-118.
- Clyde, W.C. and Gingerich, P.D., 1998. Mammalian community response to the latest Paleocene thermal maximum: An isotaphonomic study in the northern Bighorn Basin, Wyoming. *Geology*, 26: 1011-1014.
- Clyde, W.C., Khan, I.H. and Gingerich, P.D., 2003. Stratigraphic response and mammalian dispersal during initial India-Asia collision: Evidence from the Ghazij Formation, Balochistan, Pakistan. *Geology*, 31(12): 1097-1100.
- Cocconi, R., Basso, D., Brinkhuis, H., Galeotti, S., Gardin, S., Monechi, S. and Spezzaferri, S., 2000. Marine biotic signals across a late Eocene impact layer at Massignano, Italy: Evidence for long-term environmental perturbations? *Terra Nova*, 12: 258-263.
- Cocconi, R., Di-Leo, R., Galeotti, S. and Monechi, S., 1994. Integrated biostratigraphy and benthic foraminiferal faunal turnover across the Paleocene-Eocene boundary at Trabakua Pass section, northern Spain. *Palaeopelagos*, 4: 87-100.
- Cogné, J.-P. and Humler, E., 2006. Trends and rhythms in global seafloor generation rate. *Geochemistry, Geophysics, Geosystems*, 7(Q03011): doi:10.1029/2005GC001148.
- Cojan, I., Moreau, M.G. and Stott, L.E., 2000. Stable carbon isotope stratigraphy of the Paleogene pedogenic series of southern France as a basis for continental-marine correlation. *Geology*, 28(3): 259-262.
- Collinson, M.E., Hooker, J.J. and Grocke, D.R., 2003. Cobham lignite bed and pencontemporaneous macrofloras of southern England: A record of vegetation and fire across the Paleocene-Eocene Thermal Maximum. In: S.L. Wing, P.D. Gingerich, B. Schmitz and E. Thomas (Editors), *Causes and Consequences of Globally Warm Climates in the Early Paleogene*. Geological Society of America Special Paper 369. Geological Society of America, Boulder, Colorado, pp. 333-349.
- Collinson, M.E., Steart, D., Handley, L., Pancost, R., Scott, A.C., Glasspool, I., Hooker, J. and Stott, A., 2006. Fire regimes and palaeoenvironments across the onset of the Palaeocene/Eocene thermal maximum, S. England, *Climate and Biota of the Early Paleogene*, Bilbao.
- Colosimo, A.B., Bralower, T.J. and Zachos, J.C., 2005. Evidence for lysocline shoaling at the Paleocene/Eocene Thermal Maximum on Shatsky Rise, northwest Pacific. In: T.J. Bralower, I. Premoli-Silva and M.J. Malone (Editors), *Proceedings of the Ocean Drilling Program 198*, Available from World Wide Web: http://www-odp.tamu.edu/publications/198_SR/VOLUME/CHAPTERS/112.PDF, pp. 1-36.
- Corradini, D. and Biffi, U., 1988. Etude des dinokystes a la limite Messinien-Pliocene dans la coupe Cava Serredi, Toscane, Italie (Dinocysts at the Messinian-Pliocene boundary in the Cava Serredi section, Tuscany, Italy). *Bulletin des Centres de Recherches Exploration-Production Elf-Aquitaine*, 12(1): 221-236.

References

- Cramer, B.S., Aubry, M.-P., Miller, K.G., Olsson, R.K., Wright, J.D. and Kent, D.V., 1999. An exceptional chronologic, isotopic, and clay mineralogic record of the latest Paleocene thermal maximum, Bass River, NJ, ODP 174AX. *Bulletin de la Société Géologique de France*, 170: 883-897.
- Cramer, B.S. and Kent, D.V., 2005. Bolide summer: The Paleocene/Eocene thermal maximum as a response to an extraterrestrial trigger. *Palaeogeography, Palaeoclimatology, Palaeoecology*, 224(1-3): 144-166.
- Cramer, B.S., Wright, J.D., Kent, D.V. and Aubry, M.-P., 2003. Orbital climate forcing of $\delta^{13}\text{C}$ excursions in the late Paleocene-early Eocene (chrons C24n-C25n). *Paleoceanography*, 18(4): 10.1029/2003PA000909.
- Crouch, E.M., 2001. Environmental change at the time of the Paleocene-Eocene biotic turnover. *Laboratory of Palaeobotany and Palynology Contribution Series*, 14, 216 pp.
- Crouch, E.M. and Brinkhuis, H., 2005. Environmental change across the Paleocene-Eocene transition from eastern New Zealand: A marine palynological approach. *Marine Micropaleontology*, 56(3-4): 138-160.
- Crouch, E.M., Brinkhuis, H., Visscher, H., Adatte, T. and Bolle, M.-P., 2003a. Late Paleocene - early Eocene dinoflagellate cyst records from the Tethys: Further observations on the global distribution of *Apectodinium*. In: S.L. Wing, P.D. Gingerich, B. Schmitz and E. Thomas (Editors), *Causes and Consequences of Globally Warm Climates in the Early Paleogene*. Geological Society of America Special Paper 369. Geological Society of America, Boulder, Colorado, pp. 113-131.
- Crouch, E.M., Dickens, G.R., Brinkhuis, H., Aubry, M.-P., Hollis, C.J., Rogers, K.M. and Visscher, H., 2003b. The *Apectodinium* acme and terrestrial discharge during the Paleocene-Eocene thermal maximum: new palynological, geochemical and calcareous nannoplankton observations at Tawanui, New Zealand. *Palaeogeography, Palaeoclimatology, Palaeoecology*, 194: 387-403.
- Crouch, E.M., Heilmann-Clausen, C., Brinkhuis, H., Morgans, H.E.G., Rogers, K.M., Egger, H. and Schmitz, B., 2001. Global dinoflagellate event associated with the late Paleocene thermal maximum. *Geology*, 29(4): 315-318.
- Crouch, E.M. and Visscher, H., 2003. Terrestrial vegetation record across the initial Eocene thermal maximum at the Tawanui marine section, New Zealand. In: S.L. Wing, P.D. Gingerich, B. Schmitz and E. Thomas (Editors), *Causes and Consequences of Globally Warm Climates in the Early Paleogene*. Geological Society of America Special Paper 369, Boulder, Colorado, pp. 351-363.
- Dale, B., 1983. Dinoflagellate resting cysts: "Benthic plankton". In: G.A. Fryxell (Editor), *Survival Strategies of the Algae*. Cambridge University Press, Cambridge, pp. 69-136.
- Dale, B., 1996. Dinoflagellate cyst ecology: Modeling and geological applications. In: J. Jansonius and D.C. McGregor (Editors), *Palynology: Principles and Applications*. American Association of Stratigraphic Palynologists Foundation, Dallas, pp. 1249-1276.
- Dale, B., 2001. The sedimentary record of dinoflagellate cysts: looking back into the future of phytoplankton blooms. *Scientia Marina*, 65, 257-272 pp.
- Dale, B. and Dale, A.L., 1992. Dinoflagellate contributions to the deep sea. *Ocean Biocoenosis Series*, 5: 1-77.
- Dale, B. and Fjellså, A., 1994. Dinoflagellate cysts as paleoproductivity indicators: State of the art, potential and limits. In: R. Zahn, T.F. Pedersen, M.A. Kaminski and L. Labeyrie (Editors), *Carbon Cycling in the Glacial Ocean: Constraints on the Ocean's Role in Global Change*. Springer, Berlin, pp. 521-537.
- DeConto, R.M. and Pollard, D., 2003. A coupled climate-ice sheet modeling approach to the Early Cenozoic history of the Antarctic ice sheet. *Palaeogeography, Palaeoclimatology, Palaeoecology*, 198(1-2): 39-52.
- Delille, B., Harlay, J., Zondervan, I., Jacquet, S., Chou, L., Wollast, R., Bellerby, R.G.J., Frankignoulle, M., Vieira-Borges, A., Riebesell, U. and Gattuso, J.-P., 2005. Response of primary production and calcification to changes of pCO₂ during experimental blooms of the coccolithophorid *Emiliana huxleyi*. *Global Biogeochemical Cycles*, 19(GB2023): doi:10.1029/2004GB002318.
- de Vernal, A., Goyette, C. and Rodrigues, C.G., 1989. Contribution palynostratigraphique (dinokystes, pollen et spores) à la connaissance de la mer de Champlain: coupe de Saint Cezaire, Québec. *Canadian Journal of Earth Sciences*, 26: 2450-2464.
- de Vernal, A., Hillaire-Marcel, C., Turon, J.-L. and Matthiessen, J., 2000. Reconstruction of sea-surface temperature, salinity, and sea-ice cover in the northern North Atlantic during the last glacial maximum based on dinocyst assemblages. *Canadian Journal of Earth Sciences*, 37: 725-750.
- de Vernal, A., Matthiessen, J., Mudie, P.J., Rochon, A., Boessenkool, K.P., Eynaud, F., Grøsfjeld, K., Guiot, J., Hamel, D., Harland, R., Head, M.J., Kunz-Pirrung, M., Loucheur, V., Peyron, O., Pospelova, V., Radi, T., Turon, J.-L. and Voronina, E., 2001. Dinoflagellate cyst assemblages as tracers of sea-surface conditions in the northern North Atlantic, Arctic and sub-Arctic seas: the new "n-677" data base and its application for quantitative paleoceanographic reconstruction. *Journal of Quaternary Science*, 16: 681-698.

References

- de Vernal, A. and Mudie, P.J., 1992. Pliocene and Quaternary dinoflagellate cyst stratigraphy in the Labrador Sea: Paleoenvironmental implications. In: M.J. Head and J.H. Wrenn (Editors), Neogene and Quaternary Dinoflagellate Cysts and Acritarchs. American Association of Stratigraphic Palynologists Foundation, Dallas, pp. 329-436.
- de Vernal, A., Rochon, A., Turon, J.-L. and Matthiessen, J., 1998. Organic-walled dinoflagellate cysts: Palynological tracers of sea-surface conditions in middle to high latitude marine environments. *Geobios*, 30: 905-920.
- de Vernal, A., Turon, L. and Guiot, J., 1994. Dinoflagellate distribution in high-latitude marine environments and quantitative reconstruction of sea-surface salinity, temperature and seasonality. *Canadian Journal of Earth Sciences*, 31: 48-62.
- Devillers, R. and DeVernal, A., 2000. Distribution of dinoflagellate cysts in surface sediments of the northern North Atlantic in relation to nutrient content and productivity in surface waters. *Marine Geology*, 166: 103-124.
- Dickens, G.R., 2000. Methane oxidation during the late Paleocene Thermal Maximum. *Bulletin de la Société Géologique de France*, 171: 37-49.
- Dickens, G.R., 2001a. Carbon addition and removal during the Late Palaeocene Thermal Maximum: Basic theory with a preliminary treatment of the isotope record at Ocean Drilling Program Site 1051, Blake Nose. In: D. Kroon, R.D. Norris and A. Klaus (Editors), Western North Atlantic Paleogene and Cretaceous Paleocyanography, *Geol. Soc. London Spec. Publ.* 183.
- Dickens, G.R., 2001b. On the fate of past gas: What happens to methane released from a bacterially mediated gas hydrate capacitor? *Geochemistry Geophysics Geosystems*, 2.
- Dickens, G.R., 2001c. The potential volume of oceanic methane hydrates with variable external conditions. *Organic Geochemistry*, 32: 1179-1193.
- Dickens, G.R., Castillo, M.M. and Walker, J.C.G., 1997. A blast of gas in the latest Paleocene: Simulating first-order effects of massive dissociation of oceanic methane hydrate. *Geology*, 25(3): 259-262.
- Dickens, G.R., Fewless, T., Thomas, E. and Bralower, T.J., 2003. Excess barite accumulation during the Paleocene/Eocene thermal maximum: Massive input of dissolved barium from seafloor gas hydrate reservoirs. In: S.L. Wing, P.D. Gingerich, B. Schmitz and E. Thomas (Editors), Causes and Consequences of Globally Warm Climates in the Early Paleogene, Geological Society of America Special Publication, 369. Geological Society of America, Boulder, Colorado, pp. 11-23.
- Dickens, G.R. and Francis, J.M., 2003. Comment on "A case for a comet impact trigger for the Paleocene/Eocene thermal maximum and carbon isotope excursion" by D.V. Kent et al. [*Earth Planet. Sci. Lett.* 211 (2003) 13-26]. *Earth and Planetary Science Letters*, 217: 197-200.
- Dickens, G.R., O'Neil, J.R., Rea, D.K. and Owen, R.M., 1995. Dissociation of oceanic methane hydrate as a cause of the carbon isotope excursion at the end of the Paleocene. *Paleocyanography*, 10: 965-971.
- Downie, C., Hussain, M.A. and Williams, G.L., 1971. Dinoflagellate cyst and acritarch associations in the Paleogene of Southeast England. *Geoscience and Man*, 3: 29-35.
- Dupuis, C., Aubry, M.-P., Steurbaut, E., Berggren, W.A., Ouda, K., Magioncalda, R., Cramer, B.S., Kent, D.V., Speijer, R.P. and Heilmann-Clausen, C., 2003. The Dababiya Quarry section: Lithostratigraphy, clay mineralogy, geochemistry and paleontology. In: K. Ouda and M.-P. Aubry (Editors), The upper Paleocene-lower Eocene of the Upper Nile Valley: Part 1, Stratigraphy. *Micropaleontology*, v.49 (supplement 1), pp. 41-59.
- Edwards, L.E., 1989. Dinoflagellate Cysts from the Lower Tertiary Formations, Haynesville Cores, Richmond County, Virginia, *Geology and Paleontology of the Haynesville Cores - Northeastern Virginia Coastal Plain*. U.S. Geological Survey professional paper 1489-C. United States Government Printing Office, Washington, pp. 23.
- Edwards, L.E. and Andrieu, V.A., 1992. Distribution of selected dinoflagellate cysts in modern sediments. In: M.J. Head and J.H. Wrenn (Editors), Neogene and Quaternary Dinoflagellate cysts and Acritarchs. American Association of Stratigraphic Palynologists Foundation, Dallas, pp. 259-288.
- Egger, H., Fenner, J., Heilmann-Clausen, C., Rögl, F., Sachsenhofer, R. and Schmitz, B., 2003. Paleoproductivity of the northwestern Tethyan margin (Anthering section, Austria) across the Paleocene-Eocene transition. In: S.L. Wing, P.D. Gingerich, B. Schmitz and E. Thomas (Editors), Causes and Consequences of Globally Warm Climates in the Early Paleogene, Geological Society of America Special Paper 369. Geological Society of America, Boulder, Colorado, pp. 133-146.
- Egger, H., Heilmann-Clausen, C. and Schmitz, B., 2000. The Paleocene-Eocene boundary interval of a Tethyan deep-sea section and its correlation with the North Sea basin. *Bulletin de la Société Géologique de France*, 171: 207-216.
- Ellegaard, M., 2000. Variations in dinoflagellate cyst morphology under conditions of changing salinity during the last 2000 years. *Review of Palaeobotany and Palynology*, 109: 65-81.
- Emanuel, K., DesAutels, C., Holloway, C. and Korty, R., 2004. Environmental Control of

References

- Tropical Cyclone Intensity. *Journal of the Atmospheric Sciences*, 61: 843-858.
- Emerson, S. and Hedges, J.I., 1988. Processes controlling the ocean carbon content of open ocean sediments. *Paleoceanography*, 3: 621-634.
- Ernst, S.R., Guasti, E., Dupuis, C. and Speijer, R.P., 2006. Environmental perturbation in the southern Tethys across the Paleocene/Eocene boundary (Dababiya, Egypt): Foraminiferal and clay mineral records. *Marine Micropaleontology*, 60(1): 89-111.
- Eshet, Y., Almogi, L.A. and Bein, A., 1994. Dinoflagellate cysts, paleoproductivity and upwelling systems; a Late Cretaceous example from Israel. *Marine Micropaleontology*, 23(2): 231-240.
- Fairbanks, R.G., 1982. The origin of continental shelf and slope water in the New York Bight and Gulf of Maine: Evidence from $H_2^{18}O/H_2^{16}O$ ratio measurements. *Journal of Geophysical Research*, 87(5796-5808).
- Farley, K.A., 2001. Extraterrestrial helium in seafloor sediments: identification, characteristics, and accretion rate over geologic time. In: B. Peucker-Ehrinbrink and B. Schmitz (Editors), *Accretion of Extraterrestrial Matter Throughout Earth's History*. Kluwer, New York, pp. 179-204.
- Farley, K.A. and Eltgroth, S.F., 2003. An alternative age model for the Paleocene-Eocene thermal maximum using extraterrestrial 3He . *Earth and Planetary Science Letters*, 208: 135-148.
- Feely, R.A., Sabine, C.L., Lee, K., Berelson, W., Kleypas, J., Fabry, V.J. and Millero, F.J., 2004. Impact of Anthropogenic CO_2 on the $CaCO_3$ System in the Oceans. *Science*, 305(5682): 362-366.
- Fehn, U., Snyder, G. and Egeberg, P.K., 2000. Dating of Pore Waters with ^{129}I : Relevance for the Origin of Marine Gas Hydrates. *Science*, 289(5488): 2332-2335.
- Fensome, R.A., Gocht, H. and Williams, G.L., 1996a. The Eisenack Catalog of Fossil Dinoflagellates. New Series, 4. E. Schweizerbart'sche Verlagsbuchhandlung, Stuttgart, Germany, 2009-2548 pp.
- Fensome, R.A., MacRae, R.A., Moldovan, J.M., Taylor, F.J.R. and Williams, G.L., 1996b. The early Mesozoic radiation of dinoflagellates. *Paleobiology*, 22: 329-338.
- Fensome, R.A., Taylor, F.J.R., Norris, G., Sarjeant, W.A.S., Wharton, D.I. and Williams, G.L., 1993. A Classification of Fossil and Living Dinoflagellates. *Micropaleontology, Special Publication*, 7, 351p. pp.
- Firth, J.V., 1996. Upper Middle Eocene to Oligocene dinoflagellate biostratigraphy and assemblage variations in Hole 913B, Greenland Sea. In: J. Thiede, A.M. Myrhe, J.V. Firth, G.L. Johnson and W.F. Ruddiman (Editors), *Proceedings of the Ocean Drilling Program, Scientific Results*. Texas A & M University, Ocean Drilling Program, College Station, TX, United States, College Station, Texas, pp. 203-242.
- Fogg, G.E., 2002. Harmful algae - a perspective. *Harmful Algae*, 1: 1-4.
- Fricke, H.C., Clyde, W.C., O'Neil, J.R. and Gingerich, P.D., 1998. Evidence for rapid climate change in North America during the latest Paleocene thermal maximum: oxygen isotope compositions of biogenic phosphate from the Bighorn Basin (Wyoming) *Earth and Planetary Science Letters*, 160(1-2): 193-208.
- Fricke, H.C. and Wing, S.L., 2004. Oxygen isotope and paleobotanical estimates of temperature and $\delta^{18}O$ -latitude gradients over North America during the early Eocene. *American Journal of Science*, 304(7): 612-635.
- Galeotti, S., Brinkhuis, H. and Huber, M., 2004. Records of post-Cretaceous-Tertiary boundary millennial-scale cooling from the western Tethys: A smoking gun for the impact-winter hypothesis? *Geology*, 32(6): 529-532.
- Galeotti, S., Kaminski, M.A., Coccioni, R. and Speijer, R.P., 2005. High resolution deep water agglutinated foraminiferal records across the Paleocene/Eocene transition in the Contessa Road Section (Italy). In: M. Bubik and M.A. Kaminski (Editors), *Proceedings of the Sixth International Workshop on Agglutinated Foraminifera (Grzybowski Foundation Special Publication, Volume 8)*.
- Gavrilov, Y., Shcherbinina, E.A. and Oberhänsli, H., 2003. Paleocene-Eocene boundary events in the northeastern Peri-Tethys. In: S.L. Wing, P.D. Gingerich, B. Schmitz and E. Thomas (Editors), *Causes and Consequences of Globally Warm Climates in the Early Paleogene*. Geological Society of America Special Paper 369. Geological Society of America, Boulder, Colorado, pp. 147-168.
- Gervais, P., 1877. Enumeration de quelques ossements d'animaux vertebres recueillis aux environs de Reims par M. Lemoine. *Journal de Zoologie*, 6: 74-79.
- Gibbs, S.J., Bralower, T.J., Bown, P.R., Zachos, J.C. and Bybell, L.M., 2006. Shelf and open-ocean calcareous phytoplankton assemblages across the Paleocene-Eocene Thermal Maximum: Implications for global productivity gradients. *Geology*, 34(4): 233-236.
- Gibson, T.G. and Bybell, L.M., 1994. Sedimentary Patterns across the Paleocene-Eocene boundary in the Atlantic and Gulf coastal plains of the United States. *Bulletin de la Société Belge de Géologie*, 103(3-4): 237-265.
- Gibson, T.G., Bybell, L.M. and Mason, D.B., 2000. Stratigraphic and climatic implications of clay mineral changes around the Paleocene/Eocene boundary of the northeastern US

References

- margin. *Sedimentary Geology*, 134: 65-92.
- Gibson, T.G., Bybell, L.M. and Owens, J.P., 1993. Latest Paleocene lithologic and biotic events in neritic deposits of Southwestern New Jersey. *Paleoceanography*, 8(4): 495-514.
- Gingerich, P.D., 1989. New earliest Wasatchian mammalian fauna from the Eocene of Northwestern Wyoming: Composition and diversity in a rarely sampled high-floodplain assemblage. *University of Michigan Papers on Paleontology*, 28: 1-97.
- Gingerich, P.D., 2000. Paleocene-Eocene boundary and continental vertebrate faunas of Europe and North America. In: B. Schmitz, B. Sundquist and F.P. Andreasson (Editors), *Early Paleogene Warm Climates and Biosphere Dynamics*. GFF (Geologiska Föreningens Förhandlingar), Geological Society of Sweden. , Uppsala, pp. 57-59.
- Gingerich, P.D., 2006. Environment and evolution through the Paleocene-Eocene thermal maximum. *Trends in Ecology & Evolution*, 21(5): 246-253.
- Gingerich, P.D. and Clyde, W.C., 2001. Overview of mammalian biostratigraphy of the Paleocene-Eocene Fort Union and Willwood formations of the Bighorn and Clarks Fork basins. In: P.D. Gingerich (Editor), *Paleocene-Eocene Stratigraphy and Biotic Change in the Bighorn and Clarks Fork Basins, Wyoming*. University of Michigan Papers on Paleontology, 33, pp. 1-14.
- Giusberti, L., Rio, D., Agnini, C., Backman, J., Fornaciari, E., Tateo, F. and Oddone, M., submitted. An expanded marine PETM section in the Venetian Pre-Alps, Italy.
- Goodman, D.K., 1987. Dinoflagellate cysts in ancient and modern sediments. In: F.J.R. Taylor (Editor), *The Biology of Dinoflagellates*, Botanical Monographs, 21, pp. 649-722.
- Goodman, D.K. and Ford, L.N., 1983. Preliminary dinoflagellate biostratigraphy for the middle Eocene to lower Oligocene From the Southwest Atlantic Ocean. In: W.J. Ludwig and V.A. Krasheninnikov, et al. (Editors), *Initial Reports of the Deep Sea Drilling Project*, 71, pp. 859-877.
- Gradstein, F.M., Kaminski, M.A., Berggren, W.A., Kristiansen, I.L. and D'Ioro, M.A., 1994. Cenozoic biostratigraphy of the North Sea and Labrador Shelf. *Micropaleontology*, 40(Supplement): 152.
- Gradstein, F.M., Kristiansen, I.L., Loemo, L. and Kaminski, M.A., 1992. Cenozoic foraminiferal and dinoflagellate cyst biostratigraphy of the central North Sea. *Micropaleontology*, 38: 101-137.
- Gradstein, F.M., Ogg, J.G. and Smith, A.G., 2004. *A Geologic Time Scale 2004*. Cambridge University Press, Cambridge.
- Grøsfjeld, K., Larsen, E., Sejrup, H.-P., DeVernal, A., Flatebf, T., Vestbf, M., Hafliðason, H. and Aarseth, I., 1999. Dinoflagellate cysts reflecting surface-water conditions in Voldafjorden, western Norway during the last 11.300 years. *Boreas*, 28: 403-415.
- Guasti, E., Kouwenhoven, T.J., Brinkhuis, H. and Speijer, R.P., 2005. Paleocene sea-level and productivity changes at the southern Tethyan margin (El Kef, Tunisia). *Marine Micropaleontology*, 55(1-2): 1-17.
- Habib, D. and Miller, J.A., 1989. Dinoflagellate species and organic facies evidence of marine transgression and regression in the Atlantic coastal plain. *Palaeogeography, Palaeoclimatology, Palaeoecology*, 74: 23-47.
- Habib, D., Moshkovitz, S. and Kramer, C., 1992. Dinoflagellate and calcareous nannofossil response to sea-level change in Cretaceous-Tertiary boundary sections. *Geology*, 20: 165-168.
- Haeckel, E., 1894. *Systematische Phylogenie. Entwurf eines natürlichen Systems der Organismen auf Grund ihrer Stammesgeschichte. I. Systematische Phylogenie der Protisten und Pflanzen*. Reimer, Berlin, 400p. pp.
- Hallegraeff, G.M., 1993. A review of harmful algal blooms and their apparent global increase. *Phycologia*, 32: 79-99.
- Haq, B.U., Hardenbol, J. and Vail, P.R., 1987. Chronology of fluctuating sea levels since the Triassic. *Science*, 235: 1156-1167.
- Haq, B.U., Hardenbol, J. and Vail, P.R., 1988. Mesozoic and Cenozoic chronostratigraphy and cycles of sea level change. In: C.K. Wilgus and B.S. Hastings (Editors), *Sea Level Changes: An Integrated Approach*. Society of Economic Paleontologists and Mineralogists Special Publication, 42, pp. 71-108.
- Harland, R., 1973. Dinoflagellate cysts and acritarchs from the Bearpaw Formation (Upper Campanian) of southern Alberta, Canada. *Palaeontology*, 16: 665-706.
- Harland, R., 1983. Distribution maps of recent dinoflagellate in bottom sediments from the North Atlantic Ocean and adjacent seas. *Palaeontology*, 26: 321-387.
- Harland, R. and Long, D., 1996. A Holocene dinoflagellate cyst record from offshore North-east England. *Proceedings of the Yorkshire Geological Society*, 51(1): 65-74.
- Harrington, G.J., 2003. Geographic patterns in the floral response to Paleocene-Eocene warming. In: S.L. Wing, P.D. Gingerich, B. Schmitz and E. Thomas (Editors), *Causes and Consequences of Globally Warm Climates in the Early Paleogene*. Geological Society of America Special Paper 369, Boulder, Colorado, pp. 381-393.
- Harrington, G.J. and Kemp, S.J., 2001. US Gulf Coast vegetation dynamics during the latest

References

- Palaeocene. *Palaeogeography Palaeoclimatology Palaeoecology*, 167(1-2): 1-21.
- Harrington, G.J., Kemp, S.J. and Koch, P.L., 2004. Palaeocene-Eocene paratropical floral change in North America: Responses to climate change and plant immigration. *Journal of the Geological Society*, 161: 173-184.
- Head, M.J., 1996. Modern dinoflagellate cysts and their biological affinities. In: J. Jansonius and D.C. McGregor (Editors), *Palynology: principles and applications*. American Association of Stratigraphic Palynologists Foundations, Dallas, Texas, pp. 1197-1248.
- Head, M.J. and Norris, G., 1989. Palynology and dinocyst stratigraphy of the Eocene and Oligocene in ODP Leg 105, Hole 647A, Labrador Sea. In: S.K. Stewart (Editor), *Proceedings of the Ocean Drilling Program, Scientific Results 105*, pp. 515-550.
- Heil, C.A., Glibert, P.M., Al-Sarawi, M.A., Faraj, M., Behbehani, M. and Husain, M., 2001. First record of a fish-killing *Gymnodinium* sp. bloom in Kuwait Bay, Arabian Sea: chronology and potential causes. *Marine Ecology Progress Series*, 214: 15-23.
- Heilmann-Clausen, C., 1985. Dinoflagellate stratigraphy of the Uppermost Danian to Ypresian in the Viborg 1 borehole, Central Jylland, Denmark. *DGU A7*: 1-69.
- Heilmann-Clausen, C. and Egger, H., 2000. The Anthering outcrop (Austria), a key-section for correlation between Tethys and Northwestern Europe near the Paleocene/Eocene boundary. In: B. Schmitz, B. Sundquist and F.P. Andreasson (Editors), *Early Paleogene Warm Climates and Biosphere Dynamics*. GFF (Geologiska Föreningens Förhandlingar), Geological Society of Sweden. , Uppsala, pp. 69.
- Heiskanen, A.-S., 1993. Mass encystment and sinking of dinoflagellates during a spring bloom. *Marine Biology*, 116: 161-167.
- Heslop, D. and Dekkers, M.J., 2002. Spectral analysis of unevenly spaced climatic time series using CLEAN: signal recovery and derivation of significance levels using a Monte Carlo simulation. *Physics of the Earth and Planetary Interiors*, 130: 103-116.
- Higgins, J.A. and Schrag, D.P., 2006. Beyond methane: Towards a theory for the Paleocene-Eocene Thermal Maximum. *Earth and Planetary Science Letters*, 245(3-4): 523-537.
- Hilgen, F.J., 1991. Astronomical calibration of Gauss to Matuyama sapropels in the Mediterranean and implications for the Geomagnetic Polarity time Scale. *Earth and Planetary Science Letters*, 104(2-4): 226-244.
- Hilgen, F.J., Krijgsman, W., Langereis, C.G., Lourens, L.J., Santarelli, A. and Zacheriasse, W.J., 1995. Extending the astronomical (polarity) time scale into the Miocene. *Earth and Planetary Science Letters*, 136(3-4): 495-510.
- Hochuli, P.A. and Frank, S.M., 2000. Palynology (dinoflagellate cysts, spores, and pollen) and stratigraphy of the Lower Carnian Raibl Group. *Eclogae geologicae Helvetiae*, 93: 429-443.
- Hollis, C.J., Dickens, G.R., Field, B.D., Jones, C.M. and Percy Strong, C., 2005. The Paleocene-Eocene transition at Mead Stream, New Zealand: a southern Pacific record of early Cenozoic global change. *Palaeogeography, Palaeoclimatology, Palaeoecology*, 215(3-4): 313-343.
- Holroyd, P.A., Hutchison, J.H. and Strait, S.G., 2001. Turtle diversity and abundance through the lower Eocene Willwood Formation of the southern Bighorn Basin. *University of Michigan Papers on Paleontology*, 33: 97-107.
- Hooker, J.J., 1998. Mammalian faunal change across the Paleocene-Eocene transition in Europe. In: M.P. Aubry, S. Lucas and W.A. Berggren (Editors), *Late Paleocene-Early Eocene Climatic and Biotic Events in the Marine and Terrestrial Records*. Columbia University Press, New York, pp. 428-450.
- Hopmans, E.C., Weijers, J.W.H., Schefuß, E., Herfort, L., Sinninghe Damsté, J.S. and Schouten, S., 2004. A novel proxy for terrestrial organic matter in sediments based on branched and isoprenoid tetraether lipids. *Earth and Planetary Science Letters*, 224: 107-116.
- Horner, R.A., 1985. *Sea ice biota*. CRC Press, Boca Raton, 215 pp.
- Huber, M. and Sloan, L.C., 2001. Heat transport, deep waters, and thermal gradients: Coupled simulation of an Eocene Greenhouse Climate. *Geophysical Research Letters*, 28(18): 3481-3484.
- Huber, M., Brinkhuis, H., Stickley, C.E., Döös, K., Sluijs, A., Warnaar, J., Schellenberg, S.A. and Williams, G.L., 2004. Eocene circulation of the Southern Ocean: Was Antarctica kept warm by subtropical waters? *Paleoceanography*, 19(PA4026): doi:10.1029/2004PA001014.
- Huber, M., Sloan, L.C. and Shellito, C.J., 2003. Early Paleogene oceans and climate: A fully coupled modeling approach using the NCAR CCSM. In: S.L. Wing, P.D. Gingerich, B. Schmitz and E. Thomas (Editors), *Causes and Consequences of Globally Warm Climates in the Early Palaeogene*, Geological Society of America Special Paper 369. Geological Society of America, Boulder, Colorado, pp. 25-47.
- Hultberg, S.U., 1987. Palynological evidence for a diachronous low-salinity event in the C-T boundary clay at Stevns Klint, Denmark. *Journal of Micropalaeontology*, 6(2): 35-40.
- Iakovleva, A.I., Brinkhuis, H. and Cavagnetto, C., 2001. Late Palaeocene-Early Eocene dinoflagellate cysts from the Turgay Strait, Kazakhstan; correlations across ancient seaways. *Palaeogeography, Palaeoclimatology, Palaeoecology*, 172: 243-268.

References

- IPCC: Intergovernmental Panel on Climate Change, 2001. Climate Change 2001, The Scientific Basis.
- Jackson, S.T. and Overpeck, J.T., 2000. Responses of plant populations and communities to environmental changes of the late Quaternary. *Paleobiology*, 26(4 SUPPS): 194-220.
- JanDuChêne, R.E. and Adediran, S.A., 1984. Late Paleocene to Early Eocene Dinoflagellates from Nigeria. *Cahiers de Micropaléontologie*, 3-1984: 88.
- Jaramillo, C.A. and Dilcher, D.L., 2000. Microfloral diversity patterns of the late Paleocene-Eocene interval in Colombia, northern South America. *Geology*, 28(9): 815-818.
- Jarvis, I., Carson, G.A., Cooper, M.K.E., Hart, M.B., Leary, P.N., Tocher, B.A., Horne, D. and Rosenfeld, A., 1988. Microfossil assemblages and the Cenomanian-Turonian (late Cretaceous) Oceanic Anoxic Event (OAE). *Cretaceous Research*, 9: 3-103.
- Jenkyns, H.C., Forster, A., Schouten, S. and Sinninghe Damsté, J.S., 2004. High temperatures in the Late Cretaceous Arctic Ocean. *Nature*, 432(16 December 2004): 888-892.
- John, C.M., Bohaty, S., Zachos, J.C., Sluijs, A., Gibbs, S.J., Brinkhuis, H. and Bralower, T.J., in prep. Impact of the Paleocene-Eocene thermal maximum on continental margins and implications for the carbon cycle in near-shore environments.
- Kahn, A. and Aubry, M.-P., 2004. Provincialism associated with the Paleocene/Eocene thermal maximum: temporal constraint. *Marine Micropaleontology*, 52(1-4): 117-131.
- Kaiho, K., Arinobu, T., Ishiwatari, R., Morgans, H.E.G., Okada, H., Takeda, N., Tazaki, K., Zhou, G., Kajiwaru, Y., Matsumoto, R., Hirai, A., Niitsuma, N. and Wada, H., 1996. Latest Paleocene benthic foraminiferal extinction and environmental change at Tawanui, New Zealand. *Paleoceanography*, 11: 447-465.
- Kaminski, M.A., Kuhnt, W.A. and Radley, J.D., 1996. Palaeocene-Eocene deep water agglutinated foraminifera from the Namibian Flysch (Rif, Northern Morocco): their significance for the palaeoceanography of the Gibraltar Gateway. *Journal of Micropalaeontology*, 15: 1-19.
- Katz, M.E., Cramer, B.S., Mountain, G.S., Katz, S. and Miller, K.G., 2001. Uncorking the bottle: What triggered the Paleocene/Eocene thermal maximum methane release? *Paleoceanography*, 16(0): 1-14.
- Kelly, D.C., Bralower, T.J. and Zachos, J.C., 1998. Evolutionary consequences of the latest Paleocene thermal maximum for tropical planktonic foraminifera. *Palaeogeography, Palaeoclimatology, Palaeoecology*, 141: 139-161.
- Kelly, D.C., Bralower, T.J., Zachos, J.C., Premoli Silva, I. and Thomas, E., 1996. Rapid diversification of planktonic foraminifera in the tropical Pacific (ODP Site 865) during the late Paleocene thermal maximum. *Geology*, 24(423-426).
- Kelly, D.C., Zachos, J.C., Bralower, T.J. and Schellenberg, S.A., 2005. Enhanced terrestrial weathering/runoff and surface ocean carbonate production during the recovery stages of the Paleocene-Eocene thermal maximum. *Paleoceanography*, 20(PA4023): doi:10.1029/2005PA001163.
- Kenig, F., Hudson, J., D., Sinninghe Damsté, J.S. and Popp, B.N., 2004. Intermittent euxinia: Reconciliation of a Jurassic black shale with its biofacies. *Geology*, 32(5): 421-424.
- Kennett, J.P. and Stott, L.D., 1991. Abrupt deep-sea warming, palaeoceanographic changes and benthic extinctions at the end of the Palaeocene. *Nature*, 353: 225-229.
- Kent, D.V., Cramer, B.S., Lanci, L., Wang, D., Wright, J.D. and van der Voo, R., 2003. A case for a comet impact trigger for the Paleocene/Eocene thermal maximum and carbon isotope excursion. *Earth and Planetary Science Letters*, 211: 13-26.
- Kirschvink, J.L., 1980. the least-square line and plane and the analysis of paleomagnetic data. *Geophysical Journal of the Royal Astronomical Society*, 62: 699-718.
- Knox, R.W.O.B., Corfield, R.M. and Dunay, R.S. (Editors), 1996. Correlation of the Early Paleogene in Northwest Europe. Geological Society of London Special Publication, 101, London, UK.
- Koch, P.L., Clyde, W.C., Hepple, R.P., Fogel, M.L., Wing, S.L. and Zachos, J.C., 2003. Carbon and oxygen isotope records from paleosols spanning the Paleocene-Eocene boundary, Bighorn Basin, Wyoming. In: S.L. Wing, P.D. Gingerich, B. Schmitz and E. Thomas (Editors), Causes and Consequences of Globally Warm Climates in the Early Paleogene. Geological Society of America Special Paper, Boulder, Colorado, pp. 49-64.
- Koch, P.L., Zachos, J.C. and Dettman, D.L., 1995. Stable isotope stratigraphy and paleoclimatology of the Paleogene Bighorn Basin (Wyoming, USA). *Palaeogeography, Palaeoclimatology, Palaeoecology*, 115: 61-89.
- Koch, P.L., Zachos, J.C. and Gingerich, P.D., 1992. Correlation between isotope records in marine and continental carbon reservoirs near the Palaeocene/Eocene boundary. *Nature*, 358: 319-322.
- Kokinos, J.P. and Anderson, D.M., 1995. Morphological development of resting cysts in cultures of the marine dinoflagellate *Lingulodinium polyedrum* (= *L. machaerophorum*). *Palynology*, 19: 143-166.
- Koopmans, M.P., Koster, J., Van Kaam-Peters, H.M.E., Kenig, F., Schouten, S., Hartgers, W.A., de Leeuw, J.W. and Sinninghe Damsté, J.S., 1996. Diagenetic and catagenetic products of isorenieratene: Molecular indicators for photic zone anoxia. *Geochimica et*

References

- Cosmochimica Acta, 60(22): 4467-4496.
- Köthe, A., 1990. Paleogene dinoflagellates from Northwest Germany. Geologisches Jahrbuch, 118: 1-111.
- Köthe, A., Khan, A.M. and Ashraf, M., 1988. Biostratigraphy of the Surghar Range, Salt Range, Sulaiman Range and the Kohat area, Pakistan, according to Jurassic through Paleogene Calcareous Nannofossils and Paleogene Dinoflagellates. Geologisches Jahrbuch, Reihe B, 71. Bundesanstalt für Geowissenschaften und Rohstoffe und den Geologischen Landesämtern in der Bundesrepublik Deutschland, Hannover, 87 pp.
- Kouli, K., Brinkhuis, H. and Dale, B., 2001. *Spiniferites cruciformis*: a fresh water dinoflagellate cyst? Review of Palaeobotany and Palynology, 113: 273-286.
- Krause, D.W. and Maas, M.C., 1990. The biogeographic origins of late Paleocene-early Eocene mammalian immigrants to the western interior of North America. In: T.M. Bown and K.D. Rose (Editors), Dawn of the age of mammals in the northern part of the Rocky Mountain interior, North America. Geological Society of America, Boulder, pp. 71-105.
- Krutzsch, W., 1962. Die Mikroflora der Geiseltalbraunkohle, Teil III. Süßwasserdinoflagellaten aus subaquatisch gebildeten Blätterkohlenlagen des mittleren Geiseltales. Hallesches Jahrbuch für Mitteldeutsche Erdgeschichte, 4: 40-45.
- Kurtz, A., Kump, L.R., Arthur, M.A., Zachos, J.C. and Paytan, A., 2003. Early Cenozoic decoupling of the global carbon and sulfur cycles. Paleocyanography, 18(1090, doi:10.1029/2003PA000908).
- Kvenvolden, K.A., 1988. Methane hydrate - a major reservoir of carbon in the shallow geosphere? Chemical Geology, 71: 41-51.
- Kvenvolden, K.A., 1993. Gas hydrates: Geological perspective and global change. Review of Geophysics, 31: 173-187.
- Laskar, J., Robutel, P., Joutel, F., Gastineau, M., Correia, A.C.M. and Levrard, B., 2004. A long-term numerical solution for the insolation quantities of the Earth. Astronomy & Astrophysics, 428: 261-286.
- Leckie, D.A., Singh, C., Bloch, J., Wilson, M. and Wall, J., 1992. An anoxic event at the Albian-Cenomanian boundary: the Fish Scale Marker Bed, northern Alberta, Canada. Palaeogeography, Palaeoclimatology, Palaeoecology, 92: 139-166.
- LeHérissé, A., Masure, E., Al Ruwaili, M. and Massa, D., 2000. Revision of *Arpylorus antiquus* from the Silurian: The end of a myth. In: W. Wang, S. Ouyang, X. Sun and G. Yu (Editors), Abstracts 10th International Palynological Congress. National Natural Science Foundation of China, Nanjing, pp. 88.
- Lentin, J.K. and Williams, G.L., 1976. A Monograph of Fossil Peridinioid Dinoflagellate Cysts. Bedford Institute of Oceanography Report Series, BI-R-75-16. Bedford Institute of Oceanography, Dartmouth, NS, 237 pp.
- Lentin, J.K. and Williams, G.L., 1980. Dinoflagellate provincialism with emphasis on Campanian peridiniaceans. American Association of Stratigraphic Palynologists, Contribution Series, 7: 1-47.
- Lewis, J., Ellegaard, M., Hallett, R., Harding, I. and Rochon, A., 2003. Environmental control of cyst morphology in Gonyaulacoid dinoflagellates. In: K. Matsuoka, M. Yoshida and M. Iwataki (Editors), Dino7, Seventh International Conference on Modern and fossil Dinoflagellates, Abstract Volume, pp. Additional abstract.
- Lewis, J. and Hallett, R., 1997. *Lingulodinium polyedrum* (*Gonyaulax polyedra*), a blooming dinoflagellate. Oceanography and Marine Biology Annual Review, 35: 97-161.
- Lewis, J., Rochon, A. and Harding, I., 1999. Preliminary observations of cyst-theca relationships in *Spiniferites ramosus* and *Spiniferites membranaceus* (Dinophyceae). Grana, 38: 113-124.
- Lourens, L.J., Antonarakou, A., Hilgen, F.J., Hoof, A.A.M.V., Vergnaud-Grazzini, C. and Zacheriasse, W.J., 1996. Evaluation of the Plio-Pleistocene astronomical timescale. Paleocyanography, 11(4): 391-413.
- Lu, G. and Keller, G., 1993. The Paleocene-Eocene transition in the Antarctic Indian Ocean: Inference from planktic foraminifera. Marine Micropaleontology, 21(1-3): 101-142.
- MacDonald, G.J., 1990. Role of methane clathrates in past and future climates. Climatic Change, 16(3): 247-281.
- Machlus, M., Hemming, S.R., Olsen, P.E. and Christie-Blick, N., 2004. Eocene calibration of geomagnetic polarity time scale reevaluated: Evidence from the Green River Formation in Wyoming. Geology, 32: 137-140.
- MacLennan, J. and Jones, S.M., 2006. Regional uplift, gas hydrate dissociation and the origins of the Paleocene-Eocene Thermal Maximum. Earth and Planetary Science Letters, 245(1-2): 65-80.
- MacRae, R.A., Fensome, R.A. and Williams, G.L., 1996. Fossil dinoflagellate diversity, originations, and extinctions and their significance. Canadian Journal of Botany, 74: 1687-1694.
- Magioncalda, R., Dupuis, C., Smith, T., Steurbaut, E. and Gingerich, P.D., 2004. Paleocene-Eocene carbon isotope excursion in organic carbon and pedogenic carbonate: Direct

References

- comparison in a continental stratigraphic section. *Geology*, 32(7): 553-556.
- Manum, S.B., 1960. Some dinoflagellates and hystrichosphaerids from the Lower Tertiary of Spitsbergen. *Nytt Magasin for Botanikk*, 8: 17-26.
- Manum, S.B., Boulter, M.C., Gunnarsdottir, H., Ragnes, K. and Scholze, A., 1989. Eocene to Miocene palynology of the Norwegian Sea (ODP leg 104). In: O. Eldholm, J. Thiede and E. Taylor (Editors), *Proceedings of the Ocean Drilling Program, Scientific Results*, vol 104, pp. 611-661.
- Mao, S. and Mohr, B.A.R., 1995. Middle Eocene dinoflagellate paleoenvironmental and paleogeographic conclusions. *Review of Palaeobotany and Palynology*, 86: 235-263.
- Markwick, P.J., 1998. Fossil crocodylians as indicators of Late Cretaceous and Cenozoic climates: Implications for using palaeontological data in reconstructing palaeoclimate. *Palaeogeography, Palaeoclimatology, Palaeoecology*, 137: 205-271.
- Marland, G., Boden, T.A. and Andres, R.J., 2005. *Global, Regional, and National CO₂ Emissions, Trends: A Compendium of Data on Global Change*. Carbon Dioxide Information Analysis Center, Oak Ridge National Laboratory, U.S. Department of Energy, Oak Ridge, Tenn., U.S.A.
- Marret, F. and Scourse, J., 2002. Control of modern dinoflagellate cyst distribution in the Irish and Celtic seas by seasonal stratification dynamics. *Marine Micropaleontology*, 47: 101-116.
- Marshall, K.L. and Batten, D.J., 1988. Dinoflagellate cyst associations in Cenomanian-Turonian "black shale" sequences of northern Europe. *Review of Palaeobotany and Palynology*, 54: 85-103.
- Martini, E., 1971. Standard Tertiary and Quaternary calcareous nannoplankton zonation. In: A. Farinacci (Editor), *Proceedings of the II Planktonic Conference, Roma 1970*. Vol. 2. Edizioni Tecnoscienza, Rome, pp. 739-785.
- Matsumoto, R., 1995. Causes of the $\delta^{13}\text{C}$ anomalies of carbonates and a new paradigm "gas-hydrate hypothesis". *Journal of the Geological Society of Japan*, 101: 902-904.
- Matsuoka, K., 1992. Species diversity of modern dinoflagellate cysts in surface sediments around the Japanese Islands. In: M.J. Head and J.H. Wrenn (Editors), *Neogene and Quaternary Dinoflagellate Cysts and Acritarchs*. American Association of Stratigraphic Palynologists Foundation, Dallas, pp. 33-53.
- Matsuoka, K., Yoshida, M. and Iwataki, M., 2003. *Dino7, Seventh International Conference on Modern and Fossil Dinoflagellates*, Nagasaki.
- Matthiessen, J. and Brenner, W., 1996. Chlorococcalalgen und Dinoflagellatenzysten in rezenten Sedimenten des Greifswalder Boddens (südliche Ostsee). *Senckenbergiana Maritima*, 27: 33-48.
- Matthiessen, J., Zonneveld, K.A.F., DeVernal, A., Head, M.J. and Harland, R., 2005. Recent and Quaternary organic-walled dinoflagellate cysts in arctic marine environments and their paleoenvironmental significance. *Paläontologische Zeitschrift*, 79(1): 3-51.
- McKenna, M.C., 1983. Holarctic landmass rearrangement, cosmic events, and Cenozoic terrestrial organisms. *Annals of the Missouri Botanical Garden*, 70: 459-489.
- McMinn, A., 1995. Why are there no post-Paleogene dinoflagellate cysts in the Southern Ocean? *Micropaleontology*, 41(4): 383-386.
- Meng, J., Bowen, G.J., Ye, J., Koch, P.L., Ting, S., Li, Q. and Jin, X., 2004. Gomphos elkema (Gires, Mammalia) from the Erlian Basin: Evidence for the Early Tertiary Bumbanian Land Mammal Age in Nei-Mongol, China. *American Museum Novitates*, 3425: 1-25.
- Milkov, A.V., 2004. Global estimates of hydrate-bound gas in marine sediments: how much is really out there? *Earth-Science Reviews*, 66(3-4): 183-197.
- Miller, K.G., 1997. Coastal Plain Drilling And The New Jersey Sea-Level Transect. In: K.G. Miller and S.W. Snyder (Editors), *Proceedings of the Ocean Drilling Program, Scientific Results, Volume 150X*. Ocean Drilling Program, College Station, TX, , pp. 3-12.
- Miller, K.G., Fairbanks, R.G. and Mountain, G.S., 1987. Tertiary oxygen isotope synthesis, sea-level history, and continental margin erosion. *Paleoceanography*, 2: 1-19.
- Miller, K.G., Kominz, M.A., Browning, J.V., Wright, J.D., Mountain, G.S., Katz, M.E., Sugarman, P.J., Cramer, B.S., Christie-Blick, N. and Pekar, S.F., 2005a. The Phanerozoic Record of Global Sea-Level Change. *Science*, 310(5752): 1293-1298.
- Miller, K.G., Mountain, G.S., Browning, J.V., Kominz, M., Sugarman, P.J., Christie-Blick, N., Katz, M.E. and Wright, J.D., 1998a. Cenozoic global sea level, sequences, and the New Jersey transect: results from coastal plain and continental slope drilling. *Reviews of Geophysics*, 36: 569-601.
- Miller, K.G., Sugarman, P.J., Browning, J.V. et al., 1998b. *Proceedings of the Ocean Drilling Program, Initial Reports 174AX* [Online]. Available from World Wide Web: http://www-odp.tamu.edu/publications/174AX_IR/174AXTOC.HTM.
- Miller, K.G., Wright, J.D. and Browning, J.V., 2005b. Visions of ice sheets in a greenhouse world. *Marine Geology*, 217(3-4): 215-231.
- Miller, K.G., Wright, J.D. and Fairbanks, R.G., 1991. Unlocking the Ice House: Oligocene-Miocene oxygen isotopes, eustasy and margin erosion. *Journal of Geophysical Research*, 96: 6829-6848.

References

- Moldowan, J.M. and Talyzina, N.M., 1998. Biogeochemical evidence for dinoflagellate ancestors in the Early Cambrian. *Science*, 281: 1168-1170.
- Monechi, S., Angori, E. and von-Salis, K., 2000. Calcareous nannofossil turnover around the Paleocene/Eocene transition at Alamedilla (southern Spain). *Bulletin de la Société Géologique de France*, 171: 477-489.
- Moore, T.C., Rabinowitz, P.D. et al. (Editors), 1984. Initial Reports of the Deep Sea Drilling Program Leg 74. U.S. Govt. Printing Office, Washington.
- Moran, K., Backman, J., Brinkhuis, H., Clemens, S.C., Cronin, T., Dickens, G.R., Eynaud, F.d.r., Gattacceca, J.r.m., Jakobsson, M., Jordan, R.W., Kaminski, M., King, J., Koc, N., Krylov, A., Martinez, N., Matthiessen, J., McInroy, D., Moore, T.C., Onodera, J., O'Regan, M., PÅllike, H., Rea, B., Rio, D., Sakamoto, T., Smith, D.C., Stein, R., St John, K., Suto, I., Suzuki, N., Takahashi, K., Watanabe, M., Yamamoto, M., Farrell, J., Frank, M., Kubik, P., Jokat, W. and Kristoffersen, Y., 2006. The Cenozoic palaeoenvironment of the Arctic Ocean. *Nature*, 441(7093): 601-605.
- Moshkovitz, S. and Habib, D., 1993. Calcareous nannofossil and dinoflagellate stratigraphy of the Cretaceous-Tertiary boundary, Alabama, and Georgia. *Micropaleontology*, 39: 167-191.
- Mudie, P.J., Aksu, A.E. and Yasar, D., 2001. Late Quaternary dinoflagellate cysts from the Black, Marmara and Aegean Seas: variations in assemblages, morphology and paleosalinity. *Marine Micropaleontology*, 43: 155-178.
- Mudie, P.J. and Harland, R., 1996. Aquatic Quaternary. In: J. Jansonius and D.C. McGregor (Editors), *Palynology: Principles and Applications*. American Association of Stratigraphic Palynologists Foundation, Dallas, pp. 843-877.
- Mudie, P.J., Rochon, A., Aksu, A.E. and Gillespie, H., 2002. Dinoflagellate cysts, freshwater algae and fungal spores as salinity indicators in late Quaternary cores from Marmara and Black seas. In: A.E. Aksu and C. Yaltirak (Editors), *Quaternary paleoclimatic-paleoceanographic and tectonic evolution of the Marmara Sea and environs; a collection of papers dedicated to the memory of Dr. Ihsan Ketin*. *Marine Geology* 190, 1-2, pp. 203-231.
- Murray, A.E., Blakis, A., Massana, R., Strawzewski, S., Passow, U., Alldredge, A. and DeLong, E.F., 1999. A time series assessment of planktonic archaeal variability in the Santa Barbara Channel. *Aquatic Microbial Ecology*, 20(2): 129-145.
- Nehring, S., 1994a. Dinoflagellaten-Dauercysten in deutschen Küstengewässern: Vorkommen, Verbreitung und Bedeutung als Rekrutierungspotential. *Berichte des Instituts für Meereskunde der Christian-Albrechts-Universität Kiel*, 259: 1-231.
- Nehring, S., 1994b. Spatial distribution of dinoflagellate resting cysts in Recent sediments of Kiel Bight, Germany (Baltic Sea). *Ophelia*, 39: 137-158.
- Nicoll, R.S. and Foster, C.B., 1994. Late Triassic conodont and palynomorph biostratigraphy and conodont thermal maturation, North West Shelf, Australia. *Journal of Australian Geology and Geophysics*, 15: 1-101.
- Nohr-Hansen, H., 2003. Dinoflagellate cyst stratigraphy of the Palaeogene strata from the Hellefisk-1, Ikermiut-1, Kangamiut-1, Nukik-1, Nukik-2 and Qulleq-1 wells, offshore West Greenland. *Marine and Petroleum Geology*, 20(9): 987-1016.
- Norris, G., 1965. Provincialism of Callovian-Neocomian dinoflagellate cysts in the northern and southern hemispheres. *American Association of Stratigraphic Palynologists, Contribution Series*, 4: 29-35.
- Norris, R.D. and Röhl, U., 1999. Carbon cycling and chronology of climate warming during the Palaeocene/Eocene transition. *Nature*, 401: 775-778.
- Nunes, F. and Norris, R.D., 2006. Abrupt reversal in ocean overturning during the Palaeocene/Eocene warm period. *Nature*, 439(7072): 60-63.
- O'Neil, J.R., Clayton, R.N. and Mayeda, T.K., 1969. Oxygen isotope fractionation in divalent metal carbonates. *Journal of Chemical Physics*, 51: 5547-5558.
- Okada, H. and Bukry, D., 1980. Supplementary modification and introduction of code numbers to the low-latitude coccolith biostratigraphic zonation (Bukry, 1973; 1975). *Marine Micropaleontology*, 5: 321-325.
- Orr, J.C., Fabry, V.J., Aumont, O., Bopp, L., Doney, S.C., Feely, R.A., Gnanadesikan, A., Gruber, N., Ishida, A., Joos, F., Key, R.M., Lindsay, K., Maier-Reimer, E., Matear, R., Monfray, P., Mouchet, A., Najjar, R.G., Plattner, G.-K., Rodgers, K.B., Sabine, C.L., Sarmiento, J.L., Schlitzer, R., Slater, R.D., Totterdell, I.J., Weirig, M.-F., Yamanaka, Y. and Yool, A., 2005. Anthropogenic ocean acidification over the twenty-first century and its impact on calcifying organisms. *Nature*, 437(7059): 681-686.
- Ortiz, N., 1995. Differential patterns of benthic foraminiferal extinctions near the Paleocene/Eocene boundary in the North Atlantic and western Tethys. *Marine Micropaleontology*, 26: 341-359.
- Overpeck, J.T., Webb, R.S. and Webb III, T., 1992. Mapping eastern North American vegetation change of the past 18 ka: No-analogs and the future. *Geology*, 20: 1071-1074.
- Pagani, M., Pedentchouk, N., Huber, M., Sluijs, A., Schouten, S., Brinkhuis, H., Sinninghe

References

- Damsté, J.S., Dickens, G.R. and Expedition-Scientists, 2006. Arctic hydrology during global warming at the Palaeocene-Eocene thermal maximum. *Nature*, 442(7103): 671-675.
- Paillard, D., Labeyrie, L. and Yiou, P., 1996. Macintosh Program Performs time-Series Analysis. *Eos Transactions of the American Geophysical Union*, 77: 379.
- Pak, D.K. and Miller, K.G., 1992. Paleocene to Eocene benthic foraminiferal isotopes and assemblages: Implications for deep water circulation. *Paleoceanography*, 7(4): 405-422.
- Passier, H.F., Bosch, H.-J., Nijenhuis, I.A., Lourens, L.J., Bottcher, M.E., Leenders, A., Damsté, J.S.S., de Lange, G.J. and Leeuw, J.W., 1999. Sulphidic Mediterranean surface waters during Pliocene sapropel formation. *Nature*, 397(6715): 146-149.
- Patten, B.C., 1962. Species diversity in net phytoplankton of Raritan Bay. *Journal of Marine Research*, 20: 57-75.
- Peeters, F.J.C., Hoek, R.P., Brinkhuis, H., Wilpshaar, M., de Boer, P.L., Krijgsman, W. and Meulenkamp, J.E., 1998. Differentiating glacio-eustasy and tectonics; a case study involving dinoflagellate cysts from the Eocene-Oligocene of the Pindos Foreland Basin (NW Greece). *Terra Nova*, 10: 245-249.
- Peters, R.B. and Sloan, L.C., 2000. High concentrations of greenhouse gases and polar stratospheric clouds: A possible solution to high-latitude faunal migration at the latest Paleocene thermal maximum. *Geology*, 28(11): 979-982.
- Phipps Morgan, J., Reston, T.J. and Ranero, C.R., 2004. Contemporaneous mass extinctions, continental flood basalts, and 'impact signals': are mantle plume-induced lithospheric gas explosions the causal link? *Earth and Planetary Science Letters*, 217(3-4): 263-284.
- Pierrehumbert, R.T., 2002. The hydrologic cycle in deep-time climate problems. *Nature*, 419(6903): 191-198.
- Pope, K.O., Baines, K.H., Ocampo, A.C. and Ivanov, B.A., 1997. Energy, volatile production, and climatic effects of the Chicxulub Cretaceous/Tertiary impact. *Journal of Geophysical Research*, 102: 21.645-21.664.
- Popp, B.N., Laws, E.A., Bidigare, R.R., Dore, J.E., Hanson, K.L. and Wakeham, S.G., 1998. Effect of Phytoplankton Cell Geometry on Carbon Isotopic Fractionation. *Geochimica et Cosmochimica Acta*, 62(1): 69-77.
- Poulsen, N.E., Manum, S.B., Williams, G.L. and Ellegaard, M., 1996. Tertiary dinoflagellate biostratigraphy of sites 907, 908 and 909 in the Norwegian-Greenland Sea. In: R.N. Riegel (Editor), *Proceedings of the Ocean Drilling Program, Scientific Results*, 151, pp. 255-287.
- Powell, A.J., 1992. Dinoflagellate cysts of the Tertiary System. In: A.J. Powell (Editor), *A stratigraphic Index of Dinoflagellate Cysts*. British Micropalaeontological Society Publication Series, Chapman & Hall, pp. 155-251.
- Powell, A.J., Brinkhuis, H. and Bujak, J.P., 1996. Upper Paleocene - Lower Eocene dinoflagellate cyst sequence biostratigraphy of southeast England. In: R.W.O.B. Knox, R.M. Corfield and R.S. Dunay (Editors), *Correlation of the Early Paleogene in Northwest Europe*, Geological Society Special Publication, 101, pp. 145-183.
- Powell, A.J., Lewis, J. and Dodge, J.D., 1992. The palynological expressions of post-Paleogene upwelling: a review. In: C.P. Summerhayes, W.L. Prell and K.C. Emeis (Editors), *Upwelling Systems: Evolution since the Early Miocene*. The Geological Society, London, pp. 215-226.
- Pross, J., 2001a. Dinoflagellate cyst biogeography and biostratigraphy as a tool for palaeoceanographic reconstructions: An example from the Oligocene of western and northwestern Europe. In: H. Luterbacher, J. Pross and W. Wille (Editors), *Studies in Dinoflagellate Cysts in Honour of Hans Gocht*, *Neues Jahrbuch für Geologie und Paläontologie, Abhandlungen*, pp. 207-219.
- Pross, J., 2001b. Paleo-oxygenation in Tertiary epeiric seas: Evidence from dinoflagellate cysts. *Palaeogeography, Palaeoclimatology, Palaeoecology*, 166: 369-381.
- Pross, J. and Brinkhuis, H., 2005. Organic-walled dinoflagellate cysts as paleoenvironmental indicators in the Paleogene; a synopsis of concepts. *Paläontologische Zeitschrift*, 79(1): 53-59.
- Pross, J., Kotthoff, U. and Zonneveld, K.A.F., 2004. Die Anwendung organischwandiger Dinoflagellatenzysten zur Rekonstruktion von Paläoumwelt, Paläoklima und Paläozeanographie: Möglichkeiten und Grenzen. *Paläontologische Zeitschrift*, 78: 5-39.
- Pross, J. and Schmiedl, G., 2002. Early Oligocene dinoflagellate cysts from the Upper Rhine Graben (SW Germany): Paleoenvironmental and paleoclimatic implications. *Marine Micropaleontology*, 45: 1-24.
- Pujalte, V. and Schmitz, B., 2006. Abrupt climatic and sea level changes across the Paleocene-Eocene boundary, as recorded in an ancient coastal plain setting (Pyrenees, Spain), *Climate and Biota of the Early Paleogene*, Bilbao, Spain.
- Radinova, E.P., Khokhlova, I.E., Baniamovskii, V.N., Shcherbinina, E.A., Iakovleva, A.I. and

References

- Sadchikova, T.A., 2001. The Paleocene/Eocene transition in the northeastern Peri-Tethys area: Sokolovskii key section of the Turgay Passage (Kazakhstan). *Bulletin de la Société Géologique de France*, 172(2): 245-256.
- Raffi, I., Backman, J. and Pälike, H., 2005. Changes in calcareous nannofossil assemblages across the Paleocene/Eocene transition from the paleo-equatorial Pacific Ocean. *Palaeogeography, Palaeoclimatology, Palaeoecology*, 226(1-2): 93-126.
- Ravizza, G., Norris, R.N. and Blusztajn, J., 2001. An Osmium isotope excursion associated with the late Paleocene thermal maximum: Evidence of intensified chemical weathering. *Paleoceanography*, 16: 155-163.
- Reichart, G.-J. and Brinkhuis, H., 2003. Late Quaternary Protoperidinium cysts as indicators of paleoproductivity in the northern Arabian Sea. *Marine Micropaleontology*, 49(4): 303-370.
- Reichart, G.-J., Brinkhuis, H., Huiskamp, F. and Zachariasse, W.J., 2004. Hyper-stratification following glacial overturning events in the northern Arabian Sea. *Paleoceanography*, 19(2): PA2013. 10.1029/2003PA000900.
- Robert, C. and Kennett, J.P., 1994. Antarctic subtropical humid episode at the Paleocene-Eocene boundary: clay mineral evidence. *Geology*, 22(211-214).
- Roberts, D.H., Lehar, J. and Dreher, J.W., 1987. Time series analysis with CLEAN. Part I. Derivation of a spectrum. *Astronomical Journal*, 93(968-989).
- Rochon, A., DeVernal, A., Sejrup, H.-P. and Hafliðason, H., 1998. Palynological evidence of climatic and oceanographic changes in the North Sea during the last deglaciation. *Quaternary Research*, 49: 197-207.
- Rochon, A., Vernal, A., de, Turon, J.L., Mathiessen, J. and Head, M.J., 1999. Distribution of recent dinoflagellate cysts in surface sediments from the North Atlantic Ocean and adjacent seas in relation to sea-surface parameters. *American Association of Stratigraphic Palynologists Foundation Contributions Series*, 35.
- Rognon, P. and Coudé-Gaussen, G., 1996. Paleoclimates off Northwest Africa (28°-35°N) about 18,000 yr B.P. based on continental eolian deposits. *Quaternary Research*, 46: 118-126.
- Röhl, U., Bralower, T.J., Norris, G. and Wefer, G., 2000. A new chronology for the late Paleocene thermal maximum and its environmental implications. *Geology*, 28: 927-930.
- Röhl, U., Brinkhuis, H., Sluijs, A. and Fuller, M., 2004a. On the search for the Paleocene/Eocene Boundary in the Southern Ocean: Exploring ODP Leg 189 Holes 1171D and 1172D, Tasman Sea. In: N.F. Exon, M. Malone and J.P. Kennett (Editors), *The Cenozoic Southern Ocean: Tectonics, Sedimentation, and Climate Change Between Australia and Antarctica*. Geophysical Monograph Series 151, pp. 113-125.
- Röhl, U., Brinkhuis, H., Stickle, C.E., Fuller, M., Schellenberg, S.A., Wefer, G. and Williams, G.L., 2004b. Sea level and astronomically induced environmental changes in Middle and Late Eocene sediments from the East Tasman Plateau. In: N.F. Exon, M. Malone and J.P. Kennett (Editors), *The Cenozoic Southern Ocean: Tectonics, Sedimentation, and Climate Change Between Australia and Antarctica*. American Geophysical Union Geophysical Monograph Series, 151, pp. 127-151.
- Röhl, U., Norris, R.D. and Ogg, J.G., 2003. Cyclostratigraphy of upper Paleocene and lower Eocene sediments at Blake Nose Site 1051 (western North Atlantic). In: S.L. Wing, P.D. Gingerich, B. Schmitz and E. Thomas (Editors), *Causes and Consequences of Globally Warm Climates in the Early Paleogene*. Geological Society of America Special Paper 369. Geological Society of America., Boulder, Colorado, pp. 567-589.
- Röhl, U., Westerhold, T., Bralower, T.J. and Zachos, J.C., in prep. Status of the duration of the Paleocene-Eocene thermal maximum.
- Röhl, U., Westerhold, T., Monechi, S., Thomas, E., Zachos, J.C. and Donner, S., 2006. The third Early Eocene Thermal Maximum: Characteristics, Timing, and Mechanisms of the "X" Event, EGU 2006, Vienna.
- Sangiorgi, F., Capotondi, L. and Brinkhuis, H., 2002. A centennial scale organic-walled dinoflagellate cyst record of the last deglaciation in the South Adriatic Sea (central Mediterranean). *Palaeogeography, Palaeoclimatology, Palaeoecology*, 186(3-4): 199-216.
- Sangiorgi, F., Capotondi, L., Combourieu Nebout, N., Vigliotti, L., Brinkhuis, H., Giunta, S., Lotter, A.F., Morigi, C., Negri, A. and Reichart, G.J., 2003. Holocene seasonal sea surface temperature variations in the South Adriatic Sea inferred from a multi-proxy approach. *Journal of Quaternary Science*, 18(8): 723-732.
- Sarjeant, W.A.S., 1978. *Arpylorus antiquus Calandra* emend., a dinoflagellate cyst from the Upper Silurian. *Palynology*, 2: 167-179.
- Sarjeant, W.A.S., Lacalli, T. and Gaines, G., 1987. The cysts and skeletal elements of dinoflagellates: speculations on the ecologic causes for their morphology and development. *Micropaleontology*, 33: 1-36.
- Sarnthein, M., Winn, K., Duplessy, J.C. and Fontugne, M.R., 1988. Global variations of surface ocean productivity in low and middle latitudes: influence on CO₂ reservoirs of

References

- deep ocean and atmosphere during the last 21,000 years. *Paleoceanography*, 3: 361-399.
- Scheibner, C., Speijer, R.P. and Marzouk, A.M., 2005. Turnover of larger foraminifera during the Paleocene-Eocene Thermal Maximum and paleoclimatic control on the evolution of platform ecosystems. *Geology*, 33(6): 493-496.
- Schellenberg, S.A., Brinkhuis, H., Stickley, C.E., Fuller, M., Kyte, F.T. and Williams, G.L., 2004. The Cretaceous/Paleogene transition on East Tasman Plateau, southwestern Pacific. In: N.F. Exon, M. Malone and J.P. Kennett (Editors), *The Cenozoic Southern Ocean: Tectonics, Sedimentation, and Climate Change Between Australia and Antarctica*. American Geophysical Union Geophysical Monograph Series, 151.
- Schenau, S.J., Prins, M., de Lange, G.J. and Monnin, C., 2001. Barium accumulation in the Arabian Sea: Controls on barite preservation in marine sediments. *Geochimica et Cosmochimica Acta*, 65(10): 1545-1556.
- Schmidt, G.A. and Schindell, D.T., 2003. Atmospheric composition, radiative forcing, and climate change as a consequence of a massive methane release from gas hydrates. *Paleoceanography*, 18(1004): doi:10.1029/2002PA000757.
- Schmitz, B., Peucker-Ehrenbrink, B., Heilmann-Clausen, C., Åberg, G., Asaro, F. and Lee, C.-T.A., 2004. Basaltic explosive volcanism, but no comet impact, at the Paleocene-Eocene boundary: high-resolution chemical and isotopic records from Egypt, Spain and Denmark. *Earth and Planetary Science Letters*, 225: 1-17.
- Schmitz, B., Pujalte, V. and Nunez-Betelu, K., 2001. Climate and sea-level perturbations during the Incipient Eocene Thermal Maximum: evidence from siliciclastic units in the Basque Basin (Ermua, Zumaia and Trabakua Pass), northern Spain. *Palaeogeography, Palaeoclimatology, Palaeoecology*, 165(3-4): 299-320.
- Schmitz, B. and Pujalte, V., 2003. Sea-level, humidity, and land-erosion records across the initial Eocene thermal maximum from a continental-marine transect in northern Spain. *Geology*, 31(8): 689-692.
- Schouten, S., Hopmans, E.C., Forster, A., Breugel, Y.V., Kuypers, M.M.M. and Sinninghe Damsté, J.S., 2003. Extremely high sea-surface temperatures at low latitudes during the middle Cretaceous as revealed by archaeal membrane lipids. *Geology*, 31(12): 1069-1072.
- Schouten, S., Hopmans, E.C., Schefuß, E. and Sinninghe Damsté, J.S., 2002. Distributional variations in marine crenarchaeotal membrane lipids: a new tool for reconstructing ancient sea water temperatures? *Earth and Planetary Science Letters*, 204: 265-274.
- Schouten, S., Hopmans, E.C. and Sinninghe Damsté, J.S., 2004. The effect of maturity and depositional redox conditions on archaeal tetraether lipid palaeothermometry. *Organic Geochemistry*, 35: 567-571.
- Schrag, D.P., dePaolo, D.J. and Richter, F.M., 1995. Reconstructing past sea surface temperatures: Correcting for diagenesis of bulk marine carbonate. *Geochimica et Cosmochimica Acta*, 59: 2265-2278.
- Scotese, C.R., 2002. <http://www.scotese.com>, (PALEOMAP website).
- Shackleton, N.J., 1967. Oxygen isotope analyses and Pleistocene temperatures reassessed. *Nature*, 215: 15-17.
- Shellito, C.J., Sloan, L.C. and Huber, M., 2003. Climate model sensitivity to atmospheric CO₂ levels in the Early-Middle Paleogene. *Palaeogeography, Palaeoclimatology, Palaeoecology*, 193(1): 113-123.
- Shimmield, G.B., 1992. Can sediment geochemistry record changes in coastal upwelling palaeoproductivity? Evidence from northwest Africa and the Arabian Sea. In: C.P. Summerhayes, W.L. Prell and K.C. Emeis (Editors), *Upwelling Systems: Evolution since the Early Miocene*, Geological Society Special Publication, 64, pp. 29-46.
- Sinninghe Damsté, J.S. and Koster, J., 1998. A euxinic southern North Atlantic Ocean during the Cenomanian/Turonian oceanic anoxic event. *Earth and Planetary Science Letters*, 158(3-4): 165-173.
- Sinninghe Damsté, J.S., Rijpstra, W.I.C. and Reichert, G.-J., 2002. The influence of oxic degradation on the sedimentary biomarker record II. Evidence from Arabian Sea sediments. *Geochimica et Cosmochimica Acta*, 66(15): 2737-2754.
- Sinninghe Damsté, J.S., Wakeham, S.G., Kohnen, M.E.L., Hayes, J.M. and de Leeuw, J.W., 1993. A 6,000-year sedimentary molecular record of chemocline excursions in the Black Sea. *Nature*, 362: 827 - 829.
- Sloan, L.C. and Pollard, D., 1998. Polar stratospheric clouds: A high latitude warming mechanism in an ancient greenhouse world. *Geophysical Research Letters*, 25(18): 3517-3520.
- Sluijs, A., Brinkhuis, H., Stickley, C.E., Warnaar, J., Williams, G.L. and Fuller, M., 2003. Dinoflagellate cysts from the Eocene/Oligocene transition in the Southern Ocean; results from ODP Leg 189. In: N.F. Exon, J.P. Kennett and M.J. Malone (Editors), *Proceedings Ocean Drilling Program, Scientific Results*. Available from World Wide Web: http://www-odp.tamu.edu/publications/189_SR/104/104.htm. College Station, Texas, pp. 1-42.
- Smith, T. and Smith, R., 1995. Synthèse des données actuelles sur les vertébrés de la

References

- transition Paleocene-Eocene de Dormaal (Belgique). *Bulletin de la Société belge de Géologie*, 104(1-2): 119-131.
- Smith, T., Rose, K., D. and Gingerich, P., D., 2006. Rapid Asia-Europe-North America geographic dispersal of earliest Eocene primate *Teilhardina* during the Paleocene-Eocene Thermal Maximum. *Proceedings of the National Academy of Sciences of the United States of America*, 103(30): 11223-11227.
- Speijer, R.P. and Morsi, A.-M.M., 2002. Ostracode turnover and sea-level changes associated with the Paleocene-Eocene thermal maximum. *Geology*, 30(1): 23-26.
- Speijer, R.P. and Schmitz, B., 1998. A benthic foraminiferal record of Paleocene sea level and trophic/redox conditions at Gebel Aweina, Egypt. *Palaeogeography, Palaeoclimatology, Palaeoecology*, 137(1-2): 79-101.
- Speijer, R.P., Van der Zwaan, G.J. and Schmitz, B., 1996. The impact of Paleocene/ Eocene boundary events on middle neritic benthic foraminiferal assemblages from Egypt. *Marine Micropaleontology*, 28(2): 99-132.
- Speijer, R.P. and Wagner, T., 2002. Sea-level changes and black shales associated with the late Paleocene thermal maximum: Organic-geochemical and micropaleontologic evidence from the southern Tethyan margin (Egypt-Israel). *Geological Society of America Special Paper*, 356: 533-549.
- Spero, H.J., 1987. Symbiosis in the planktonic foraminifer, *Orbulina universa*, and the isolation of its symbiotic dinoflagellate, *Gymnodinium beii* sp. nov. *Journal of Phycology*, 23: 307-317.
- Spero, H.J., Bijma, J., Lea, D.W. and Bemis, B.E., 1997. Effect of seawater carbonate concentration on foraminiferal carbon and oxygen isotopes. *Nature*, 390(6659): 497-500.
- Spieß, V., 1990. Cenozoic magnetostratigraphy of Leg 113 drill sites, Maud Rise, Weddell Sea, Antarctica. In: P.F. Barker, J.P. Kennett and e. al (Editors), *Proceedings of the Ocean Drilling Project, Initial Reports 113*, pp. 261-315.
- Sprangers, M., Dammers, N., Brinkhuis, H., Weering, T.C.E., van and Lotter, A.F., 2004. Modern organic-walled dinoflagellate cyst distribution offshore NW Iberia; tracing the upwelling system. *Review of Palaeobotany and Palynology*, 128(1-2): 97-106.
- Steineck, P.L. and Thomas, E., 1996. The latest Paleocene crisis in the deep sea: Ostracode succession at Maud Rise, Southern Ocean. *Geology*, 24(7): 583-586.
- Steurbaut, E., Magioncalda, R., Dupuis, C., Van Simaey, S., Roche, E. and Roche, M., 2003. Palynology, paleoenvironments, and organic carbon isotope evolution in lagoonal Paleocene-Eocene boundary settings in North Belgium. In: S.L. Wing, P. Gingerich, B. Schmitz and E. Thomas (Editors), *Causes and consequences of Globally Warm Climates in the Early Paleogene*, Geological Society of America Special Paper 369. Geological Society of America, Boulder, Colorado, pp. 291-317.
- Stickley, C.E., Brinkhuis, H., Schellenberg, S.A., Sluijs, A., Röhl, U., Fuller, M., Grauert, M., Huber, M., Warnaar, J. and Williams, G.L., 2004. Timing and nature of the deepening of the Tasmanian Gateway. *Paleoceanography*, 19(PA4027): doi:10.1029/2004PA001022.
- Stoll, H.M., 2005. Limited range of interspecific vital effects in coccolith stable isotopic records during the Paleocene-Eocene thermal maximum. *Paleoceanography*, 20(PA1007): doi:10.1029/2004PA001046.
- Stover, L.E., 1977. Oligocene and early Miocene dinoflagellates from Atlantic corehole 5/5b, Blake Plateau. In: W.C. Elsik (Editor), *Contributions of Stratigraphic Palynology 1; Cenozoic palynology*. American Association of Stratigraphic Palynologists Contribution Series 5A, pp. 66-89.
- Stover, L.E., Brinkhuis, H., Damassa, S.P., de Verteuil, L., Helby, R.J., Monteil, E., Partridge, A., Powell, A.J., Riding, J.B., Smelror, M. and Williams, G.L., 1996. Mesozoic-Tertiary dinoflagellates, acritarchs and prasinophytes. In: J. Jansonius and D.C. McGregor (Editors), *Palynology: Principles and Applications*. American Association of Stratigraphic Palynologists Foundation, Dallas, pp. 641-750.
- Stover, L.E. and Hardenbol, J., 1994. Dinoflagellates and depositional sequences in the Lower Oligocene (Rupelian) Boom clay formation, Belgium. *Bulletin van de Belgische Vereniging voor Geologie / Bulletin de la Société belge de Géologie*, T. 102(1-2): 5-77.
- Suess, E., 1980. Particulate organic carbon flux in the oceans - surface productivity and oxygen utilization. *Nature*, 288: 260-263.
- Svensen, H., Planke, S., Mørch-Sørensen, A., Jamveit, B., Myklebust, R., Eidem, T.R. and Rey, S.S., 2004. Release of methane from a volcanic basin as a mechanism for initial Eocene global warming. *Nature*, 429: 542-545.
- Tappan, H. and Loeblich, A.R.J., 1971. Geobiologic implications of fossil phytoplankton evolution and time-space distribution. In: R.M. Kosanke and A.T. Cross (Editors), *Symposium on Palynology of the Late Cretaceous and Early Tertiary*. Geological Society of America Special Paper, 127, pp. 247-340.
- Targarona, J., Warnaar, J., Boessenkool, K.P., Brinkhuis, H. and Canals, M., 2000. Recent dinoflagellate cyst distribution in the north Canary Basin, NW Africa. *Grana*, 38(2/3): 170-178.

References

- Tauxe, L., Pick, T. and Kok, Y.S., 1995. Relative paleointensity in sediments: a pseudo-Thellier approach. *Geophysical Research Letters*, 22(2885-2888).
- Taylor, D.L. and Seliger, H.H., 1979. *Toxic Dinoflagellate Blooms*. Elsevier, North Holland, 505 pp.
- Taylor, F.J.R. (Editor), 1987. *The biology of dinoflagellates*. Botanical Monographs, 21. Blackwell Scientific Publications, London, 785 pp.
- Thomas, D.J., 2004. Evidence for deep-water production in the North Pacific Ocean during the early Cenozoic warm interval. *Nature*, 430(6995): 65-68.
- Thomas, D.J. and Bralower, T.J., 2005. Sedimentary trace element constraints on the role of North Atlantic Igneous Province volcanism in late Paleocene-early Eocene environmental change. *Marine Geology*, 217(3-4): 233-254.
- Thomas, D.J., Bralower, T.J. and Zachos, J.C., 1999. New evidence for subtropical warming during the late Paleocene thermal maximum: Stable isotopes from Deep Sea Drilling Project Site 527, Walvis Ridge. *Paleoceanography*, 14: 561-570.
- Thomas, D.J., Zachos, J.C., Bralower, T.J., Thomas, E. and Bohaty, S., 2002. Warming the fuel for the fire: Evidence for the thermal dissociation of methane hydrate during the Paleocene-Eocene thermal maximum. *Geology*, 30(12): 1067-1070.
- Thomas, E., 1989. Development of Cenozoic deep-sea benthic foraminiferal faunas in Antarctic waters. *Geological Society London Special Publication*, 47: 283-296.
- Thomas, E., 1998. Biogeography of the late Paleocene benthic foraminiferal extinction. In: M.-P. Aubry, S.G. Lucas and W.A. Berggren (Editors), *Late Paleocene-early Eocene climatic and biotic events in the marine and terrestrial records*. Columbia University Press, New York, pp. 214-243.
- Thomas, E., 2003. Extinction and food at the seafloor: A high-resolution benthic foraminiferal record across the Initial Eocene Thermal Maximum, Southern Ocean Site 690. In: S.L. Wing, P.D. Gingerich, B. Schmitz and E. Thomas (Editors), *Causes and Consequences of Globally Warm Climates in the Early Paleogene*. Geological Society of America Special Paper 369. Geological Society of America, Boulder, Colorado, pp. 319-332.
- Thomas, E., in press. Cenozoic mass extinctions in the deep sea: what perturbs the largest habitat on earth? *Geological Society of America Special Paper*.
- Thomas, E. and Shackleton, N.J., 1996. The Palaeocene-Eocene benthic foraminiferal extinction and stable isotope anomalies. In: R.W.O.B. Knox, R.M. Corfield and R.E. Dunay (Editors), *Correlation of the Early Paleogene in Northwestern Europe*, Geological Society London Special Publication, 101, pp. 401-441.
- Thomas, E. and Zachos, J.C., 2000. Was the late Paleocene thermal maximum a unique event? *Geologiska Föreningens i Stockholm Förhandlingar (GFF; Transactions of the Geological Society in Stockholm)*, 122: 169-170.
- Thomas, E., Zachos, J.C. and Bralower, T.J., 2000. Deep-sea environments on a warm Earth: Latest Paleocene-early Eocene. In: B.T. Huber, K. MacLeod and S.L. Wing (Editors), *Warm Climates in Earth History*. Cambridge University Press, pp. 132-160.
- Ting, S., Bowen, G.J., Koch, P.L., Clyde, W.C., Wang, Y., Wang, Y. and McKenna, M.C., 2003. Biostratigraphic, chemostratigraphic, and magnetostratigraphic study across the Paleocene/Eocene boundary in the Hengyang Basin, Hunan, China. In: S.L. Wing, P.D. Gingerich, B. Schmitz and E. Thomas (Editors), *Causes and Consequences of Globally Warm Climates in the Early Paleogene*. Geological Society of America Special Paper 369, Boulder, Colorado, pp. 521-535.
- Torricelli, S., Knezaurek, G. and Biffi, U., 2006. Sequence biostratigraphy and paleoenvironmental reconstruction in the Early Eocene Figols Group of the Tremp-Graus Basin (south-central Pyrenees, Spain). *Palaeogeography, Palaeoclimatology, Palaeoecology*, 232(1): 1-35.
- Tremolada, F. and Bralower, T.J., 2004. Nannofossil assemblage fluctuations during the Paleocene-Eocene Thermal Maximum at Sites 213 (Indian Ocean) and 401 (North Atlantic Ocean): palaeoceanographic implications. *Marine Micropaleontology*, 52(1-4): 107-116.
- Trench, R., 1987. Non-parasitic symbioses. In: F.J.R. Taylor (Editor), *The Biology of Dinoflagellates*, Botanical Monographs, pp. 530-570.
- Tripati, A. and Elderfield, H., 2005. Deep-Sea Temperature and Circulation Changes at the Paleocene-Eocene Thermal Maximum. *Science*, 308(5730): 1894-1898.
- Tripati, A., Zachos, J., Marincovich, J., Louie and Bice, K., 2001. Late Paleocene Arctic coastal climate inferred from molluscan stable and radiogenic isotope ratios. *Palaeogeography, Palaeoclimatology, Palaeoecology*, 170(1-2): 101-113.
- Tripati, A.K. and Elderfield, H., 2004. Abrupt hydrographic changes in the equatorial Pacific and subtropical Atlantic from foraminiferal Mg/Ca indicate greenhouse origin for the thermal maximum at the Paleocene-Eocene Boundary. *Geochemistry, Geophysics, Geosystems*, 5(2): doi:10.1029/2003GC000631.
- Turon, J.-L., 1981. *Le palyonplancton dans l'environnement actuel de l'Atlantique Nord oriental. Evolution climatique et hydrologique depuis le Dernier Maximum Glaciaire*. Thèse d'Etat Thesis, Université de Bordeaux, 313 pp.

References

- van Mourik, C.A. and Brinkhuis, H., 2000. Data report: Organic Walled Dinoflagellate Cyst Biostratigraphy of the Latest Middle to Late Eocene at Hole 1053A (Subtropical Atlantic Ocean). In: D. Kroon, R.D. Norris and A. Klaus (Editors), Proceedings of the Ocean Drilling Program, Scientific Results, 171B. Available from World Wide Web: http://www-odp.tamu.edu/publications/171B_SR/chap_06/chap_06.htm. College Station, Texas.
- van Mourik, C.A., Brinkhuis, H. and Williams, G.L., 2001. Middle to Late Eocene organic walled dinoflagellate cysts from ODP Leg 171B, offshore Florida. In: D. Kroon, Norris, R.D. and A. Klaus (Editors), Western North Atlantic Palaeogene and Cretaceous Palaeoceanography. Geological Society London Special Publications, 183, pp. 225-251.
- van Simaey, S., Brinkhuis, H., Pross, J., Williams, G.L. and Zachos, J.C., 2005. Arctic dinoflagellate migrations mark the stongest Oligocene glaciations. *Geology*, 33(9): 709-712.
- Vandenbergh, N., Brinkhuis, H. and Steurbaut, E., 2003. The Eocene/Oligocene boundary in the North Sea area: a sequence stratigraphic approach. In: D.R. Prothero, L.C. Ivany and E.A. Nesbitt (Editors), From Greenhouse to Icehouse: the Marine Eocene Oligocene Transition. Columbia University Press, New York, pp. 419-437.
- Varadi, F., Bunnegar, B. and Ghil, M., 2003. Successive refinements in long-term integrations of planetary orbits. *Astrophysical Journal*, 592(1.1): 620-630.
- Versteegh, G.J.M., 1994. Recognition of cyclic and non-cyclic environmental changes in the Mediterranean Pliocene: A palynological approach. *Marine Micropaleontology*, 23: 147-183.
- Versteegh, G.J.M. and Zonneveld, K.A.F., 1994. Determination of (palaeo-)ecological preferences of dinoflagellates by applying Detrended and Canonical Correspondence analysis to Late Pliocene dinoflagellate cyst assemblages of the south Italian Singha section. *Review of Palaeobotany and Palynology*, 84: 181-199.
- Versteegh, G.J.M. and Zonneveld, K.A.F., 2002. Use of selective degradation to separate preservation from productivity. *Geology*, 30(7): 615-618.
- Vonhof, H.B., Brinkhuis, H., Van der Hoeven, M., Smit, J., Montanari, A. and Nederbragt, A.J., 2000. Global cooling accelerated by early late Eocene impacts. *Geology*, 28: 687-690.
- Wakeham, S.G. and Canuel, E.A., 1988. Organic geochemistry of particulate matter in the eastern tropical North Pacific Ocean: Implications for particle dynamics. *Journal of Marine Research*, 46: 183-213.
- Wakeham, S.G. and Lee, C., 1993. Production, transport, and alteration of particulate organic matter in the marine water column. In: M. Engel and S. Macko (Editors), *Organic Geochemistry*. Plenum Press, New York, pp. 145-169.
- Wall, D. and Dale, B., 1974. Dinoflagellates in late Quaternary deep-water sediments of Black Sea. In: E.T. Degens and D.A. Ross (Editors), *The Black Sea - Geology, Chemistry and Biology*, American Association of Petroleum Geologists Memoir, pp. 364-380.
- Wall, D., Dale, B. and Harada, K., 1973. Descriptions of new fossil dinoflagellates from the late Quaternary of the Black Sea. *Micropaleontology*, 19: 18-31.
- Wall, D., Dale, B., Lohmann, G.P. and Smith, W.K., 1977. The environmental and climatic distribution of dinoflagellate cysts in the North and South Atlantic and adjacent seas. *Marine Micropaleontology*, 30: 319-343.
- Wefer, G., Berger, W.H., Bijma, J. and Fischer, G., 1999. Clues to ocean history: a brief overview of proxies. In: G. Fischer and G. Wefer (Editors), *Use of Proxies in Paleoceanography*, Examples from the South Atlantic. Springer, Berlin, pp. 1-68.
- Weijers, J.W.H., Schouten, S., van der Linden, M., van Geel, B. and Sinninghe Damsté, J.S., 2004. Water table related variations in the abundance of intact archaeal membrane lipids in a Swedish peat bog. *FEMS Microbiol. Lett.*, 239: 51-56.
- Westerhold, T., Röhl, U., Laskar, J., Raffi, I., Bowles, J., Lourens, L.J. and Zachos, J.C., submitted. On the duration of Magnetochrons C24r and C25n, and the timing of early Eocene global warming events: Implications from the OCP Leg 208 Walvis Ridge depth transect. *Paleoceanography*.
- Williams, G.L., 1977. Dinoflagellate cysts, their classification, biostratigraphy and palaeoecology. In: A.T.S. Ramsay (Editor), *Oceanic Micropaleontology*. Academic Press, London, pp. 1231-1325.
- Williams, G.L., Brinkhuis, H., Pearce, M.A., Fensome, R.A. and Weegink, J.W., 2004. Southern Ocean and global dinoflagellate cyst events compared; index events for the Late Cretaceous-Neogene. In: N.F. Exon, J.P. Kennett and M.J. Malone (Editors), Proceedings Ocean Drilling Program, Scientific Results. Available from World Wide Web: http://www-odp.tamu.edu/publications/189_SR/107/107.htm. College Station, Texas, pp. 1-98.
- Wilpsaar, M., Santarelli, A., Brinkhuis, H. and Visscher, H., 1996. Dinoflagellate cysts and mid-Oligocene chronostratigraphy in the central Mediterranean region. *Journal of the Geological Society, London*, 153: 553-561.
- Wing, S.L., 1998. Late Paleocene-early Eocene floral and climate change in the Bighorn Basin, Wyoming. In: M.-P. Aubry, S.G. Lucas and W.A. Berggren (Editors), *Late Paleocene-early Eocene climatic and biotic events in the marine and terrestrial records*. Columbia

References

- University Press, New York, pp. 380-400.
- Wing, S.L. and Harrington, G.J., 2001. Floral response to rapid warming in the earliest Eocene and implications for concurrent faunal change. *Paleobiology*, 27(3): 539-563.
- Wing, S.L., Harrington, G.J., Bowen, G.J. and Koch, P.L., 2003. Floral change during the Initial Eocene Thermal Maximum in the Powder River Basin, Wyoming. In: S.L. Wing, P.D. Gingerich, B. Schmitz and E. Thomas (Editors), *Causes and Consequences of Globally Warm Climates in the Early Paleogene*. Geological Society of America Special Paper 369, Boulder, Colorado, pp. 425-440.
- Wing, S.L., Harrington, G.J., Smith, F.A., Bloch, J.I., Boyer, D.M. and Freeman, K.H., 2005. Transient Floral Change and Rapid Global Warming at the Paleocene-Eocene Boundary. *Science*, 310(5750): 993-996.
- Wolff, T., Grieger, B., Hale, W., Dürkoop, A., Mulitza, S., Pätzold, J. and Wefer, G., 1999. On the reconstruction of paleosalinities. In: G. Fischer and G. Wefer (Editors), *Use of Proxies in Paleoceanography, Examples from the South Atlantic*. Springer, Berlin, pp. 207-228.
- Wrenn, J.H. and Beckmann, S.W., 1982. Maceral, total organic carbon, and palynological analyses of Ross Ice Shelf Project site J9 cores. *Science*, 216: 187-189.
- Wrenn, J.H. and Hart, G.F., 1988. Paleogene Dinoflagellate Cyst Biostratigraphy of Seymour Island, Antarctica. In: R.M. Feldmann and M.O. Woodburne (Editors), *Geology and Paleontology of Seymour Island, Antarctic Peninsula*, *Memoirs of the Geological Society of America*, Boulder, Colorado, pp. 321-447.
- Wuchter, C., Schouten, S., Boschker, H.T.S. and Sinninghe Damsté, J.S., 2003. Bicarbonate uptake by marine Crenarchaeota. *FEMS Microbiology Letters*, 219(2): 203-207.
- Wuchter, C., Schouten, S., Coolen, M.J.L. and Sinninghe Damsté, J.S., 2004. Temperature-dependent variation in the distribution of tetraether membrane lipids of marine Crenarchaeota: Implications for TEX₈₆ paleothermometry. *Paleoceanography*, 19(PA402): doi:10.1029/2004PA001041.
- Xu, W., Lowell, R.P. and Peltzer, E.T., 2001. Effect of seafloor temperature and pressure variations on methane flux from a gas hydrate layer: Comparison between current and late Paleocene climate conditions. *Journal of Geophysical Research*, 106(26,413-26,423).
- Zachos, J., Pagani, M., Sloan, L., Thomas, E. and Billups, K., 2001. Trends, rhythms, and aberrations in global climate 65 Ma to present. *Science*, 292: 686-693.
- Zachos, J.C., Kroon, D., Blum, P. et al. 2004. Proceedings of the Ocean Drilling Program, Initial Reports, 208. Available from World Wide Web: http://www-odp.tamu.edu/publications/208_IR/208ir.htm.
- Zachos, J.C., Lohmann, K.C., Walker, J.C.G. and Wise, S.W., 1993. Abrupt climate change and transient climates during the Palaeogene: A marine perspective. *Journal of Geology*, 101: 191-213.
- Zachos, J.C., Stott, L.D. and Lohmann, K.C., 1994. Evolution of early Cenozoic marine temperatures. *Paleoceanography*, 9(353-387).
- Zachos, J.C., Wara, M.W., Bohaty, S., Delaney, M.L., Petrizzo, M.R., Brill, A., Bralower, T.J. and Premoli Silva, I., 2003. A transient rise in tropical sea surface temperature during the Paleocene-Eocene thermal maximum. *Science*, 302: 1151-1154.
- Zevenboom, D., 1996. Late Oligocene-early Miocene dinoflagellate cysts from the Lemme-Carrosio section (NW Italy); biostratigraphy and palaeoenvironmental interpretation. In: F.F. Steininger, S. Iaccarino and F. Cati (Editors), *In search of the Paleogene-Neogene boundary; Part 3, The global stratotype section and point, the GSSP for the base of the Neogene (the Paleogene-Neogene boundary)*. *Giornale di Geologia* 58, 1-2, pp. 81-93.
- Zevenboom, D., Brinkhuis, H. and Visscher, H., 1994. Dinoflagellate cysts palaeoenvironmental analysis of the Oligocene/ Miocene transition in northwest and central Italy, Miocene stratigraphy of Italy and adjacent regions. *Giornale di Geologia*, 56, 1, Bologna, Italy, pp. 155-169.