

A VARIATION OF MIXED HODGE STRUCTURE FOR A SPECIAL CASE OF APPELL'S F_4

JAN STIENSTRA

dedicated to Mr Taniguchi on the occasion of his ninetieth birthday

Abstract

We present a variation of mixed Hodge structure of geometric origin for Appell's system of differential equations of type F_4 with parameters $0, \frac{1}{2}, 1, 1$ and variables τ_1^2, τ_2^2 .

1 Appell's F_4

In [1] Appell introduced the hypergeometric series of type F_4 with parameters $\alpha, \beta, \gamma, \gamma'$ and variables u_1, u_2 :

$$F_4(\alpha, \beta, \gamma_1, \gamma_2, u_1, u_2) = \sum_{m, n \geq 0} \frac{(\alpha, m+n)(\beta, m+n)}{(\gamma_1, m)(\gamma_2, n)(1, m)(1, n)} u_1^m u_2^n.$$

Here $(\lambda, k) = \lambda \cdot (\lambda + 1) \cdots (\lambda + k - 1)$. It is one solution for the system of hypergeometric differential equations, with $D_i = u_i \frac{\partial}{\partial u_i}$, $i = 1, 2$,

$$D_i(D_i + \gamma_i - 1)Z = u_i(D_1 + D_2 + \alpha)(D_1 + D_2 + \beta)Z.$$

In this paper we want to study Appell's system of differential equations for $(\alpha, \beta, \gamma, \gamma') = (0, \frac{1}{2}, 1, 1)$ and $u_1 = \tau_1^2, u_2 = \tau_2^2$: with $\mathcal{D}_i = \tau_i \frac{\partial}{\partial \tau_i}$, $i = 1, 2$,

$$\mathcal{D}_i^2 Z = \tau_i^2 (\mathcal{D}_1 + \mathcal{D}_2)(\mathcal{D}_1 + \mathcal{D}_2 + 1)Z. \quad (1)$$

This system turns out to be quite interesting, although from the point of view of Appell's function F_4 it looks somewhat degenerate. Set

$$\begin{aligned} \Delta &= (1 + \tau_1 + \tau_2)(1 + \tau_1 - \tau_2)(1 - \tau_1 + \tau_2)(1 - \tau_1 - \tau_2) \\ \Gamma &= 1 - \tau_1^2 - \tau_2^2 \\ \Lambda_{\pm} &= ((2\tau_1^2 \pm \Gamma) \tau_2 d\tau_1 \pm (2\tau_2^2 \pm \Gamma) \tau_1 d\tau_2) / (2\tau_1 \tau_2) \end{aligned}$$

Let $Z_{\pm} = (\mathcal{D}_1 Z \pm \mathcal{D}_2 Z)/2$ and $Z_3 = (\mathcal{D}_1 Z + \mathcal{D}_2 Z + 2\mathcal{D}_1 \mathcal{D}_2 Z)/2\Gamma$. Noticing that the differential equations can be rewritten as $\mathcal{D}_i^2 Z = 2\tau_i^2 Z_3$, one easily deduces the following Pfaffian system equivalent to (1)

$$d \begin{bmatrix} Z \\ Z_- \\ Z_+ \\ Z_3 \end{bmatrix} = \begin{bmatrix} 0 & \frac{d(\tau_1/\tau_2)}{\tau_1/\tau_2} & \frac{d(\tau_1\tau_2)}{\tau_1\tau_2} & 0 \\ 0 & 0 & \frac{d(\tau_1/\tau_2)}{2\tau_1/\tau_2} & \Lambda_- \\ 0 & 0 & -\frac{d(\tau_1\tau_2)}{2\tau_1\tau_2} & \Lambda_+ \\ 0 & 0 & -\frac{\Lambda_+}{\Delta} & \frac{d(\tau_1\tau_2)}{2\tau_1\tau_2} - \frac{d\Delta}{\Delta} \end{bmatrix} \begin{bmatrix} Z \\ Z_- \\ Z_+ \\ Z_3 \end{bmatrix} \quad (2)$$

2 Variation of mixed Hodge structure

We will associate with (2) a variation of mixed Hodge structure $(\mathbf{V}_{\mathbf{Z}}, W, \mathcal{F})$ of geometric origin [6]. In this section we describe this VMHS abstractly. In the subsequent sections it will be realized in geometry.

The base space \mathcal{B} is the complement in the projective plane \mathbf{P}^2 of the complete quadrilateral with equation

$$t_1 t_2 (t_0 + t_1 + t_2)(t_0 + t_1 - t_2)(t_0 - t_1 + t_2)(t_0 - t_1 - t_2) = 0 \quad (3)$$

The construction of the vector bundle with connection works completely within the algebraic geometry of schemes over $\mathbf{Z}[\frac{1}{2}]$. The vector bundle \mathcal{V} underlying the VMHS has rank 4 and is generated by global sections $|\omega_1|, |\omega_2|, |\omega_3|, |\omega_4|$. The weight filtration $W_0\mathcal{V} \subset W_1\mathcal{V} \subset W_2\mathcal{V} \subset W_3\mathcal{V} \subset W_4\mathcal{V}$ is

$$0 \subset \mathcal{O}_{\mathcal{B}} |\omega_3| \oplus \mathcal{O}_{\mathcal{B}} |\omega_4| \subset W_1\mathcal{V} \oplus \mathcal{O}_{\mathcal{B}} |\omega_2| = W_3\mathcal{V} \subset \mathcal{V}.$$

The Hodge filtration $\mathcal{F}^0\mathcal{V} \supset \mathcal{F}^1\mathcal{V} \supset \mathcal{F}^2\mathcal{V} \supset \mathcal{F}^3\mathcal{V}$ is given by

$$\mathcal{V} \supset \mathcal{O}_{\mathcal{B}} |\omega_1| \oplus \mathcal{O}_{\mathcal{B}} |\omega_2| \oplus \mathcal{O}_{\mathcal{B}} |\omega_3| \supset \mathcal{O}_{\mathcal{B}} |\omega_1| \supset 0.$$

The vector bundle carries a connection compatible with the weight filtration

$$\nabla : \mathcal{V} \rightarrow \Omega_{\mathcal{B}/\mathbf{Z}}^1 \otimes \mathcal{V}.$$

With respect to the basis $|\omega_1|, |\omega_2|, |\omega_3|, |\omega_4|$ the connection is given by

$$\nabla \underline{\omega} = L \underline{\omega} \quad (4)$$

where L is the connection matrix from (2) and $\underline{\omega} = (|\omega_1|, |\omega_2|, |\omega_3|, |\omega_4|)^t$.

For the local system on \mathcal{B} we must pass to analytic geometry and view \mathcal{B} as a complex analytic space. The local system $\mathbf{V}_{\mathbf{Z}}$ on \mathcal{B} and its relation with \mathcal{V} are most easily described over the universal covering space $\tilde{\mathcal{B}}$ of \mathcal{B} . Let $q : \tilde{\mathcal{B}} \rightarrow \mathcal{B}$ be the covering projection. Then $\tilde{\mathbf{V}}_{\mathbf{Z}} = q^* \mathbf{V}_{\mathbf{Z}}$ is a constant sheaf on $\tilde{\mathcal{B}}$, equal to a free \mathbf{Z} -module with basis $\mathbf{e}_1, \mathbf{e}_2, \mathbf{e}_3, \mathbf{e}_4$. The weight filtration $W_0 \tilde{\mathbf{V}}_{\mathbf{Z}} \subset W_1 \tilde{\mathbf{V}}_{\mathbf{Z}} \subset W_2 \tilde{\mathbf{V}}_{\mathbf{Z}} \subset W_3 \tilde{\mathbf{V}}_{\mathbf{Z}} \subset W_4 \tilde{\mathbf{V}}_{\mathbf{Z}}$ is

$$0 \subset \mathbf{Z} \mathbf{e}_3 \oplus \mathbf{Z} \mathbf{e}_4 \subset W_1 \tilde{\mathbf{V}}_{\mathbf{Z}} \oplus \mathbf{Z} \mathbf{e}_2 = W_3 \tilde{\mathbf{V}}_{\mathbf{Z}} \subset \tilde{\mathbf{V}}_{\mathbf{Z}}.$$

There is an isomorphism which is compatible with the weight filtrations

$$\mathcal{I} : \mathcal{V} \otimes_{\mathcal{O}_{\mathcal{B}}} \mathcal{O}_{\tilde{\mathcal{B}}} \xrightarrow{\cong} \tilde{\mathbf{V}}_{\mathbf{Z}} \otimes_{\mathbf{Z}} \mathcal{O}_{\tilde{\mathcal{B}}}.$$

With respect to the bases $\underline{\omega}$ and $\underline{\mathbf{e}}$ it is described by a matrix $\mathcal{P} = (\mathcal{P}_{ij})$ of analytic functions on $\tilde{\mathcal{B}}$:

$$\mathcal{I} \underline{\omega} = \mathcal{P} \underline{\mathbf{e}}. \quad (5)$$

Here $\underline{\mathbf{e}}$ is the column vector $(\mathbf{e}_1, \mathbf{e}_2, \mathbf{e}_3, \mathbf{e}_4)^t$. The frame $\underline{\mathbf{e}}$ is horizontal:

$$\nabla \mathbf{e}_i = 0, \quad i = 1, 2, 3, 4. \quad (6)$$

From (4)-(6) one sees that the columns of \mathcal{P} satisfy the differential equation (2) and that $\mathcal{P}_{11}, \mathcal{P}_{12}, \mathcal{P}_{13}, \mathcal{P}_{14}$ span the solution space of Appell's system (1) over \mathbf{C} . This only involves the \mathbf{C} -vector space $\tilde{\mathbf{V}}_{\mathbf{Z}} \otimes \mathbf{C}$.

Fix a base point $b \in \mathcal{B}$. The natural action of the fundamental group $\pi_1 = \pi_1(\mathcal{B}, b)$ on \mathcal{B} induces a representation ν of π_1 on the sheaf of functions $\mathcal{O}_{\tilde{\mathcal{B}}}$. Let $\mathbf{1}$ denote the trivial representation of π_1 on \mathcal{V} . The monodromy representation μ of π_1 on $\tilde{\mathbf{V}}_{\mathbf{Z}}$ is such that \mathcal{I} is actually an isomorphism of representations $\mathcal{I} : \mathbf{1} \otimes \nu \simeq \mu \otimes \nu$. The integral structure $\tilde{\mathbf{V}}_{\mathbf{Z}}$ comes in if one wants the monodromy representation to take values in $GL_4(\mathbf{Z})$.

3 A family of elliptic curves minus four points

This section realizes $(W_1 \mathcal{V} \subset W_2 \mathcal{V}, \nabla)$ in the de Rham cohomology of a family of elliptic curves minus four points.

Consider in $\mathbf{P}^2 \times \mathbf{P}^1 \times \mathbf{P}^1$ over $\mathbf{Z}[\frac{1}{2}]$ the subscheme \mathcal{E} defined by

$$2t_0 x_1 y_1 x_2 y_2 + t_1 (x_1^2 + y_1^2) x_2 y_2 + t_2 x_1 y_1 (x_2^2 + y_2^2) = 0.$$

Take in \mathbf{P}^2 the complete quadrilateral $\boxed{\times}$ defined by (3) and in $\mathbf{P}^1 \times \mathbf{P}^1$ the quadrilateral $\square : x_1 y_1 x_2 y_2 = 0$. Let p and p' be the projections from $\mathbf{P}^2 \times \mathbf{P}^1 \times \mathbf{P}^1$ onto \mathbf{P}^2 and $\mathbf{P}^1 \times \mathbf{P}^1$, respectively. We define

$$\mathcal{B} = \mathbf{P}^2 \setminus \boxed{\times}, \quad \mathcal{A} = \mathcal{E} \cap p^{-1}(\mathcal{B}), \quad \mathcal{S} = \mathcal{A} \cap p'^{-1}(\square), \quad \mathcal{A}^\circ = \mathcal{A} \setminus \mathcal{S}.$$

Then $p : \mathcal{A} \rightarrow \mathcal{B}$ is a family of elliptic curves over the base \mathcal{B} and \mathcal{S} consists of the four sections (00) , (0∞) , $(\infty 0)$ and $(\infty\infty)$.

On \mathcal{A}° we use the inhomogeneous coordinates $\xi_1 = x_1/y_1$, $\xi_2 = x_2/y_2$. For the computations showing that the Gauss-Manin connection on a subspace of the de Rham cohomology of $\mathcal{A}^\circ/\mathcal{B}$ matches the lower right hand 3×3 -block of (2) it suffices to work on $\mathcal{B} \setminus \{t_0 = 0\}$ and use the inhomogeneous coordinates $\tau_1 = t_1/t_0$, $\tau_2 = t_2/t_0$. In these coordinates \mathcal{A}° is described by the equation

$$\mathcal{A}^\circ : \quad 2 + \tau_1(\xi_1 + \xi_1^{-1}) + \tau_2(\xi_2 + \xi_2^{-1}) = 0.$$

Consider the three differential forms on \mathcal{A}/\mathcal{B} :

$$\begin{aligned} \omega_3 &= \frac{-1}{\tau_1(\xi_1 - \xi_1^{-1})} \frac{d\xi_2}{\xi_2} \\ \omega_4 &= \frac{2(\xi_1 + \xi_1^{-1})}{\tau_1^2(\xi_1 - \xi_1^{-1})^3} \frac{d\xi_2}{\xi_2} \\ \omega_2 &= \frac{\tau_1(\xi_1 + \xi_1^{-1}) - \tau_2(\xi_2 + \xi_2^{-1})}{2\tau_1(\xi_1 - \xi_1^{-1})} \frac{d\xi_2}{\xi_2}. \end{aligned}$$

These are forms of the first, second and third kind respectively. Their cohomology classes $|\omega_3|$, $|\omega_4|$, $|\omega_2|$ are elements of the de Rham cohomology group $\mathbf{H}^1(\mathcal{A}, \Omega_{\mathcal{A}/\mathcal{B}}^\bullet(\log(\mathcal{S})))$, hypercohomology of the complex of differential forms on \mathcal{A} relative to \mathcal{B} with at most logarithmic singularities along \mathcal{S} . The 3-dimensional subspace spanned by these classes can be characterized as the +1 eigenspace of the involution σ on $\mathbf{H}^1(\mathcal{A}, \Omega_{\mathcal{A}/\mathcal{B}}^\bullet(\log(\mathcal{S})))$ induced by the involution σ of $\mathcal{A}^\circ/\mathcal{B}$ defined by

$$\sigma(\xi_1) = \xi_1^{-1}, \quad \sigma(\xi_2) = \xi_2^{-1}.$$

$\mathbf{H}^1(\mathcal{A}, \Omega_{\mathcal{A}/\mathcal{B}}^\bullet(\log(\mathcal{S})))^{\sigma=1}$ carries a Gauss-Manin connection ∇ [3, §4]. One can show (the proofs will appear elsewhere)

$$\nabla \begin{pmatrix} |\omega_2| \\ |\omega_3| \\ |\omega_4| \end{pmatrix} = L \begin{pmatrix} |\omega_2| \\ |\omega_3| \\ |\omega_4| \end{pmatrix}$$

where L is the lower right hand 3×3 block of the matrix in (2). It is well known that the vector bundle with connection $(\mathbf{H}^1(\mathcal{A}, \Omega_{\mathcal{A}/\mathcal{B}}^\bullet(\log(\mathcal{S})))^{\sigma=1}, \nabla)$ underlies a variation of mixed Hodge structure on \mathcal{B} [6]. The local system of this VMHS is $\mathbf{R}^1 p_* \mathbf{Z}_{\mathcal{A}^\circ}$, where $p^\circ : \mathcal{A}^\circ \rightarrow \mathcal{B}$ is the restriction of p and $\mathbf{Z}_{\mathcal{A}^\circ}$ is the constant sheaf \mathbf{Z} on \mathcal{A}° .

We use the following notation emphasizing the weight filtration

$$\begin{aligned} W_1 \mathcal{V} &= \mathbf{H}^1(\mathcal{A}, \Omega_{\mathcal{A}/\mathcal{B}}^\bullet), & W_2 \mathcal{V} &= \mathbf{H}^1(\mathcal{A}, \Omega_{\mathcal{A}/\mathcal{B}}^\bullet(\log \mathcal{S}))^{\sigma=1} \\ W_1 \mathbf{V}_{\mathbf{Z}} &= \mathbf{R}^1 p_* \mathbf{Z}_{\mathcal{A}}, & W_2 \mathbf{V}_{\mathbf{Z}} &= (\mathbf{R}^1 p_* \mathbf{Z}_{\mathcal{A}^\circ})^{\sigma=1} \end{aligned}$$

This VMHS is an extension of the Tate Hodge structure $\mathbf{Z}(-1)$, viewed as a constant VHS on \mathcal{B} , by the VHS associated with the family $\mathcal{A} \rightarrow \mathcal{B}$ of complete elliptic curves.

4 A rank 4 VMHS; Bloch's regulator

There is an obvious and simple way to extend the rank 3 bundle $W_2 \mathcal{V}$ to a rank 4 bundle \mathcal{V} with connection giving the full 4×4 connection matrix of (2); namely

$$\mathcal{V} = W_2 \mathcal{V} \oplus \mathcal{O}_{\mathcal{B}} \cdot |\omega_1|$$

with connection ∇ inducing the Gauss-Manin connection on $W_2 \mathcal{V}$ and such that for the global section $|\omega_1|$ of \mathcal{V}

$$\nabla |\omega_1| = \frac{d(\tau_1/\tau_2)}{\tau_1/\tau_2} |\omega_2| + \frac{d(\tau_1\tau_2)}{\tau_1\tau_2} |\omega_3|.$$

In order to get a rank 4 VMHS we must also extend the local system. For this we use Bloch's regulator map and some K-theory.

Algebraic K-theory [5] constructs for every scheme \mathcal{X} abelian groups $K_i(\mathcal{X})$, $i \geq 0$. For a commutative ring R with 1 one sets $K_i(R) = K_i(\text{Spec } R)$. Two invertible elements u, v of R determine an element $\{u, v\}$ of $K_2(R)$ called Steinberg symbol. It satisfies $\{u^{-1}, v^{-1}\} = \{u, v\}$ [4]. In particular one has

$$\{\xi_1, \xi_2\} \in K_2(\mathcal{A}^\circ)^{\sigma=1}.$$

Bloch's regulator map [2] for an open Riemann surface \mathcal{X} is defined on the subgroup of $K_2(\mathcal{X})$ generated by the Steinberg symbols and takes values in $\mathbf{H}^1(\mathcal{X}, \mathbf{C}^*)$. The construction applies to Steinberg symbols in K_2 of the fibres of the family $\mathcal{A}^\circ/\mathcal{B}$, but the formulas in [2] can easily be re-interpreted so that they work for $\{\xi_1, \xi_2\}$ in $K_2(\mathcal{A}^\circ)$. For this we need the multiplicative

de Rham complex on \mathcal{A} relative to \mathcal{B} with at most logarithmic singularities along \mathcal{S} :

$$\Omega_{\mathcal{A}/\mathcal{B}}^{\times\bullet}(\log(\mathcal{S})) = [j_* \mathcal{O}_{\mathcal{A}^\circ}^* \xrightarrow{d \log} \Omega_{\mathcal{A}/\mathcal{B}}^1(\log(\mathcal{S}))].$$

Here j is the inclusion $\mathcal{A}^\circ \hookrightarrow \mathcal{A}$. Let \mathcal{A}°_+ resp. $\mathcal{A}^\circ_- \subset \mathcal{A}^\circ$ be the inverse image of $\mathbf{C} \setminus (-\infty, 0]$ resp. $\mathbf{C} \setminus [0, \infty)$ with respect to the function ξ_1 . Let \log_+ resp. \log_- be a branch of the logarithm on $\mathbf{C} \setminus (-\infty, 0]$ resp. $\mathbf{C} \setminus [0, \infty)$. Then

$$\begin{aligned} \xi_2^{(\log_-(\xi_1) - \log_+(\xi_1))/2\pi i} & \text{ on } \mathcal{A}^\circ_+ \cap \mathcal{A}^\circ_- \\ \frac{-1}{2\pi i} \log_\pm(\xi_1) \frac{d\xi_2}{\xi_2} & \text{ on } \mathcal{A}^\circ_\pm \end{aligned} \quad (7)$$

gives a Čech cocycle for $\Omega_{\mathcal{A}/\mathcal{B}}^{\times\bullet}(\log(\mathcal{S}))$ and hence represents an element $r\{\xi_1, \xi_2\}$ of its \mathbf{H}^1 . Formula (7) shows that $r\{\xi_1, \xi_2\}$ is invariant under the involution σ :

$$r\{\xi_1, \xi_2\} \in \mathbf{H}^1(\mathcal{A}, \Omega_{\mathcal{A}/\mathcal{B}}^{\times\bullet}(\log(\mathcal{S})))^{\sigma=1}.$$

One can immitate the Katz-Oda construction for the Gauss-Manin connection and construct a map

$$\mathbf{H}^1(\mathcal{A}, \Omega_{\mathcal{A}/\mathcal{B}}^{\times\bullet}(\log(\mathcal{S}))) \xrightarrow{\nabla} \Omega_{\mathcal{B}/\mathbf{C}}^1 \otimes \mathbf{H}^1(\mathcal{A}, \Omega_{\mathcal{A}/\mathcal{B}}^\bullet(\log(\mathcal{S}))) \quad (8)$$

which composes with the map, coming from the exponential sequence,

$$\mathbf{H}^1(\mathcal{A}, \Omega_{\mathcal{A}/\mathcal{B}}^\bullet(\log(\mathcal{S}))) \rightarrow \mathbf{H}^1(\mathcal{A}, \Omega_{\mathcal{A}/\mathcal{B}}^{\times\bullet}(\log(\mathcal{S}))), \quad (9)$$

to the usual Gauss-Manin connection on $\mathbf{H}^1(\mathcal{A}, \Omega_{\mathcal{A}/\mathcal{B}}^\bullet(\log(\mathcal{S})))$. One can easily compute $\nabla r\{\xi_1, \xi_2\}$ in (8):

$$2\pi i \nabla r\{\xi_1, \xi_2\} = -\frac{d\xi_1}{\xi_1} \frac{d\xi_2}{\xi_2} = \frac{d(\tau_1/\tau_2)}{\tau_1/\tau_2} \omega_2 + \frac{d(\tau_1\tau_2)}{\tau_1\tau_2} \omega_3 = \nabla|\omega_1|. \quad (10)$$

Now we need the universal covering space $\tilde{\mathcal{B}}$ of \mathcal{B} . Let $\tilde{\mathcal{A}}, \tilde{\mathcal{S}}, \tilde{\mathcal{A}}^\circ$ be the fibre products over \mathcal{B} of $\tilde{\mathcal{B}}$ with $\mathcal{A}, \mathcal{S}, \mathcal{A}^\circ$, respectively. The involution σ lifts to $\tilde{\mathcal{A}}^\circ/\tilde{\mathcal{B}}$. The obvious analogue of (7) defines an element $\tilde{r}\{\xi_1, \xi_2\}$ in $\mathbf{H}^1(\tilde{\mathcal{A}}, \Omega_{\tilde{\mathcal{A}}/\tilde{\mathcal{B}}}^{\times\bullet}(\log(\tilde{\mathcal{S}})))^{\sigma=1}$. It can be shown (details will appear elsewhere) that in the analogue of (9) $\tilde{r}\{\xi_1, \xi_2\}$ can be lifted to an element R in $\mathbf{H}^1(\tilde{\mathcal{A}}_{\tilde{\mathcal{B}}}, \Omega_{\tilde{\mathcal{A}}/\tilde{\mathcal{B}}}^\bullet(\log(\tilde{\mathcal{S}})))^{\sigma=1}$. Define

$$e_1 = \frac{1}{(2\pi i)^2} |\omega_1| - \frac{1}{2\pi i} R \quad (11)$$

This is a section of the vector bundle $\tilde{\mathcal{V}} = \mathcal{V} \otimes \mathcal{O}_{\tilde{\mathcal{B}}}$. Because of (10) \mathbf{e}_1 is a horizontal section:

$$\nabla \mathbf{e}_1 = 0.$$

The vector bundles $W_1\mathcal{V}$, $W_2\mathcal{V}$ pull back to the vector bundles $W_1\mathcal{V} \otimes \mathcal{O}_{\tilde{\mathcal{B}}}$, $W_2\mathcal{V} \otimes \mathcal{O}_{\tilde{\mathcal{B}}}$ on $\tilde{\mathcal{B}}$. The local systems $W_1\mathbf{V}_{\mathbf{Z}}$, $W_2\mathbf{V}_{\mathbf{Z}}$ pull back to constant sheaves $W_1\tilde{\mathbf{V}}_{\mathbf{Z}}$, $W_2\tilde{\mathbf{V}}_{\mathbf{Z}}$ on $\tilde{\mathcal{B}}$. Choosing a base point $b \in \mathcal{B}$ we have

$$\begin{aligned} W_1\tilde{\mathbf{V}}_{\mathbf{Z}} &= H^1(\mathcal{A}_b, \mathbf{Z}) \\ W_2\tilde{\mathbf{V}}_{\mathbf{Z}} &= H^1(\mathcal{A}^\circ_b, \mathbf{Z})^{\sigma=1} \end{aligned}$$

where \mathcal{A}_b and \mathcal{A}°_b are the fibres over b . We can choose any basis $\mathbf{e}_3, \mathbf{e}_4$ for the free rank 2 \mathbf{Z} -module $W_1\tilde{\mathbf{V}}_{\mathbf{Z}}$ and extend to a basis $\mathbf{e}_2, \mathbf{e}_3, \mathbf{e}_4$ for the free rank 3 \mathbf{Z} -module $W_2\tilde{\mathbf{V}}_{\mathbf{Z}}$ so that under the period isomorphism from Hodge theory $W_2\mathcal{V} \otimes \mathcal{O}_{\tilde{\mathcal{B}}} \simeq W_2\tilde{\mathbf{V}}_{\mathbf{Z}} \otimes \mathcal{O}_{\tilde{\mathcal{B}}}$ we have

$$|\omega_2| \equiv 2\pi i \mathbf{e}_2 \pmod{W_1\mathcal{V} \otimes \mathcal{O}_{\tilde{\mathcal{B}}}}$$

Define

$$\tilde{\mathbf{V}}_{\mathbf{Z}} = \mathbf{Z}\mathbf{e}_1 \oplus W_3\tilde{\mathbf{V}}_{\mathbf{Z}} \subset \tilde{\mathcal{V}}$$

The fundamental group $\pi_1 = \pi_1(\mathcal{B}, b)$ acts on $\tilde{\mathcal{B}}$, $\tilde{\mathcal{A}}$, $\tilde{\mathcal{A}}^\circ$, the various de Rham complexes and their cohomology. $\tilde{r}\{\xi_1, \xi_2\}$ is the image of $r\{\xi_1, \xi_2\}$ under the map on cohomology induced by the covering map $q: \tilde{\mathcal{B}} \rightarrow \mathcal{B}$. So it is invariant under the action of π_1 . Note also that $|\omega_1|$ is π_1 -invariant. Therefore, for $\phi \in \pi_1$,

$$\phi \mathbf{e}_1 - \mathbf{e}_1 = \frac{1}{2\pi i} (R - \phi R) \in W_3\tilde{\mathbf{V}}_{\mathbf{Z}}$$

We see that

for the given choice of $\mathbf{e}_1, \mathbf{e}_2, \mathbf{e}_3, \mathbf{e}_4$ and $\tilde{\mathbf{V}}_{\mathbf{Z}}$ the monodromy representation takes values in $GL_4(\mathbf{Z})$:

$$\pi_1(\mathcal{B}, b) \rightarrow \text{Aut}_{\mathbf{Z}}(\tilde{\mathcal{V}}) \simeq GL_4(\mathbf{Z}).$$

This completes the construction of the rank 4 variation of mixed Hodge structure for Appell's system of differential equations of type F_4 with parameters $0, \frac{1}{2}, 1, 1$ and variables τ_1^2, τ_2^2 .

References

- [1] Appell, P.- Kampé de Fériet, J. *Fonctions hypergéométriques et hypersphériques; polynômes d'Hermite*, Paris 1926: Gauthier-Villars
- [2] Bloch, S. *The dilogarithm and extensions of Lie algebras*, in: Lecture Notes in Math. 854 (Springer Verlag 1981) 1-23
- [3] Katz, N. *Nilpotent connections and the monodromy theorem: applications of a result of Turrittin*, Publ. Math. IHES 39 (1970) 175-232
- [4] Milnor, J. *Introduction to algebraic K-theory*, Annals of Math. Studies 72, Princeton: Princeton University Press, 1971
- [5] Quillen, D. *Higher algebraic K-theory I*, in: Lecture Notes in Math. 341 (Springer Verlag 1973) 85-147
- [6] Steenbrink, J.- Zucker, S. *Variation of mixed Hodge structure I*, Invent. Math. 80 (1985) 489-542

Acknowledgements. I would like to thank the Taniguchi Foundation for its support and the organizers of the Workshop and the Symposium on "Special Differential Equations" for creating an interesting and stimulating atmosphere.

Jan Stienstra
Mathematisch Instituut der Rijksuniversiteit te Utrecht
Postbus 80.010, 3508 TA Utrecht, The Netherlands