

MOTIVIC MEASURES

EDUARD LOOIJENGA

1. INTRODUCTION

An n -jet of an arc in an algebraic variety is a one parameter Taylor series of length n in that variety. To be precise, if the variety X is defined over the algebraically closed field k , then it is a $k[[t]]/(t^{n+1})$ -valued point of X . The set of such n -jets are the closed points of a variety $\mathcal{L}_n(X)$ also defined over k and the arc space of X , $\mathcal{L}(X)$, is the projective limit of these. Probably Nash [24] was the first to study arc spaces in a systematic fashion (the paper in question was written in 1968). He concentrated on arcs based at a given point of X and observed that to each irreducible component of this ‘provariety’ there corresponds in an injective manner an irreducible component of the preimage of this point in any resolution of X . He asked the (still unanswered) question how to identify these components on a given resolution. The renewed interest in arc spaces has a different origin, however. Batyrev [4] proved that two connected projective complex manifolds with trivial canonical bundle which are birationally equivalent must have the same Betti numbers. This he showed by first lifting the data to a situation over a discrete valuation ring with finite residue field and then exploiting a p -adic integration technique. (Such a p -adic integration approach to problems in complex algebraic geometry had also been used by Denef and Loeser [12] in their work on topological zeta functions attached to singular points of complex varieties.) When Kontsevich learned of Batyrev’s result he saw how this proof could be made to work in a complex setting using arc spaces. The new proof also gave more: equality of Hodge numbers, and even an isomorphism of Hodge structures with rational coefficients. The underlying technique, now going under the name of *motivic integration*, has led to an avalanche of applications. These include new (so-called *stringy*) invariants of singularities, a complex analogue of the Igusa zeta function, a motivic version of the Thom-Sebastiani property and the motivic McKay correspondence. Some of these were covered in a recent talk by Reid [26] in this seminar.

The idea is simple if we keep in mind an analogous, more classical situation. Consider the case of a complete discrete valuation ring (R, m) with finite residue field F . There is a Haar measure on the Boolean algebra consisting of the cosets of powers of m that takes the value 1 on R (so it is also a probability measure). This induces one on a suitable Boolean algebra of subsets of the set of R -valued points of any scheme that is flat of pure dimension and of finite type over $\text{Spec}(R)$. Associated to this measure is a function that essentially counts the number of ‘points’ in each reduction modulo m^k : the *Igusa zeta function*, introduced by Weil, and intensively studied by Igusa, Denef and Loeser (and reported on by Denef in this seminar [11]). A missing case was that of equal characteristic zero: $\mathcal{O} = k[[t]]$, $k \supset \mathbb{Q}$. The proposal of Kontsevich is to give \mathcal{O} a measure that takes values in a Grothendieck ring of k -varieties in which the class of the affine line, \mathbb{L} , is invertible: the value on

the ideal (t^n) is then simply \mathbb{L}^{-n} (or \mathbb{L}^{1-n} , which is sometimes more convenient). If \mathcal{X} is a suitable \mathcal{O} -scheme, then we obtain a measure on the set of sections as before, but now with values in this Grothendieck ring. The corresponding zeta function is a very fine bookkeeping device, for it does its counting in a ring that is huge. There is no a priori reason to restrict to the case of equal characteristic, for Kontsevich's idea makes sense for any complete discrete valuation ring. Indeed, with little extra effort the material in Sections 2,3 and 9 can be generalized to that context.

This report concerns mainly work of Denef and Loeser. Some of their results are presented here somewhat differently, and this is why more proofs are provided than one perhaps expects of the write up of a seminar talk. References to the sources are in general given after the section titles, rather than in the statements of theorems.

I thank Jan Denef for inviting me for a short visit to Leuven to discuss the material exposed here. I am also indebted to Maxim Kontsevich and especially to Jan Denef for comments on previous versions, from which this text has greatly benefitted (though remaining errors are my responsibility only). This applies in particular to the motivic Thom-Sebastiani theorem and a word of explanation is in order here. In the original version I had introduced (albeit somewhat implicitly) a binary operator on a certain Grothendieck ring of motives, called here quasi-convolution. Quasi-convolution is almost associative, but not quite, and since I thought this to be a serious defect, I passed to the universal associative quotient. But in a recent overview, Denef and Loeser [19] noted that there is no need for this: the property one wants (which is another than associativity) holds already without passing that to quotient. As this no longer justifies its introduction, I thought it best to take advantage of their observation and rewrite things accordingly.

2. THE ARC SPACE AND ITS MEASURE [14], [23]

Throughout the talk we fix a complete discrete valuation ring \mathcal{O} whose residue field k is assumed to be algebraically closed and of characteristic zero. The spectrum of \mathcal{O} is denoted \mathbb{D} with generic point \mathbb{D}^\times and closed point o . A uniformizing parameter is often denoted by t so that $\mathcal{O} = k[[t]]$. The assumption that k be algebraically closed is for convenience only: in most situations this restriction is unnecessary or can be avoided.

The symbol \mathbb{N} stands for the set of nonnegative integers.

The Grothendieck ring of varieties. Consider the *Grothendieck ring* $K_0(\mathcal{V}_k)$ of reduced k -varieties: this is the abelian group generated by the isomorphism classes of such varieties, subject to the relations $[X - Y] = [X] - [Y]$, where Y is a closed in X . The product over k turns it into a ring. Note that if we restrict ourselves to smooth varieties we get the same ring: the reason is that every k -variety X admits a stratification (i.e., a filtration by closed subschemes $X = X^0 \supset X^1 \supset \dots \supset X^{d+1} = \emptyset$ such that $X^k - X^{k+1}$ is smooth) and that any two such admit a common refinement. The latter property implies that $[X] := \sum_k [X^k - X^{k+1}]$ is unambiguously defined. In fact, $K_0(\mathcal{V}_k)$ is generated by the classes of complete nonsingular varieties, for any smooth variety U admits a completion \overline{U} by adding a normal crossing divisor and then $[U] = \sum (-1)^i [\overline{U}^i]$, where \overline{U}^i stands for the normalization of the codimension i skeleton of the resulting stratification. Włodarczyk's weak factorization theorem (in the form of the main theorem of [1]) can be used to show that relations of the following simple type suffice: if X is smooth

projective and $\tilde{X} \rightarrow X$ is obtained by blowing up a smooth closed subvariety $Y \subset X$ with exceptional divisor \tilde{Y} , then $[\tilde{X}] - [\tilde{Y}] = [X] - [Y]$.

We denote the class of the affine line \mathbb{A}^1 by \mathbb{L} and we write M_k for the localization $K_0(\mathcal{V}_k)[\mathbb{L}^{-1}]$. Recall that a subset of a variety X is called *constructible* if it is a finite union of (locally closed) subvarieties. Any constructible subset C of X defines an element $[C] \in M_k$. The constructible subsets of X form a Boolean algebra and so we obtain in a tautological manner a M_k -valued measure μ_X defined on this Boolean algebra. More generally, a morphism $f : Y \rightarrow X$ defines on that same algebra an M_k -valued measure $f_*\mu_Y$: assign to a constructible subset of X its preimage in Y .

The ring M_k is interesting, big, and hard to grasp. Fortunately, there are several characteristics of M_k (i.e., ring homomorphisms from M_k to a ring) that are well understood. We describe some of these in decreasing order of complexity under the assumption that k is a subfield of \mathbb{C} . The first example is the Grothendieck ring $K_0(\text{HS})$ of the category of Hodge structures. A *Hodge structure* consists of a finite dimensional \mathbb{Q} -vector space H , a finite bigrading $H \otimes \mathbb{C} = \bigoplus_{p,q \in \mathbb{Z}} H^{p,q}$ such that $H^{p,q}$ is the complex conjugate of $H^{q,p}$ and each *weight summand*, $\bigoplus_{p+q=m} H^{p,q}$, is defined over \mathbb{Q} . There are evident notions of tensor product and morphism of Hodge structures so that we get an abelian category HS with tensor product. The Grothendieck construction produces a group $K_0(\text{HS})$, elements of which are representable as a formal difference of Hodge structures $[H] - [H']$ and $[H] = [H']$ if and only if H and H' are isomorphic. The tensor product makes it a ring.

For every complex variety X , the cohomology with compact supports, $H_c^r(X; \mathbb{Q})$, comes with a natural finite increasing filtration $W_\bullet H_c^r(X; \mathbb{Q})$, the *weight filtration*, such that the associated graded $\text{Gr}_\bullet^W H_c^r(X; \mathbb{Q})$ underlies a Hodge structure having $\text{Gr}_m^W H_c^r(X; \mathbb{Q})$ as weight m summand. We assign to X the *Hodge characteristic*¹

$$\chi_h(X) := \sum_r (-1)^r [H_c^r(X; \mathbb{Q})] \in K_0(\text{HS})$$

If $Y \subset X$ is closed subvariety, then the exact sequence

$$\cdots \rightarrow H_c^r(X - Y) \rightarrow H_c^r(X) \rightarrow H_c^r(Y) \rightarrow H_c^{r+1}(X - Y) \rightarrow \cdots$$

is compatible in a strong sense with the Hodge data. This implies the additivity property $\chi_h(X) = \chi_h(X - Y) + \chi_h(Y)$. For the affine line \mathbb{A}^1 , $H_c^r(\mathbb{A}^1; \mathbb{Q})$ is nonzero only for $r = 2$; the cohomology group $H_c^2(\mathbb{A}^1; \mathbb{Q})$ is one-dimensional and of type $(1, 1)$. So $\chi_h(\mathbb{A}^1)$ (usually denoted as $\mathbb{Q}(-1)$) is invertible. It follows that χ_h factorizes over M_k . If we only care for dimensions, then we compose with the ring homomorphism $K_0(\text{HS}) \rightarrow \mathbb{Z}[u, u^{-1}, v, v^{-1}]$, $[H] \mapsto \sum_{p,q} \dim(H^{p,q}) u^p v^q$, to get the *Hodge number characteristic* $\chi_{\text{hn}} : M_k \rightarrow \mathbb{Z}[u, u^{-1}, v, v^{-1}]$. It takes \mathbb{L} to uv . The *weight characteristic* $\chi_{\text{wt}} : M_k \rightarrow \mathbb{Z}[w, w^{-1}]$ is obtained if we go further down along the map $\mathbb{Z}[u, u^{-1}, v, v^{-1}] \rightarrow \mathbb{Z}[w, w^{-1}]$ that sends both u and v to w . Evaluating the latter at $w = 1$ gives the ordinary² Euler characteristic $\chi_{\text{top}} : M_k \rightarrow \mathbb{Z}$.

In the spirit of this discussion is the following question raised by Kapranov [22]:

¹As all our characteristics are compactly supported we omit the otherwise desirable subscript c from the notation.

²A complex algebraic variety can be compactified within its homotopy type by giving it a topological boundary that is stratifiable into strata of odd dimension. This boundary has zero Euler characteristic, hence the compactly supported Euler characteristic of the variety is its ordinary Euler characteristic.

Question 2.1. Let X be a variety over k . If $\sigma_n(X) \in M_k$ denotes the class of its n th symmetric power, is then

$$Z_X(T) := 1 + \sum_{n=1}^{\infty} \sigma_n(X) T^n \in M_k[[T]]$$

a rational function in the sense that it determines an element in a suitable localization of $M_k[[T]]$? (Since the logarithmic derivative Z'/Z defines an additive map $M_k \rightarrow M_k[[T]]$, we may restrict ourselves here to the case of a smooth variety.) Does it satisfy a functional equation when X is smooth and complete? Kapranov shows that the answer to both questions is yes in case $\dim(X) \leq 1$.

A measure on the space of sections. Let us call a \mathbb{D} -variety a separated reduced scheme that is flat and of finite type over \mathbb{D} and whose closed fiber is reduced. Given a \mathbb{D} -variety \mathcal{X}/\mathbb{D} with closed fiber X , then the set of its sections up to order n , \mathcal{X}_n , is the set of closed points of a k -variety (also denoted \mathcal{X}_n) naturally associated to \mathcal{X} . It is obtained from \mathcal{X} modulo \mathfrak{m}^{n+1} essentially by Weil restriction of scalars [20]. So $\mathcal{X}_0 = X$. The set \mathcal{X}_∞ of sections of $\mathcal{X} \rightarrow \mathbb{D}$ is the projective limit of these and is therefore the set of closed points of a provariety. If \mathcal{X}/\mathbb{D} is of the form $X \times \mathbb{D} \rightarrow \mathbb{D}$, with X a k -variety, then we are dealing with the space of n -jets (of curves) on X and the *arc space* of X , here denoted by $\mathcal{L}_n(X)$ resp. $\mathcal{L}(X)$.

For $m \geq n$ we have a forgetful morphism $\pi_n^m : \mathcal{X}_m \rightarrow \mathcal{X}_n$. (When $n = 0$, we shall often write π_X^m, π_X instead of π_0^m, π_0 .) A fiber of π_n^{n+1} lies in an affine space over the Zariski tangent space of the base point. In case X is smooth, it is in fact an affine space over the tangent space of the base point: π_n^{n+1} has then the structure of a torsor over the tangent bundle. A theorem of Greenberg [21] asserts that there exists a constant c such that the image of π_n equals the image of π_n^{cn} . So $\pi_n(\mathcal{X}_\infty)$ is constructible.

The goal is to define a measure on an interesting algebra of subsets of \mathcal{X}_∞ in such a way that its direct image under π_X is the tautological measure μ_X when X is smooth. (This will lead us to deviate from the definition of Denef-Loeser and Batyrev by a factor \mathbb{L}^d and to adopt the one used in [26] instead.) For this we assume that \mathcal{X} is of pure relative dimension d and we say that a subset A of \mathcal{X}_∞ is *stable* if for some $n \in \mathbb{N}$ we have

- $\pi_n(A)$ is constructible in \mathcal{X}_n and $A = \pi_n^{-1}\pi_n(A)$,
- for all $m \geq n$ the projection $\pi_{m+1}(A) \rightarrow \pi_m(A)$ is a piecewise trivial fibration (that is, trivial relative to a decomposition into subvarieties) with fiber an affine space of dimension d .

The second condition is of course superfluous in case \mathcal{X}/\mathbb{D} is smooth. It is clear that $\dim \pi_m(A) - md$ is independent of the choice of $m \geq n$; we call this the (*virtual*) *dimension* $\dim A$ of A . The same is true for the class $[\pi_m(A)]\mathbb{L}^{-md} \in M_k$; we denote that class by $\tilde{\mu}_{\mathcal{X}}(A)$. The collection of stable subsets of \mathcal{X} is a Boolean ring (i.e., is closed under finite union and difference) on which $\tilde{\mu}_{\mathcal{X}}$ defines a finite additive measure. A theorem of Denef-Loeser (see Theorem 9.1) ensures that there are plenty of stable sets.

In order to extend the measure to a bigger collection of interesting subsets of \mathcal{X}_∞ we need to complete M_k . Given $m \in \mathbb{Z}$, let $F_m M_k$ be the subgroup of M_k spanned by the $[Z]\mathbb{L}^{-r}$ with $\dim Z \leq m + r$. This is a filtration of M_k as a ring: $F_m M_k \cdot F_n M_k \subset F_{m+n} M_k$. So the separated completion of M_k with respect to this

filtration,

$$\hat{M}_k := \varprojlim M_k / F_m M_k \quad (m \rightarrow -\infty \text{ in this limit}),$$

to which we will refer as the *dimensional completion*, is also a ring. The kernel of the natural map $M_k \rightarrow \hat{M}_k$ is $\cap_m F_m M_k$, of course. It is not known whether this is zero³. In case $k \subset \mathbb{C}$, the Hodge characteristic extends to this completion:

$$\chi_h : \hat{M}_k \rightarrow \hat{K}_0(\text{HS}).$$

Here $\hat{K}_0(\text{HS})$ is defined in a similar way as \hat{M}_k with ‘dimension’ replaced by ‘weight’. The assertion follows from the fact that the weights in the compactly supported cohomology of a variety of dimension d are $\leq 2d$. Likewise we can extend the characteristics counting Hodge numbers or weight numbers (with values Laurent power series in the reciprocals of their variables). This does not apply to the Euler characteristic, but in many cases of interest the weight characteristic gives a rational function in w that has no pole at $w = 1$. Its value there is then a good substitute.

We will be mostly concerned with the composite of $\tilde{\mu}_{\mathcal{X}}$ and the completion map, for it is this measure that we shall extend. We call this the *motivic measure* on \mathcal{X} and denote it by $\mu_{\mathcal{X}}$. Let us say that a subset $A \subset \mathcal{X}_{\infty}$ is *measurable* if for every (negative) integer m there exist a stable subset $A_m \subset \mathcal{X}_{\infty}$ and a sequence $(C_i \subset \mathcal{X}_{\infty})_{i=0}^{\infty}$ of stable subsets such that the symmetric difference $A \Delta A_m$ is contained in $\cup_{i \in \mathbb{N}} C_i$ with $\dim C_i < m$ for all i and $\dim C_i \rightarrow -\infty$, for $i \rightarrow \infty$.

Proposition 2.2. *The measurable subsets of \mathcal{X}_{∞} make up a Boolean subring and $\mu_{\mathcal{X}}$ extends as a measure to this ring by*

$$\mu_{\mathcal{X}}(A) := \lim_{m \rightarrow -\infty} \mu_{\mathcal{X}}(A_m).$$

In particular, the above limit exists in \hat{M}_k and its value only depends on A .

The proof is based on

Lemma 2.3. *Let \mathcal{X}/\mathbb{D} be of pure dimension and $A \subset \mathcal{X}_{\infty}$ a stable subset. If $\mathcal{C} = \{C_i\}_{i=1}^{\infty}$ is a countable covering of A by stable subsets with $\dim C_i \rightarrow -\infty$ as $i \rightarrow \infty$, then A is covered by a finite subcollection of \mathcal{C} .*

Proof. Let $n \in \mathbb{N}$ be such that $A = \pi_n^{-1} \pi_n(A)$. Suppose that A is not covered by a finite subcollection of \mathcal{C} . Choose $k \in \mathbb{N}$ such that $\dim C_i < -(n+2)d$ for $i > k$ and let $u_{n+1} \in \pi_{n+1}(A \setminus \cup_{i \leq k} C_i)$. We have $\pi_{n+1}^{-1} u_{n+1} \subset A$. This set is not covered by a finite subcollection of \mathcal{C} , for clearly $\pi_{n+1}^{-1}(u_{n+1})$ is not covered by $\{C_i\}_{i \leq k}$ and for $i > k$, $C_i \cap \pi_{n+1}^{-1}(u)$ is of positive codimension in $\pi_{n+1}^{-1}(u)$.

With induction we find a sequence $\{u_m \in \mathcal{L}_m(X)\}_{m > n}$ so that for all $m > n$ u_{m+1} lies over u_m and $\pi^{-1}(u_m)$ is not covered by a finite subcollection of \mathcal{C} . The sequence defines an element $u \in \mathcal{X}$. Since $\pi_n(u) \in \pi_n(A)$, we have $u \in A$ and so $u \in C_i$ for some i . But if C_i is stable at level $m > n$, then $\pi_m^{-1}(u_m) \subset C_i$, which contradicts a defining property of u_m . \square

For $k = \mathbb{C}$, the condition $\lim_{i \rightarrow \infty} \dim C_i = -\infty$ is unnecessary, for we may then use the Baire property of \mathbb{C} instead [5].

³This issue is avoided if we work with the adic completion $\mathbb{Z}((L^{-1})) \otimes_{\mathbb{Z}[L]} K_0(\mathcal{V}_k)$ instead, but in practice this is too small. Nevertheless, it seems that in all applications we are dealing with elements lying in the localization $\mathbb{Q}(L) \otimes_{\mathbb{Z}[L]} K_0(\mathcal{V}_k)$.

Proof of 2.2. Suppose we have another solution $A\Delta A'_m \subset \cup_{i \in \mathbb{N}} C'_i$ with A'_m and C'_i stable, $\dim(C'_i) < m$ for all i and $\dim C'_i \rightarrow -\infty$ as $i \rightarrow \infty$. It is enough to prove that the dimension of the stable set $A_m\Delta A'_m$ is $< m$. Since $A_m\Delta A'_m \subset \cup_{i \in \mathbb{N}} (C_i \cup C'_i)$, Lemma 2.3 applies and we find that $A_m\Delta A'_m \subset \cup_{i \leq N} (C_i \cup C'_i)$ for some N . Since every term has dimension $< m$, this is also true for $A_m\Delta A'_m$. \square

So a countable union of stable sets $A = \cup_{n \in \mathbb{N}} A_n$ with $\lim_{n \rightarrow \infty} \dim A_n = -\infty$ is measurable and $\mu_{\mathcal{X}}(A) = \lim_{n \rightarrow \infty} \mu_{\mathcal{X}}(\cup_{k \leq n} A_k)$.

Remark 2.4. Given a \mathbb{D} -variety \mathcal{X} , then for any $d \in \mathbb{N}$ there is a d -measure $\mu_{\mathcal{X}}^d$ that induces $\mu_{\mathcal{Y}}$ on \mathcal{Y}_{∞} for any \mathbb{D} -subvariety \mathcal{Y} of pure dimension d . We expect this measure to extend to a much bigger collection of subsets of \mathcal{X} so that if $f : \mathcal{X} \rightarrow \mathcal{S}$ is a dominant \mathbb{D} -morphism of pure relative dimension d , then every fiber of $f_* : \mathcal{X}_{\infty} \rightarrow \mathcal{S}_{\infty}$ is $\mu_{\mathcal{X}}^d$ -measurable.

Here is a sample of the results of Denef and Loeser on the rationality of Poincaré series [14].

Theorem 2.5. *Let X be a k -variety. Then $\sum_{n=0}^{\infty} \mu_{\mathcal{X}}(\pi_n \mathcal{L}(X)) T^n \in M_k[[T]]$ is a rational expression in T with each factor in the denominator of the form $1 - \mathbb{L}^a T^b$ where $a \in \mathbb{Z}$ and b is a positive integer.*

We will not discuss its proof, since this theorem is not used in what follows. Denef and Loeser derive this by means of Kontsevich's transformation rule discussed below, which is applied to a suitable projective resolution \mathcal{X} , and a theorem about semialgebraic sets, due to Pas [25]. It is likely that this theorem still holds for the space of sections of any \mathbb{D} -variety.

3. THE TRANSFORMATION RULE [23], [14], [16]

We describe two results that are at the basis of the theory. The proofs are relegated to Section 9.

Proposition 3.1. *For a \mathbb{D} -variety \mathcal{X}/\mathbb{D} of pure dimension, the preimage of any constructible subset under $\pi_n : \mathcal{X}_{\infty} \rightarrow \mathcal{X}_n$ is measurable. In particular, \mathcal{X}_{∞} is measurable. If $\mathcal{Y} \subset \mathcal{X}$ is nowhere dense, then \mathcal{Y}_{∞} is of measure zero.*

For \mathcal{X}/\mathbb{D} of pure relative dimension we have the notion of an integrable function $\Phi : \mathcal{X}_{\infty} \rightarrow \hat{M}_k$: this requires the fibers of Φ to be measurable and the sum $\sum_a \mu_{\mathcal{X}}(\Phi^{-1}(a))a$ to converge, i.e., there are at most countably many nonzero terms $(\mu_{\mathcal{X}}(\Phi^{-1}(a_i))a_i)_{i \in \mathbb{N}}$ and we have $\mu_{\mathcal{X}}(\Phi^{-1}(a_i))a_i \in F_{m_i} \hat{M}_k$ with $\lim_{i \rightarrow \infty} m_i = -\infty$. The motivic integral of Φ is then by definition the value of this series:

$$\int \Phi d\mu_{\mathcal{X}} = \sum_i \mu_{\mathcal{X}}(\Phi^{-1}(a_i))a_i.$$

We have a similar notion for maps with values in topological \hat{M}_k -modules. An important example arises from an ideal $\mathcal{I} \subset \mathcal{O}_{\mathcal{X}}$: such an ideal defines a function $\text{ord}_{\mathcal{I}} : \mathcal{X}_{\infty} \rightarrow \mathbb{N} \cup \{\infty\}$ by assigning to $\gamma \in \mathcal{X}_{\infty}$ the multiplicity of $\gamma^* \mathcal{I}$. The condition $\text{ord}_{\mathcal{I}} \gamma = n$ only depends on the n -jet of γ and this defines a constructible subset $C_n \subset \mathcal{X}_n$. Hence the fibers of $\text{ord}_{\mathcal{I}}$ are measurable. We shall see that the function

$$\mathbb{L}^{-\text{ord}_{\mathcal{I}}} : \mathcal{X}_{\infty} \rightarrow \hat{M}_k$$

is integrable.

There is a beautiful transformation rule for motivic integrals under modifications. Let $H : \mathcal{Y} \rightarrow \mathcal{X}$ be a morphism of \mathbb{D} -varieties of pure dimension d . We define the *Jacobian ideal* $\mathcal{J}_H \subset \mathcal{O}_{\mathcal{Y}}$ of H as 0th Fitting ideal of $\Omega_{\mathcal{Y}/\mathcal{X}}$. This has the nice property that its formation commutes with base change. The following theorem generalizes an unpublished theorem of Kontsevich.

Theorem 3.2. *Let $H : \mathcal{Y} \rightarrow \mathcal{X}$ be a \mathbb{D} -morphism of pure dimensional \mathbb{D} -varieties with \mathcal{Y}/\mathbb{D} smooth. If A is a measurable subset of \mathcal{Y}_{∞} with $H|_A$ injective, then HA is measurable and $\mu_{\mathcal{X}}(HA) = \int_A \mathbb{L}^{-\text{ord}_{\mathcal{J}_H}} d\mu_{\mathcal{Y}}$.*

4. THE BASIC FORMULA [14]

A relative Grothendieck ring. It is convenient to be able to work in a relative setting. Given a variety S , denote by $K_0(\mathcal{V}_S)$ the Grothendieck ring of S -varieties and by M_S its localization with respect \mathbb{L} . The ring M_S can be dimensionally completed as usual. Notice that an element of M_S defines a M_k -valued measure on the Boolean algebra of constructible subsets of S . Often measures are naturally represented this way. For instance, the preceding shows that for all $n \in \mathbb{N}$, the direct image of $\mu_{\mathcal{X}}$ on \mathcal{X}_n is given by an element $\mu_{\mathcal{X},n} \in \hat{M}_{\mathcal{X}_n}$. (Notice that $\mu_{\mathcal{X},n}$ is then the direct image of $\mu_{\mathcal{X},n+1}$.)

A morphism $f : S' \rightarrow S$ induces a ring homomorphism $f^* : M_S \rightarrow M_{S'}$. This makes $M_{S'}$ a M_S -module. We also have a direct image $f_* : M_{S'} \rightarrow M_S$ that is a homomorphism of M_S -modules. Notice that f itself defines an element $[f] \in M_S$; this is also the image of $1 \in M_{S'}$ under f_* .

There are corresponding characteristics. For instance, the ordinary Euler characteristic χ_{top} becomes a ring homomorphism from M_S to the Grothendieck ring of constructible \mathbb{Q} -vector spaces on S . This ring is generated by direct images of irreducible local systems of \mathbb{Q} -vector spaces over smooth irreducible subvarieties Z of S . (A better choice is to take the intersection cohomology sheaf in S of this local system along Z ; this has the advantage that it only depends on the generic point of Z .)

Similarly, the Hodge characteristic χ_{h} takes values in a ring $K_0(\text{HS}_S)$ that is generated by variations of Hodge structures over a smooth subvariety of S . The homomorphisms f^* and f_* persist on this level: $f : S' \rightarrow S$ induces homomorphisms $f^* : K_0(\text{HS}_S) \rightarrow K_0(\text{HS}_{S'})$ and $f_* : K_0(\text{HS}_{S'}) \rightarrow K_0(\text{HS}_S)$.

The basic computation. A case of interest is when the base variety is $(\mathbb{N} \times \mathbb{G}_m)^r$. This fails to be finite type, but that is of no consequence and we identify $\hat{M}_{(\mathbb{N} \times \mathbb{G}_m)^r}$ with $\hat{M}_{\mathbb{G}_m^r}[[T_1, \dots, T_r]]$ in the obvious way.

We use a uniformizing parameter of \mathcal{O} to define

$$\text{ac} : \mathcal{L}(\mathbb{A}^1) - \{0\} \rightarrow \mathbb{N} \times \mathbb{G}_m,$$

by assigning to γ its order $\text{ord}(\gamma)$ resp. the first nonzero coefficient of γ (ac stands for *angular component*). Integration along ac sends a \hat{M}_k -valued measure on $\mathcal{L}(\mathbb{A}^1)$ to an element of $\hat{M}_{\mathbb{G}_m}[[T]]$. The prime example is when this measure is given by a regular function $f : \mathcal{X} \rightarrow \mathbb{A}^1$ on a \mathbb{D} -variety \mathcal{X} of pure relative dimension: this induces a map $f_* : \mathcal{X}_{\infty} \rightarrow \mathcal{L}(\mathbb{A}^1)$ and we then define

$$\text{ac}_f : \mathcal{X}_{\infty} \xrightarrow{f_*} \mathcal{L}(\mathbb{A}^1) \xrightarrow{\text{ac}} \mathbb{N} \times \mathbb{G}_m,$$

so that $[\text{ac}_f] \in \hat{M}_{\mathbb{G}_m}[[T]]$. More generally, given a morphism $f = (f_1, \dots, f_r) : \mathcal{X} \rightarrow \mathbb{A}^r$, we abbreviate

$$\text{ac}_{X,f} := (\pi_X, \text{ac}_{f_1}, \dots, \text{ac}_{f_r}) : \mathcal{X}_\infty \rightarrow X \times (\mathbb{N} \times \mathbb{G}_m)^r.$$

So $[\text{ac}_{X,f}] \in \hat{M}_{X \times \mathbb{G}_m^r}[[T_1, \dots, T_r]]$.

Conventions 4.1. If E is a simple normal crossing hypersurface on a smooth k -variety Y , then we adhere to the following notation throughout the talk: $(E_i)_{i \in \text{irr}(E)}$ denotes the collection of irreducible components of E (so these are all smooth by assumption) and for any subset $I \subset \text{irr}(E)$, E_I° stands for the locus of $p \in \tilde{X}$ with $p \in E_i$ if and only if $i \in I$. (With this convention, $E_\emptyset^\circ = Y - E$.) We denote the complement of the zero section of the normal bundle of E_i by U_{E_i} (so this is a \mathbb{G}_m -bundle over E_i) and U_I designates the fiber product of the bundles $U_{E_i}|_{E_I^\circ}$, $i \in I$ (a \mathbb{G}_m^I -bundle whose total space has the same dimension as Y).

If \mathcal{E} is a simple normal crossing hypersurface on a \mathbb{D} -variety \mathcal{Y}/\mathbb{D} with \mathcal{Y} smooth, then we shall always assume that its union with the closed fiber Y has also normal crossings. The notational conventions are as above to the extent that restriction or intersection with Y is indicated by switching from calligraphic to roman font (e.g., $E_i = \mathcal{E}_i \cap Y$). If Y is smooth, then we may identify $\text{irr}(E)$ with a subset of $\text{irr}(\mathcal{E})$. (An equality if \mathcal{E} has no component in Y .)

The following proposition accounts for many of the rationality assertions in [14].

Proposition 4.2. *Let \mathcal{X}/\mathbb{D} be a \mathbb{D} -variety of pure relative dimension and $H : \mathcal{Y} \rightarrow \mathcal{X}$ a resolution of singularities. Let \mathcal{E} be a simple normal crossing hypersurface on \mathcal{Y} that has no irreducible component in the closed fiber Y . Assume that the Jacobian ideal \mathcal{J}_H of H is principal and has divisor $\sum_i (\nu_i - 1)\mathcal{E}_i$ (so $\nu_i \geq 1$). Let for $\rho = 1, \dots, r$, $f_\rho : \mathcal{X} \rightarrow \mathbb{A}^1$ be a regular function such that $f_\rho H$ has zero divisor $\sum_i N_{i,\rho}\mathcal{E}_i$ and put $N_i := (N_{i,1}, \dots, N_{i,r}) \in \mathbb{N}^r$, $i \in \text{irr}(E)$. Then*

$$[\text{ac}_{X,f}] = \sum_{I \subset \text{irr}(E)} [U_I/X \times \mathbb{G}_m^r] \prod_{i \in I} (\mathbb{L}^{\nu_i} T^{-N_i} - 1)^{-1} \text{ in } \hat{M}_{X \times \mathbb{G}_m^r}[[T_1, \dots, T_r]],$$

where $U_I \rightarrow X \times \mathbb{G}_m^r$ has first component projection onto $E_I^\circ \subset X$ followed by the restriction of H and second component induced by fH .

Proof. Given $m \in \mathbb{N}^{\text{irr}(E)}$, consider the set $\mathcal{Y}(m)$ of $\gamma \in \mathcal{Y}_\infty$ with order m_i along \mathcal{E}_i . So for $\gamma \in \mathcal{Y}(m)$ we have $\text{ord}_{\mathcal{J}_H}(\gamma) = \sum_i m_i(\nu_i - 1)$ and $\text{ord}_{f_\rho H}(\gamma) = \sum_i m_i N_{\rho,i}$. If $\text{supp}(m) \subset \text{irr}(E)$ is the support of m , then we have a natural projection $e_m : \mathcal{Y}(m) \rightarrow U_{\text{supp}(m)}$. Its composite with the morphism $U_{\text{supp}(m)} \rightarrow X \times \mathbb{G}_m^r$ is a restriction of $\text{ac}_{X,fH} := (\pi_X H, \text{ac}_{f_1 H}, \dots, \text{ac}_{f_r H}) : \mathcal{Y}_\infty \rightarrow X \times (\mathbb{N} \times \mathbb{G}_m)^r$ with \mathbb{N}^r -component $\sum_i m_i N_i$. In other words,

$$[\text{ac}_{X,fH} |_{\mathcal{Y}(m)}] = [U_{\text{supp}(m)}/X \times \mathbb{G}_m^r] \mathbb{L}^{-\sum_i m_i} T^{\sum_i m_i N_i}.$$

So the transformation formula 3.2 yields

$$\begin{aligned} [\text{ac}_{X,f}] &= \sum_{m \in \mathbb{N}^{\text{irr}(E)}} [U_{\text{supp}(m)}/X \times \mathbb{G}_m^r] \prod_{i \in \text{supp}(m)} (\mathbb{L}^{-m_i - m_i(\nu_i - 1)} T^{m_i N_i}) \\ &= \sum_{I \subset \text{irr}(E)} [U_I/X \times \mathbb{G}_m^r] \prod_{i \in I} \frac{\mathbb{L}^{-\nu_i} T^{N_i}}{1 - \mathbb{L}^{-\nu_i} T^{N_i}}. \end{aligned}$$

□

If we drop the assumption that \mathcal{E} has no irreducible component in Y , then the above formula must be somewhat modified: now each irreducible component of Y contributes with an expression of the above form times a monomial in \mathbb{L}^{-1} and T_1, \dots, T_r .

Corollary 4.3. *In the situation of 4.2, the class of $(\pi_X, \text{ord}_f) : \mathcal{X}_\infty \rightarrow X \times \mathbb{N}^r$ in $\hat{M}_X[[T_1, \dots, T_r]]$ equals*

$$\sum_{I \subset \text{irr}(E)} [E_I^\circ/X] \prod_{i \in I} \frac{\mathbb{L} - 1}{\mathbb{L}^{\nu_i T^{-N_i}} - 1}.$$

In particular, the direct image of μ_X on X is represented by

$$\sum_{I \subset \text{irr}(E)} [E_I^\circ/X] \prod_{i \in I} [\mathbb{P}^{\nu_i - 1}]^{-1}.$$

Proof. Since U_I is a \mathbb{G}_m^I -bundle over E_I° , the class of the projection $U_I \rightarrow X$ is $(\mathbb{L} - 1)^{|I|}$ times the class of $E_I^\circ \rightarrow X$. \square

This corollary shows that \mathcal{X}_∞ is measurable so that the measurable subsets of \mathcal{X}_∞ form in fact a Boolean algebra. It also implies that the Hodge number characteristic of \mathcal{X}_∞ is an element of $\mathbb{Q}[u, v][[(uv)^N - 1]^{-1} \mid N = 1, 2, \dots]$ on which the Euler characteristic takes the value $\sum_{I \subset \text{irr}(E)} \chi_{\text{top}}(E_I^\circ) \prod_{i \in I} \nu_i^{-1}$.

Remark 4.4. We can also express the direct image of μ_X on X in terms of the closed subvarieties E_I : if $\text{irr}'(E)$ denotes the set of $i \in \text{irr}(E)$ with $\nu_i \geq 2$, then

$$\sum_{I \subset \text{irr}'(E)} (-\mathbb{L})^{|I|} [E_I/X] \prod_{i \in I} \frac{[\mathbb{P}^{\nu_i - 2}]}{[\mathbb{P}^{\nu_i - 1}]}.$$

All varieties appearing in this expression are proper over X and nonsingular. So it gives rise to an element of a complex cobordism ring of X localized away from the classes of the complex projective varieties. This class, and the values that various genera take on it, might deserve closer study.

5. THE MOTIVIC NEARBY FIBER [13], [18]

An equivariant Grothendieck ring. Let G be an affine algebraic group. We consider varieties X with *good* G -action, where ‘good’ means that every orbit is contained in an affine open subset. For instance, a representation of G on a k -vector space V is good. For a fixed variety S with G -action, we define the Grothendieck group $K_0^G(\mathcal{V}_S)$ as generated by isomorphism types of S -varieties with good G -action modulo the usual equivalence relation (defined by pairs) and the relation that declares that every finite dimensional representation ρ of G has the same class as the trivial representation of the same degree (i.e., $\mathbb{L}^{\deg(\rho)}$).

In case the action on S is trivial, the product makes $K_0^G(\mathcal{V}_S)$ a $K_0(\mathcal{V}_S)$ -algebra. If moreover G is finite abelian, then assigning to a variety X with good G -action its G -orbit space $\bar{X} := G \backslash X$ augments this as a $K_0(\mathcal{V}_S)$ -module:

$$K_0^G(\mathcal{V}_S) \rightarrow K_0(\mathcal{V}_S), \quad a \mapsto \bar{a}.$$

(Not as an algebra, for the orbit space of a product is in general not the product of orbit spaces.) That this is well-defined follows from the lemma below. (We do not know whether this holds for arbitrary finite G .)

Lemma 5.1. *Let be given a representation of a finite abelian group G on a k -vector space V of finite dimension n . Then the class of \overline{V} in $K_0(\mathcal{V}_k)$ is \mathbb{L}^n .*

Proof. Let $V = \bigoplus_{\chi \in \hat{G}} V_\chi$ be the eigenspace decomposition of the G -action. Given a subset $I \subset \hat{G}$, denote by V_I the set of vectors in V whose V_χ -component is nonzero if and only if $\chi \in I$. We have a natural projection $\overline{V_I} \rightarrow \prod_{\chi \in I} \mathbb{P}(V_\chi)$. This has the structure of a torus bundle, the torus in question being a quotient of \mathbb{G}_m^I by a finite subgroup. So the class of $\overline{V_I}$ in M_k is $(\mathbb{L} - 1)^{|I|}$ times the class of $\prod_{\chi \in I} \mathbb{P}(V_\chi)$. Since V_I has also that structure, the classes of $\overline{V_I}$ and V_I in M_k coincide. Hence the same is true for \overline{V} and V . \square

Similarly we can form $M_S^G := K_0^G(\mathcal{V}_S)[\mathbb{L}^{-1}]$ and its dimensional completion. The class of an S -variety Z/S with G -action in M_S^G or \hat{M}_S^G is denoted by $[Z/S; G]$. If G is abelian and acts trivially on S , then we have corresponding augmentations taking values in M_S and its completion.

There are corresponding characteristics in case $k \subset \mathbb{C}$. For instance, the ordinary Euler characteristic defines a ring homomorphism from M_k^G to the Grothendieck ring $K_0^G(\mathbb{Q})$ of finite dimensional representations of G over \mathbb{Q} and more generally, we have a ring homomorphism χ_{top}^G from M_S^G to the Grothendieck ring of constructible sheaves with G -action on S , $K_0^G(\mathbb{Q}_S)$. Similarly, there is a Hodge character $\chi_h^G : M_S^G \rightarrow K_0^G(\text{HS}_S)$.

The case $G = \hat{\mu}$. We will mostly (but not exclusively) be concerned with the case when G is a group of roots of unity. We have the Grothendieck ring $M_S^{\hat{\mu}}$ of varieties with a topological action of the procyclic group $\hat{\mu} = \varprojlim \mu_n$ (such an action factorizes through a finite quotient μ_n). The inverse automorphism of $\hat{\mu}$, $\zeta \mapsto \zeta^{-1}$, defines an involution $*$ in $M_S^{\hat{\mu}}$.

The group of continuous characters of $\hat{\mu}$ is naturally isomorphic with \mathbb{Q}/\mathbb{Z} , with the involution $*$ acting as multiplication by -1 ; the projection $\hat{\mu} \rightarrow \mu_n$ followed by the inclusion $\mu_n \subset \mathbb{G}_m$ corresponds to $\frac{1}{n} \pmod{\mathbb{Z}}$. In other words, $K_0^{\hat{\mu}}(\mathbb{C}) \cong \mathbb{Z}[e^\alpha \mid \alpha \in \mathbb{Q}/\mathbb{Z}]$. For every positive integer n there is a rational irreducible representation χ_n of μ_n , namely the field $\mathbb{Q}(\mu_n)$, regarded as \mathbb{Q} -vector space. These make up an additive basis of $K_0^{\hat{\mu}}(\mathbb{Q})$. The image of χ_n in $\mathbb{Z}[\mathbb{Q}/\mathbb{Z}]$ is $\sum_{(k,n)=1} e^{k/n}$, which allows us to regard $K_0^{\hat{\mu}}(\mathbb{Q})$ as a subring of $\mathbb{Z}[e^\alpha \mid \alpha \in \mathbb{Q}/\mathbb{Z}]$.

The so-called mapping torus construction gives rise to an M_k -linear map

$$M_k^{\hat{\mu}} \rightarrow M_{\mathbb{G}_m}$$

with the property that composition with the direct image homomorphism $M_{\mathbb{G}_m} \rightarrow M_k$ is $(\mathbb{L} - 1)$ times the augmentation $M_k^{\hat{\mu}} \rightarrow M_k$. It is defined as follows. If X is a variety with good μ_n , then its *mapping torus* is the étale locally trivial fibration $\mathbb{G}_m \times^{\mu_n} X \rightarrow \mathbb{G}_m$ whose total space is the orbit space of the μ_n -action on $\mathbb{G}_m \times X$ defined by $\zeta(\lambda, x) = (\lambda\zeta^{-1}, \zeta x)$ and for which the projection is induced by $(\lambda, x) \mapsto \lambda^n$. Notice that the fiber over $1 \in \mathbb{G}_m$ can be identified with X and that the monodromy is given by the action of μ_n on X . The projection on the second factor induces a morphism $\mathbb{G}_m \times^{\mu_n} X \rightarrow \mathbb{G}_m \rightarrow \overline{X}$ that has the structure of a piecewise \mathbb{G}_m -bundle. So the image of $\mathbb{G}_m \times^{\mu_n} X$ in M_k is $(\mathbb{L} - 1)[\overline{X}]$. If $m = kn$ is a positive multiple of n and we let μ_m act on X via $\mu_m \rightarrow \mu_n$, then $(\lambda, x) \mapsto (\lambda^k, x)$ identifies the two fibrations over \mathbb{G}_m and so we have a map as

asserted. This generalizes at once to the case where we have a base variety with trivial $\hat{\mu}$ -action.

Aut(\mathbb{D})-equivariance. The automorphism group $\text{Aut}(\mathbb{D})$ can be identified with the group of formal power series $k[[t]]$ with nonzero constant term where the group law is given by substitution. It acts on the arc space of any k -variety by composition: $h(\gamma) := \gamma h^{-1}$. If the variety is of pure dimension, then this action is free outside negligible subset. Clearly, a morphism of k -varieties induces an $\text{Aut}(\mathbb{D})$ -equivariant map between their arc spaces. Since we end up with more than just a $\text{Aut}(\mathbb{D})$ -invariant measure on an arc space, it is worthwhile to explicate this structure by means of a definition. If \mathbb{D}_n denotes the subscheme of \mathbb{D} defined by the ideal (t^{n+1}) , then $\text{Aut}(\mathbb{D}_n)$ (which has the same underlying variety as the group of units of $k[[t]]/(t^{n+1})$) acts naturally on $\mathcal{L}_n(X)$. For $n \geq 1$, the kernel of $\text{Aut}(\mathbb{D}_{n+1}) \rightarrow \text{Aut}(\mathbb{D}_n)$ can be identified with \mathbb{G}_a . Its action is trivial on $(\pi_1^{n+1})^{-1}(0)$ and free on the complement $(\pi_1^{n+1})^{-1}(TX - \{0\})$. By choosing a constructible section of the latter we lift the direct image homomorphism $(\pi_n^{n+1})_*$ to a map

$$M_{\mathcal{L}_{n+1}(X)}^{\text{Aut}(\mathbb{D}_{n+1})} \rightarrow M_{\mathcal{L}_n(X)}^{\text{Aut}(\mathbb{D}_n)}.$$

The result is easily seen to be independent of this choice.

Definition 5.2. An *equivariant motivic measure* on $\mathcal{L}(X)$ is a collection $\lambda = (\Lambda_n \in \hat{M}_{\mathcal{L}_n(X)}^{\text{Aut}(\mathbb{D}_n)})_{n=1}^\infty$, so that Λ_n is the direct image of Λ_{n+1} for all n .

It is clear that such a collection determines an \hat{M}_k -valued measure on the stable subsets. The definition is so devised that the measure $\mu_{\mathcal{L}(X)}$ constructed earlier comes from an equivariant motivic measure.

This notion is of particular interest when the variety in question is a smooth curve C and we are given a closed point $o \in C$. An $\text{Aut}(\mathbb{D})$ -orbit in $\mathcal{L}(C, o)$ is given by a positive integer n that may also take the value ∞ . If n is finite, then this orbit projects onto the set of nonzero elements of $(\pi_{n-1}^n)^{-1}(0) \cong T_{C,o}^{\otimes n}$. The group $\text{Aut}(\mathbb{D}_n)$ acts on the latter orbit through $\text{Aut}(\mathbb{D}_1) \cong \mathbb{G}_m$ with $\mu_n \subset \mathbb{G}_m$ as isotropy group. So the value of λ on a fiber over $T_{C,o}^{\otimes n} - \{0\}$ is naturally an element λ_n of $M_k^{\mu_n}$. We call the generating series

$$\lambda(T) := \sum_{n=1}^{\infty} \lambda_n T^n$$

the *zeta function* of λ . It is not hard to verify that this series determines λ completely. This is particularly so if we view λ as a \hat{M}_k -valued measure on $\mathcal{L}(C, o)$. For instance, its value on the preimage in $\mathcal{L}(C, o)$ of a constructible subset A of $\mathcal{L}_m(C, o)$ consisting of order n -arcs (with $n \leq m$) is $\mathbb{L}^{n-m}[A]\bar{\lambda}_n$. Notice that the series $\sum_{n=1}^{\infty} (\mathbb{L} - 1)\bar{\lambda}_n$ converges to the full integral of λ .

A motivic zeta function. Given a pure dimensional variety X and a flat morphism $X \rightarrow \mathbb{A}^1$, let $X_0 := f^{-1}(0)$ and denote by f the restriction $(X, X_0) \rightarrow (\mathbb{A}^1, 0)$. Then the direct image of $\mu_{\mathcal{L}(X, X_0)}$ (regarded as an equivariant measure) on $X_0 \times \mathcal{L}(\mathbb{A}^1, 0)$ is then also equivariant. We will (perhaps somewhat ambiguously) refer to this measure as the direct image of the motivic measure of $\mathcal{L}(X, X_0)$ on

$X_0 \times \mathcal{L}(\mathbb{A}^1, 0)$. Its zeta function is denoted by

$$S(f) = \sum_{n=1}^{\infty} S_n(f) T^n \in \hat{M}_{X_0}^{\hat{\mu}}[[T]].$$

We now assume that X is smooth and connected. The smoothness of X ensures that the preimage in $\mathcal{L}(X, X_0)$ of a stable subset of $\mathcal{L}(\mathbb{A}^1, 0)$ of level n is stable of level n , so that $S_n(f)$ already is defined as an element of M_{X_0} (but we shall not bring out the distinction in our notation). The series $S(f)$ can be computed from an embedded resolution of the zero set of f , $H : Y \rightarrow X$ of X , as in 4.1. We assume here that the preimage E of X_0 is a simple normal crossing hypersurface that contains the exceptional set. Let m be a positive integer that is divided by all the coefficients N_i of the divisor (f) on the irreducible components of E . If we make a base change of $\tilde{f} := fH$ over the m th power map $\mathbb{A}^1 \rightarrow \mathbb{A}^1$ and normalize, then we get a μ_m -covering $\tilde{Y} \rightarrow Y$. Let \tilde{E}_I° be a connected component of the preimage of E_I° in \tilde{Y} . The restriction $\tilde{E}_I^{\circ} \rightarrow E_I^{\circ}$ is unramified, and has μ_m -stabilizer of \tilde{E}_I° as its Galois group. The latter is easily seen to be the subgroup μ_{N_I} , where $N(I) := \gcd\{N_i \mid i \in I\}$. This defines

$$[\tilde{E}_I^{\circ}/Y; \mu_{N(I)}] \in M_Y^{\mu_{N(I)}}.$$

This element lies over X_0 if I is nonempty, an assumption we make from now on. We wish to compare it with $U_I(1) \subset U_I$, the fiber over 1 of the projection $U_I \rightarrow \mathbb{G}_m$ induced by \tilde{f} . This projection has weights $(N_i)_{i \in I}$ relative to the \mathbb{G}_m^I -action and so $\prod_{i \in I} \mu_{N_i} \subset \mathbb{G}_m^I$ preserves $U_I(1)$. This finite group contains a monodromy action by $\mu_{N(I)}$: write $N(I) = \sum_{i \in I} \alpha_i N_i$ and embed \mathbb{G}_m in \mathbb{G}_m^I by $t \mapsto (t^{\alpha_i})_{i \in I}$ (since the $(\alpha_i)_{i \in I}$ are relatively prime, this is an embedding indeed). Notice that the projection $U_I \rightarrow \mathbb{G}_m$ is homogeneous of degree N_I relative to the action of this one parameter subgroup. This implies that $\mu_{N(I)} \subset \mathbb{G}_m$ may serve as monodromy group. (There are a priori several choices for this action, but they are all E_I° -isomorphic.)

Lemma 5.3. *In $M_{Y_0}^{\hat{\mu}}$ we have $[U_I(1)/Y_0; \mu_{N(I)}] = (\mathbb{L} - 1)^{|I|-1} [\tilde{E}_I^{\circ}/Y_0; \mu_{N(I)}]$.*

Proof. One verifies that the Stein factorization of the projection $U_I(1) \rightarrow E_I^{\circ}$ has $\tilde{E}_I^{\circ} \rightarrow E_I^{\circ}$ as finite factor with $U_I(1) \rightarrow \tilde{E}_I^{\circ}$ being an algebraic torus bundle of rank $|I| - 1$. In view of Lemma 5.1 the equivariant class of the latter is $(\mathbb{L} - 1)^{|I|-1}$ times the equivariant class of the base. The lemma follows. \square

Much of the work of Denef-Loeser on motivic integration centers around the following

Theorem 5.4. *The following identity holds in $M_{X_0}^{\hat{\mu}}[[T]]$:*

$$S(f) = \sum_{\emptyset \neq I \subset \text{irr}(E)} (\mathbb{L} - 1)^{|I|-1} [\tilde{E}_I^{\circ}/X_0; \mu_{N(I)}] \prod_{i \in I} (\mathbb{L}^{\nu_i} T^{-N_i} - 1)^{-1}.$$

Proof. Start with the identity of Proposition 4.2 (with $r = 1$). Omit at both sides the constant terms (on the right this amounts to summing over nonempty I only), and restrict the resulting identity to the fiber over $1 \in \mathbb{G}_m$. If we take into account the monodromies and use Lemma 5.3, we get the asserted identity, at least if we take our coefficients in $M_{X_0}^{\hat{\mu}}$. Inspection of the proof shows that this actually holds in $M_{X_0}^{\hat{\mu}}[[T]]$. \square

So the expression at the righthand side is independent of the resolution, something that is not at all evident a priori. Since it lies in the $M_{X_0}^{\hat{\mu}}$ -subalgebra of $M_{X_0}^{\hat{\mu}}[[T]]$ generated by the fractions $(L^\nu T^{-N} - 1)^{-1}$ with $\nu, N > 0$, $S(f)$ has a value at $T = \infty$:

$$S(f)|_{T=\infty} = - \sum_{\emptyset \neq I \subset \text{irr}(E)} (1 - \mathbb{L})^{|I|-1} [\tilde{E}_I^\circ / X_0; \mu_{N(I)}]$$

Comparison with ordinary monodromy. The element $-S(f)|_{T=\infty}$ has an interpretation in terms of the nearby cycle sheaf of f as we shall now explain.

Suppose first that $k = \mathbb{C}$. Let $\widetilde{X - X_0} \rightarrow X - X_0 \subset X$ be the pull-back along f of the universal covering $\exp : \mathbb{C} \rightarrow \mathbb{C}^\times \subset \mathbb{C}$. Take the full direct image of the constant sheaf $\mathbb{Q}_{\widetilde{X - X_0}}$ on X and restrict to X_0 : this defines ψ_f as an element of the derived category of constructible sheaves on X_0 . Let $\sigma : \widetilde{X - X_0} \rightarrow \widetilde{X - X_0}$ be a generator of the covering transformation that induces in \mathbb{C} translation over $-2\pi\sqrt{-1}$. This generator has the property that its action in ψ_f is the monodromy.

Let $H : Y \rightarrow X$ be a resolution as in 4.1. In the same way, $\psi_{\tilde{f}}$ is defined as an element of the derived category of constructible sheaves on the zero set Y_0 of \tilde{f} . The full direct image of $\psi_{\tilde{f}}$ on X_0 is equal to ψ_f .

An elementary calculation shows that the stalk of $\psi_{\tilde{f}}$ at a point of E_I° is the cohomology of N_I copies of a real torus of dimension $N_I - 1$. More precisely, the restriction of $\psi_{\tilde{f}}$ to E_I° is naturally representable as the full direct image of the constant sheaf on $U_I(1)$ (an algebraic torus bundle of dimension $N_I - 1$ over \tilde{E}_I°) under the projection $U_I(1) \rightarrow E_I^\circ$. We have a canonical isomorphism $H^k(\mathbb{G}_m^r; \mathbb{Q}) \cong H_c^{k+r}(\mathbb{G}_m^r; \mathbb{Q})$ and hence the Euler characteristic $\sum_k (-1)^k [H^k(\mathbb{G}_m^r; \mathbb{Q})]$ in $K_0(HS)$ is $(-1)^r$ times the Euler characteristic $\sum_k (-1)^k [H_c^k(\mathbb{G}_m^r; \mathbb{Q})]$. In other words, it is the value of χ_h on $(1 - \mathbb{L})^r$. Hence, if Z is a subvariety of E_I° with preimage \tilde{Z} in \tilde{E}_I° , then $\sum_k (-1)^k [H_c^k(Z; \psi_{\tilde{f}})]$ is the value of χ_h on $(1 - \mathbb{L})^{|I|-1} [\tilde{Z}; \mu_{N(I)}]$. This shows that ψ_f and $-S(f)|_{T=\infty}$ have the same Hodge characteristic. We therefore put

$$[\psi_f] := -S(f)|_{T=\infty} = \sum_{\emptyset \neq I \subset \text{irr}(E)} (1 - \mathbb{L})^{|I|-1} [\tilde{E}_I^\circ / X_0; \mu_{N(I)}].$$

We refer to $[\psi_f]$ as the *nearby cycle class* of f along X_0 . Its component in the augmentation submodule,

$$[\phi_f] := [\psi_f] - \overline{[\psi_f]} \in M_{X_0}^{\hat{\mu}},$$

is by definition the *vanishing cycle class* of f .

Let S be a variety with trivial $\hat{\mu}$ -action. Given a S -variety Z with a good topological $\hat{\mu}$ -action, then for any positive integer n the fixed point locus of $\ker(\hat{\mu} \rightarrow \mu_n)$ in Z is a S -variety which inherits a good μ_n -action. This defines a homomorphism of M_S -algebras

$$\text{Tr}_n : M_S^{\hat{\mu}} \rightarrow M_S^{\mu_n}.$$

If $\sigma \in \hat{\mu}$ generates a dense subgroup of $\hat{\mu}$, then the fixed point locus of $\ker(\hat{\mu} \rightarrow \mu_n)$ is also the fixed point locus of σ^n . In case $k \subset \mathbb{C}$, a Lefschetz fixed point formula (applied to a partition of Z by orbit type) implies that $\chi_h[Z^{\sigma^n}]$ equals the trace of

σ^n in $\chi_h[Z]$. So we may then think of $Tr_n[Z]$ as the motivic trace of σ^n . This is why the following proposition is a motivic version of a result of A'Campo [2].

Proposition 5.5 (see Denef-Loeser [18]). *The series $S(f)$ and $\sum_{n=1}^{\infty} Tr_n[\psi_f]T^n$ in $M_{X_0}^{\mu_n}[[T]]$ are congruent modulo $\mathbb{L} - 1$.*

Proof. The monodromy σ acts on \tilde{E}_I° as a covering transformation of order N_I . So σ^n has no fixed point if N_I does not divide n and is equal to all of \tilde{E}_I° otherwise. It follows from formula for the nearby cycle class that

$$Tr_n[\psi_f] = \sum_{I \subset \text{irr}(E), N_I | n} (1 - \mathbb{L})^{|I|-1} [\tilde{E}_I^\circ / X_0].$$

So

$$\begin{aligned} \sum_{n=1}^{\infty} Tr_n[\psi_f]T^n &= \sum_{n=1}^{\infty} \sum_{I \subset \text{irr}(E), N_I | n} (1 - \mathbb{L})^{|I|-1} [\tilde{E}_I^\circ / X_0] T^n \\ &= \sum_{\emptyset \neq I \subset \text{irr}(E)} (1 - \mathbb{L})^{|I|-1} [\tilde{E}_I^\circ / X_0] \sum_{k \geq 1} T^{kN_I} \\ &= \sum_{\emptyset \neq I \subset \text{irr}(E)} (1 - \mathbb{L})^{|I|-1} [\tilde{E}_I^\circ / X_0] \frac{T^{N_I}}{1 - T^{N_I}}. \end{aligned}$$

If we reduce modulo $(\mathbb{L} - 1)$ only the terms with I a singleton remain. Theorem 5.4 shows that this has the same reduction modulo $(\mathbb{L} - 1)$ as $S(f)$. \square

6. THE MOTIVIC ZETA FUNCTION OF DENEFF-LOESER [13]

This function is a motivic analogue of Igusa's local zeta function. It captures slightly less than the function $S(f)$, but has the virtue that it is defined in greater generality. First we introduce two homomorphisms of Grothendieck rings.

An arrow $M_S^{\mu_{rn}} \rightarrow M_S^{\mu_n}$ is defined by assigning to a variety with good μ_{rn} -action its orbit space with respect to the subgroup $\mu_r \subset \mu_{rn}$ (with a residual action of μ_n). The totality of these arrows forms a projective system whose limit we denote by $M_S(\hat{\mu})$. This is not the same as $M_S^{\hat{\mu}}$, but there is certainly a natural ring homomorphism

$$\rho : M_S^{\hat{\mu}} \rightarrow M_S(\hat{\mu}).$$

It is given by assigning to a variety X with good $\hat{\mu}$ -action, the system $(X_n)_n$, where X_n is the orbit space of X by the kernel of $\hat{\mu} \rightarrow \mu_n$ endowed with the residual action of μ_n .

We next define the *Kummer map*

$$M_{S \times \mathbb{G}_m} \rightarrow M_S(\hat{\mu}), \quad [f] \mapsto [f]^{1/\infty}.$$

Given a S -variety Y and a morphism $f : Y \rightarrow \mathbb{G}_m$, then for every positive integer n , let $f^{1/n} : Y(f^{1/n}) \rightarrow \mathbb{G}_m$ be the pull-back of f over the n th power map $[n] : \mathbb{G}_m \rightarrow \mathbb{G}_m$. So $Y(f^{1/n})$ is the hypersurface in $\mathbb{G}_m \times Y$ defined by $f(z) = u^n$. The projection of $Y(f^{1/n}) \rightarrow Y$ is a μ_n -covering and thus defines an element $[f]^{1/n}$ of $M_S^{\mu_n}$. Notice that $Y(f^{1/n})$ is the orbit space of $Y(f^{1/nr})$ relative to the subgroup $\mu_r \subset \mu_{rn}$. Hence the $[f]^{1/n}$'s define an element $[f]^{1/\infty} \in M_S(\hat{\mu})$.

The following lemma is a straightforward exercise.

Lemma 6.1. *The composition of the mapping torus construction and the Kummer map is equal to $(\mathbb{L} - 1)\rho$.*

For \mathcal{X} a smooth \mathbb{D} -variety of pure relative dimension d , define the *Denef-Loeser zeta function* by

$$I(f) := \mathbb{L}^{-d} \sum_{n=0}^{\infty} [\text{ac}_{f,n}]^{1/\infty} \mathbb{L}^{-sn} \in M_X(\hat{\mu})[[\mathbb{L}^{-s}]],$$

where \mathbb{L}^{-s} is just a variable with a suggestive notation. Then Corollary 4.3 and Lemma 6.1 yield

Theorem 6.2. *The following identity holds in $M_X(\hat{\mu})[[\mathbb{L}^{-s}]]$:*

$$I(f) = \mathbb{L}^{-d} \sum_{I \subset \text{irr}(E)} \rho[\tilde{E}_I^\circ/X; \mu_{N(I)}] \prod_{i \in I} \frac{\mathbb{L} - 1}{\mathbb{L}^{\nu_i + sN_i} - 1}.$$

Putting $\mathbb{L} = 1$. Consider the $\mathbb{Z}[L, L^{-1}]$ -subalgebra S of $\mathbb{Q}(L, L^{-s})$ generated by the rational functions $(L - 1)(L^{n+sN} - 1)^{-1}$, $n, N \geq 1$. The spectrum of S contains the generic point of the exceptional divisor of the blow up of $(1, 1)$ in $\mathbb{G}_m \times \mathbb{A}^1$. The corresponding specialization is the evaluation homomorphism $S \rightarrow \mathbb{Q}(s)$ which sends $(L - 1)(L^{n+sN} - 1)^{-1}$ to $(n + sN)^{-1}$. According to Theorem 6.2, $I(f)$ lies in $S \otimes_{\mathbb{Z}[L, L^{-1}]} M_X(\hat{\mu})$. Evaluation at $\mathbb{L} = 1$ yields

$$I(f)|_{\mathbb{L}=1} = \sum_{I \subset \text{irr}(D)} \rho[\tilde{E}_I^\circ/X; \mu_{N(I)}] \prod_{i \in I} \frac{1}{\nu_i + sN_i} \in M_X(\hat{\mu})/(\mathbb{L} - 1) \otimes_{\mathbb{Z}} \mathbb{Q}(s).$$

This is the motivic incarnation of the topological zeta function considered earlier by Denef and Loeser in [12]. At the time the resolution independence of this function was established using Theorem 6.3 below.

Comparison with Igusa's p -adic zeta function. Suppose we are given a complete discrete valuation ring (R, m) of characteristic zero whose residue field $F = R/m$ has finite cardinality q . Then R contains all the $(q - 1)$ st roots of unity μ_{q-1} and this group projects isomorphically onto F^\times . Let K be the quotient field of R . If we choose a uniformizing parameter $\pi \in m - m^2$, then the collection $(\zeta\pi^k)_{\zeta \in \mu_{q-1}, k \in \mathbb{Z}}$ is a system of representatives of $K^\times/(1 + m)$. Define

$$\text{ac}^s : K \rightarrow \mathbb{Z}[\mu_{q-1}][q^{-s}]$$

by assigning to $u \in \zeta\pi^k + m^{k+1}$ the value ζq^{-ks} and 0 to 0. (Here q^{-s} is just the name of a variable; the righthand side can be more canonically understood as the group algebra of $K^\times/(1 + m)$.) There is a natural (additive) Haar measure μ on the the Boolean ring of subsets of K generated by the cosets of powers of m that takes the value 1 on R . It takes values in $\mathbb{Z}[q^{-1}]$. Given an $f \in R[x_1, \dots, x_d]$ whose reduction mod m is nonzero, then its *Igusa local zeta function* is defined by

$$Z(f) := \int_{R^m} \text{ac}^s f(x) d\mu(x),$$

where R^m is endowed with the product measure. We regard this as an element of $\mathbb{Q}[\mu_{q-1}][[q^{-s}]]$: the coefficient of ζq^{-ns} is the volume of $f^{-1}(\zeta\pi^n + m^{n+1})$. (It is customary to let s be a complex number—the series then converges in a right half plane—and to compose with a complex character $\mu_{q-1} \rightarrow \mathbb{C}^\times$.)

Let us write \mathcal{X} for $\text{Spec}(R[x_1, \dots, x_d])$ and regard f as a morphism $\mathcal{X} \rightarrow \mathbb{A}_R^1$ over $\text{Spec}(R)$. Suppose we have an embedded resolution $H : \mathcal{Y} \rightarrow \mathcal{X}$ of the zero locus of f over $\text{Spec}(R)$ with a simple normal crossing hypersurface \mathcal{E} relative to $\text{Spec}(R)$ (so no irreducible component in the closed fiber). Then we get an embedded resolution of the closed fiber $Y \rightarrow X$ with simple normal crossing divisor E . Make a base change of $fH : \mathcal{Y} \rightarrow \mathbb{A}_R^1$ over the $(q-1)$ st power map $[q-1] : \mathbb{A}_R^1 \rightarrow \mathbb{A}_R^1$ and normalize; this gives a μ_{q-1} -covering $\tilde{\mathcal{Y}} \rightarrow \mathcal{Y}$. We now get a covering $\hat{E}_I^\circ \rightarrow E_I^\circ$ defined over F with Galois group $\mu_{N_q(I)}$, where $N_q(I) := \gcd(q-1, (N_i)_{i \in I})$ over F in much the same way as before. The $\mu_{N_q(I)}$ -set $\hat{E}_I^\circ(F)$ determines an element

$$\#[\hat{E}_I^\circ; \mu_{q-1}] \in \mathbb{Q}[\mu_{N_q(I)}] \subset \mathbb{Q}[\mu_{q-1}],$$

where the last inclusion is defined by the surjection $\mu_{q-1} \rightarrow \mu_{N_q(I)}$. Denef proved earlier [10] the following analogue of 6.2:

Theorem 6.3 (Denef). *In this situation we have*

$$Z(f) = q^{-d} \sum_{I \subset \text{irr}(E)} \#[\hat{E}_I^\circ; \mu_{q-1}] \prod_{i \in I} \frac{q-1}{q^{\nu_i + s N_i} - 1},$$

where ν_i and N_i have the usual meaning.

As appears from 6.2, $Z(f)$ is what we get from the value of $I(f_{\bar{K}})$ on $X_0(\bar{K})$ (with \bar{K} an algebraic closure of K) if we replace classes in $M_{\bar{K}}$ by the number of F -rational points in their F -counterparts (so that we substitute q for \mathbb{L}) and pass from $\hat{\mu}$ to μ_{q-1} . This should be understood on a more conceptual level that involves a Grothendieck ring $M_{\text{Spec}(R)}^{\mu_{q-1}}$ which specializes to both $M_{\text{Spec}(\bar{K})}^{\mu_{q-1}}$ and $\mathbb{Q}[\mu_{q-1}][[q^{-s}]]$, and avoids resolution.

7. MOTIVIC CONVOLUTION [15]

Join and quasi-convolution. Consider the Fermat curve J_n in \mathbb{G}_m^2 defined by $u^n + v^n = 1$. Notice that it is invariant under the subgroup $\mu_n^2 \subset \mathbb{G}_m^2$. If d is a positive divisor of n , then the μ_d^2 -orbit space of J_n is $J_{n/d}$. In particular, the μ_n^2 -orbit space of J_n is J_1 , an affine line less two points. Given varieties X and Y with good μ_n -action, then we have the variety with $\mu_n \times \mu_n$ -action

$$J_n(X, Y) := J_n \times^{(\mu_n \times \mu_n)} (X \times Y).$$

(If a group G acts well on varieties A and B , then $A \times^G B$ stands for quotient of $A \times B$ by the equivalence relation $(ga, b) \sim (a, gb)$ with G acting well on it by $g[a, b] := [ga, b] = [a, gb]$.) Let μ_n act on $J_n(X, Y)$ diagonally: $\zeta[(u, v), (x, y)] := [(\zeta u, \zeta v), (x, y)]$. The natural map $J_n(X, Y) \rightarrow J_1$ is étale locally trivial. If Y has trivial μ_n -action, then $J_n(X, Y) = J_n(X, pt) \times Y$ and the variety $J_n(X, pt)$ can be identified with $(\mathbb{G}_m - \{\mu_n\}) \times^{\mu_n} X$. The latter has the structure of a piecewise \mathbb{G}_m -bundle over \overline{X} from which a copy of X has been removed. Similarly, the natural projection of $\overline{J_n(X, Y)} \rightarrow \overline{X} \times \overline{Y}$ is a piecewise \mathbb{G}_m -bundle from which a copy of $X \times Y$ has been removed.

The construction is perhaps better understood in terms of the fibrations over \mathbb{G}_m defined by the mapping torus construction. Recall that for a variety X with μ_n -action, its mapping torus $\mathbb{G}_m \times^{\mu_n} X$ fibers over \mathbb{G}_m by $[\lambda, x] \mapsto \lambda^n$ with $\{1\} \times X$

mapping to the fiber over 1. The monodromy is the given μ_n -action on X . If Y is another variety with μ_n -action, then the composite

$$(\mathbb{G}_m \times^{\mu_n} X) \times (\mathbb{G}_m \times^{\mu_n} Y) \longrightarrow \mathbb{G}_m \times \mathbb{G}_m \subset \mathbb{G}_a \times \mathbb{G}_a \xrightarrow{+} \mathbb{G}_a$$

is a fibration over \mathbb{G}_m . The fiber over $1 \in \mathbb{G}_a$ is identified as $J_n(X, Y)$ and the monodromy is the given μ_n -action on $J_n(X, Y)$ defined above.

Clearly, $J_n(X, Y) \cong J_n(Y, X)$. If m is a divisor of n and the action of μ_n on X and Y is through μ_m , then $J_m(X, Y) = J_n(X, Y)$. So this induces a binary operation, the *join*

$$J : M_k^{\hat{\mu}} \times M_k^{\hat{\mu}} \rightarrow M_k^{\hat{\mu}}.$$

The preceding discussion shows that the join is commutative and bilinear over M_k and that (i) $J(a, 1) = (\mathbb{L} - 1)\bar{a} - a$ and (ii) $\overline{J(a, b)} = (\mathbb{L} - 1)\bar{a}\bar{b} - \overline{ab}$, where we recall that $a \in M_k^{\hat{\mu}} \mapsto \bar{a} \in M_k$ is the augmentation defined by ‘passing to the orbit space’. This suggests to define another binary operation $*$, the *quasi-convolution*, on $M_k^{\hat{\mu}}$ by:

$$a * B := -J(a, b) + (\mathbb{L} - 1)\overline{ab}.$$

The quasi-convolution is commutative and bilinear over M_k , whereas the properties (i) and (ii) come down to

- (i) 1 is a unit for $*$: $a * 1 = a$ (and hence $a * \bar{b} = \overline{a\bar{b}}$) and
- (ii) $\overline{a * b} = \overline{ab}$.

Neither the join nor the quasi-convolution is associative, but we do have:

- (iii) $a * (b * c) - (\mathbb{L} - 1)\overline{a(b * c)} + (\mathbb{L} - 1)^2\overline{a\bar{b}\bar{c}}$ is symmetric in a, b and c ,

which shows that the quasi-convolution is associative modulo elements of M_k . This property is seen as follows. Let J_n^2 denotes the Fermat surface in \mathbb{G}_m^3 defined by $u^n + v^n + w^n = 1$ and consider the morphism

$$J_n \times J_n \rightarrow J_n^{(2)}, \quad ((u_1, v_1), (u_2, v_2)) \mapsto (u, v, w) = (u_1, v_1 u_1, v_1 u_2).$$

This morphism is equivariant with respect to the action of μ_n on $J_n \times J_n$ that is diagonal on the first factor and trivial on the second and the diagonal action μ_n on $J_n^{(2)}$. It also factorizes over the orbit space of $J_n \times J_n$ with respect to the μ_n action defined by $\zeta((u_1, v_1), (u_2, v_2)) = ((u_1, \zeta^{-1}v_1), (\zeta u_2, \zeta v_2))$. One easily verifies that this identifies the orbit space for this action with in $J_n^{(2)} - K_n$, where $K_n \subset J_n^{(2)}$ is defined by $u^n = 1$. A choice of an n th root α of -1 , identifies K_n with $\mu_n \times \mathbb{G}_m \times \mu_n$ via $(u, v, w) \mapsto (u, v, \alpha w/v)$. The μ_n^3 -action on K_n carries in an obvious manner to $\mu_n \times \mathbb{G}_m \times \mu_n$.

It follows from these observations that if X, Y, Z are varieties with good μ_n -action, then $J_n^{(2)} \times^{\mu_n^3} X \times Y \times Z$ decomposes as a μ_n -variety into two pieces that can be identified with $J_n(X, J_n(Y, Z))$ and $X \times (\mathbb{G}_m \times^{\mu_n} (Y \times Z))$ respectively. The factor $\mathbb{G}_m \times^{\mu_n} (Y \times Z)$ has the structure of a \mathbb{G}_m -bundle over $\bar{Y} \times \bar{Z}$. Passing now to $M_k^{\hat{\mu}}$ we find that

$$J(a, J(b, c)) + (\mathbb{L} - 1)\overline{abc}$$

is symmetric in a, b, c and this is equivalent to property (iii) above.

Join and quasi-convolution extend to $\hat{M}_k^{\hat{\mu}}$ and admit relative variants.

Formation of the spectrum. Join and quasi-convolution also descend to the Grothendieck ring $K_0^{\hat{\mu}}(HS)$ of Hodge structures with $\hat{\mu}$ -action. We need:

Lemma 7.1 (Shioda-Katsura, [27]). *Given $(\alpha, \beta) \in (\mathbb{Q}/\mathbb{Z})^2$, then for every common denominator n of α and β , the Hodge type of the eigenspace $I_{\alpha, \beta}$ of $\mu_n \times \mu_n$ in $H_c^1(J_n)$ with character $(\alpha, \beta) \in (n^{-1}\mathbb{Z}/\mathbb{Z})^2$ is independent of n and we have $\dim I_{\alpha, \beta} = 1$ for $(\alpha, \beta) \neq (0, 0)$ and $\dim I_{0,0} = 2$. If $\alpha \in \mathbb{Q}/\mathbb{Z} \mapsto \tilde{\alpha} \in [0, 1[$ is the obvious section, then*

$$I_{\alpha, \beta} \text{ is of Hodge type } \begin{cases} (0, 1) & \text{if } \alpha \neq 0 \neq \beta \text{ and } 0 < \tilde{\alpha} + \tilde{\beta} < 1, \\ (1, 0) & \text{if } 1 < \tilde{\alpha} + \tilde{\beta} < 2 \text{ and} \\ (0, 0) & \text{otherwise: } \alpha = 0 \text{ or } \beta = 0 \text{ or } \alpha + \beta = 0. \end{cases}$$

The only other nonzero group is $H_c^2(J_n)$, which is isomorphic to $\mathbb{Q}(-1)$ and has trivial character $(0, 0)$.

Corollary 7.2. *If $H, H' \in K_0^{\hat{\mu}}(HS)$, then*

$$H * H' = H_0 \otimes H'_0 + \sum_{\alpha \neq 0} H_{\alpha} \otimes H'_{-\alpha}(-1) + \sum_{\alpha + \beta \neq 0} H_{\alpha} \otimes H'_{\beta} \otimes I_{\alpha, \beta}.$$

Anderson [3] investigated Hodge structures with $\hat{\mu}$ -action using a notion of a *fractional Hodge structure*. For us such a structure will consist of a complex vector space V defined over \mathbb{Q} with a complex decomposition $V = \bigoplus_{p, q \in \mathbb{Q}: p+q \in \mathbb{Z}} V^{p, q}$ such that $V^{q, p}$ is the complex conjugate of $V^{p, q}$ and $\bigoplus_{p+q=n} V^{p, q}$ is defined over \mathbb{Q} for every $n \in \mathbb{Z}$. They form an abelian category $HS(\mathbb{Q})$ with tensor product. Anderson associates to a Hodge structure H with $\hat{\mu}$ -action a fractional Hodge structure $\sigma(H)$ whose underlying vector space is H , leaves the bidegrees on H_0 unaltered and increases the bidegrees of H_{α} by $(\tilde{\alpha}, 1 - \tilde{\alpha})$ if $\alpha \neq 0$. We shall refer to this operation as the *formation of the spectrum*. It defines an additive functor and hence a homomorphism of groups $\text{sp} : K_0^{\hat{\mu}}(HS) \rightarrow K_0(HS(\mathbb{Q}))$. This is not a ring homomorphism, but Corollary 7.2 shows that sp takes quasi-convolution to the tensor product:

$$\text{sp}(H * H') = \text{sp}(H) \otimes \text{sp}(H').$$

Convolution. In what follows we need the (additive) group structure on the affine line, so we write \mathbb{G}_a instead of \mathbb{A}^1 . We have a bijection $\mathcal{L}(\mathbb{G}_a, 0) \cong \mathfrak{m}$, defined by assigning to $\gamma \in \mathcal{L}(\mathbb{G}_a, 0)$ the pull-back of the standard coordinate on \mathbb{G}_a .

Let $\lambda = (\Lambda_n)_n$ and $\lambda' = (\Lambda'_n)_n$ be equivariant measures on $\mathcal{L}(\mathbb{G}_a, 0)$. Then $\lambda \times \lambda' := (\Lambda_n \times \Lambda'_n)_{n=1}^{\infty}$ defines a measure on the algebra of stable subsets of $\mathcal{L}(\mathbb{G}_a, 0)^2$ (that is, preimages of constructible subset of some truncation $\mathcal{L}_n(\mathbb{G}_a, 0)^2$). For instance, if $C \subset \mathcal{L}_n(\mathbb{G}_a, 0)^2$ is constructible and consists of pairs of truncated arcs of fixed order (k, l) (with $k, l \leq n$), then the value of $\lambda \times \lambda'$ on the preimage of C in $\mathcal{L}(\mathbb{G}_a, 0)^2$ is $\lambda_k \lambda'_l [C] \mathbb{L}^{-2n}$.

The direct image of $\lambda \times \lambda'$ under the addition morphism $\text{add} : \mathbb{G}_a \times \mathbb{G}_a \rightarrow \mathbb{G}_a$, $\lambda * \lambda' := (\mathcal{L}_n(\text{add})(\Lambda_n \times \Lambda'_n))_{n=1}^{\infty}$, is an equivariant measure on $\mathcal{L}(\mathbb{G}_a, 0)$, called the *convolution* of λ and λ' .

Lemma 7.3. *The zeta function of $\lambda * \lambda'$ is determined by those of λ and λ' :*

$$(\lambda * \lambda')_n = -(\lambda_n * \lambda'_n) + (\mathbb{L} - 1) \sum_{i \leq n} \mathbb{L}^{i-n} \overline{\lambda_i \lambda'_i} + (\mathbb{L} - 1) \sum_{i > n} (\lambda_n \overline{\lambda'_i} + \overline{\lambda_i} \lambda'_n).$$

Proof. The preimage of $t^n + \mathfrak{m}^{n+1}$ in $\mathfrak{m} \times \mathfrak{m}$ under $\mathcal{L}(\text{add})$ decomposes into the following pieces: $(t^n + \mathfrak{m}^{n+1}) \times \mathfrak{m}^{n+1}$, $\mathfrak{m}^{n+1} \times (t^n + \mathfrak{m}^{n+1})$ and for $i = 1, \dots, n$ the preimage $\tilde{C}_{n,i}$ of the subset $C_{n,i} \subset ((\mathfrak{m}^i - \mathfrak{m}^{i+1})/\mathfrak{m}^{n+1})^2$ of pairs $(\alpha_i t^i + \dots + \alpha_n t^n, \beta_i t^i + \dots + \beta_n t^n)$ with $\alpha_k + \beta_k = 0$ for $k = i, \dots, n-1$ and $\alpha_n + \beta_n = 1$. We must evaluate $\lambda \times \lambda'$ on each of these (relative to the diagonal μ_n -action). The first piece gives $\lambda_n \sum_{i>n} (\mathbb{L} - 1) \overline{\lambda}_i$ and the second the same expression with λ and λ' interchanged. Since $[C_{n,i}] = [(\mathfrak{m}^i - \mathfrak{m}^{i+1})/\mathfrak{m}^{n+1}] = (\mathbb{L} - 1) \mathbb{L}^{n-i}$, we find that for $i < n$, the value of $\lambda \times \lambda'$ on $\tilde{C}_{n,i}$ equals $(\mathbb{L} - 1) \mathbb{L}^{i-n} \overline{\lambda}_i \overline{\lambda}'_i$ (the action of μ_n is trivial here). Notice that $C_{n,n}$ is embedded in $(\mathfrak{m}^n - \mathfrak{m}^{n+1}/\mathfrak{m}^{n+1})^2$ as J_1 in \mathbb{G}_m^2 . From the above discussion one sees that $\lambda \times \lambda'$ takes on this set the value $J(\lambda_n, \lambda'_n)$. If we substitute the defining equation for $*$, the Lemma follows. \square

This lemma suggests a notion of a convolution operator for series

$$\lambda(T) = \sum_{n=1}^{\infty} \lambda_n T^n \in \hat{M}_k^{\hat{\mu}}[[T]]$$

with the property that the mass $(\mathbb{L} - 1) \sum_{n=1}^{\infty} \overline{\lambda}_n$ converges.

For a $\mathbb{Z}[L, L^{-1}]$ -module M we set

$$M\langle T \rangle := M[T] \left[\frac{1}{T^N - L^\nu} \mid \nu \in \mathbb{Z}, N = 1, 2, 3, \dots \right].$$

Expanding the denominators $(1 - T^N L^{-\nu})^{-1}$ in T embeds $M\langle T \rangle$ in $M[[T]]$ and expanding $(1 - T^{-N} L^\nu)^{-1}$ in T^{-1} embeds $M\langle T \rangle$ in $M[[T^{-1}]]\langle T \rangle$.

According to Theorem 5.4, $S(f) \in \hat{M}^{\hat{\mu}}\langle T \rangle$.

Theorem 7.4 (Abstract Thom-Sebastiani property). *Let λ and λ' be equivariant measures on $\mathcal{L}(\mathbb{G}_a, 0)$ whose zeta functions lie in $\hat{M}^{\hat{\mu}}\langle T \rangle$. Then $\lambda * \lambda'$ has this property, too. If moreover λ and λ' have zero mass and zeta functions converging at $T = \infty$, then $\lambda * \lambda'\langle T \rangle$ has these properties as well and $(\lambda * \lambda')(\infty) = \lambda(\infty) * \lambda'(\infty)$.*

Corollary 7.5. *Let X and Y be smooth connected varieties and $f : X \rightarrow \mathbb{G}_a$, $g : Y \rightarrow \mathbb{G}_a$ nonconstant morphisms with zero fibers X_0 and Y_0 . Let $f * g : X \times Y \rightarrow \mathbb{G}_a$ be defined by $(f * g)(x, y) := f(x) + g(y)$. Then the restriction of $[\phi_{f * g}]$ to $X_0 \times Y_0$ and the exterior $*$ -product $[\phi_f] * [\phi_g] \in M_{X_0 \times Y_0}$ coincide.*

If we apply the Hodge number characteristic followed by formation of the spectrum, then we recover the Thom-Sebastiani property for the spectrum, proved earlier by Varchenko in case f and g have isolated singularities and by M. Saito [28] in general.

For the proof of Theorem 7.4 we need the following

Lemma 7.6. *Let M and N be $\mathbb{Z}[L, L^{-1}]$ -modules and let $a \in M\langle T \rangle$ and $b \in N\langle T \rangle$ both be zero at $T = 0$ and regular at $T = \infty$. If $\sum_{k>0} a_k T^k$ resp. $\sum_{k>0} b_k T^k$ are their expansions at 0, then $\sum_{k>0} (a_k \otimes b_k) T^k$ is the expansion at zero of a $c \in (M \otimes_{\mathbb{Z}[L, L^{-1}]} N)\langle T \rangle$ whose value at $T = \infty$ equals $-a(\infty) \otimes b(\infty)$.*

Proof. It is easy to see that it suffices to prove this for $M = N = \mathbb{Z}[L, L^{-1}]$. The idea of the proof in this case is inspired by a paper of Deligne [9]. Fix for the moment $L \in \mathbb{C} - \{0\}$. Let $r_0 > 0$ be a radius of convergence for the two expansions. Let $T \in \mathbb{C}$ be such that $|T| < r_0^2$ and choose $|T|/r_0 < r < r_0$. Consider the integral

$$c(T) := \frac{1}{2\pi\sqrt{-1}} \int_{|\tau|=r} a(T/\tau) b(\tau) \frac{d\tau}{\tau}.$$

On the circle of integration the expansions converge uniformly and absolutely and so

$$c(T) = \frac{1}{2\pi\sqrt{-1}} \int_{|\tau|=r} \sum_{k,l \in \mathbb{N}} a_k T^k b^l \tau^{k-l} \frac{d\tau}{\tau}.$$

Since summation and integration may be interchanged, only the terms with $k = l$ remain and hence $c(T) = \sum_{k \in \mathbb{N}} a_k b_k T^k$. If P_a resp. P_b denotes the set of poles of a resp. b , then the integrand has polar set $TP_a^{-1} \cup P_b$ (there is no pole in 0 or ∞) and the poles enclosed by the circle of integration are those in TP_a^{-1} . By the theory of residues, $-c(T)$ must then be equal to the sum of the residues of the integrand at P_b . This description no longer requires $|T| < r_0^2$ and defines an analytic extension of c to the complement of $P_a P_b$. This extension is easily seen to be meromorphic at $P_a P_b$. To compute its behavior at ∞ , we note that $a(T/\tau)$ converges for $T \rightarrow \infty$ on a neighborhood of P_b absolutely (with all its derivatives) to the constant function $a(\infty)$. So as $T \rightarrow \infty$, $-c(T)$ tends to the sum of the residues of $a(\infty)b(\tau)\tau^{-1} d\tau$ at P_b . This sum is opposite to the residue at the remaining pole ∞ , hence equal to $a(\infty)b(\infty)$. In particular, c is a rational function with polar set contained in $P_a P_b$.

Assume now that $a, b \in \hat{R}$. A pole of an element of R in $\mathbb{C}^\times \times \mathbb{C}$ satisfies an equation $T^N = L^\nu$ for certain integers $N > 0$, $\nu \geq 0$. A product of such poles satisfies a similar equation, and this implies that a product of c and a finite set of polynomials of the form $T^N - L^\nu$ is in $\mathbb{C}[L, L^{-1}, T]$. Since the expansion of c at $T = 0$ has integral coefficients, this product lies in $\mathbb{Z}[L, L^{-1}, T]$. \square

Proof of Theorem 7.4. We start with the convolution formula 7.3. It says that

$$\begin{aligned} (\lambda * \lambda')(T) = & \\ - \sum_{n>0} \lambda_n * \lambda_n T^n + (\mathbb{L} - 1) \sum_{0 < i \leq n} \overline{\lambda_i \lambda'_i} \mathbb{L}^{i-n} T^n + (\mathbb{L} - 1) \sum_{i > n > 0} (\lambda_n \overline{\lambda'_i} + \overline{\lambda_i} \lambda'_n) T^n. \end{aligned}$$

We now assume that λ and λ' are massless so that $(\mathbb{L} - 1) \sum_{i > n} \overline{\lambda_i} = -(\mathbb{L} - 1) \sum_{i=1}^n \overline{\lambda_i}$ and similarly for λ' . We then have

$$\begin{aligned} (\lambda * \lambda')(T) = & - \sum_{n>0} \lambda_n * \lambda'_n T^n + (\mathbb{L} - 1) \sum_{0 < i \leq n} \overline{\lambda_i \lambda'_i} \mathbb{L}^{i-n} T^n + \\ & + (\mathbb{L} - 1) \sum_{n>0} \overline{\lambda_n \lambda'_n} T^n - (\mathbb{L} - 1) \sum_{0 < i \leq n} (\lambda_n \overline{\lambda'_i} + \overline{\lambda_i} \lambda'_n) T^n. \end{aligned}$$

We consider each series on the right separately. By Lemma 7.6, $-\sum_{n>0} \lambda_n * \lambda'_n T^n$ is in the in $\hat{M}_k^{\hat{\mu}} \langle T \rangle$ with value at ∞ equal to $\lambda(\infty) * \lambda'(\infty)$. We also have

$$(\mathbb{L} - 1) \sum_{0 < i \leq n} \overline{\lambda_i \lambda'_i} \mathbb{L}^{i-n} T^n = (\mathbb{L} - 1) \sum_{i>0} \sum_{k \geq 0} \overline{\lambda_i \lambda'_i} \mathbb{L}^{-k} T^{k+i} = \frac{\mathbb{L} - 1}{1 - \mathbb{L}^{-1} T} \sum_{i>0} \overline{\lambda_i \lambda'_i} T^i.$$

By the same 7.6 the righthand side is in $\hat{M}_k^{\hat{\mu}} \langle T \rangle$ and takes the value zero at ∞ . Since

$$\sum_{0 < i \leq n} \overline{\lambda_i} T^n = -(T - 1)^{-1} \sum_{i>0} \overline{\lambda_i} T^i$$

is in $\hat{M}_k^\mu\langle T \rangle$ with value zero at ∞ it follows from 7.6 that the same is true for $(\mathbb{L} - 1) \sum_{0 < i \leq n} (\bar{\lambda}_i \lambda'_n) T^n$. Likewise for $(\mathbb{L} - 1) \sum_{0 < i \leq n} (\lambda_n \bar{\lambda}'_i) T^n$. So $(\lambda * \lambda')(T)$ is in $\hat{M}_k^\mu\langle T \rangle$ and has value $\lambda(\infty) * \lambda'(\infty)$ at ∞ . \square

8. THE MCKAY CORRESPONDENCE [6], [16], [26]

Suppose a group G of finite order m acts well and effectively on a smooth connected variety U of dimension d . This defines an orbifold $p : U \rightarrow U_G$ with underlying variety $G \backslash U$. Let us write X for the orbifold U_G . We also fix a primitive m th root of unity ζ_m .

Let $g \in G$ and let U^g be its fixed point set in U . The action of g in the normal bundle of U^g decomposes that bundle into a direct sum of eigensubbundles

$$\nu_{U/U^g} = \bigoplus_{k=1}^{m-1} \nu_g^k,$$

where ν_g^k has eigenvalue ζ_m^k . We like to think of ν_g^k as the pull-back of a fractional bundle on a subvariety of X whose virtual rank is k/m times that of ν_g^k . A more formal discussion involves the extension $M_X[\mathbb{L}^{1/m}]$ of M_X obtained by adjoining an m th root of \mathbb{L} . To be precise, let $w(g) := \sum_k \frac{k}{m} \text{rk}(\nu_g^k)$, considered as locally constant function $U^g \rightarrow m^{-1}\mathbb{Z}$, and let $\mathbb{L}_{U^g}^{w(g)}$ be the element of $M_{U^g}[\mathbb{L}^{1/m}] \subset M_U[\mathbb{L}^{1/m}]$ that this defines. Then $\sum_{g \in G} \mathbb{L}_{U^g}^{w(g)}$ is the image under p^* of

$$W(X) = \sum_{[g] \in \text{conj}(G)} \sum_{i \in \pi_0(U^g)} [(G_i \backslash U_i^g)/X] \mathbb{L}^{w_i} \in M_X[\mathbb{L}^{1/m}].$$

Here U_i^g is the connected component of U^g labeled by i , G_i is the G -stabilizer of this component, and w_i the value of $w(g)$ on U_i^g . The sum is over a system of representatives of the conjugacy classes of G and can be rewritten as one over the orbifold strata of X (see Reid [26]): the decomposition of U into connected strata by orbit type (a stratum is a connected component of the locus of points with given G -stabilizer) induces a partition of X into orbifolds and $W(X)$ has the form $\sum_S [S] W_S$, where the sum is over the orbifold strata, and W_S is a polynomial in $\mathbb{L}^{1/m}$. We will see that $W(X)$ can be understood as the class of an obstruction bundle for lifting arcs in X to arcs in U .

The McKay correspondence identifies $W(X)$ in terms of a resolution of X :

Theorem 8.1 (Batyrev [6], Denef-Loeser [16]). *Let $H : Y \rightarrow X$ be a resolution of the orbifold X whose exceptional divisor E has simple normal crossings. With the usual meaning of E_I° and with ν_i^* as defined below we have the following identity in $\hat{M}_X[\mathbb{L}^{1/m}]$:*

$$W(X) = \sum_{I \subset \text{irr}(D)} [E_I^\circ/X] \prod_{i \in I} \frac{\mathbb{L} - 1}{\mathbb{L}^{\nu_i^*} - 1}.$$

The statement does not involve arc spaces, but the proof does. It could well be that the identity is already valid in $M_X[\mathbb{L}^{1/m}]$. The relative simplicity of the lefthand side has implications for the righthand side, one of which is that all the ‘non-Tate’ material in a fiber of H must cancel out in the sum. For that same reason the lefthand side is hardly affected if we apply the weight character relative to X to it, that is, if we take the image of $W(X)$ in the Grothendieck ring of constructible $\mathbb{Z}((w^{-1/m}))$ -modules on X : just substitute w^2 for \mathbb{L} .

We first seek an orbifold measure on $\mu_{\mathcal{L}(X)}^{\text{orb}}$ on $\mathcal{L}(X)$ with the property that for every G -invariant measurable $A \subset \mathcal{L}(U)$ we have

$$\mu_{\mathcal{L}(X)}^{\text{orb}}(p_*A) := \overline{\mu_{\mathcal{L}(U)}(A)},$$

where the righthand side should be interpreted as follows: think of $\mu_{\mathcal{L}(U)}(A)$ as an element of \hat{M}_k^G , and then let $\overline{\mu_{\mathcal{L}(U)}(A)}$ be the image of $\mu_{\mathcal{L}(U)}(A)$ under the augmentation $\hat{M}_k^G \rightarrow \hat{M}_k$. Since $p_* : \mathcal{L}(U) \rightarrow \mathcal{L}(X)$ need not be surjective, this will not characterize the orbifold measure a priori. But it suggests how to define it: suppose that the Jacobian ideal \mathcal{J}_p has constant order e along A . Then the usual measure of $\mathcal{L}(X)$ pulled back to A is $\mathbb{L}^{-e} \mu_{\mathcal{L}(U)}|_A$. We therefore want the orbifold measure restricted to $p_*(A)$ to be the restriction of $\mathbb{L}^e \mu_{\mathcal{L}(X)}$. This can be done as follows. Let r be a positive integer such that $(\Omega_U^d)^{\otimes r}$ descends to an invertible sheaf $\omega_X^{(r)}$ on X . (So for every $u \in U$, G_u acts on the tangent space $T_u U$ with determinant an r th root of unity.) There is a natural homomorphism $(\Omega_X^d)^{\otimes r} \rightarrow \omega_X^{(r)}$ whose kernel is the torsion of $(\Omega_X^d)^{\otimes r}$. The image of this homomorphism has the form $\mathcal{I}^{(r)} \omega_X^{(r)}$ for an ideal $\mathcal{I}^{(r)}$. We set

$$\mu_{\mathcal{L}(X)}^{\text{orb}} := \mathbb{L}^{\text{ord}_{\mathcal{I}^{(r)}}/r} \mu_{\mathcal{L}(X)}.$$

It is a measure that takes values in $\hat{M}_k[\mathbb{L}^{1/r}]$.

Lemma 8.2. *The pull-back of $\mu_{\mathcal{L}(X)}^{\text{orb}}$ under p^* is a measure that assigns to any G -invariant measurable subset A of $\mathcal{L}(U)$ the image of A under the augmentation map $\hat{M}_k^G \rightarrow \hat{M}_k$.*

Proof. If we apply p^* to the identity $(\Omega_X^d)^{\otimes r}/\text{tors} = \mathcal{I}^{(r)} \omega_X^{(r)}$ we get $\mathcal{J}_p^r (\Omega_U^d)^{\otimes r} = p^*(\mathcal{I}^{(r)}) \omega_U^{\otimes r}$. Since $\Omega_U^d = \omega_U$, it follows that $p^*(\mathcal{I}^{(r)}) = \mathcal{J}_p^r$. So $\mu_{\mathcal{L}(X)}^{\text{orb}}$ pulls back under p^* to $\mathbb{L}^{-\text{ord}_{\mathcal{J}_p} + p^*(\mathcal{I}^{(r)})/r} \mu_{\mathcal{L}(U)} = \mu_{\mathcal{L}(U)}$. The rest is left to the reader. \square

The following lemma describes the direct image of $\mu_{\mathcal{L}(X)}^{\text{orb}}$ on X in terms of a resolution of X : let $Y \rightarrow X$ be a resolution of singularities with simple normal crossing divisor E . We have $H^* \omega_X^{(r)} = \tilde{\mathcal{I}}^{(r)} \omega_Y^{\otimes r}$ for some fractional ideal $\tilde{\mathcal{I}}^{(r)}$ on Y . It is known that the multiplicity m_i of E_i in this ideal is $> -r$. So $\nu_i^* := 1 + m_i/r$ is positive. Entirely analogous to the proof of Theorem 4.2 one derives:

Lemma 8.3. *The direct image of $\mu_{\mathcal{L}(X)}^{\text{orb}}$ on X is represented by the class*

$$\sum_{I \subset \text{Irr}(E)} [E_I^\circ/X] \prod_{i \in I} \frac{\mathbb{L} - 1}{\mathbb{L}^{\nu_i^*} - 1} \in \hat{M}_X[\mathbb{L}^{1/r}].$$

Let $\mathcal{L}'(X)$ be the set of arcs in X not contained in the discriminant of $p : U \rightarrow X$. This is a subset of full measure. We decompose $\mathcal{L}'(X)$ according to the ramification behavior of $p : U \rightarrow X$. Let $[m] : \mathbb{D} \rightarrow \mathbb{D}$ be the m th power map and denote the parameter of the domain by $t^{1/m}$. We regard ζ_m (through its action on the domain) as generator of the Galois group of $[m]$. For $\gamma \in \mathcal{L}'(X)$, $\gamma[m]$ lifts to a morphism $\tilde{\gamma} : \mathbb{D} \rightarrow \tilde{X}$ and this lift is unique up to conjugation with G . Given the lift, there is a $g \in G$ such that $g\tilde{\gamma} = \tilde{\gamma}\zeta_m$. Its conjugacy class $[g]$ in G only depends on γ . This conjugacy class determines the isomorphism type of the G -covering over γ : if m' is the order of g , then $\gamma^*(p)$ is isomorphic to $G \times^{(g)} \mathbb{D} \rightarrow \mu_{m'} \backslash \mathbb{D}$, with g acting on \mathbb{D} as multiplication by $\zeta^{m/m'}$. Notice that $\tilde{\gamma}(o)$ is in the fixed point set U^g . The

‘fractional lifts’ $\tilde{\gamma}$ that so arise are like arcs in the total space of the normal bundle $\oplus_k \nu_g^k$ of U^g (based at the zero section) which in the ν_g^k -direction develop as $t^{k/m}$ times a power series in t .

Denote the set of arcs in $\mathcal{L}'(X)$ belonging to the conjugacy class of $[g]$ of g by $\mathcal{L}(X, [g])$. The McKay correspondence now results from:

Lemma 8.4. *The subset $\mathcal{L}(X, [g])$ is measurable for $\mu_{\mathcal{L}(X)}^{\text{orb}}$ and the restriction of $\mu_{\mathcal{L}(X)}^{\text{orb}}$ to this subset is represented by the class $[(G_g \backslash U^g)/X] \mathbb{L}^{w(g^{-1})} \in M_X[\mathbb{L}^{1/m}]$, where G_g is the G -stabilizer of U^g .*

The proof is a calculation which we only discuss in a heuristic fashion. The elements of $\mathcal{L}(X, [g])$ correspond to G_g -orbits of fractional lifts as described above. In view of our definition of orbifold measure, we need to argue that these fractional lifts are represented by the element $\mathbb{L}_{U^g}^{w(g^{-1})}$. If r_1, \dots, r_{m-1} are positive integers, then the arcs in $\oplus_k \nu_g^k$ of U^g based at the zero section and which in the ν_g^k -direction have order r_k make up a constructible subset of $\mathcal{L}(\oplus_k \nu_g^k)$ whose class is easily seen to be equal to $\mathbb{L}_{U^g}^w$, with $w = \sum_k (1 - r_k) \text{rk}(\nu_g^k)$. The fact is that this also holds for the fractional values $r_k = k/m$. So in that case we have $w = \sum_k (1 - k/m) \text{rk}(\nu_g^k) = w(g^{-1})$.

9. PROOF OF THE TRANSFORMATION RULE [14]

Let \mathcal{X}/\mathbb{D} be a \mathbb{D} -variety of pure relative dimension d . The d th Fitting ideal of $\Omega_{\mathcal{X}/\mathbb{D}}$ defines the locus where \mathcal{X} fails to be smooth over \mathbb{D} ; we denote that ideal by $\mathcal{J}(\mathcal{X}/\mathbb{D})$. Locally this ideal is obtained as follows: if \mathcal{X} is given as a closed subset of $(\mathbb{A}^{d+l})_{\mathbb{D}}$, then $\mathcal{J}_{\mathcal{X}/\mathbb{D}}$ is the restriction to \mathcal{X} of the ideal generated by the determinants $\det((\partial f_j / \partial x_{i_k})_{j,k=1}^l)$, where f_1, \dots, f_l are taken from the ideal $I_{\mathcal{X}} \subset \mathcal{O}[x_1, \dots, x_{d+l}]$ defining \mathcal{X} and $1 \leq i_1 < \dots < i_l \leq d+l$.

Let $\gamma \in \mathcal{X}_{\infty}$ be such that $\gamma^* \mathcal{J}(\mathcal{X}/\mathbb{D})$ has finite order e . This implies that γ maps \mathbb{D}^{\times} to the part $(\mathcal{X}/\mathbb{D})_{\text{reg}}$ where \mathcal{X} is smooth over \mathbb{D} . In particular, $\gamma^* \Omega_{\mathcal{X}/\mathbb{D}}$ is a \mathcal{O} -module of rank d . Since the formation of a Fitting ideal commutes with base change, the d th Fitting ideal of $\gamma^* \Omega_{\mathcal{X}/\mathbb{D}}$ will be \mathfrak{m}^e . This means that the torsion of $\gamma^* \Omega_{\mathcal{X}/\mathbb{D}}$ has length e .

It is clear that $\text{Der}_{\mathcal{O}}(\mathcal{O}_{\mathcal{X}, \gamma(o)}, \mathcal{O}) \cong \text{Hom}_{\mathcal{O}}(\gamma^* \Omega_{\mathcal{X}/\mathbb{D}}, \mathcal{O})$ is a free \mathcal{O} -module of rank d (where \mathcal{O} is a $\mathcal{O}_{\mathcal{X}, \gamma(o)}$ -module via γ^*). The fiber over o , $\text{Hom}_{\mathcal{O}}(\gamma^* \Omega_{\mathcal{X}/\mathbb{D}}, \mathcal{O}) \otimes_{\mathcal{O}} k$, is d -dimensional subspace of the Zariski tangent space $T_{\mathcal{X}, \gamma(o)}$, which we shall denote by $\hat{T}_{\mathcal{X}, \gamma}$. Any \mathcal{O} -homomorphism $\gamma^* \Omega_{\mathcal{X}/\mathbb{D}} \rightarrow \mathcal{O}/\mathfrak{m}^{n+1}$ that kills the torsion lifts to a \mathcal{O} -homomorphism $\gamma^* \Omega_{\mathcal{X}/\mathbb{D}} \rightarrow \mathcal{O}$. This is automatic when $n \geq e$ and so $\hat{T}_{\mathcal{X}, \gamma}$ only depends on the e -jet of γ . The space $\hat{T}_{\mathcal{X}, \gamma}$ has a simple geometric interpretation: it is the ‘limiting position’ of the tangent space along the fibers of \mathcal{X}/\mathbb{D} at the generic point of $\gamma(\mathbb{D})$ in the closed point $\gamma(o)$.

If $\gamma' : \mathbb{D} \rightarrow \mathcal{X}$ has the same n -jet as γ , then γ^* and γ'^* differ by a homomorphism $\mathcal{O}_{\mathcal{X}, \gamma(o)} \rightarrow \mathfrak{m}^{n+1}$. The reduction modulo $\mathfrak{m}^{2(n+1)}$ of this homomorphism is a \mathcal{O} -derivation, i.e., defines an element of $\text{Hom}_{\mathcal{O}}(\gamma^* \Omega_{\mathcal{X}/\mathbb{D}}, \mathfrak{m}^{n+1}/\mathfrak{m}^{2(n+1)})$. Its reduction modulo \mathfrak{m}^{n+2} will lie in $\hat{T}_{\mathcal{X}, \gamma} \otimes \mathfrak{m}^{n+1}/\mathfrak{m}^{n+2}$, provided that $n \geq e$. The next lemma shows that every element of this k -vector space so arises.

Lemma 9.1. *Assume that $n \geq e$. The fiber of $\pi_{n+1}\mathcal{X}_\infty \rightarrow \pi_n\mathcal{X}_\infty$ over $\pi_n(\gamma)$ is an affine space with translation space $\hat{T}_{X,\gamma} \otimes_k \mathfrak{m}^{n+1}/\mathfrak{m}^{n+2}$. This defines an affine space bundle of rank d over the locus of $\pi_n\mathcal{X}_\infty$ defined by $\text{ord}_{\mathcal{J}(\mathcal{X}/\mathbb{D})} \leq n$.*

Proof. Assume that \mathcal{X} is given as a closed subset of $(\mathbb{A}^{d+l})_{\mathbb{D}}$ as above. There exist $f_1, \dots, f_l \in I_{\mathcal{X}}$ and $1 \leq i_1 < \dots < i_l \leq d+l$ such that the Jacobian matrix $\det((\partial f_j / \partial x_{i_k})_{j,k=1}^l)$ has order e along γ , whereas for any other matrix thus formed the order is $\geq e$. By means of a coordinate change we may arrange that

$$\gamma^* df_j \equiv t^{e_j} dx_j \pmod{t^{e_j+1}(dx_{j+1}, \dots, dx_{d+l})}, \quad j = 1, \dots, l,$$

so that $e = \sum_j e_j$. The subspace of \mathbb{A}_k^{d+l} spanned by the last d basis vectors is then just $\hat{T}_{X,\gamma}$.

We investigate which $u_0 \in k^{d+l}$ appear as the constant coefficient of an $u \in \mathcal{O}^{d+l}$ with the property that $\gamma + t^{n+1}u \in \mathcal{X}_\infty$. We first do this for the complete intersection defined by f_1, \dots, f_l . This complete intersection contains \mathcal{X} and the irreducible component that contains the image of γ lies in \mathcal{X} . So we want $f_j(\gamma + t^{n+1}u) = 0$ for $j = 1, \dots, l$. By expanding at γ this amounts to identities of the form

$$t^{n+1}D_\gamma f_j(u) + t^{2(n+1)}F_j(u) = 0, \quad j = 1, \dots, l,$$

with $D_\gamma f_j$ the derivative of f_j at γ and $F_j \in \mathcal{O}[x_1, \dots, x_{d+l}]$. Equivalently:

$$t^{-e_j}D_\gamma f_j(u) + t^{n+1-e_j}F_j(u) = 0, \quad j = 1, \dots, l.$$

All the terms are regular and the reduction modulo t yields the j th unit vector in k^{d+l} . Hensel's lemma says that a solution u exists if and only if u_0 solves this set of equations modulo t . This just means that $u_0 \in \hat{T}_{X,\gamma}$. In particular, we see that for all $k \in \mathbb{N}$, $\pi_{n+k}\pi_n^{-1}\pi_n(\gamma)$ is isomorphic to an affine space and hence is irreducible. This implies that all elements of $\pi_n^{-1}\pi_n(\gamma)$ map to the same irreducible component of the common zero locus of f_1, \dots, f_l . It follows that $\pi_n^{-1}\pi_n(\gamma) \subset \mathcal{X}_\infty$. The last assertion is easy. \square

Proof of Proposition 3.1. Suppose that \mathcal{X} is of pure relative dimension d . Let C_e denote the subset of \mathcal{X}_∞ defined by $\text{ord}_{\mathcal{J}(\mathcal{X}/\mathbb{D})} = e$. It is clear that $C_e = \pi_e^{-1}\pi_e(C_e)$. It follows from Greenberg's theorem [21] that $\pi_e(C_e)$ is constructible. Hence C_e is stable by Lemma 9.1. We have $\cup_e C_e = \mathcal{X}_\infty - (\mathcal{X}_{\text{sing}})_\infty$. In view of Lemma 2.3 it now suffices to see that $\dim \pi_e(C_e) - de \rightarrow -\infty$ as $e \rightarrow \infty$. This is not difficult. \square

Let $H : \mathcal{Y} \rightarrow \mathcal{X}$ be a \mathbb{D} -morphism of \mathbb{D} -varieties of pure relative dimension d . Recall that the Jacobian ideal \mathcal{J}_H of H is the 0th Fitting ideal of $\Omega_{\mathcal{Y}/\mathcal{X}}$. Suppose $\gamma \in \mathcal{Y}_\infty$ is such that \mathcal{J}_H has finite order e along γ . Then γ resp. $H\gamma$ maps the generic point \mathbb{D}^\times to $(\mathcal{Y}/\mathbb{D})_{\text{reg}}$ resp. $(\mathcal{X}/\mathbb{D})_{\text{reg}}$. We have an exact sequence of \mathcal{O} -modules

$$(H\gamma)^*\Omega_{\mathcal{X}/\mathbb{D}} \rightarrow \gamma^*\Omega_{\mathcal{Y}/\mathbb{D}} \rightarrow \gamma^*\Omega_{\mathcal{Y}/\mathcal{X}} \rightarrow 0.$$

The base change property of Fitting ideals implies that the length of $\gamma^*\Omega_{\mathcal{Y}/\mathcal{X}}$ must be e . So if $\gamma^*\Omega_{\mathcal{Y}/\mathbb{D}}$ is torsion free and $n \geq e$, then the kernel of the map

$$D_\gamma^{(n)} : \text{Hom}_{\mathcal{O}}(\gamma^*\Omega_{\mathcal{Y}/\mathbb{D}}, \mathcal{O}/\mathfrak{m}^{n+1}) \rightarrow \text{Hom}_{\mathcal{O}}((\gamma H)^*\Omega_{\mathcal{X}/\mathbb{D}}, \mathcal{O}/\mathfrak{m}^{n+1})$$

induced by the derivative of H is contained in $\text{Hom}_{\mathcal{O}}(\gamma^*\Omega_{\mathcal{Y}/\mathbb{D}}, \mathfrak{m}^{n+1-e}/\mathfrak{m}^{n+1})$, can be identified with $\text{Hom}_{\mathcal{O}}(\gamma^*\Omega_{\mathcal{Y}/\mathcal{X}}, \mathcal{O}/\mathfrak{m}^{n+1})$, and is of length e . The proof of Theorem 3.2 now rests on the

Key lemma 9.2. *Suppose \mathcal{Y}/\mathbb{D} smooth and let $A \subset \mathcal{Y}_\infty$ be a stable subset of level l : $A = \pi_l^{-1}\pi_l(A)$. Assume that $H|_A$ is injective and that $\text{ord}_{\mathcal{J}_H}|_A$ is constant equal to $e < \infty$. If $n \geq \sup\{2e, l + e, \text{ord}_{\mathcal{J}(\mathcal{X}/\mathbb{D})}|_{HA}\}$, then $H_n : \pi_n A \rightarrow H_n \pi_n A$ has the structure of affine-linear bundle of dimension e .*

Proof. Let $\gamma \in A$ and put $x := \gamma(o)$, $y := H(x)$. Suppose $\gamma' \in A$ is such that $H\gamma'$ and $H\gamma$ have the same n -jet. We first show that γ and γ' have the same $(n - e)$ -jet. We do this by constructing a $\gamma_1 \in \mathcal{Y}_\infty$ (by successive approximation) with the same $(n - e)$ -jet as γ and with $H\gamma_1 = H\gamma'$. Since $n - e \geq l$, we will have $\gamma_1 \in A$ and our injectivity assumption then implies $\gamma_1 = \gamma'$.

The difference $(H\gamma)^* - (H\gamma')^*$ defines a \mathcal{O} -derivation $\mathcal{O}_{\mathcal{X},x} \rightarrow \mathfrak{m}^{n+1}/\mathfrak{m}^{2(n+1)}$ over γ^* and hence a $\tilde{v} \in \text{Hom}_{\mathcal{O}}((\gamma H)^*\Omega_{\mathcal{X}/\mathbb{D}}, \mathfrak{m}^{n+1}/\mathfrak{m}^{2n+2})$. Since $n \geq \text{ord}_{H\gamma} \mathcal{J}(\mathcal{X}/\mathbb{D})$, this element annihilates the torsion of $(\gamma H)^*\Omega_{\mathcal{X}/\mathbb{D}}$. This is then also true for its reduction modulo \mathfrak{m}^{n+2} and it follows from the fact that $n \geq e$ that this reduction is of the form $D_\gamma^{(n+1)}(u)$ for some $u \in \text{Hom}_{\mathcal{O}}(\gamma^*\Omega_{\mathcal{Y}/\mathbb{D}}, \mathfrak{m}^{n-e+1}/\mathfrak{m}^{n+2})$. Regard u as a \mathcal{O} -derivation $\mathcal{O}_{\mathcal{Y},y} \rightarrow \mathfrak{m}^{n-e+1}/\mathfrak{m}^{n+2}$ and let $\gamma_1 \in \mathcal{Y}_\infty$ be such that $\gamma_1^* - \gamma^*$ represents u . Then $\pi_{n-e}(\gamma_1) = \pi_{n-e}(\gamma)$ and $\pi_{n+1}(H\gamma_1) = \pi_{n+1}(H\gamma)$. Replace γ by γ_1 and continue with induction on n .

So $(\gamma')^* - \gamma^*$ defines a \mathcal{O} -derivation $\mathcal{O}_{\mathcal{X},x} \rightarrow \mathfrak{m}^{n-e+1}/\mathfrak{m}^{2(n-e+1)}$ and hence a \mathcal{O} -derivation $\mathcal{O}_{\mathcal{X},x} \rightarrow \mathfrak{m}^{n-e+1}/\mathfrak{m}^{n+1}$ (because $n \geq 2e$). The latter is zero if and only if $\pi_n(\gamma') = \pi_n(\gamma)$. This proves that the fiber of $H_n|_{\pi_n A}$ through $\pi_n(\gamma)$ is an affine space over the kernel of $D_\gamma^{(n)}$, $\text{Hom}_{\mathcal{O}}(\gamma^*\Omega_{\mathcal{Y}/\mathcal{X}}, \mathcal{O}/\mathfrak{m}^{n+1})$, which has length e .

The last assertion is easy. \square

Proof of 3.2. It is enough to prove this for A stable. In that case the theorem follows in a straightforward manner from Lemma 9.2. \square

REFERENCES

- [1] D. Abramovich, K. Karu, K. Matsuki, J. Włodarczyk, *Torification and Factorization of Birational Maps*, 30 p., [arXiv.org/abs/math/9904135](https://arxiv.org/abs/math/9904135).
- [2] N. A'Campo, *La fonction zêta d'une monodromie*, *Comment. Math. Helv.* **50** (1975), 233–248.
- [3] G. Anderson, *Cyclotomy and an extension of the Taniyama group*, *Comp. Math.* **57** (1986), 153–217.
- [4] V. Batyrev, *Birational Calabi–Yau n -folds have equal Betti numbers*, in *New trends in algebraic geometry*, Klaus Hulek *et al.*, eds., CUP, 1999, 1–11.
- [5] V. Batyrev, *Stringy Hodge numbers of varieties with Gorenstein canonical singularities*, in *Integrable systems and algebraic geometry (Kobe/Kyoto, 1997)*, 1–32, World Sci. Publishing, River Edge, NJ, 1998.
- [6] V. Batyrev, *Non-Archimedean integrals and stringy Euler numbers of log-terminal pairs*, *J. Eur. Math. Soc.* **1** (1999), 5–33.
- [7] V. Batyrev and D. Dais, *Strong McKay correspondence, string-theoretic Hodge numbers and mirror symmetry*, *Topology* **35** (1996), 901–929.
- [8] A. Craw, *An introduction to motivic integration*, 23 pp., [arXiv.org/abs/math/9911179](https://arxiv.org/abs/math/9911179).
- [9] P. Deligne, *Intégration sur un cycle évanescant*, *Invent. Math.* **76** (1984), no. 1, 129–143.
- [10] J. Denef, *Local zeta functions and Euler characteristics*, *Duke Math. J.* **63** (1991), no. 3, 713–721.
- [11] J. Denef, *Report on Igusa's local zeta function*, in *Séminaire Bourbaki, volume 1990/91*, exposé 741, *Astérisque* **201-202-203** (1991), 359–386.
- [12] J. Denef, F. Loeser, *Caractéristiques d'Euler-Poincaré, fonctions zêta locales et modifications analytiques*, *J. Amer. Math. Soc.* **5** (1992), 705–720.
- [13] J. Denef, F. Loeser, *Motivic Igusa zeta functions*, *J. Algebraic Geom.*, **7** (1998), 505–537.

- [14] J. Denef, F. Loeser, *Germes of arcs on singular algebraic varieties and motivic integration*, Invent. Math., **135** (1999), 201–232.
- [15] J. Denef, F. Loeser, *Motivic exponential integrals and a motivic Thom-Sebastiani Theorem*, Duke Math. J., **99** (1999), 285–309.
- [16] J. Denef, F. Loeser, *Motivic integration, quotient singularities and the McKay correspondence*, 20 p., [arXiv.org/abs/math/9903187](https://arxiv.org/abs/math/9903187).
- [17] J. Denef, F. Loeser, *Definable sets, motives and P-adic integrals*, 45 p., [arXiv.org/abs/math/9910107](https://arxiv.org/abs/math/9910107).
- [18] J. Denef, F. Loeser, *Lefschetz numbers of the monodromy and truncated arcs*, 10 p., [arXiv.org/abs/math/0001105](https://arxiv.org/abs/math/0001105).
- [19] J. Denef, F. Loeser, *Geometry on arc spaces of algebraic varieties*, 22 p., [arXiv.org/abs/math/0006050](https://arxiv.org/abs/math/0006050).
- [20] M. Greenberg, *Schemata over local rings*, Ann. Math. **73** (1961), 624–648.
- [21] M. Greenberg, *Rational points in discrete valuation rings*, Publ. Math. I.H.E.S. **31** (1966), 59–64.
- [22] M. Kapranov, *The elliptic curve in the S-duality theory and Eisenstein series for Kac-Moody groups*, 41 pp., [arXiv.org/abs/math/0001005](https://arxiv.org/abs/math/0001005).
- [23] M. Kontsevich, Lecture at Orsay (December 7, 1995).
- [24] J. Nash Jr., *Arc structure of singularities*, Duke Math. J., **81** (1995), 31–38.
- [25] J. Pas, *Uniform p-adic cell decomposition and local zeta functions*, J. f. d. reine u. angew. Math. **399** (1989), 137–172.
- [26] M. Reid, *La correspondance de McKay*, in *Séminaire Bourbaki* (en anglais), exposé 867, Novembre 1999, 20 p., in this volume and at [arXiv.org/abs/math/9911195](https://arxiv.org/abs/math/9911195).
- [27] T. Shioda, T. Katsura, *On Fermat varieties*, Tohoku math. J. **31** (1979), 97–115.
- [28] J. Steenbrink, *The spectrum of hypersurface singularities*, in *Théorie de Hodge, Luminy 1987*, Astérisque **179-180** (1989), 163–184.
- [29] A. Varchenko, *Asymptotic Hodge structure in the vanishing cohomology*, Math. USSR Izvestija **18** (1982), 469–512.
- [30] W. Veys, *The topological zeta function associated to a function on a normal surface germ*, Topology **38** (1999), no. 2, 439–456.
- [31] W. Veys, *Zeta functions and ‘Kontsevich invariants’ on singular varieties*, 27 pp., [arXiv.org/abs/math.AG/0003025](https://arxiv.org/abs/math.AG/0003025).

MATHEMATISCH INSTITUUT, UNIVERSITEIT UTRECHT, PO BOX 80.010, NL-3508 TA UTRECHT, NEDERLAND

E-mail address: looiijeng@math.uu.nl