

**Larsen C ice shelf
meteorology**

P. Kuipers Munneke et al.

This discussion paper is/has been under review for the journal The Cryosphere (TC).
Please refer to the corresponding final paper in TC if available.

Near-surface climate and surface energy budget of Larsen C ice shelf, Antarctic Peninsula

P. Kuipers Munneke¹, M. R. van den Broeke¹, J. C. King², T. Gray², and C. H. Reijmer¹

¹Institute for Marine and Atmospheric Research, Utrecht University, Utrecht, The Netherlands

²British Antarctic Survey, National Environmental Research Council, Cambridge, UK

Received: 20 September 2011 – Accepted: 29 September 2011 – Published: 12 October 2011

Correspondence to: P. Kuipers Munneke (p.kuipersmunneke@uu.nl)

Published by Copernicus Publications on behalf of the European Geosciences Union.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Abstract

Data collected by two automatic weather stations (AWS) on the Larsen C ice shelf, Antarctica, between 22 January 2009 and 1 February 2011 are analyzed and used as input for a model that computes the surface energy budget (SEB), including melt energy. The two AWSs are separated by about 70 km in the north-south direction, and both the near-surface meteorology and the SEB show similarities, although small differences in all components (most notably the melt flux) can be seen. The impact of subsurface absorption of shortwave radiation on melt and snow temperature is significant, and discussed. In winter, longwave cooling of the surface is entirely compensated by a downward turbulent transport of sensible heat. In summer, the positive net radiative flux is compensated by melt, and quite frequently by upward turbulent diffusion of heat and moisture, leading to sublimation and weak convection over the ice shelf. The month of November 2010 is highlighted, when strong westerly flow over the Antarctic Peninsula led to a dry and warm föhn wind over the ice shelf, resulting in warm and sunny conditions. Under these conditions the increase in shortwave and sensible heat fluxes is larger than the reduction of net longwave and latent heat fluxes, providing energy for significant melt.

1 Introduction

Over the past 50 years, the Antarctic Peninsula has seen an atmospheric warming that is much larger than the global average (Turner et al., 2005). In relation to this, Antarctic Peninsula ice shelves, which are the floating extensions of glaciers originating in the mountains of the Antarctic Peninsula, have retreated everywhere in this region (Cook and Vaughan, 2010). The collapse of Larsen A and B ice shelves (Rott et al., 1996) have become iconic images for the rapid climate change in this region. The disappearance of ice shelves has caused acceleration and thinning of the glaciers previously feeding them (Rignot et al., 2004; Scambos et al., 2004; Rott et al., 2011). Whether the shrinkage of the ice shelves is driven mainly by atmospheric warming or by

Larsen C ice shelf meteorology

P. Kuipers Munneke et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Larsen C ice shelf meteorology

P. Kuipers Munneke et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

changes in oceanic circulation underneath is a subject of lively scientific discourse, but it is certain that atmospheric warming has profound effects on the state of the snow and firn layer covering the ice shelves (Holland et al., 2011). It has been hypothesized that an increase in snowmelt could enhance hydrofracturing of surface crevasses (Scambos et al., 2000; Van den Broeke, 2005), influencing the stability of ice shelves.

Obviously, it is important to study the way in which the atmosphere interacts with the snow-covered ice-shelf surface. It is therefore somewhat surprising that no detailed accounts of the surface energy budget (SEB), which represents the energy transfer between the atmosphere and the snow (including melt), over Antarctic Peninsula ice shelves are available in the scientific literature. In order to better understand the influence of near-surface climate on the condition of the snowpack, a network of six year-round operating automatic weather stations (AWS) has been installed since 2008 on the Larsen C ice shelf in a collaboration between the British Antarctic Survey (BAS, UK), the University of Colorado (USA), the Centro de Estudios Científicos (CECS, Chile) and the Institute for Marine and Atmospheric research at Utrecht University (UU/IMAU, The Netherlands). In February 2011, an additional UU/IMAU AWS was installed on Scar Inlet, the remainder of the Larsen B ice shelf after its collapse in 2002. Here, we present the first two years of data from the two UU/IMAU AWSs on the Larsen C ice self, and give an overview of the near-surface climate and the SEB at these locations. In Sect. 2, we present the data and methods central to this paper, and in Sect. 3 we provide an analysis of the near-surface meteorology and SEB. A discussion of these results and general conclusions are presented in Sect. 4.

2 Data and methods

The data collected by the two AWSs described in this paper are used to drive a model that solves the surface energy budget. In this section, we describe the AWS sites, their instrumentation, the way in which we corrected the raw data, and we briefly discuss the surface energy budget model.

2.1 AWS sites and instrumentation

AWS 14 is the northerly UU/IMAU AWS at $67^{\circ}00.8' \text{ S } 61^{\circ}28.8' \text{ W}$ (see map in Fig. 1), at an elevation of 40 m a.s.l. The other station, AWS 15, is located about 70 km to the SSW at $67^{\circ}34.3' \text{ S } 62^{\circ}07.5' \text{ W}$ (also at 40 m a.s.l.). Both stations are situated roughly 125 km from the grounding line to the west, and 55 km from the ice shelf front to the east. The stations are equipped with a GPS antenna, data from which indicate that both stations are moving eastward by about 400 m yr^{-1} . The surface at both locations is homogeneous and flat, with a surface slope less than 0.1° . The stations were installed in January 2009, serviced in January 2010, and serviced and raised in January 2011.

The following quantities are measured by both AWSs: air temperature (T_a) and relative humidity (RH) by a Vaisala HMP35AC. The temperature sensor is naturally-ventilated. Air pressure (p) is measured by a Vaisala PTB101B. Wind speed (v) and direction are observed using a Young 05103, and the radiative fluxes (shortwave fluxes (SW_{\downarrow} and SW_{\uparrow}), longwave fluxes (LW_{\downarrow} and LW_{\uparrow})) are measured by a naturally-ventilated Kipp and Zonen CNR1 radiometer. The tilt of the AWS mast in two perpendicular directions is observed using homemade inclinometers, and snow temperature (T_{sn}) is observed by thermistor strings at initial depths of 0.25, 0.50, 1.0, 2.0, 3.0, 4.0, 5.0, 7.0, 10.0, and 15.0 m. During the 2011 visit, new snow thermistors were installed at 0.05, 0.10, 0.20, 0.40, and 0.80 m depth. Additionally, surface height is measured using a sonic height ranger. The initial height of the wind and air temperature sensors was 3.80 m above the snow. All quantities are sampled every 6 min, from which hourly means are stored on a Campbell CR10X datalogger. An exception to this is the air pressure, which is sampled once per hour.

2.2 AWS data treatment

Data coverage of AWS 14 is continuous between 20 January 2009 and 1 April 2011. Hourly data were recovered from the data logger on 16 January 2011, and data between 16 January and 1 April 2011 have been obtained by interpolating station data

TCD

5, 2665–2697, 2011

Larsen C ice shelf meteorology

P. Kuipers Munneke et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



received from the ARGOS satellite network. Data coverage of AWS 15 is also continuous between 20 January 2009 and 28 January 2011. Since then, the ARGOS antenna malfunctioned, so that no data are available after that date.

Before the AWS data can be used as input for the SEB model, some corrections to the raw data are necessary. Many of these corrections have been described before in detail by Van den Broeke et al. (2004); Kuipers Munneke et al. (2009); Giesen et al. (2009). First of all, the shortwave fluxes are corrected for sensor tilt using the tilt sensors of the AWS. When the wind speed sensor is rimed, data from one AWS are used to fill the gap of the other AWS. This is justified because wind speed variations at the two sites are well-correlated and mean wind speeds are similar (see Fig. 2 and Table 1). When both vanes are rimed (occurring in three separate periods of 2, 5, and 8 days during winter), we assume a constant wind speed of 1 m s^{-1} . Relative humidity is corrected with respect to sublimation over ice. Since the air temperature screen is not ventilated to save energy, an overestimation of air temperature occurs on calm, sunny days. A correction is applied that is a function of wind speed and the sum of the incoming and reflected shortwave fluxes (Smeets, 2006).

2.3 Surface energy budget model

The SEB model used in this study is identical to the one used in Kuipers Munneke et al. (2009), including the computation of subsurface absorption of solar radiation. For details, we refer to the above-mentioned paper, but we will reiterate the concept of the model here. The model uses ρ , RH, T_a , \mathbf{v} , SW_{\downarrow} , SW_{\uparrow} , and LW_{\downarrow} as input. The sensible heat flux H_{sen} and latent heat flux H_{lat} are calculated using a bulk method, applied to the air layer between the sensor level and the snow surface. The ground heat flux G is calculated using a multi-layer snowpack that allows for melting, refreezing, and percolation of meltwater. The actual value of G is computed using the vertically integrated temperature change of the snowpack (Kuipers Munneke et al., 2009). The

Larsen C ice shelf meteorology

P. Kuipers Munneke et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[I◀](#)[▶I](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

SEB equation is:

$$SW_{\downarrow} + SW_{\uparrow} + Q + LW_{\downarrow} + LW_{\uparrow} + H_{\text{sen}} + H_{\text{lat}} + G = M, \quad (1)$$

where M is the sum of surface and subsurface melt. Q is the amount of shortwave radiation absorbed below the surface. All fluxes are defined as positive when directed towards the surface. It should be noted that Eq. (1) is cast in such a way that it covers the energy budget of the entire snowpack, including the subsurface. This is done because we take into account absorption of radiation below the surface.

The model iterates until a skin temperature is found for which the energy budget is closed. This skin temperature yields LW_{\uparrow} , which can be verified against the observed LW_{\uparrow} to assess the performance of the SEB model. In this way, a closed and consistent surface energy budget is found.

The scalar roughness length is taken as 0.11 mm, derived from the direct measurements of turbulent fluxes. Roughness lengths for temperature and humidity are calculated from the roughness length for momentum using the relations by Andreas (1987). Longwave emissivity of the snowpack is taken as 0.98. The snow grain size for the subsurface radiation penetration is taken as 100 μm . The model snow density profile for both AWSs is derived from the mean of more than 60 snow pits dug in January 2011 around AWS 14, and taken constant over the entire period. The error associated with this assumption is small (Van As et al., 2005; Kuipers Munneke et al., 2009). The temperature of the model snowpack is initialized using thermistor string observation at the start of the observation period.

3 Results

In this section, we will discuss the results of the SEB model described in Sect. 2.3. First, we will discuss the general near-surface meteorological conditions in Sect. 3.1. In Sect. 3.2, we present the different components of the SEB. In Sects. 3.3–3.5, we

Larsen C ice shelf meteorology

P. Kuipers Munneke et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



zoom in on some particular aspects of the SEB and their relation to the near-surface meteorology on Larsen C ice shelf.

3.1 Near-surface meteorology

Here, we give an overview of the near-surface meteorology of both AWS sites for the measurement period. Due to the short period under consideration, this overview should not be seen as a site climatology, but rather as a description of the typical meteorological conditions encountered during these two years. Since the sensor height above the surface is changing due to snow accumulation, we interpolate air temperature and humidity to the 2-m level using vertical profiles that emerge from the bulk method for turbulent fluxes described above. Likewise, wind speed is extrapolated to 10 m above the surface.

The mean 2-m (nominal) air temperature at AWS 14 (15) is -15.5 (-15.8) $^{\circ}\text{C}$ (see Table 1 and Fig. 2). In summer, the melting surface keeps the air temperature close to 0°C , and the temporal variability is reduced. From autumn to spring, temperature variability is much larger, as the air temperature is governed mainly by advection. During advection of warm air, near-surface temperature can rise close to the melting point even in winter, like on 14 July 2010 (-0.6°C at AWS 14) or 13 September 2010 (1.5°C). The lowest temperature recorded during this period was -46.9 (-45.4) $^{\circ}\text{C}$ at AWS 14 (15) on 12 (10) July 2009.

Specific humidity exhibits a strong annual cycle, with one order of magnitude higher values during the summer months. In general, the air over the Larsen C ice shelf is very moist, with a mean relative humidity of almost 95 % at both stations (Table 1).

The mean wind speed is 4.5 (4.3) m s^{-1} at AWS 14 (15) for this period. These values are among the lowest of coastal Antarctica – Larsen C is sheltered from the westerlies by the Antarctic Peninsula mountains, its flat surface does not generate katabatic flow, and it is not or weakly under the influence of outflow of remotely generated cold air from the continent through the barrier wind mechanism. In Fig. 3, wind roses are shown for both stations. Interestingly, south is the dominant wind direction at AWS 15,

Larsen C ice shelf meteorology

P. Kuipers Munneke et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



while at AWS 14 it is more SSW/SW. This turning of the wind results from cyclonic flow around a climatological low-pressure area over the Weddell Sea. The strongest winds also blow from these dominant directions, resulting from the above-mentioned cyclonic flow in combination with the topographic constraint of the mountains to the west. The weakest winds are blowing from the NW and E at both stations.

We computed cloud optical thickness using an algorithm that relates the observed SW_{\downarrow} to the theoretical clear-sky value of SW_{\downarrow} (Fitzpatrick et al., 2004), extrapolated to the winter months using a good correlation between cloud optical thickness and the longwave radiation balance in summer (Kuipers Munneke et al., 2010). The cloud cover at AWS 15 is generally optically thicker than at AWS 14 (Fig. 4). At both locations, cloud cover is optically thinnest in late winter and spring, when the atmosphere is coldest and sea-ice cover over the Weddell Sea reduces evaporation of sea water.

3.2 Surface energy budget

The SEB model performance is tested by comparing the observed surface temperature (computed with LW_{\uparrow}) to the surface temperature found by the model to close the surface energy budget (Fig. 5). Based on hourly (daily) means, the difference is 0.44 (0.44) K for AWS 14, with a RMS difference of 1.40 (1.08) K. At AWS 15, the difference based on hourly (daily) means is 0.19 (0.19) K, with a RMS difference of 1.48 (1.10) K.

Monthly mean values of all SEB components are presented in Fig. 6 for both locations. Seasonal and annual averages of all SEB components are shown in Table 2. At both sites, there is a clear annual cycle in all the SEB components. In winter, SW_{net} becomes almost zero (JJA: 1.0 W m^{-2}). The surface starts cooling, so that LW_{\uparrow} decreases and the net longwave budget becomes less negative (JJA: -9.8 W m^{-2}). The latent heat flux becomes small in winter (JJA: 1.0 W m^{-2}), since the low air temperature reduces near-surface specific moisture gradients, inhibiting effective heat loss through sublimation. In the winter of 2009, the sensible heat flux is also low or slightly positive, indicating that there is some vertical transport of warmer air towards the surface

Larsen C ice shelf meteorology

P. Kuipers Munneke et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

through turbulent motion. In the winter of 2010, the sensible heat flux attains higher values, which is caused by frequent advection of warmer air (see air temperature record in Fig. 2). This means that more heat is transported to the surface. The resulting increased surface temperature decreases the net longwave radiation, restoring the surface energy balance.

In summer, net shortwave radiation is the most prominent source of energy – it is balanced by a net longwave cooling (DJF value: -21.1 W m^{-2}), and by a negative latent heat flux (DJF: -9.6 W m^{-2}) indicating a steady sublimation. Moreover, a slightly negative sensible heat (DJF: -5.0 W m^{-2}) is visible in summer, indicating that the surface layer becomes unstable and convective in nature. We will discuss this feature in more detail in Sect. 3.4. The negative ground heat flux (DJF: -2.1 W m^{-2}) indicates that the snowpack is heating up. A considerable amount of energy is consumed by melting (DJF: 8.7 W m^{-2}).

There are some small differences in the SEB of the two locations. First of all, the net shortwave radiation is somewhat larger at AWS 14 than at AWS 15. This is caused by a somewhat higher albedo at AWS 15 (0.88 averaged over the entire period compared to 0.85 at AWS 14). The higher albedo is caused by the thicker cloud cover, which in itself also is a cause for reduced incoming shortwave radiation. In a relative sense, the largest difference between the two locations is the melt flux: averaged over the year, the melt energy flux at AWS 14 is more than twice that at AWS 15. The reduced melt is also a cause for a higher albedo at AWS 15. The lower net radiative flux at AWS 15 results in slightly less negative ground and latent heat fluxes in summer.

At both locations, November 2010 is characterized by a relatively strongly negative net longwave flux, a large net shortwave flux, and a remarkably high melt flux. It turns out that an unusually persistent westerly föhn wind descended from the Antarctic Peninsula over the ice shelf for much of the month. In order to illustrate the effect of such föhn winds on the SEB and near-surface meteorology, we give special attention to this period in Sect. 3.5.

**Larsen C ice shelf
meteorology**

P. Kuipers Munneke et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



3.3 Radiation penetration, snow temperature, and melt

The role of shortwave radiation penetration in the snowpack is to deliver energy below the surface, enabling a more rapid transfer of energy into the snow (Brandt and Warren, 1993; Kuipers Munneke et al., 2009). This effect is more pronounced in ice than in snow (Van den Broeke et al., 2008). Shortwave radiation penetration depends on the size of the snow grains: the larger the grains, the more photons are scattered in the forward direction (i.e. into the snowpack), leading to more absorption of shortwave radiation inside the snowpack. Inclusion of this process into the SEB has two important consequences: first, the uppermost snow layers heat up more quickly in summer. Second, the absorption of shortwave radiation in layers below the surface allows for a larger contribution of subsurface melting. The partitioning of melt into surface melt and subsurface melt will change when shortwave radiation penetration is allowed. Moreover, the total melt (the sum of surface and subsurface melt) increases.

To illustrate these points, we show snow temperature at 0.5 m (Fig. 7) and melt fluxes (Fig. 8) for three model settings (at AWS 14): no radiation penetration, and radiation penetration with 0.1 and 0.5 mm radius snow grains. The effect on snow temperature at 0.5 m depth is clearly visible (Fig. 7): when more radiation penetration into the snowpack is allowed, the snow remains at the melting point for a longer period. A comparison with observed snow temperature (interpolated to 0.5 m) weakly suggests that radiation penetration with a snow grain size of 0.1 mm is the most appropriate choice: without radiation penetration, the temperature drops below melting two weeks too early, while with stronger radiation penetration, the melting point is attained too long. In the remainder of March 2011, the observed signal deviates somewhat from the modeled ones (both in gradient and shape) which is likely due to a simplified representation of snow density and refreezing in the model.

The implication for the melt budget is clearly observed in Fig. 8. Without accounting for radiation penetration, the total melt (green solid line) equals the surface melt, since subsurface melt is zero. Radiation penetration leads to smaller surface melt (dashed lines), but the increase in subsurface melt (dotted lines) is larger than the decrease

Larsen C ice shelf meteorology

P. Kuipers Munneke et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



in surface melt, resulting in a net higher total melt (solid lines). We have attempted to corroborate these results with the sonic height ranger for the period 4–16 January 2011 (black line in Fig. 8), assuming a density of the melted snow of 400 kg m^{-3} . We found a reasonable agreement, although a discrepancy with modeled melt remains.

The comparison is difficult since the sonic height ranger itself is not anchored to a fixed reference surface, nor is it known which part of the snowpack is actually melting (low-density surface snow or high-density subsurface snow and ice lenses). Moreover, the height change signal also includes densification and accumulation. This has a large impact on the conversion from height change to a melt rate, and thus the melt signal derived from observations has quite a large uncertainty. Therefore, it cannot be concluded from the comparison in Fig. 8 which setting for radiation penetration is the most adequate to simulate the SEB at this site. On the positive side, the comparison shows that the model computes a melt signal that corresponds well to the observed height change, both in magnitude and in variability.

3.4 Convection and a ceasing temperature inversion

An interesting feature of the summertime SEB at both locations is the frequent occurrence of a negative sensible heat flux. This indicates that the surface layer, normally characterized by a temperature inversion in the polar regions, breaks down and gives way to convective turbulent motion. Convection occurs so frequently that the monthly-mean sensible heat flux becomes negative in the summer months (Fig. 6). We study this phenomenon in more detail in this section.

A typical Antarctic boundary layer is characterized by a stable stratification and a temperature deficit at the surface. In winter, this temperature deficit is driven by a radiation deficit at the surface. In response, sensible heat is transported towards the surface by turbulent motion in the atmosphere. In summer, a temperature inversion can still exist, but then it is usually limited to nighttime, or driven by advection of warm air over a melting surface, the temperature of which is limited to the melting point. This situation also leads to a sensible heat flux directed towards the surface.

Larsen C ice shelf meteorology

P. Kuipers Munneke et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Larsen C ice shelf meteorology

P. Kuipers Munneke et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



At the Larsen C ice shelf, we find that heat is advected away from the surface during the summer, indicative of a convective surface layer. This has also been observed on the Brunt ice shelf at Halley Station (King et al., 2006). In Fig. 9, we zoom in on a period of four days in January 2011 illustrating the typical circumstances under which this convection develops. During this period, manual observations and radiosonde data are available from a temporary research camp at the AWS 14 site. In the upper panel of Fig. 9, we see that during daytime, the surface temperature exceeds the 2-m temperature by up to 2–3 °C. There is little wind, blowing from the S to E. This wind advects humid and cloudy air from the Weddell Sea: manual weather observations made during this period reveal a mean cloud cover of 7.7 oktas ($n = 32$) with a cloud base at 1–1.5 km above the surface. The cloud cover results in only a slightly negative net longwave flux, making the sum of net longwave and shortwave fluxes on the order of 50–70 W m⁻². This radiative heating of the snow surface is compensated by melting and/or by upward turbulent fluxes of heat and moisture (sublimation). The vertical extent of this convective layer is determined when plotting a vertical profile of potential temperature θ obtained by a radiosonde (Fig. 10). Negative values of $d\theta/dz$ are a sign of an unstable boundary layer, and extend up to only 50 m above the surface. Theoretically, an air parcel can be lifted to the level at which the potential temperature equals the surface potential temperature, which is at about 1150 m, which is the possible depth of convection.

3.5 November 2010: strong westerlies and intense melting

In the two-year time series discussed in this paper, November 2010 stands out as a peculiar month, with low net longwave radiation, high net shortwave radiation, and significant melt occurring already in the austral spring. Using the regional atmospheric model RACMO2 at 27-km horizontal resolution (Lenaerts et al., 2011), which is forced at the lateral boundaries by ERA-Interim reanalysis data, we found persistent westerly winds flowing over the Antarctic Peninsula for a period of 15 days (2–18 November 2010). Moderate westerlies impinging on the Antarctic Peninsula mountains have been shown

to cause a northerly flow on the eastern side of the Peninsula (Orr et al., 2004), advecting warm air over the ice shelves. When the westerly flow becomes even stronger, the air flows over the mountains, causing a föhn effect on their lee side. Under these conditions, the local meteorological conditions at Larsen C ice shelf are characterized by cloudless skies and a north-to-westerly advection of dry and warm air.

To illustrate the effect of this warm northerly flow and westerly föhn winds, we zoom in at the period 10–25 November 2010. This period shows two very contrasting regimes: between 10 and 18 November, the flow was predominantly from the north and west; between 19 and 25 November, the wind was from the south and east. The near-surface wind and 850 hPa geopotential heights typical for these contrasting regimes are illustrated in Fig. 11. In Fig. 12, we show the SEB components at AWS 14 between 10 and 25 November 2010, along with the observed daily mean wind direction. Additionally, Table 3 shows the means of meteorological variables and SEB components for these two contrasting periods. In the first half of this period, strong westerly flow west of the peninsula travels over the mountains onto the ice shelves. The descending motion east of the mountains warms and dries the air. The resulting clear-sky conditions are reflected in the SEB as low net longwave radiation (-49.9 W m^{-2}), and high net shortwave radiation (63.1 W m^{-2}). Since the snow surface is at the melting point during the day and the air temperature is above 0°C , the sensible heat flux is significantly positive (16.5 W m^{-2}). On the other hand, the dry air allows for significant sublimation, reflected in the SEB as a negative latent heat flux (-13.0 W m^{-2}). This result is consistent with the anticorrelation of sensible and latent heat fluxes found over the Larsen Ice Shelf using airborne observations of the stable boundary layer (King et al., 2008). Overall, the increase of the shortwave and sensible heat fluxes outweighs the decrease of the longwave and latent heat fluxes. As a result, a relatively large amount of energy is available for snow melt (29.6 W m^{-2}).

From 19 November onwards, the atmospheric flow comes from the east and south, bringing cold and cloudy air in from the Weddell Sea and colder regions to the south. This is immediately reflected in the SEB: net shortwave fluxes are much smaller, and

Larsen C ice shelf meteorology

P. Kuipers Munneke et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



sensible heat fluxes are now negative (-6.8 W m^{-2}), indicating weak convection over the ice shelf. The negative sensible heat flux is explained by the advection of cold air over a warm and wet snowpack that first needs to refreeze before it can start to cool.

4 Conclusions

Two years of data collected by the two AWSs on Larsen C ice shelf offer a preliminary insight into the SEB typical for this region. The time series are too short to present a climatology, but several interesting aspects of the SEB can be highlighted. First of all, the importance of modelling subsurface absorption of solar radiation and its implications on total melt have been discussed. However, it remains difficult to determine the best settings for this part of the model. Perhaps a seasonally varying density profile and snow grain size would improve the simulation of the snowpack. A radical step would be to incorporate snow densification and snow metamorphism into the SEB model, but we did not pursue this for the present study.

Also, we have shown that calm and cloudy conditions dominate in summer, and can give rise to long continuous periods of daytime surface melt, as well as daytime convection in the surface layer. On quite a number of days in summer, the air temperature at 2 m above the surface remains below 0°C while the surface is melting. Interestingly, this means that air temperature measured at some distance above the surface, is not necessarily a good indicator of melt. Models or algorithms that depend solely on air temperature input for the computation of melt (like positive degree-day models) will likely underestimate the amount of melt when the air temperature input is recorded well above the surface. This stresses the need not only to model the full SEB, but also to measure the SEB with an AWS, including the individual radiation components.

Although the analysis is limited to only two locations, the topographic homogeneity makes that the results presented here are representative for a large area. Especially in the north-south direction, there are some notable climate gradients over Larsen C ice shelf, however: in several surface mass balance components (Lenaerts et al., 2011), in

Larsen C ice shelf meteorology

P. Kuipers Munneke et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



5 firm air conditions (Holland et al., 2011), and in number of melt days (Tedesco and Monaghan, 2009). These gradients are a result of gradients in the SEB components, and ultimately of gradients in meteorological variables like temperature and cloud cover. Already between AWS 14 and 15, separated by a mere 70 km, differences in melt and other SEB components can be seen. It would be interesting and useful to compare observations from other locations to get insight in the spatial variability of near-surface climate and SEB. To that end, BAS and UU/IMAU extended the coverage of the Antarctic Peninsula ice shelves northward by installing an AWS at Scar Inlet, the remains of the Larsen B ice shelf (AWS 17, shown on the map in Fig. 1) in February 2011.

10 Lastly, we illustrated the important effect that westerly flow crossing the Antarctic Peninsula mountain range has on the SEB and most importantly on the melt flux. Over the past 50 years, the Antarctic Peninsula has seen a large temperature increase (Turner et al., 2005), linked to a positive trend of the Southern Annular Mode (SAM, the principal mode of variability in the atmospheric circulation of the Southern Hemisphere extratropics, Marshall, 2003). Given the marked increase in westerly flow during this period (Marshall, 2002), the melt volume has likely increased over these 50 years. Since the late 1980s however, melt trends have been decreasing in the Antarctic Peninsula (Tedesco, 2009; Kuipers Munneke et al., 2011). The future trend of temperature and melt is uncertain, but continued observation and modelling is crucial to understand the repercussions of future climate change on the surface energy budget, and ultimately on the stability of the ice shelves in the Antarctic Peninsula.

25 *Acknowledgements.* We would like to thank Ian Hey, Ian Potten, Nick Alford (BAS) and Wim Boot (IMAU) for help with installation and maintenance of the two IMAU Automatic Weather Stations. Henk Snellen and Marcel Portanger are thanked for help with the development of the Automatic Weather Stations and data transfer. Phil Anderson (BAS) is thanked for his contribution in the 2011 field experiment at the AWS 14 site. This research is partly funded by NWO/ALW grant 818.01.016. The radiosonde measurements presented in Fig. 10 were carried out as part of the project “Orographic Flows and the Climate of the Antarctic Peninsula” (OFCAP), funded by the UK Natural Environmental Research Council (NERC).

Larsen C ice shelf meteorology

P. Kuipers Munneke et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



References

- Andreas, E. L.: A theory for the scalar roughness and the scalar transfer coefficients over snow and sea ice, *Bound.-Lay. Meteorol.*, 38, 159–184, 1987. 2670
- van As, D., van den Broeke, M. R., Reijmer, C. H., and van de Wal, R. S. W.: The summer surface energy balance of the high Antarctic Plateau, *Bound.-Lay. Meteorol.*, 115, 289–317, 2005. 2670
- Brandt, R. E. and Warren, S. G.: Solar-heating rates and temperature profiles in Antarctic snow and ice, *J. Glaciol.*, 39(131), 99–110, 1993. 2674
- van den Broeke, M. R.: Strong surface melting preceded collapse of Antarctic Peninsula ice shelf, *Geophys. Res. Lett.*, 32, L12815, 2005. 2667
- van den Broeke, M. R., van As, D., Reijmer, C. H., and van de Wal, R. S. W.: Assessing and improving the quality of unattended radiation observations in Antarctica, *J. Atmos. Ocean. Tech.*, 21, 1417–1431, 2004. 2669
- van den Broeke, M., Smeets, P., Ettema, J., van der Veen, C., van de Wal, R., and Oerlemans, J.: Partitioning of melt energy and meltwater fluxes in the ablation zone of the west Greenland ice sheet, *The Cryosphere*, 2, 179–189, doi:10.5194/tc-2-179-2008, 2008. 2674
- Cook, A. J. and Vaughan, D. G.: Overview of areal changes of the ice shelves on the Antarctic Peninsula over the past 50 years, *The Cryosphere*, 4, 77–98, doi:10.5194/tc-4-77-2010, 2010. 2666
- Fitzpatrick, M. F., Brandt, R. E., and Warren, S. G.: Transmission of solar radiation by clouds over snow and ice surfaces: a parameterization in terms of optical depth, solar zenith angle, and surface albedo, *J. Climate*, 17, 266–275, 2004. 2672
- Giesen, R. H., Andreassen, L. M., van den Broeke, M. R., and Oerlemans, J.: Comparison of the meteorology and surface energy balance at Storbreen and Midtdalsbreen, two glaciers in southern Norway, *The Cryosphere*, 3, 57–74, doi:10.5194/tc-3-57-2009, 2009. 2669
- Holland, P. R., Corr, H. F. J., Pritchard, H. D., Vaughan, D. G., Arthern, R. J., Jenkins, A., and Tedesco, M.: The air content of Larsen Ice Shelf, *Geophys. Res. Lett.*, 38, L10503, doi:10.1029/2011GL047245, 2011. 2667, 2679
- King, J. C., Argentini, S. A., and Anderson, P. S.: Contrasts between the summertime surface energy balance and boundary layer structure at Dome C and Halley stations, Antarctica, *J. Geophys. Res.*, 111, D02105, 2006. 2676
- King, J. C., Lachlan-Cope, T. A., Ladkin, R. S., and Weiss, A.: Airborne measurements in

Larsen C ice shelf meteorology

P. Kuipers Munneke et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Larsen C ice shelf meteorology

P. Kuipers Munneke et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



the stable boundary layer over the Larsen Ice Shelf, Antarctica, Bound.-Lay. Meteorol., 127, 413–428, doi:10.1007/s10546-008-9271-4, 2008. 2677

Kuipers Munneke, P., van den Broeke, M. R., Reijmer, C. H., Helsen, M. M., Boot, W., Schneebeli, M., and Steffen, K.: The role of radiation penetration in the energy budget of the snowpack at Summit, Greenland, The Cryosphere, 3, 155–165, doi:10.5194/tc-3-155-2009, 2009. 2669, 2670, 2674

Kuipers Munneke, P., Reijmer, C. H., and van den Broeke, M. R.: Assessing the retrieval of cloud properties from radiation measurements over snow and ice, Int. J. Climatol., 31, 756–769, doi:10.1002/joc.2114, 2010. 2672

Kuipers Munneke, P., van den Broeke, M. R., Picard, G., Lenaerts, J. T. M., and van Meijgaard, E.: Estimating Antarctic snowmelt volume and its recent trend, in prep., 2011. 2679
Lenaerts, J. T. M., van den Broeke, M. R., van de Berg, W. J., van Meijgaard, E., and Kuipers Munneke, P.: A new high-resolution surface mass balance map of Antarctica (1989–2009) based on regional climate modelling, Geophys. Res. Lett., in review, 2011. 2676, 2678

Marshall, G. J.: Analysis of recent circulation and thermal advection in the Northern Antarctic Peninsula, Int. J. Climatol., 22, 1557–1567, 2002. 2679

Marshall, G. J.: Trends in the Southern Hemisphere annular mode from observations and reanalyses, J. Climate, 16, 4134–4143, 2003.

Orr, A., Cresswell, D., Marshall, G. J., Hunt, J. C. R., Sommeria, J., Wang, C. G., and Light, M.: A “low-level” explanation for the recent large warming trend over the Western Antarctic Peninsula involving blocked winds and changes in zonal circulation, Geophys. Res. Lett., 31, L06204, doi:10.1029/2003GL019160, 2004. 2677

Rignot, E., Casassa, G., Gogineni, P., Krabill, W., Rivera, A., and Thomas, R.: Accelerated ice discharge from the Antarctic Peninsula following the collapse of Larsen B ice shelf, Geophys. Res. Lett., 31, L18401, doi:10.1029/2004GL020697, 2004. 2666

Rott, H., Skvarça, P., and Nagler, T.: Rapid collapse of Northern Larsen Ice Shelf, Antarctica, Science, 271, 788–792, 1996. 2666

Rott, H., Müller, F., Nagler, T., and Floricioiu, D.: The imbalance of glaciers after disintegration of Larsen-B ice shelf, Antarctic Peninsula, The Cryosphere, 5, 125–134, doi:10.5194/tc-5-125-2011, 2011. 2666

Scambos, T. A., Hulbe, C., Fahnestock, M., and Bohlander J.: The link between climate warming and break-up of ice shelves in the Antarctic Peninsula, J. Glaciol., 46, 516–530, 2000. 2667

Scambos, T. A., Bohlander, J. A., Shuman, C. A., and Skvarça, P.: Glacier acceleration and thinning after ice shelf collapse in the Larsen B embayment, Antarctica, *Geophys. Res. Lett.*, 31, L18402, doi:10.1029/2004GL020670, 2004. 2666

5 Smeets, C. J. P. P.: Assessing unspirated temperature measurements using a thermocouple and a physically based model, in: *The mass budget of Arctic glaciers*, IASC Working group on Arctic Glaciology, 99–101, Utrecht, The Netherlands, 2006. 2669

Tedesco, M.: Assessment and development of snowmelt retrieval algorithms over Antarctica from K-band spaceborne brightness temperature (1979–2008), *Remote Sens. Environ.*, 113, 979–997, 2009. 2679

10 Tedesco, M. and Monaghan, A. J.: An updated Antarctic melt record through 2009 and its linkages to high-latitude and tropical climate variability, *Geophys. Res. Lett.*, 36, L18502, 2009. 2679

15 Turner, J., Colwell, S. R., Marshall, G. J., Lachlan-Cope, T. A., Carleton, A. M., Jones, P. D., Lagun, V., Reid, P. A., and Iagovkina, S.: Antarctic climate change during the last 50 years, *Int. J. Climatol.*, 25, 279–294, 2005. 2666, 2679

Larsen C ice shelf meteorology

P. Kuipers Munneke et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Larsen C ice shelf meteorology

P. Kuipers Munneke et al.

Table 1. Means of meteorological variables 1 February 2009–31 January 2011 (28 January 2011 for AWS 15). ANN = average over all months.

| AWS 14 | | ANN | DJF | MAM | JJA | SON |
|-------------------------|--------------------|-------|-------|-------|-------|-------|
| Air temperature (2 m) | °C | −15.5 | −4.2 | −20.1 | −24.0 | −13.4 |
| Specific humidity (2 m) | g kg ^{−1} | 1.33 | 2.52 | 0.84 | 0.60 | 1.41 |
| Relative humidity (2 m) | % | 94.8 | 90.6 | 98.9 | 96.3 | 93.3 |
| Surface temperature | °C | −16.1 | −4.0 | −20.7 | −25.2 | −14.2 |
| Wind speed | m s ^{−1} | 4.5 | 4.4 | 3.9 | 4.3 | 5.4 |
| Air pressure | hPa | 985.1 | 986.4 | 985.5 | 988.7 | 979.8 |
| AWS 15 | | ANN | DJF | MAM | JJA | SON |
| Air temperature (2 m) | °C | −15.8 | −4.5 | −20.2 | −24.0 | −13.7 |
| Specific humidity (2 m) | g kg ^{−1} | 1.30 | 2.49 | 0.82 | 0.60 | 1.36 |
| Relative humidity (2 m) | % | 94.7 | 90.9 | 98.0 | 96.3 | 93.3 |
| Surface temperature | °C | −16.7 | −4.6 | −21.0 | −25.4 | −14.8 |
| Wind speed | m s ^{−1} | 4.3 | 4.5 | 3.6 | 4.0 | 5.0 |
| Air pressure | hPa | 985.1 | 986.7 | 985.5 | 988.5 | 979.7 |

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[I◀](#)
[▶I](#)
[◀](#)
[▶](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)


Larsen C ice shelf meteorology

P. Kuipers Munneke et al.

Table 2. Means of SEB components 1 February 2009–31 January 2011 for AWS 14, in W m^{-2} . ANN = average over all months.

| AWS 14 | ANN | DJF | MAM | JJA | SON |
|----------------------|-------|-------|------|------|-------|
| Net shortwave | 17.8 | 38.0 | 6.2 | 1.0 | 26.5 |
| Subsurface shortwave | 4.1 | 9.2 | 1.4 | 0.2 | 5.8 |
| Net longwave | -15.7 | -21.1 | -8.6 | -9.8 | -23.3 |
| Sensible heat flux | 1.5 | -5.0 | 0.3 | 7.1 | 3.6 |
| Latent heat flux | -3.2 | -9.6 | -0.2 | 1.0 | -4.0 |
| Ground heat flux | 0.0 | -2.1 | 3.4 | 1.5 | -2.8 |
| Melt energy | 2.8 | 8.7 | 0.0 | 0.0 | 2.5 |

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[I◀](#)
[▶I](#)
[◀](#)
[▶](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)


Larsen C ice shelf meteorology

P. Kuipers Munneke et al.

Table 3. Means of SEB components and meteorological variables for the two contrasting periods 10–18 November, and 19–25 November 2010 for AWS 14.

| AWS 14 | | 10–18 Nov | 19–25 Nov |
|-------------------------|-------------------|-----------|-----------|
| Air temperature (2 m) | °C | −0.4 | −5.6 |
| Relative humidity (2 m) | % | 79 | 92 |
| Wind speed | m s ^{−1} | 6.0 | 5.6 |
| Net shortwave | | 63.1 | 37.8 |
| Subsurface shortwave | | 13.3 | 8.8 |
| Net longwave | | −49.9 | −25.0 |
| Sensible heat flux | W m ^{−2} | 16.5 | −6.8 |
| Latent heat flux | | −13.0 | −11.1 |
| Ground heat flux | | −7.8 | −3.2 |
| Melt energy | | 29.6 | 0.9 |

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Larsen C ice shelf meteorology

P. Kuipers Munneke et al.




Fig. 1. Map of the Antarctic Peninsula central section. UU/IMAU AWS network locations shown in green; partner AWS network shown in orange; Rothera Research Station shown in blue.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion






Fig. 2. Daily-averaged air temperature, specific humidity, air pressure, and wind speed for the measurement period. AWS 14 data are shown in black with circles, AWS 15 is shown in red.

Larsen C ice shelf meteorology

P. Kuipers Munneke et al.

Title Page

Abstract Introduction

Conclusions References

Tables Figures

◀ ▶

◀ ▶

Back Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Larsen C ice shelf meteorology

P. Kuipers Munneke et al.




Fig. 3. Wind roses of daily mean wind speed and direction for the period 1 February 2009–31 January 2011, for AWS 14 (left) and 15 (right). The percentages refer to the frequency of occurrence.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**Larsen C ice shelf
meteorology**

P. Kuipers Munneke et al.




Fig. 4. Monthly means of cloud optical thickness (see text) at AWS 14 (black) and 15 (red).

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

**Larsen C ice shelf
meteorology**

P. Kuipers Munneke et al.




Fig. 5. Model performance check by comparing observed values of surface temperature to computed values, for **(a)** AWS 14; and **(b)** AWS 15. Red dots are hourly values, whereas black dots are daily-mean values.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[I◀](#)[▶I](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Larsen C ice shelf meteorology

P. Kuipers Munneke et al.




Fig. 6. Monthly means of surface energy budget components (in $W m^{-2}$) for (a) AWS 14; and (b) AWS 15.

Discussion Paper | Discussion Paper | Discussion Paper | Discussion Paper | Discussion Paper

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**Larsen C ice shelf
meteorology**

P. Kuipers Munneke et al.




Fig. 7. Comparison between observed and modeled snow temperature at 0.5 m depth at AWS 14, for different model settings of radiation penetration. Observed snow temperature (interpolated to 0.5 m) shown in black; model without radiation penetration in green; model with radiation penetration (0.1 mm snow grain radius) in blue; model with radiation penetration (0.5 mm snow grain radius) in red.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[I◀](#)[▶I](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Larsen C ice shelf
meteorology

P. Kuipers Munneke et al.




Fig. 8. Comparison of observed cumulative height change and modeled cumulative melt at AWS 14, for different model settings of radiation penetration. Observed height change (converted to mm w.e. assuming a density of 400 kg m^{-3}) shown in black; model without radiation penetration in green; model with radiation penetration (0.1 mm snow grain radius) in blue; model with radiation penetration (0.5 mm snow grain radius) in red. Dotted lines indicate subsurface melt; dashed lines indicate surface melt; and solid lines indicate total melt (= surface + subsurface melt).

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Larsen C ice shelf
meteorology

P. Kuipers Munneke et al.




Fig. 9. Meteorological conditions (upper panel) and selected SEB components (lower panel) during a four-day period showing typical daytime convection.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)




Fig. 10. Vertical profile of potential temperature (in K) from a radiosonde launched on 22 January 2011 at 15:19 LT (= 18:19 UTC). The horizontal dashed line indicates the height at which $d\theta/dz$ changes sign, while the vertical dashed line illustrates that possible convection extends up to 1150 m, the height at which potential temperature equals surface potential temperature.

**Larsen C ice shelf
meteorology**

P. Kuipers Munneke et al.




Fig. 11. Surface wind speed and direction (vectors) and the height of the 850 hPa level from RACMO2, for **(a)** 15 November 2010 and **(b)** 19 November 2010. The location of AWS 14 is denoted with the black dot.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**Larsen C ice shelf
meteorology**

P. Kuipers Munneke et al.




Fig. 12. Selected SEB components between 10 and 26 November 2010. Daily-mean wind direction is shown at the bottom of the panel.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)