



## Nationwide age references for sitting height, leg length, and sitting height/height ratio, and their diagnostic value for disproportionate growth disorders

A M Fredriks, S van Buuren, W J M van Heel, R H M Dijkman-Neerincx, S P Verloove-Vanhorick and J M Wit

*Arch. Dis. Child.* 2005;90;807-812; originally published online 29 Apr 2005;  
doi:10.1136/adc.2004.050799

---

Updated information and services can be found at:  
<http://adc.bmjournals.com/cgi/content/full/90/8/807>

---

*These include:*

### References

This article cites 14 articles, 2 of which can be accessed free at:  
<http://adc.bmjournals.com/cgi/content/full/90/8/807#BIBL>

4 online articles that cite this article can be accessed at:  
<http://adc.bmjournals.com/cgi/content/full/90/8/807#otherarticles>

### Rapid responses

You can respond to this article at:  
<http://adc.bmjournals.com/cgi/eletter-submit/90/8/807>

### Email alerting service

Receive free email alerts when new articles cite this article - sign up in the box at the top right corner of the article

---

### Topic collections

Articles on similar topics can be found in the following collections

[Other Rheumatology](#) (1705 articles)  
[Other Pediatrics](#) (1857 articles)

---

### Notes

---

To order reprints of this article go to:  
<http://www.bmjournals.com/cgi/reprintform>

To subscribe to *Archives of Disease in Childhood* go to:  
<http://www.bmjournals.com/subscriptions/>

## ORIGINAL ARTICLE

# Nationwide age references for sitting height, leg length, and sitting height/height ratio, and their diagnostic value for disproportionate growth disorders

A M Fredriks, S van Buuren, W J M van Heel, R H M Dijkman-Neerincx, S P Verloove-Vanhorick, J M Wit



*Arch Dis Child* 2005;90:807–812. doi: 10.1136/adc.2004.050799

**Aims:** To obtain age references for sitting height (SH), leg length (LL), and SH/H ratio in the Netherlands; to evaluate how SH standard deviation score (SDS), LL SDS, SH/H SDS, and SH/LL SDS are related to height SDS; and to study the usefulness of height corrected SH/H cut-off lines to detect Marfan syndrome and hypochondroplasia.

**Methods:** Cross-sectional data on height and sitting height were collected from 14 500 children of Dutch origin in the age range 0–21 years. Reference SD charts were constructed by the LMS method. Correlations were analysed in three age groups. SH/H data from patients with Marfan syndrome and genetically confirmed hypochondroplasia were compared with height corrected SH/H references.

**Results:** A positive association was observed between H SDS, SH SDS, and LL SDS in all age groups. There was a negative correlation between SH/H SDS and height SDS. In short children with a height SDS  $< -2$  SDS, a cut-off limit of  $+2.5$  SD leads to a more acceptable percentage of false positive results. In exceptionally tall children, a cut-off limit of  $-2.2$  SDS can be used. Alternatively, a nomogram of SH/H SDS versus H SDS can be helpful. The sensitivity of the height corrected cut-off lines for hypochondroplasia was 80% and for Marfan syndrome only 30%.

**Conclusions:** In exceptionally short or tall children, the dependency of the SH/H ratio (SDS) on height SDS has to be taken into consideration in the evaluation of body proportions. The sensitivity of the cut-off lines for hypochondroplasia is fair.

See end of article for authors' affiliations

Correspondence to:  
Prof. Dr J M Wit, Dept of Paediatrics, J6S, Leiden University Medical Center, PO Box 9600, 2300 RC Leiden, Netherlands; [j.m.wit@lumc.nl](mailto:j.m.wit@lumc.nl)

Accepted 5 February 2005  
Published Online First  
29 April 2005

In the diagnostic work-up of children with exceptionally short or tall stature, the visual inspection and objective measurement of body proportions can give important clues.<sup>1–3</sup> The usual method of judging body proportions of children is to calculate the ratio between sitting height and height (SH/H) or sitting height and leg length (SH/LL) and compare this with age references. Sitting height can also be used as a proxy of statural growth if height cannot be measured, for example because of lower limb deformities.

In short children, most chondrodystrophic syndromes (skeletal dysplasias) are characterised by short limbs. In contrast to achondroplasia, hypochondroplasia can be difficult to diagnose. Hypochondroplasia is an autosomal dominant condition characterised by a disproportionate short stature, with relatively short legs, micromelia, macrocrania, and lumbar lordosis, linked to N540K mutations in the FGFR3 gene. Other conditions, such as Down's syndrome and Turner's syndrome can also present with abnormal body proportions. On the other hand, some other syndromes associated with short stature present with a relatively short trunk. In tall children, it is important to diagnose Marfan syndrome, gonadotropin deficiency, and Klinefelter's syndrome, because of the clinical consequences. Marfan syndrome is an autosomal dominant disorder of connective tissue characterised by a disproportionate tall stature and relatively long legs. Thus, measuring body proportions provides vital diagnostic information in the work-up of growth disorders.

It is generally known that tall children have relatively long legs and vice versa.<sup>4,5</sup> Therefore, we conjecture that the interpretation of SH/H ratio should not only be based on age references, but also on height. This would theoretically improve the specificity of the cut-off lines. However, no such conditional references are available. There is also no

information available about the sensitivity of the usual cut-off lines of normality ( $\pm 2$  SDS), either corrected for age only or after an additional correction for height, in detecting the most frequent disproportionate growth disorders.


In this paper we present age references of SH, LL, and SH/H ratio for Dutch children, and show their relation with height. In addition, we compare SH/H of children with known Marfan syndrome and known hypochondroplasia with the new references in order to determine whether the usual cut-off limits in the reference charts are appropriate for detecting these disorders.

## METHODS

### Subjects

Cross-sectional data on height and sitting height were collected in the Fourth Dutch Growth Study in 1996 and 1997. A total of 14 500 children (7482 boys and 7018 girls) of Dutch origin in the age range 0–21 years were included. Sitting height was measured in 6877 boys and 6202 girls. Children with known growth disorders and those on medication known to interfere with growth were not included in the sample. Details have been described elsewhere.<sup>6</sup> The sample was nationally representative. Separately, we collected growth data of children with Marfan syndrome: four boys (of 3, 6, 9, and 13 years) and six girls (of 8 (n = 4), 12, and 16 years). Through the Laboratory of Clinical Genetics and referring physicians, we anonymously gathered data on individuals with DNA confirmed hypochondroplasia: seven children (three boys, of 4, 6, and 12 years; and four girls, of 1, 6, 10, and 12 years), and three adults (one man of 41 years; and two women 24 and 43 years). In addition we gathered data

**Abbreviations:** H, height; LL, leg length; SDS, standard deviation score; SH, sitting height



**Figure 1** Sitting height and leg length for age for (A) Dutch boys and (B) Dutch girls aged 0–21 years, indicating the 0,  $\pm 1$ ,  $\pm 2$ ,  $\pm 2.5$  SD lines.

on the family of a 10 year old girl with a confirmed HCH mutation in the *FGFR3* gene that caused a mild hypochondroplasia. None of these patients had been treated with any relevant medication at the time of measurement.

### Measurements

Length of infants, until 2 years of age, was measured to the nearest 0.1 cm in the supine position, fully extended with their heels in contact with a baseboard. Crown-rump length, a measure of trunk length, which is conceptually similar to sitting height in older children, was measured until 2 years of age while the child was lying in supine position on a measuring table. After the thighs were placed in a vertical plane, the footboard was pulled against the buttocks. From 2 years of age onward, standing height was measured to the nearest 0.1 cm by using a calibrated microtoise. Sitting height was measured by bringing the horizontal bar of the microtoise into the most superior midline of the head while the child was sitting in erect position on a flat stool or box. Arching of the back was avoided as much as possible by applying upward pressure to the mastoid processes while the child breathed deeply and held its breath during the measurement. The difference between crown-rump length and sitting height was on average +0.4 cm at 2 years of age. For the ratio crown-rump length/length and sitting height/height the difference was on average +0.03. Leg length was obtained by subtracting sitting height from height. The difference between length and crown-rump length is a corresponding estimate of leg length in infants.

### Statistical analysis

References for SH, LL, and SH/H for age were constructed with the LMS method. The distribution of the data is summarised by three spline curves, the L, M, and S, that vary in time: the Box-Cox transformation power that


converts data to normality and minimises the skewness of the dataset (L), the median (M), and the coefficient of variation (S).<sup>7</sup> The choice of the smoothing factors for the L, M, and S curves was made by creating local detrended QQ plots.<sup>8</sup> The associations between SH SDS, LL SDS, SH/H SDS, SH/LL SDS, and height SDS were calculated by (multiple) regression analyses and studied for three age groups: 0–<5 y (I), 5–<12.5 y (II), and 12.5–<21 y (III). Two strategies were used to find the optimal cut-off values for height SDS and SH/H SDS. First, an ellipse was drawn around 95% of the data points in the scatter plot of SH/H SDS against H SDS, and points that were located outside the ellipse were classified as unusual. The second method was to select H SDS <–2 or >+2 first, and within that group, we classified all points as unusual that were located at least 2 SDS units away from the regression line of SH/H SDS given H SDS.

### RESULTS

Reference SD charts for sitting height (SH) and leg length (LL) (fig 1), and sitting height/height (SH/H) for age (fig 2) were constructed for boys and girls aged 0–21 years.<sup>9</sup>

The corresponding L, M, and S data are shown in table 1. In infants SH represents 68% of the length, decreasing to 57% at 3 years of age for both sexes. During puberty, sitting height represents 52% of the height. Between 10 and 15 years a growth spurt in leg length is observed. The ratio SH/LL decreases from a mean of 2.10 in the first year to 1.05 in boys and 1.11 in girls at 20 years of age.

Table 2 shows the association between body proportions and height SDS. As expected, for both SH SDS and LL SDS a strong positive association with height SDS was found in all age groups. The correlations between SH/H (or SH/LL) SDS and height SDS were all negative and statistically significant ( $p < 0.001$ ).



**Figure 2** Sitting height/height ratio for age for (A) Dutch boys and (B) Dutch girls aged 0–21 years, indicating the 0,  $\pm 1$ ,  $\pm 2$ , 2.5 SD lines.

This is illustrated in fig 3 which presents a scatter plot of SH/H SDS versus height SDS. The equiprobable ellipse around 95% of the points shows a tendency towards decreasing SH/H SDS with increasing height SDS. Conversely, shorter children have higher SH/H ratios, thus relatively shorter legs. Data points located inside the ellipse may be considered as normal. Figure 4 shows the ellipse, the regression line, and two lines at 2 SDS units away of the regression line. This figure can be used as a nomogram to assess for a given height SDS the normal range of SH/H SDS.

To explore if this nomogram is a useful tool to distinguish patients with Marfan syndrome from constitutionally tall children, or patients with hypochondroplasia from idiopathic short stature, one SH/H observation per patient from these groups of patients was plotted in the figure. The purpose was to find cut-off limits that detect disproportion. Only in 3 of 10 patients with Marfan syndrome was SH/H located below the conditional  $-2$  SD line, so this cut-off criterion has a sensitivity of only 30%. In 4 of 10 patients SH/H SDS was below the unconditional  $-2$  SD line. The ellipse criterion performed better: 6 of 10 patients with Marfan syndrome were located outside the ellipse. When the conditional  $-2$  SD line is taken as diagnostic criterion, the likelihood ratio of a positive test (LR+) is  $0.3/0.02 = 15$ , and the likelihood ratio of a negative test (LR-)  $0.7/0.98 = 0.7$ . With regard to hypochondroplasia, a total of 8 out of 10 cases were located above the conditional  $+2$  SDS line, corresponding to a sensitivity of 80%. This results in an LR+ of 40 and an LR- of 0.2. We observed that here the ellipse also performed better: all hypochondroplasia cases were located outside the ellipse. Figure 5 shows sitting height/height data of the members of a family with HCH due to an FGFR3 gene mutation on the maternal side. The sitting height/height index of the index case and three of her female relatives is shown in fig 5A, and of her brother and uncle in fig 5B. The HCH mutation in the FGFR3 gene (in codon 540: substitution

of asparagine by serine) caused a mild hypochondroplasia with a variable expression pattern. All affected family members had short stature (height  $< -2$  SDS) and a mild increased sitting height/height index, indicating a disproportionate short stature with relatively short legs.

The negative correlation between SH/H SDS and H SDS signifies that for short or tall children the usual cut-off limits for body proportions ( $\pm 2.0$  SDS) would result in considerable percentages of children who would be considered as disproportionate. This is shown in table 3. If one would strive for a specificity of about 98%, the cut-off limit of SH/H SDS for short children would be  $+2.5$  SDS, and for tall children  $-2.2$ .

## DISCUSSION

This study provides new reference charts for Dutch children for SH, LL, and SH/H in relation to age. The SH/H ratio changed from 0.68 infancy to 0.52 in adolescence, indicating that in the prepubertal years growth occurs more in the limbs than in the trunk. This is also shown by the decreasing SH/LL ratio from 2.10 to 1.08 at 10 years of age. The use of a ratio might be misleading when two ratios might be equal while the nominator and denominator might be different. This effect is even stronger when a change in the nominator automatically leads to a change in the denominator, for example by using SH/LL ratio. To minimise this risk, we chose sitting height/height for age reference charts.

During the past two centuries in the Netherlands, as well as in many more industrialised countries, a positive secular growth change has been observed.<sup>6</sup> Various studies have shown that the positive secular change is mainly due to increase in leg length rather than in trunk length.<sup>1 4 10–13</sup> Tanner reported that between the 1950s and 1980s Japanese height increased solely due to change in leg length. Sitting height showed no increase, so the trunk/leg proportions changed much more towards the proportions of North Europeans, though their final height was still 1 SD lower.<sup>14</sup>

**Table 1** L, M, and S values for sitting height (SH) (cm), leg length (LL) (cm), and sitting height/height (SH/H) ratio for boys and girls of Dutch origin in the age range 0–21 years

Age (y)	SH			LL			SH/H		
	L	M	S	L	M	S	L	M	S
<b>Boys</b>									
0.25	3.072	41.691	0.045	1.061	19.376	0.080	5.220	0.682	0.034
0.50	3.087	45.669	0.044	0.936	22.136	0.078	5.220	0.671	0.034
0.75	3.092	48.035	0.044	0.824	24.605	0.077	5.220	0.660	0.034
1.0	3.087	49.710	0.044	0.729	26.922	0.075	5.220	0.648	0.034
2.0	3.030	53.599	0.046	0.536	35.711	0.071	5.220	0.604	0.034
3.0	2.971	55.952	0.046	0.513	42.289	0.067	5.220	0.572	0.033
4.0	2.886	58.741	0.047	0.530	47.110	0.064	5.220	0.555	0.033
5.0	2.768	61.831	0.047	0.558	51.291	0.062	5.220	0.546	0.031
6.0	2.624	64.916	0.047	0.588	55.143	0.061	5.220	0.541	0.030
7.0	2.474	67.784	0.046	0.617	58.826	0.060	5.220	0.536	0.029
8.0	2.331	70.446	0.047	0.644	62.405	0.060	5.220	0.531	0.028
9.0	2.212	72.642	0.047	0.672	65.720	0.059	5.220	0.525	0.027
10.0	2.115	74.502	0.048	0.703	68.825	0.059	5.220	0.520	0.026
11.0	2.023	76.435	0.048	0.742	71.947	0.059	5.220	0.516	0.025
12.0	1.931	78.893	0.049	0.791	75.332	0.059	5.220	0.513	0.025
13.0	1.868	81.971	0.050	0.851	79.180	0.059	5.220	0.509	0.025
14.0	1.871	85.482	0.050	0.910	82.832	0.058	5.220	0.508	0.026
15.0	1.925	88.994	0.047	0.957	85.618	0.057	5.220	0.510	0.026
16.0	1.985	91.555	0.044	0.990	87.476	0.057	5.220	0.512	0.027
17.0	2.024	93.017	0.042	1.007	88.456	0.057	5.220	0.513	0.027
18.0	2.047	93.854	0.041	1.013	88.821	0.057	5.220	0.515	0.028
19.0	2.059	94.311	0.040	1.014	88.885	0.057	5.220	0.516	0.028
20.0	2.062	94.432	0.040	1.016	88.954	0.057	5.220	0.516	0.029
21.0	2.060	94.654	0.040	1.019	89.144	0.057	5.220	0.513	0.029
<b>Girls</b>									
0.25	1.510	40.669	0.045	0.707	18.895	0.079	4.720	0.683	0.036
0.50	1.445	44.742	0.045	0.673	21.518	0.077	4.720	0.672	0.036
0.75	1.386	46.962	0.046	0.650	23.999	0.075	4.720	0.660	0.036
1.0	1.336	48.719	0.046	0.641	26.364	0.074	4.720	0.648	0.036
2.0	1.234	52.428	0.048	0.719	35.261	0.069	4.720	0.601	0.036
3.0	1.194	54.615	0.049	0.785	42.145	0.066	4.720	0.568	0.035
4.0	1.164	57.761	0.049	0.764	46.801	0.063	4.720	0.553	0.034
5.0	1.147	61.155	0.050	0.713	50.650	0.061	4.720	0.546	0.032
6.0	1.144	64.146	0.050	0.660	54.452	0.060	4.720	0.541	0.031
7.0	1.153	66.810	0.050	0.625	58.309	0.059	4.720	0.536	0.029
8.0	1.176	69.573	0.050	0.624	62.011	0.058	4.720	0.531	0.028
9.0	1.217	72.287	0.049	0.662	65.409	0.058	4.720	0.526	0.027
10.0	1.277	74.730	0.049	0.738	68.658	0.057	4.720	0.522	0.028
11.0	1.357	77.178	0.048	0.850	71.969	0.057	4.720	0.519	0.027
12.0	1.471	80.100	0.047	0.982	75.247	0.056	4.720	0.516	0.027
13.0	1.610	83.126	0.045	1.101	77.896	0.056	4.720	0.516	0.027
14.0	1.735	85.498	0.043	1.175	79.464	0.056	4.720	0.518	0.027
15.0	1.823	87.060	0.041	1.209	80.153	0.055	4.720	0.522	0.027
16.0	1.883	88.108	0.040	1.225	80.491	0.055	4.720	0.524	0.027
17.0	1.907	88.544	0.039	1.234	80.657	0.055	4.720	0.524	0.027
18.0	1.917	88.707	0.039	1.238	80.749	0.055	4.720	0.524	0.027
19.0	1.933	88.994	0.039	1.242	80.827	0.055	4.720	0.525	0.027
20.0	1.953	89.347	0.038	1.248	80.956	0.055	4.720	0.526	0.027
21.0	1.969	89.627	0.038	1.253	81.057	0.055	4.720	0.526	0.027


Secular trend may explain part of the difference we observed between our study and the Oosterwolde study, a previous (regional) Dutch growth study including sitting height measurements and performed in 1980 and 1990.<sup>15</sup> We found that our reference lines for SH for age and SH/H ratio for age were usually lower than the Oosterwolde study.

**Table 2** Correlations between SH SDS, LL SDS, SH/H SDS, SH/LL SDS, and height SDS in three age groups

Correlation (r)	Age group		
	0–<5 y	5–<12.5 y	12.5–21 y
SH SDS–H SDS	0.61	0.63	0.80
LL SDS–H SDS	0.50	0.69	0.87
SH/H SDS–H SDS	–0.16	–0.23	–0.23
SH/LL SDS–H SDS	–0.15	–0.22	–0.24
SH SDS–LL SDS	–0.36	–0.08	0.40

Despite the fact that the Oosterwolde sample consisted of relatively tall children from the northern part of the Netherlands, the 1997 Dutch population was even taller. The Oosterwolde study showed that in 10 years (1980–90) the increase in height was more pronounced than the increase in sitting height, so the major secular change must have been in the legs.<sup>1</sup> In the three previous national Dutch growth studies no data on body proportions were collected, so we cannot comment on the secular trend with respect to body proportions. Our present data on sitting height, leg length, and height reference values in the Netherlands are higher than in Denmark, the UK, and Sweden,<sup>16</sup> illustrating our earlier observations<sup>6</sup> that the Dutch population is probably still the tallest in the world (mean height for men 184.0 cm, for women 170.6 cm).


We have shown that in short children a cut-off of 2.5 SDS is better than a cut-off of 2 SDS and that in tall children a cut-off limit of –2.2 SDS can be used. However, we think that instead of using fixed cut-off limits, one can better plot



**Figure 3** A scatter plot of SH/H SDS versus height SDS.

individual observations on the diagram of SH/H SDS versus H SDS. The sensitivity of the conditional  $\pm 2$  SD cut-off limits for detecting hypochondroplasia and Marfan syndrome on the reference chart was studied by comparing body proportions of these two patient groups to the reference population. Based on the values of the positive and negative likelihood ratios of the conditional cut-off limit, the diagnostic value of assessing body proportions for hypochondroplasia is good. For Marfan syndrome, the LR+ is high, but the LR- is not much lower than one, suggesting that normal body proportions do not exclude Marfan syndrome. We can speculate that the major secular change that has affected leg length in particular has led to the relative lack of utility of the standards in the detection of the Marfan individuals, and that this may not be the case in other countries such as the UK. Besides tall and disproportionate stature, there are other defined characteristics to allow diagnosis of Marfan syndrome, such as arachnodactyly, joint laxity, hernias, scoliosis and chest deformations, myopia, dislocation or poor fixation of the lens, and a high arched plate. For both patient groups the equiprobable ellipse is a better criterion to detect growth disorders than the  $\pm 2$  SD lines method. Further investigations on larger groups of patients are necessary to further validate the clinical usefulness of abnormal body proportions for the detection of these and other growth disorders.

Eveleth and Tanner<sup>17</sup> reported that differences in body proportions are genetically controlled and different for European, African, and Oriental populations (Caucasians have tall stature with long legs, in contrast to Orientals). With better environmental circumstances, relatively longer



**Figure 4** A nomogram to assess for a given height SDS the normal range of SH/H SDS, indicating the ellipse, the regression line, and two lines at 2 SDS units away of the regression line. Triangles represent Marfan syndrome, circles represent hypochondroplasia.


legs appear in all ethnic groups. In fact, monitoring leg length might even be a better tool for reflection of environmental improvements than height. Abused children, who have relatively short legs, showed a significant recovery of leg length after social interventions.<sup>18</sup> In our study on body stature, mean height was related to geographical region, family size, and educational level of the parents and the child.<sup>6</sup> In the present study geographical region was only a significant predictor in the youngest group (data not shown). No significant differences were found for educational level or gender.

One of the problems in assessing body proportions is that errors in SH measurement are easily made, which can lead to considerable inter-observer variation. We did not study the inter-observer variance for sitting height measurements, but in the Fels Longitudinal Study the mean absolute inter-observer difference was 0.5 cm (SD 0.3 cm) for crown-rump length and 0.3 cm (SD 0.2 cm) for sitting height.<sup>19</sup>

In conclusion, new reference charts for sitting height, leg length, and body proportion are presented. There was a statistically significant negative correlation between SH/LL and SH/H and height. For practical purposes, in an exceptionally short child a SH/H ratio below +2.5 SDS and in a tall child a SH/H ratio above -2.2 should still be considered normal. The nomogram for SH/H SDS versus H SDS is a useful tool in the work-up of children with growth disorders and provides an objective basis for recognising disproportionate growth.

**Table 3** Percentages of short children (height SDS below -1.5 or -2.0) with an SH/H SDS  $>+2.0$  or  $+2.5$ , and percentages of tall children (height SDS above +1.5 or +2.0) with an SH/H SDS  $<-2.0$  or  $-2.5$  SDS

Short children			Tall children		
SH/H SDS	H SDS $<-2.0$	H SDS $<-1.5$	SH/H SDS	H SDS $>+1.5$	H SDS $>+2.0$
$>+2.0$	6.8%	3.6%	$<-2.0$	3.5%	5.2%
$>+2.5$	4.3%	2.5%	$<-2.5$	1.1%	1.7%



**Figure 5** SH/H data of the members of a family, with an HCH mutation in the *FGFR3* gene on the maternal side, that caused a mild hypochondroplasia with a variable expression pattern. (A) SH/H index for the index case, her cousin, her mother, her aunt, and grandmother. (B) SH/H index for the brother and an uncle of the index case.

### What is already known on this topic

- Measuring body proportions can give important clues in the work-up of growth disorders
- Tall children have relatively long legs, and short children relatively short legs

### What this study adds

- Up-to-date age references for sitting height (SH), leg length (LL), and SH/height (H) in Northern European children, adolescents, and young adults
- A nomogram of SH SDS versus height SDS
- An estimate of cut-off limits of SH/H SDS in short and tall children at a specificity of 98%
- An estimate of the sensitivity of the cut-off limits of the nomogram to detect hypochondroplasia and Marfan syndrome

### ACKNOWLEDGEMENTS

This study was carried out in cooperation with the Well Baby Clinics and Municipal Health Services and was financially supported by The Ministry of Health, Welfare and Sports, Netherlands Organisation for Health Research and Development, Nutricia Nederland BV, and Pfizer BV. We are grateful to Prof. Dr E Bakker, Mrs Y Hilhorst-Hofstee, and Mrs A Neary (Dept of Clinical Genetics, Leiden), Dr CTRM Schrandt-Stumpel and Dr A Plomp (Dept of Clinical Genetics, Maastricht), Dr M Jansen (Dept of Paediatrics, Utrecht), Dr EVM Vermeulen (Dept of Paediatrics, Zevenaar), Dr CJAM van der Burgt (Dept of Clinical Genetics, Nijmegen), and Dr CM Aalfs (Dept of Clinical Genetics, Amsterdam) for providing data of patients with hypochondroplasia. We are also grateful to Drs L Rozendaal (Dept of Paediatrics, Leiden) for providing data on patients with Marfan syndrome.

### Authors' affiliations

A M Fredriks, W J M van Heel, J M Wit, Department of Paediatrics, Leiden University Medical Center, Leiden, Netherlands

S van Buuren, S P Verloove-Vanhorick, Child Health Division, TNO Quality of Life, Leiden, Netherlands

R H M Dijkman-Neerincx, Department of Paediatrics, Hospital Rijnstate, Arnhem, Netherlands

Competing interests: none declared

### REFERENCES

- 1 Gerver WJ, De Bruin R. Relationship between height, sitting height and subischial leg length in Dutch children: presentation of normal values. *Acta Paediatr* 1995;**84**:532-5.
- 2 Cheng JC, Leung SS, Lau J. Anthropometric measurements and body proportions among Chinese children. *Clin Orthop* 1996;**323**:22-30.
- 3 Herber SM, Milner RD. Sitting heights in Sheffield, 1985: have standards changed? *Acta Paediatr Scand* 1987;**76**:818-23.
- 4 Yun DJ, Yun DK, Chang YY, et al. Correlations among height, leg length and arm span in growing Korean children. *Ann Hum Biol* 1995;**22**:443-58.
- 5 Brinkers JM, Lamore PJ, Gevers EF, et al. The effect of oestrogen treatment on body proportions in constitutionally tall girls. *Eur J Pediatr* 1994;**153**:237-4.
- 6 Fredriks AM, van Buuren S, Burgmeijer RJ, et al. Continuing positive secular growth change in the Netherlands 1955-1997. *Pediatr Res* 2000;**47**:316-23.
- 7 Cole TJ, Green PJ. Smoothing reference centile curves: the LMS method and penalized likelihood. *Stat Med* 1992;**11**:1305-19.
- 8 van Buuren S, Fredriks M. Worm plot: a simple diagnostic device for modelling growth reference curves. *Stat Med* 2001;**20**:1259-77.
- 9 *Growthcharts. 1997 (2002) Dutch Growth Study*. Bohn Stafleu van Loghum, Houten.
- 10 Udjus LG. Anthropometrical changes in Norwegian men in the twentieth century. *Universitetsforlaget* 1964.
- 11 Bogin B, Smith P, Orden AB, et al. Rapid change in height and body proportions of Maya American children. *Am J Hum Biol* 2002;**14**:753-61.
- 12 Dangour AD, Schilg S, Hulse JA, et al. Sitting height and subischial leg length centile curves for boys and girls from Southeast England. *Ann Hum Biol* 2002;**29**:290-305.
- 13 Tanner JM. Principles of growth standards. *Acta Paediatr Scand* 1990;**79**:963-7.
- 14 Tanner JM, Hayashi T, Preece MA, et al. Increase in length of leg relative to trunk in Japanese children and adults from 1957 to 1977: comparison with British and with Japanese Americans. *Ann Hum Biol* 1982;**9**:411-23.
- 15 Gerver WJM, De Bruin R. *Paediatric morphometrics, a reference manual*. Maastricht: University Press Maastricht, 2001.
- 16 Hertel NT, Scheike T, Juul A, et al. [Body proportions of Danish children. Curves for sitting height ratio, subischial length and arm span]. *Ugeskr Laeger* 1995;**157**:6876-81.
- 17 Eveleth PB, Tanner JM. *Worldwide variation in human growth*. Cambridge: Cambridge University Press, 2000:186.
- 18 Wales JK, Herber SM, Taitz LS. Height and body proportions in child abuse. *Arch Dis Child* 1992;**67**:632-5.
- 19 Roche AF, Shumei SS. *Human growth, assessment and interpretation*. Cambridge: Cambridge University Press, 2003.