

## SEPARATION AND QUANTITATION OF LEUKOTRIENES BY REVERSED-PHASE HIGH-PERFORMANCE LIQUID CHROMATOGRAPHY

Jan Verhagen, Pauline Walstra, Gerrit A. Veldink, Johannes F.G. Vliegthart,

Department of Bio-Organic Chemistry, State University of Utrecht,  
Croesestraat 79, NL-3522 AD UTRECHT, The Netherlands,  
(reprint requests to JV),

and Pieter L.B. Bruynzeel,

Department of Pulmonary Diseases, University Hospital,  
Catharijnesingel 101, NL-3511 GV UTRECHT, The Netherlands

### ABSTRACT

A rapid and sensitive reversed-phase HPLC procedure is reported which allows the simultaneous separation and quantitation of LTC<sub>4</sub>, 11*t*-LTC<sub>4</sub>, LTD<sub>4</sub>, LTB<sub>4</sub>, 12*epi*,6*t*,8*c*-LTB<sub>4</sub>, 6*t*-LTB<sub>4</sub> + 12*epi*,6*t*-LTB<sub>4</sub>, two trihydroxy-eicosatetraenoic acids tentatively identified as 20-OH-LTB<sub>4</sub> and 20-OH,12*epi*,6*t*,8*c*-LTB<sub>4</sub> and several not yet identified 15-series leukotrienes produced by the cytosol of porcine polymorphonuclear leukocytes.

### INTRODUCTION

Unsaturated fatty acids containing three or more methylene-interrupted *cis* double bonds can be dioxygenated by lipoxygenases to yield mono- and dihydroperoxy fatty acids. Bild et al. (1) have reported on the double dioxygenation of all-*cis*-5,8,11,14-eicosatetraenoic acid (ETE, arachidonic acid) by soybean lipoxygenase-1. This reaction has subsequently been studied by Van Os et al. (2) who showed that two major products were formed, both containing two hydroperoxide groups, which were identified as 8,15-DHPETE and 5,15-DHPETE. The 8,15-DHPETE has three conjugated double bonds and hence has a characteristic triene UV-spectrum with absorption maxima at 259, 268.5 and 279 nm. It was shown that all chiral centers have the S-configuration (2).

In animal systems, lipoxygenases also produce oxygenated unsaturated fatty acids, which can be enzymically converted into leukotrienes. The latter compounds have pronounced biological activities.

In the presence of glutathione or L-cysteine the sulfur-containing leukotrienes C<sub>4</sub> and D<sub>4</sub> can be formed which were shown to be potent bronchoconstrictors (3). Therefore these leukotrienes may play an important role as mediators in immediate hypersensitivity reactions (4)

LTB<sub>4</sub>, which is formed from LTA<sub>4</sub> by enzymic hydrolysis, and to a lesser degree 20-OH-LTB<sub>4</sub>, formed from LTB<sub>4</sub> through  $\omega$ -hydroxylation, show a strong chemotactic behaviour towards human neutrophils (5) and induce degranulation of human neutrophils (6). These physiological activities suggest an important role for these leukotrienes in inflammation reactions.

Upon incubation of arachidonic acid with leukocyte preparations, usually a complex mixture of leukotrienes with different structural features is formed. The composition of the mixture depends on the species used as well as on the incubation conditions i.e., the concentration of Ca<sup>2+</sup>-ions and the presence of Ca-ionophore and glutathione. The simultaneous presence of sulfidopeptide-leukotrienes, various DHETE's and THETE's then constitutes a serious analytical problem. Previously published RP-HPLC methods did not resolve LTB<sub>4</sub> and 12*epi*,6*t*,8*c*-LTB<sub>4</sub> (7,8). Recently, Masters and McMillan (Annual Symposium on Leukotrienes, London, 1982) presented a RP-HPLC procedure which allows the separation of LTB<sub>4</sub> and 12*epi*,6*t*,8*c*-LTB<sub>4</sub> but these authors did not simultaneously measure sulfidopeptide-leukotrienes and 15-series leukotrienes.

In this paper we present a rapid, quantitative and sensitive method for a complete analysis of sulfidopeptide-leukotrienes, 5- and 15-series DHETE's, and THETE's in a single isocratic RP-HPLC run.

#### MATERIALS AND METHODS

**MATERIALS:** Synthetic LTB<sub>4</sub>, LTC<sub>4</sub> and LTD<sub>4</sub> were a generous gift of Dr J. Rokach, Merck-Frosst Laboratories, Pointe-Claire/Dorval, Quebec, Canada. Arachidonic acid (purity >99%) was obtained from Fluka AG, Switzerland. Reduced glutathione, Ca-ionophore A23187 and PGB<sub>2</sub> were purchased from Sigma Chemical Company, St. Louis, Mo, U.S.A.. Human blood was obtained from healthy volunteers. Porcine blood came from the local slaughterhouse. All solvents (including water) were of HPLC quality and purchased from Baker, Deventer, The Netherlands. Dextran grade A (MW 200,000 - 275,000) was from BDH, Poole, U.K.. Ficoll-Paque was obtained from Pharmacia Fine Chemicals AB, Uppsala, Sweden.

**METHODS:** Human and porcine polymorphonuclear leukocytes (PMNL's) were isolated as follows: platelet-rich plasma was removed by centrifugation at 275 g for 15 min. After collection of the buffy coat, leukocytes were separated from erythrocytes by dextran sedimentation. PMNL's were separated from lymphocytes and monocytes by centrifugation on Ficoll-Paque at 650 g for 45 min. The remaining erythrocytes were removed by ammoniumchloride lysis at 0°C and subsequent centrifugation. The purity of the cell preparations was better than 97%. The porcine PMNL cytosol was prepared by ultrasonication of a PMNL suspension (in Dulbecco's salt solution, pH 7.4) at 0°C for 2 x 1.5 min at 20 kHz and centrifugation of the homogenate at 27,000 g for 10 min. Suspensions of human PMNL's in Dulbecco's salt solution (pH 7.4, ~ 3x10<sup>7</sup> cells/ml) or porcine PMNL cytosol (from ~ 4x10<sup>7</sup> cells/ml) were then incubated during 10 min at 37°C and pH 7.4 with 80  $\mu$ M arachidonic acid, 20  $\mu$ M A23187, 1 mM (extra) CaCl<sub>2</sub> and, in case of human PMNL's, 5 mM glutathione. The reaction was stopped by the addition of ice to the solution, followed by centrifugation at 46,000 g for 10 min at 4°C. The supernatant was then applied to a C18 reversed-phase extraction column

(Baker, Deventer, The Netherlands). The adsorbed leukotrienes were eluted with ~ 3 ml methanol.

RP-HPLC was carried out on Nucleosil 5C18 column (250 x 4.6 mm, Chrompack, Middleburg, The Netherlands) attached to a Perkin Elmer Series 1 pump fitted with a Rheodyne injector (20  $\mu$ l sample loop) and a Perkin-Elmer LC85 detector. The solvent system was tetrahydrofuran-methanol-water-acetic acid (25/30/45/0.1, by volume) which was brought to pH 5.5 with ammoniumhydroxide. The aqueous phase contained 0.1% EDTA to prevent binding to the column of cations, which could lead to excessive retention of sulfidopeptide leukotrienes (cf. 8). A flow rate of 0.9 ml/min was maintained and the effluent was monitored at 280 nm. Peak areas were determined with a Hewlett-Packard 3390A integrator and related to the internal standard PGB<sub>2</sub> using molar absorption coefficients at 280 nm of 28,650 for PGB<sub>2</sub> and 40,000 for the leukotrienes.

## RESULTS AND DISCUSSION

Incubation of human PMNL's with arachidonic acid in the presence of Ca<sup>2+</sup>, Ca-ionophore A23187 and glutathione leads to the formation of LTC<sub>4</sub>, 11*t*-LTC<sub>4</sub>, LTB<sub>4</sub>, 12*epi*,6*t*,8*c*-LTB<sub>4</sub>, a racemic mixture of non-enzymically formed 6*t*-LTB<sub>4</sub> and 12*epi*,6*t*-LTB<sub>4</sub> and two trihydroxy-eicosatetraenoic acids, identified as 20-OH-LTB<sub>4</sub> and 20-OH,12*epi*,6*t*,8*c*-LTB<sub>4</sub> (9). The relative amounts of the leukotrienes were found to be dependent on the incubation conditions as well as on the PMNL preparation. The sulfidopeptide leukotrienes LTC<sub>4</sub> and 11*t*-LTC<sub>4</sub> are only formed if glutathione or L-cysteine is also present. The relative amounts of 20-OH-LTB<sub>4</sub> and 20-OH,12*epi*,6*t*,8*c*-LTB<sub>4</sub> were found to be strongly dependent on the length of the incubation period (Masters and McMillan, Annual Symposium on Leukotrienes, London, 1982). In the absence of Ca<sup>2+</sup> and Ca-ionophore, considerable amounts of 15-series leukotrienes are synthesized (10). The production of the double dioxygenation product 12*epi*,6*t*,8*c*-LTB<sub>4</sub> and its  $\omega$ -hydroxylated metabolite 20-OH,12*epi*,6*t*,8*c*-LTB<sub>4</sub> strongly increases when the PMNL preparation is contaminated with thrombocytes (11). To get a proper insight into the absolute quantities of the leukotrienes formed under well-defined conditions, we developed a rapid and sensitive RP-HPLC assay that allows the simultaneous separation and quantitation of all the leukotrienes mentioned above. A major disadvantage of the solvent systems generally employed so far in the RP-HPLC analysis of leukotrienes is that LTB<sub>4</sub> and the double dioxygenation product 12*epi*,6*t*,8*c*-LTB<sub>4</sub> are not resolved. Unfortunately, the solvent system proposed by Masters and McMillan (Annual Symposium on Leukotrienes, London, 1982) though considerably improving the resolution of LTB<sub>4</sub> and 12*epi*,6*t*,8*c*-LTB<sub>4</sub>, does not satisfactorily separate the sulfidopeptide leukotrienes and the 15-series leukotrienes from the 5-series DHETE's. Suitable retention times for LTC<sub>4</sub> and LTD<sub>4</sub> can be obtained by changing the pH of the mobile phase because the retention of these leukotrienes depends strongly on the pH of the solvent system, due to the presence of 3 and 2 carboxyl groups in these compounds, respectively. Furthermore, increasing the tetrahydrofuran content improves the resolution of LTB<sub>4</sub> and 12*epi*,6*t*,8*c*-LTB<sub>4</sub> but deteriorates the separation of LTB<sub>4</sub> and the non-enzymically formed DHETE's. Careful adjustment of the composition of the mobile phase resulted in a solvent system that allows the separation and quantitation of all leukotrienes mentioned above within a reasonable period of time (~ 40 min).

Fig. 1 shows the RP-HPLC chromatogram of an artificial mixture of leukotrienes, prepared by mixing appropriate amounts of synthetic  $\text{LTB}_4$ ,  $\text{LTC}_4$  and  $\text{LTD}_4$  with various leukotriene samples produced by human PMNL's.




Fig. 1 RP-HPLC chromatogram of synthetic  $\text{LTB}_4$ ,  $\text{LTC}_4$ ,  $\text{LTD}_4$  and the leukotrienes produced by human PMNL's.  $\text{PGB}_2$  was added as an internal standard. Absorbance ranges: 0 - 7 min 0.08 AUFS; 7 - 50 min 0.04 AUFS

The 12*epi*,6*t*,8*c*- $\text{LTB}_4$ , the THETE's and the non-enzymically formed DHETE's were identified by their characteristic UV-spectra and by comparing their retention times on RP-HPLC and SP-HPLC with values obtained from the literature with various solvent systems. Although 6*t*- $\text{LTB}_4$  and 12*epi*,6*t*- $\text{LTB}_4$  are not completely resolved with the mobile phase used, no information is lost since these leukotrienes are formed from  $\text{LTA}_4$  by non-enzymic hydrolysis in 1 : 1 ratio. Porcine PMNL cytosol produces from arachidonic acid, under the conditions described in Materials and Methods, 15-series leukotrienes and 5-series leukotrienes, which are also formed by human PMNL's. A RP-HPLC chromatogram of a sample of leukotrienes prepared by mixing synthetic  $\text{LTB}_4$ ,  $\text{LTC}_4$  and  $\text{LTD}_4$  with the leukotrienes produced by porcine PMNL cytosol and human PMNL's is shown in Fig. 2.




Fig. 2. RP-HPLC chromatogram of synthetic LTB<sub>4</sub>, LTC<sub>4</sub>, LTD<sub>4</sub>, the leukotrienes produced by human PMNL's and the leukotrienes produced by porcine PMNL cytosol. PGB<sub>2</sub> was added as an internal standard. Absorbance range: 0.02 AUFS

Although a full structural analysis of the x-marked leukotrienes (Fig. 2) has not yet been completed, these leukotrienes belong to the 15-series since the RP-HPLC chromatogram of the leukotrienes produced by porcine lymphocytes from 15(S)-HPETE shows peaks with identical retention times and UV-spectra (unpublished results).

This RP-HPLC assay is now routinely used by us to measure leukotriene production of PMNL's purified from human blood. Since the detection limit for leukotrienes was found to be below 10 ng in this analysis and only one HPLC-run is needed to quantitate all leukotrienes, relatively small blood samples (<20 ml) are sufficient to study the leukotriene synthesis by human blood PMNL's.

#### Abbreviations used:

DHPETE: dihydroperoxy-eicosatetraenoic acid, DHETE: dihydroxy-eicosatetraenoic acid, THETE: trihydroxy-eicosatetraenoic acid.

12*epi*,6*t*,8*c*-LTB<sub>4</sub> is sometimes referred to elsewhere as 5S,12S-DHETE.

## ACKNOWLEDGEMENTS

The authors wish to thank Mrs M.L. Hamelink, Mr F. van der Lecq and Mr J. Schuring for technical assistance. This investigation was supported in part by the Netherlands Foundation for Chemical Research (SON) with financial aid from the Netherlands Organization for the Advancement of Pure Research (ZWO) and by grant 82.18 from the Netherlands Asthma Fund.

## REFERENCES

1. Bild GS, Ramadoss CS, Lim S, Axelrod B. Double dioxygenation of arachidonic acid by soybean lipoxygenase-1. *Biochem. Biophys. Res. Commun.* 74:949, 1977
2. Van Os CPA, Rijke-Schilder GPM, Van Halbeek H, Verhagen J, Vliegenthart JFG. Double dioxygenation of arachidonic acid by soybean lipoxygenase-1. Kinetics and regio/stereo specificities of the reaction steps. *Biochim. Biophys. Acta* 663:177, 1981
3. Dahlén SE, Hedqvist P, Hammarström S, Samuelsson B. Leukotrienes are potent constrictors of human bronchi. *Nature* 228:484, 1980
4. Borgeat P, Sirois P. Leukotrienes: a major step in understanding of immediate hypersensitivity reactions. *J. Med. Chem.* 24:121, 1981
5. Palmblad J, Udén A-M, Lindgren J-A, Radmark O, Hansson G, Malmsten CL. Effects of novel leukotrienes on neutrophil migration. *FEBS Lett.* 144:81, 1982
6. Feinmark SJ, Lindgren JA, Claesson H-E, Malmsten C, Samuelsson B. Stimulation of human leukocyte degranulation by leukotriene B<sub>4</sub> and its  $\omega$ -oxidized metabolites. *FEBS Lett.* 136:141, 1981
7. Mathews WR, Rokach J, Murphy RC. Analysis of leukotrienes by high-pressure liquid chromatography. *Anal. Biochem.* 118:96, 1981
8. Metz SA, Hall ME, Harper TW, Murphy RC. Rapid extraction of leukotrienes from biologic fluids and quantitation by high-performance liquid chromatography. *J. Chromatogr.* 233:193, 1982
9. Hansson G, Lindgren JA, Dahlén SE, Hedqvist P, Samuelsson B. Identification and biological activity of novel  $\omega$ -oxidized metabolites of leukotriene B<sub>4</sub> from human leukocytes. *FEBS Lett.* 130:107, 1981
10. Jubiz W, Radmark O, Lindgren JA, Malmsten C, Samuelsson B. Novel leukotrienes: products formed by initial oxygenation of arachidonic acid at C-15. *Biochem. Biophys. Res. Commun.* 99:976, 1981
11. Borgeat P, Fruteau de Lacroix B, Picard S, Drapeau J, Vallerand P, Corey EJ. Studies on the mechanism of formation of the 5S,12S-dihydroxy-6,8,10,14(E,Z,E,Z)-icosatetraenoic acid in leukocytes. *Prostaglandins* 23:713, 1982