

Flipping your Lid

Hee-Kap Ahn, Prosenjit Bose

Jurek Czyzowicz, Nicolas Hanusse

Evangelos Kranakis, Pat Morin

UU-CS-2001-13

May, 2001

Flipping your Lid

Hee-Kap Ahn* Prosenjit Bose† Jurek Czyzowicz×
Nicolas Hanusse‡ Evangelos Kranakis† Pat Morin†

*Hong Kong University of Science and Technology ▷ (pastel@cs.ust.hk)

†Carleton University ▷ ({jit,kranakis,morin}@scs.carleton.ca)

‡Université Bordeaux ▷ (hanusse@fabri.u-bordeaux.fr)

×Université du Québec à Hull ▷ (jurek@uqah.quebec.ca)

Abstract

Given a polygon P , a *flipturn* involves reflecting a pocket p of P through the midpoint of the lid of p . In 1973, Joss and Shannon (published in Grünbaum (1995)) showed that any polygon on n vertices will become convex after a sequence of at most $(n - 1)!$ flipturns. They conjectured that this bound was not tight, and that $n^2/4$ flipturns would always be sufficient. In this work, we show that any polygon on n vertices will be convex after any sequence of at most $n(n - 3)/2$ flipturns.

1 Introduction

Given a simple polygon P , a *flipturn* involves reflecting a pocket p of the convex hull of P through the midpoint of the convex hull edge defining p . See Fig. 1 for an example. In this paper, we study the number of flipturns required to convexify a polygon.

In studying this problem there are actually two questions that arise. One can consider the optimization problem of determining the minimum number of carefully chosen flipturns required to convexify any polygon, where the flipturns are chosen carefully so as to minimize this quantity. One can also consider the problem of determining the maximum number of flipturns required to convexify any polygon, where the flipturns are performed arbitrarily.




Figure 1: An example of a flipturn.

Dubins *et al* [2] show that the minimum number of floipturns required to convexify any simple lattice polygon (A lattice polygon is a polygon in which all edges have length 1 and are either horizontal or vertical.) on n vertices is at most $n - 4$.

Surprisingly, the more general case of arbitrary polygons was studied as early as 1973 when Joss and Shannon (see Grünbaum [3]) showed that the maximum number of floipturns required to convexify any polygon is at most $(n - 1)!$. They conjecture that this bound is not tight and that $n^2/4$ floipturns always suffices.

Biedl [1] has found an example where a sequence of $\Omega(n^2)$ carefully chosen floipturns are required to convexify a polygon. However, the same polygon can be convexified using a different sequence of $O(n)$ floipturns. Thus, the $\Omega(n^2)$ is only a lower bound on the maximum number of floipturns required to convexify a polygon.

Grünbaum and Zaks [4] showed that even non-simple polygons can be convexified with a finite sequence of floipturns. For a survey of these and other results on flipping polygons, see the paper by Toussaint [6].

In this paper we show that any simple polygon P with n vertices will be convexified after *any* sequence of at most $n(n - 3)/2$ floipturns, i.e., the maximum number of floipturns required to convexify any polygon is at most $n(n - 3)/2$. More generally, any polygon for which the slopes of the edges take on at most s different values will be convexified after at most $n(s - 1)/2 - s$ floipturns. In Section 2 we give some definitions. Section 3 presents our proof. Section 4 summarizes and concludes with open problems.

2 Preliminaries

Let P be a simple polygon whose vertices in counterclockwise order are v_0, \dots, v_{n-1} , and let the edges of P be oriented counterclockwise so that $e_i = (v_{i-1}, v_i)$.¹ A *pocket* $p = (v_i, \dots, v_j)$ of P is a subchain of P such that v_i and v_j are on the convex hull of P and v_k is not on the convex hull of P for all $i < k < j$. A *lid* (v_i, v_j) is the line segment joining the two endpoints of a pocket (v_i, \dots, v_j) .

In our proof, there is a special degenerate case that must be treated carefully. Let (v_i, v_j) be a lid of P . Let l be the line containing v_i and v_j and let v_k be the first vertex at or following v_j such that v_{k+1} is not contained in l . Then we call (v_i, \dots, v_k) a *modified pocket* of P and the segment (v_i, v_k) is called a *modified lid* of P . Modified pockets and lids are equivalent to standard pockets and lids except when convex hull edges have the same slope as edges of P . Fig. 2 illustrates modified pockets.

Let $p = (v_i, \dots, v_k)$ be a modified pocket of P . Then a *floipturn* $f_{i,k}(P)$ of the polygon P transforms P into a new polygon P' by reflecting all edges of p through the midpoint of the modified lid (v_i, v_k) . Equivalently, $f_{i,k}(P)$ rotates the modified pocket $p = (v_i, \dots, v_k)$ 180 degrees about the midpoint of the lid (v_i, v_k) .

¹Here and henceforth, all subscripts will be taken mod n .




Figure 2: The pocket v_i, \dots, v_j and modified pocket v_i, \dots, v_k in (a) a non-degenerate case and (b) a degenerate case.

Let $dir(e_i)$ be the direction of an edge of P , measured as the angle, in radians, between a right oriented horizontal ray and e_i . Let $S = \bigcup_{i=0}^{n-1} \{dir(e_i), -dir(e_i)\}$, i.e., the set of all directions and their negations used by edges of P . We will label the directions in S as d_0, \dots, d_{m-1} in increasing order. For two directions d_i and d_j in S we define the *discrete angle* between d_i and d_j , as $\overline{Z}d_i d_j = (j-i) \bmod m$, i.e., one plus the number of other directions in S between d_i and d_j as we rotate d_i in the counterclockwise direction.

For a vertex v_i of P incident on edges e_i and e_{i+1} we define the *weight* of v_i as

$$w(v_i) = \begin{cases} \overline{Z}dir(e_i) dir(e_{i+1}) & \text{if } v_i \text{ is convex} \\ \overline{Z}dir(e_{i+1}) dir(e_i) & \text{if } v_i \text{ is reflex} \end{cases}$$

We define the weight of P as $w(P) = \sum_{i=0}^{n-1} w(v_i)$. See Fig. 3 for an example.

For ease of notation, we define the variable s as $|S/2|$, which is exactly the number of distinct slopes used by supporting lines of edges of P . From these definitions, it is clear that $w(v_i) \leq s - 1$ and therefore $w(P) \leq n(s - 1)$.




Figure 3: A polygon for which $|S| = 8$ labelled with its vertex weights.

3 Proof of the Main Theorem

In this section we prove our main theorem by showing that the weight of P decreases by at least 2 after every flipturn. We start with the following simple lemma.

Lemma 1. *For any convex polygon P , we have $w(P) = 2s$.*




Figure 4: Four cases in the proof of Lemma 2. Arrows indicate the directions of the edges in P before performing the flipturn.

Proof. Consider the circle of all directions. The weight of a vertex v_i is the number of elements in S contained in the circular interval $I_i = [\text{dir}(e_{i-1}), \text{dir}(e_i))$. Since P is a polygon, $\bigcup_{i=0}^{n-1} I_i$ is the interval $[0, 2\pi)$. Therefore, each element of S contributes at least one to $w(P)$ so $w(P) \geq 2s$. Since P is convex, e_0, \dots, e_{n-1} are ordered in decreasing order of direction, therefore no two intervals I_i and I_j , $i \neq j$ overlap. Thus, each element of S contributes at most one to $w(P)$, so $w(P) \leq 2s$. \square

Consider a modified pocket p of P , and without loss of generality assume that the modified lid of p is parallel to the x -axis. Let v_i and v_j be the left and right vertices of the modified lid of p . Let r and b be the weight of v_i and v_j , respectively, before performing a flipturn on p and let r' and b' be the weight of the v_i and v_j , respectively, after performing the flipturn.

Lemma 2. $r + b - r' - b' \geq 2$

Proof. Let $d_w = \text{dir}(e_{i-1})$, $d_x = \text{dir}(e_i)$, $d_y = \text{dir}(e_{j-1})$, and $d_z = \text{dir}(e_j)$. To aid in understanding the problem, we place v_i and v_j at the same point and draw the four edges incident on v_i and v_j along with their extensions. There are now four cases to consider, depending on the order of d_w , d_x , d_y , and d_z . These four cases are illustrated in Fig. 4.

When viewed this way, it is clear that in each of the four cases $r + b - r' - b' = 2\alpha$, where $\alpha = \min\{d_w, d_y\} - \max\{d_x, d_z\}$. Since the discrete angles between edges of P are non-negative integers, all that remains to show is that $\alpha \neq 0$. In order to have $\alpha = 0$, the two edges defining α must both be pointing in the same direction in P before performing the flipturn. Thus, with the condition $\alpha = 0$ we obtain one of the four situations depicted in Fig. 5. However, in each of these situations, (v_i, v_j) is not a modified lid. We conclude that $\alpha \neq 0$. \square




Figure 5: Four situations corresponding to cases in the proof of Lemma 2.

Theorem 1. *Any simple polygon on n vertices is convexified after any sequence of at most $n(s - 1)/2 - s$ flipturns.*

Proof. This follows immediately from the following three facts. (1) Initially, the weight of P is at most $n(s - 1)$. (2) The weight of P once it is convexified will be $2s$. (3) During a flipturn, the only weights that change are the weights of the two vertices of the modified lid being flipped. Therefore, by Lemma 2 the weight of P decreases by at least 2 after every flipturn. \square

Strengthening the result of Joss and Shannon [3], we immediately obtain the following corollary by taking $s = n$.

Corollary 1. *Any simple polygon on n vertices is convexified after any sequence of at most $n(n - 3)/2$ flipturns.*

As for the result of Dubins *et al* [2] we take $s = 2$ and obtain the following.

Corollary 2. *Any simple lattice polygon on n vertices is convexified after any sequence of at most $n/2 - 2$ flipturns.*

Indeed, Corollary 2 is the best bound possible. This is because the weight of any vertex in a lattice polygon P is at most 1, thus the decrease in the weight of P during a flipturn is at most 2. Therefore $n/2 - 2$ flipturns are necessary to convexify any simple lattice polygon with n corners.

4 Conclusions

Table 1 summarizes the results obtained in this paper and compares them to the previous best known results. The columns labelled Min (respectively, Max) refer to the minimum (respectively, maximum) number of flipturns required. The first row of the table shows the previously known lower bounds, the second row shows the previously known upper bounds and the third row shows the new upper bounds obtained in this work.

In looking at this table, an obvious open problem is that of closing the gap between the linear lower bound and the quadratic upper bound on the minimum number of flipturns required to convexify an arbitrary polygon.

	Arbitrary Polygons		Lattice polygons	
	Min	Max	Min	Max
PLB	$n/2 - 2$	$\Omega(n^2)$ [1]	$n/2 - 2$ [2]	$n/2 - 2$ [2]
PUB	$(n - 1)!$ [3]	$(n - 1)!$ [3]	$n - 4$ [2]	2.6382^n [3, 5]
NUB	$n(n - 3)/2$	$n(n - 3)/2$	$n/2 - 2$	$n/2 - 2$

Table 1: Summary of previous and new results.

References

- [1] T. Biedl. Personal Communication, 1999.
- [2] L. E. Dubins, A. Orlicsky, J. A. Reeds, and L. A. Shepp. Self-avoiding random loops. *IEEE Transactions on Information Theory*, 34(6):1509–1516, 1988.
- [3] B. Grünbaum. How to convexify a polygon. *Geombinatorics*, 5:24–30, 1995.
- [4] B. Grünbaum and J. Zaks. Convexification of polygons by flips and flip-turns. Technical Report 6/4/98, Department of Mathematics, University of Washington, 1998.
- [5] A. J. Guttmann. On two-dimensional self-avoiding random walks. *Journal of Physics A*, 17:455–468, 1984.
- [6] G. Toussaint. The Erdős–Nagy theorem and its ramifications. In *Proceedings of the 11th Canadian Conference on Computational Geometry (CCCG'99)*, 1999. Available online at http://www.cs.ubc.ca/conferences/CCCG/elec_proc/elecproc.html.