

The relevance of non-generic events in scale space models

Arjan Kuijper
Institute of Information and Computing Sciences
Utrecht University
3584 CH Utrecht, the Netherlands

Luc Florack
Department of Biomedical Engineering
Eindhoven University of Technology
NL-5600 MB Eindhoven, The Netherlands

December 19, 2001

Abstract

In order to investigate the deep structure of Gaussian scale space images, one needs to understand the behaviour of spatial critical points under the influence of blurring. We show how the mathematical framework of catastrophe theory can be used to describe and model the behaviour of critical point trajectories when various different types of generic events, *viz.* annihilations and creations of pairs of spatial critical points, (almost) coincide. Although such events are non-generic in mathematical sense, they are not unlikely to be encountered in practice. Furthermore the behaviour leads to the observation that fine-to-coarse tracking of critical points doesn't suffice, since trajectories can form closed loops in scale space. The modelling of the trajectories include these loops. We apply the theory to an artificial image and a simulated MR image and show the occurrence of the described behaviour.

1 Introduction

The presence of structures of various sizes in an image requires image analysis tools capable of dealing with multiple levels of resolution. Various multi-scale paradigms have been developed [64], giving rise to several new topics of interest. For instance, can specific properties at some scale be related to similar properties at other scales? And if so, what can be said about the way such properties change over scale.

1.1 Scale Space History

The concept of Gaussian, or linear scale space has been introduced in the English image literature by Witkin [72] and Koenderink [42]. They showed that the natural way to represent an image at finite resolution is by convolving it with a Gaussian, thus obtaining a smoothed image at a scale determined by the bandwidth. Weickert *et al.* [70, 71] showed that the concept of scale space was introduced in Japan about twenty years earlier [30, 59]. Due to the fact that these papers were in Japanese, they remained unnoticed in the Western world. In their papers Weickert *et al.* give an overview of the several axioms leading to the paradigm of linear scale space. This approach has led to the formulation of various invariant expressions – expressions that are independent of the coordinates – that capture certain features in an image at distinct levels of scale [14, 16, 18, 19, 20, 21, 22, 39]. Such invariant features are potential candidates that one would like to trace over scale and the topological changes of which one would like to investigate, *e.g.* splitting, merging, creation or vanishing. Nowadays, (properties of) scale spaces are widely used in image analysis, segmentation, clustering and statistical analysis [32, 28, 55, 41]. Introductions to scale space can be found in several books [16, 51, 66].

1.2 Deep Structure

In this paper we focus on linear, or Gaussian, scale space. This has the advantage that each scale level only requires the choice of an appropriate scale and that the image intensity at that level follows linearly from any previous level. It is therefore possible to trace the evolution of certain image entities over scale. The exploitation of various scales simultaneously has been referred to as *deep structure* by Koenderink [42]. It pertains to the dynamic change of the image from highly detailed –including noise – to highly smoothed. Furthermore, it may be expected that large structures “live” longer than small structures (a reason that Gaussian blur is used to suppress noise). The image together with its blurred version was called “primal sketch” by Lindeberg [49, 50, 51]. Since multi-scale information can be ordered, one obtains a hierarchy representing the subsequent simplification of the image with increasing scale. In one dimensional images critical points can only vanish. Investigation of these locations has been done by several authors [8, 10, 31, 35, 36, 69, 75, 76, 77]. Higher dimensional images are more complicated as we will discuss below.

1.3 Related Work

An essentially unsolved problem in the investigation of deep structure is how to establish meaningful links across scales. A well-defined and user-independent method is obtained by linking points that satisfy some topological constraint. Thus maxima are linked to maxima, etc. This yields so-called critical curves. This approach has been used in 2-D images by various authors [27, 48, 65]. They linked extrema, but noticed that sometimes new extrema emerged, disrupting a good linking.

This creation of new extrema in scale space has been studied in detail by Damon, [11, 12, 13], proving that these creations are generic in images of dimension larger than

one. That means that they are not some kind of artifact, introduced by noise or numerical errors, but that they are to be expected in any typical case. This was somewhat counterintuitive, since blurring seemed to imply that structure could only disappear, thus suggesting that only annihilations could occur. Damon, however, showed that both annihilations and creations are generic catastrophes. Furthermore, in [11] he gave a complete list of local perturbations of these generic events. Johansen [34] derived the same results by investigating the behaviour of critical curves in scale space. Griffin [25] investigated critical point events in an affine scale space.

Whereas Damons results were stated theoretically, application of these results were reported in *e.g.* [26, 29, 43, 48, 49]. Kalitzin [37] gave artificial examples to show that the methodology of winding numbers (used for detecting critical and degenerated points) was able to detect more complicated catastrophes. Also the generic catastrophes for specified features in an image have been studied [15, 40, 58, 56, 57, 61, 62, 63].

The main outcome of the investigation of the generic results is that in order to be able to use the topological approach one necessarily needs to take into account both the annihilation and creation events. This has been done in previous work by Kuijper *et al.* [44, 45, 46, 47].

1.4 Aim

In images the location of critical points can be found up to the numerical precision of the image. The same holds for the location of catastrophe points in scale space. So although the appearance of catastrophe events can be uniquely separated in annihilations or creations of pairs of critical points, due to *e.g.* numerical limitations, (almost) symmetries in the image, or coarse sampling also indistinguishable compounds of these annihilation and creation events can be found in practise. In this way a couple of nearby generic events may well look like a single, non-generic one.

In this paper we describe these so-called non-generic catastrophes in scale space. The investigation is based on the description of the evolution of critical points in scale space, called (scale space) critical curves, in the neighbourhood of the catastrophe point(s). The compounds of generic events can be modelled using descriptions of “Catastrophe Theory”. Obviously, the models obey the property that assuming infinite precision, in non-generic compounds the generic events can be distinguished.

Furthermore we investigate the appearance of creations as described by these models in more detail and explain why they are, albeit generic, rarely found, probably the reason for current applications to simply ignore them.

The paper is organised as follows: In section 2 relevant theory on Gaussian scale space and catastrophe theory is given, as well as the way to combine them. Non-generic catastrophe models in scale space in generic coordinates and their impact on the critical curves are discussed in section 3. We give some applications in section 4 and end with a summarising discussion in section 5.

2 Theory

We define a Gaussian scale space in section 2.1. The topological change as the scale varies, is called Catastrophe Theory. A summary of relevant theory is presented in section 2.2. The embedding of Catastrophe Theory in scale space and the generic events are discussed in section 2.3.

2.1 Gaussian Scale Space

Definition 1 $L(\mathbf{x})$ denotes an arbitrary n -dimensional image. We will refer to this image as the initial image.

Definition 2 $L(\mathbf{x}; t)$ denotes the $(n + 1)$ -dimensional Gaussian scale space image of $L(\mathbf{x})$.

The Gaussian scale space image is obtained by convolution of an initial image with a normalised Gaussian kernel $G(\mathbf{x}; t)$ of zero mean and standard deviation $\sqrt{2t}$:

$$L(\mathbf{x}; t) = (G * L)(\mathbf{x}; t) \stackrel{\text{def}}{=} \int \frac{1}{\sqrt{4\pi t}^n} e^{-\frac{|\mathbf{x}-\mathbf{y}|^2}{4t}} L(\mathbf{y}) d\mathbf{y} .$$

Differentiation is now well-defined. By using multi-index notation for $\alpha = (\alpha_1, \dots, \alpha_{n+1})$, $\alpha_k \in N$, take $|\alpha| = \sum_{k=1}^{n+1} \alpha_k$ and $\partial^\alpha = \partial_1^{\alpha_1} \partial_2^{\alpha_2} \dots \partial_{n+1}^{\alpha_{n+1}}$, and derivatives of the image up to arbitrary order $|\alpha|$ at any scale t are given by

$$\partial^\alpha L = \partial^\alpha (G * L) = (\partial^\alpha G) * L.$$

That is, an arbitrary derivative of the image is obtained by the convolution of the initial image with the corresponding derivative of a Gaussian. Consequently, $L(\mathbf{x}; t)$ satisfies the diffusion equation:

$$\partial_t L(\mathbf{x}; t) = \Delta L(\mathbf{x}; t) . \quad (1)$$

Here $\Delta L(\mathbf{x}; t)$ denotes the Laplacean.

Definition 3 Spatial critical points, i.e. saddles and extrema (maxima or minima), at a certain scale t_0 are defined as the points at fixed scale t_0 where the spatial gradient vanishes: $\nabla L(\mathbf{x}; t_0) = 0$. We will refer to these points as spatial critical points.

The type of a spatial critical point is given by the eigenvalues of the Hessian H , the matrix with the second order spatial derivatives, evaluated at its location.

Definition 4 The Hessian matrix at a certain scale t_0 is defined by $H \stackrel{\text{def}}{=} \nabla \nabla^T L(\mathbf{x}; t_0)$, where each element of H is given by

$$H_{i,j} = \frac{\partial^2}{\partial x_i \partial x_j} L(\mathbf{x}; t).$$

The trace of the Hessian equals the Laplacean. For maxima (minima) all eigenvalues of the Hessian are negative (positive). At a spatial saddle point H has both negative and positive eigenvalues.

Since $L(\mathbf{x}; t)$ is a smooth function in $(\mathbf{x}; t)$ -space, spatial critical points are part of a one dimensional manifold in scale space by virtue of the implicit function theorem.

Definition 5 A critical curve is a one-dimensional manifold in scale space on which $\nabla L(\mathbf{x}; t) = 0$.

Consequently, the intersection of all critical curves in scale space with a plane of certain fixed scale t_0 yields the spatial critical points of the image at that scale.

2.2 Catastrophe Theory

The spatial critical points of a function with non-zero eigenvalues of the Hessian are called *Morse critical points*. The *Morse Lemma* states that at these points the qualitative properties of the function are determined by the quadratic part of the Taylor expansion of this function. This part can be reduced to the *Morse canonical form* by a slick choice of coordinates.

If at a spatial critical point the Hessian degenerates, so that at least one of the eigenvalues is zero, the type of the spatial critical point cannot be determined.

Definition 6 The catastrophe points of $L(\mathbf{x}; t_0)$ are defined as the points where both the spatial gradient and the determinant of the Hessian vanish: $\nabla L(\mathbf{x}; t_0) = 0$ and $\det H(\mathbf{x}; t_0) = 0$.

The term catastrophe was introduced by Thom [67, 68]. It denotes a (sudden) qualitative change in an object as the parameters on which this object depends change smoothly. This behaviour was already known by the terms perestroika, bifurcation and metamorphosis. The name catastrophe theory was suggested by Zeeman [74] to unify singularity theory, bifurcation theory and their applications and gained wide popularity. A thorough mathematical treatment on singularity theory can be found in the work of Arnol'd [2, 3, 4, 5, 6, 7]. More pragmatic introductions and applications are widely published, e.g. [9, 23, 24, 52, 60, 73, 74].

The catastrophe points are also called *non-Morse critical points*, since a higher order Taylor expansion is essentially needed to describe the qualitative properties. Although the dimension of the variables is arbitrary, the *Thom Splitting Lemma* states that one can split up the function in a Morse and a non-Morse part. The latter consists of variables representing the k “bad” eigenvalues of the Hessian that become zero. The Morse part contains the $n - k$ remaining variables. Consequently, the Hessian contains a $(n - k) \times (n - k)$ sub-matrix representing a Morse function. It therefore suffices to study the part of k variables. The canonical form of the function at the non-Morse critical point thus contains two parts: a Morse canonical form of $n - k$ variables, in terms of the quadratic part of the Taylor series, and a non-Morse part. The latter can be put into canonical form called the *catastrophe germ*, which is obviously a polynomial of degree 3 or higher.

Since the Morse part does not change qualitatively under small perturbations, it is not necessary to further investigate this part. The non-Morse part, however, does change. Generally the non-Morse critical point will split into a non-Morse critical point, described by a polynomial of lower degree, and Morse critical points, or even exclusively into Morse critical points. This event is called a *morsification*. So the non-Morse part contains the catastrophe germ and a perturbation that controls the morsifications.

Then the general form of a Taylor expansion $f(\mathbf{x})$ at a non-Morse critical point of an n dimensional function can be written as (*Thom's Theorem*):

$$f(\mathbf{x}; \lambda) = CG(x_1, \dots, x_k) + PT(x_1, \dots, x_k; \lambda_1, \dots, \lambda_l) + \sum_{i=k+1}^n \epsilon_i x_i^2, \quad (2)$$

where $CG(x_1, \dots, x_k)$ denotes the catastrophe germ, $PT(x_1, \dots, x_k; \lambda_1, \dots, \lambda_l)$ the perturbation germ with an l -dimensional space of parameters, and in the Morse part $\epsilon_i = \pm 1$.

The so-called simple real singularities in 2D have catastrophe germs given by the two infinite series

- $A_k^\pm \stackrel{\text{def}}{=} \pm x^{k+1}, k \geq 1$. The germs A_k^+ and A_k^- are equivalent for $k = 1$ and k even.
- $D_k^\pm \stackrel{\text{def}}{=} x^2 y \pm y^{k-1}, k \geq 4$, which we will rewrite to $x^{k-1} \pm xy^2$ for notational convenience.

Furthermore three exceptional singularities occur:

- $E_6 \stackrel{\text{def}}{=} x^3 \pm y^4$,
- $E_7 \stackrel{\text{def}}{=} x^3 + xy^3$, and
- $E_8 \stackrel{\text{def}}{=} x^3 + y^5$.

2.3 Catastrophes and Scale Space

In Definition 6, the number of equations defining the catastrophe point equals $n + 1$ and therefore it is over-determined with respect to the n spatial variables. Consequently, catastrophe points are generically not found in typical images. In scale space, however, the number of variables equals $n + 1$ and catastrophes occur as isolated points.

Although the list of catastrophes starts very simple, it is not trivial to apply it directly to scale space by assuming that scale is just one of the perturbation parameters. As Damon [11, 13] points out: “There are significant problems in trying to directly apply Morse theory to solutions of the heat equation. First, it is not clear that generic solutions to the heat equation must be generic in the Morse sense. Second, standard models for Morse critical points and their annihilation and creation do not satisfy the heat equation. How must these models be modified? Third, there is the question of what constitutes generic behaviour. This depends on what notion of local equivalence one uses between solutions to the heat equation.”

For example, in one-dimensional images the A_2 catastrophe reduces to $x^3 + \lambda x$. It describes the change from a situation with two critical points (a maximum and a minimum) for $\lambda < 0$ to a situation without critical points for $\lambda > 0$. See *e.g.* Figure in Section for an example of such an annihilation sequence. This event can occur in two ways. The extrema are annihilated for increasing λ , but the opposite – creation of two extrema for decreasing λ – is also possible.

In scale space, however, there is an extra constraint: the germ has to satisfy the diffusion equation. Thus the catastrophe germ x^3 implies an extra term $6xt$. On the other hand, the perturbation term is given by $\lambda_1 x$, so by taking $\lambda_1 = 6t$ scale plays the role of the perturbing parameter. This gives a directionality to the perturbation parameter, in the sense that the only remaining possibility for this A_2 -catastrophe in one-dimensional images is an annihilation. So the Fold catastrophe is adjusted such that it satisfies the heat equation, but this adjustment only allows annihilations. However, it does not imply that only annihilations are generic in scale space. In higher dimensional images also the opposite – *i.e.* a A_2 catastrophe describing the creation of a pair of critical points – is possible. Then the perturbation $\lambda = -6t$ with increasing t requires an additional term of the form $-6xy^2$ in order to satisfy the diffusion equation as we will see.

The transfer of the catastrophe germs to scale space has been made by many authors, [11, 12, 13, 17, 33, 35, 36, 37, 38, 43, 45, 46, 47, 49, 51], among whom Damon's account is probably the most rigorous. He showed that the only generic morsifications in scale space are the aforementioned A_2 catastrophes describing *annihilations* and *creations* of pairs of critical points. These two points have opposite sign of the determinant of the Hessian before annihilation and after creation. All other events are compounds of such events. It is however possible that one may not be able to distinguish these generic events, *e.g.* due to numerical limitations, coarse sampling, or (almost) symmetries in the image. For instance, one may find at some scale three nearby critical points, *e.g.* two extrema and a saddle, and at the subsequent scale only one extremum. Obviously, one pair of critical points is annihilated, but one may not be able to identify the annihilating extremum at the former scale. This is illustrated in Figure 1. On the left the critical paths are shown, together with a grey area representing the uncertainty in determining the catastrophe location. On the right the non-generic model of this event is displayed.

Definition 7 *The scale space catastrophe germs are defined by*

$$\begin{aligned} f^A(\mathbf{x}; t) &\stackrel{\text{def}}{=} x_1^3 + 6x_1 t + Q(\mathbf{x}; t), \\ f^C(\mathbf{x}; t) &\stackrel{\text{def}}{=} x_1^3 - 6x_1 t - 6x_1 x_2^2 + Q(\mathbf{x}; t). \end{aligned}$$

The quadratic term $Q(\mathbf{x}; t)$ is defined

$$Q(\mathbf{x}; t) \stackrel{\text{def}}{=} \sum_{i=2}^n \epsilon_i (x_i^2 + 2t),$$

where $\sum_{i=2}^n \epsilon_i \neq 0$ and $\epsilon_i \neq 0 \forall i$.




Figure 1: Left: Annihilation of two critical points in the neighbourhood of a third critical point. The grey area represents the uncertainty in determining the catastrophe. Right: Non-generic representation and model of this event.

Note that the scale space catastrophe germs f^A and f^C , and the quadratic term Q satisfy the diffusion equation. The germs f^A and f^C correspond to the two qualitatively different A_2 catastrophes at the origin, an annihilation and a creation respectively. From Definition 7 it is obvious that annihilations occur in any dimension, but creations require at least 2 dimensions. Consequently, in 1D signals only annihilations occur. Furthermore, for images of arbitrary dimension and less than three vanishing eigenvalues of the Hessian at a degenerated point, it suffices to investigate the 2D case due to the Splitting Lemma.

2.3.1 The Annihilation Germ

Spatial critical points at any scale t for f^A follow directly from $\nabla f^A(\mathbf{x}; t) = 0$:

$$\begin{cases} 3x_1^2 = -6t \\ 2\epsilon_i x_i = 0, & i \geq 2 \end{cases}$$

Then the critical curve is parametrised by $(\pm\sqrt{-2t}, 0, \dots, 0; t)$, $t \leq 0$. At the origin a catastrophe takes place. The determinant of the Hessian is given by $\det H = cx_1$, with the constant $c = 3 \cdot 2^n \prod_{i=2}^n \epsilon_i$. So two critical points with opposite sign approach the origin as t increases to zero and meet transversally to the Hessian zero-crossing. Note that $\text{tr } H = 6x_1 + \sum_{i=2}^n 2\epsilon_i$, which is generically non-zero at catastrophe points. This explains the constraints on the ϵ_i in Definition 7.

2.3.2 The Creation Germ

The creation germ is a bit more complicated. Spatial critical points at any scale t for f^C follow from $\nabla f^C(\mathbf{x}; t) = 0$:

$$\begin{cases} 3x_1^2 - 6x_2^2 = 6t \\ 2x_2(\epsilon_2 - 6x_1) = 0 \\ 2\epsilon_i x_i = 0, & i \geq 3 \end{cases}$$

Since we look in the neighbourhood of the origin, we take $x_2 = 0$. Then the critical curve is parametrised by $(\pm\sqrt{2t}, 0, \dots, 0; t)$, $t \geq 0$. At the origin a catastrophe takes

place. The determinant of the Hessian is given by $\det H = cx_1(\epsilon_2 - 6x_1) - 12cx_2^2$, with the constant $c = 3 \cdot 2^n \prod_{i=3}^n \epsilon_i$, so two critical points with opposite sign leave the origin as t increases from zero, again transversally to the Hessian zero-crossing. Note that this catastrophe is a A_2 catastrophe, since it describes the creation of two critical points, although there is a striking resemblance to the description of the D_4^- catastrophe. Furthermore, the description of the catastrophe is essentially local: If t is taken too large, the (non-generic) degeneration of the Hessian at $x_1 = \frac{1}{6}\epsilon_2$ has to be taken into account. We will elaborate on these items in Section 3.

3 Scale space catastrophe models

In this section we describe how the following catastrophes (with nicknames and perturbation germs) can be used to model events in $(2 + 1)$ -dimensional scale space. The catastrophes describe in canonical coordinates how critical curves pass the origin yielding compounds of annihilations and / or creations of pairs of critical points.

- A_2 Fold catastrophe: $x^3 + \lambda_1 x \pm y^2$.
- A_3 Cusp catastrophe: $\pm x^4 + \lambda_1 x + \lambda_2 x^2 \pm y^2$.
- D_4^+ Hyperbolic Umbilic catastrophe, $x^3 + xy^2 + \lambda_1 x + \lambda_2 y + \lambda_3 y^2$.
- D_4^- Elliptic Umbilic catastrophe: $x^3 - xy^2 + \lambda_1 x + \lambda_2 y + \lambda_3 y^2$.
- D_5^\pm Parabolic Umbilic catastrophe: $x^4 \pm xy^2 + \lambda_1 x + \lambda_2 y + \lambda_3 y^2 + \lambda_4 x^2$.
- D_6^+ Second Hyperbolic Umbilic catastrophe: $x^5 + xy^2 + \lambda_1 x + \lambda_2 y + \lambda_3 x^2 + \lambda_4 y^2 + \lambda_5 x^3$.
- D_6^- Second Elliptic Umbilic catastrophe: $x^5 - xy^2 + \lambda_1 x + \lambda_2 y + \lambda_3 x^2 + \lambda_4 y^2 + \lambda_5 x^3$.

The \pm signs at the A_3 and D_5 denote ‘dual’ possibilities with similar geometry.

Again we emphasise that this list not a complete list as can be found in [11], containing all the relevant mathematical details. However, the germs mentioned above are related to the perturbations of the generic annihilation and creation given in [11]. We will see that although most of these catastrophes are non-generic, they may still be relevant for modelling compounds of generic events that one is not capable of, or willing to, segregate as such. Recall, for example, Figure 1.

The germs in this list are adjusted such that they satisfy the heat equation. Furthermore, by choosing the perturbation terms non-zero and adjusting them in the same way, descriptions of critical curves in scale space are obtained. These critical curves only contain the generic Fold annihilation(s) and/or creation(s).




Figure 2: Critical paths of the Cusp catastrophe. The catastrophe is located at the dot. a) $\lambda_1 = 1$. b) $\lambda_1 = 0$. c) $\lambda_1 = -1$. Note that if the perturbation is very small, these three distinct cases may very well be confused.

3.1 A_2 Fold catastrophe

The Fold catastrophe in scale space is given by

$$L(x, y; t) = x^3 + 6xt + \delta(y^2 + 2t),$$

where $\delta = \pm 1$. Critical curves and the catastrophe point follow from

$$\begin{cases} L_x = 3x^2 + 6t \\ L_y = 2\delta y \\ \det(H) = 12\delta x. \end{cases}$$

One can verify that at the origin a saddle and an extremum (a minimum if $\delta = 1$, a maximum if $\delta = -1$) moving in the $y = 0$ plane meet and annihilate while increasing the scale parameter t .

3.2 A_3 Cusp catastrophe

The Cusp catastrophe germ is given by x^4 . Its scale space addition is $12x^2t + 12t^2$. The perturbation term contains two terms: $\lambda_1 x + \lambda_2 x^2$. Obviously, scale takes the role of λ_2 . Taking the dual Cusp gives the same geometry by changing the sign of λ_1 , or by setting $x = -x$. The scale space Cusp catastrophe germ with perturbation is thus defined by

$$L(x, y; t) = x^4 + 12x^2t + 12t^2 + \lambda_1 x + \delta(y^2 + 2t),$$

with $\delta = \pm 1$. Again, the critical curves and the catastrophe point follow from

$$\begin{cases} L_x = 4x^3 + 24xt + \lambda_1 \\ L_y = 2\delta y \\ \det(H) = 24\delta(x^2 + 2t). \end{cases}$$

Morsification by the perturbation $\lambda_1 \neq 0$ yields one Fold catastrophe and one regular critical curve, see Figure 2a,c. The differences in behaviour depending on the sign of δ is studied in detail in [44]. It suffices here to note that if $\lambda_1 = 0$, at the origin three critical points transform to one critical point while increasing scale, see Figure 2b.

One can verify that the A_k , $k > 3$ catastrophes describes the (non-generic) simultaneous annihilations of critical points in one dimension under the influence of blurring, albeit in more complicated appearances.



Figure 3: Critical paths of the D_4^+ -catastrophe. a) Unperturbed. b) Perturbed. Again, if the perturbation is small we may not be able to tell which configuration is the actual one.

3.3 D_4^\pm Umbilic catastrophes

The D_4^\pm Umbilic catastrophe germs are given by $x^3 + \delta xy^2$, where $\delta = \pm 1$. The scale space addition is $(6 + 2\delta)xt$, yielding $x^3 + xy^2 + 8xt$ for the Hyperbolic Umbilic catastrophe, and $x^3 - xy^2 + 4xt$ for the Elliptic Umbilic catastrophe. The perturbation contains three terms: $\lambda_1 x + \lambda_2 y + \lambda_3 y^2$. Obviously, scale takes the role of λ_1 .

3.3.1 D_4^+ Hyperbolic Umbilic catastrophe

The scale space D_4^+ Hyperbolic Umbilic catastrophe germ with perturbation is thus defined by

$$L(x, y; t) = x^3 + xy^2 + 8xt + \lambda_3(y^2 + 2t) + \lambda_2 y .$$

The critical curves and catastrophe points follow from

$$\begin{cases} L_x &= 3x^2 + 8t + y^2 \\ L_y &= 2xy + 2\lambda_3 y + \lambda_2 \\ \det(H) &= 12x(x + \lambda_3) - 4y^2. \end{cases}$$

In the unperturbed situation four critical points exist for each $t < 0$ on the x - and y -axes. At $t = 0$ the four critical curves annihilate simultaneously at the origin, see Figure 3a. Taking perturbation into account, the curves are separated into two critical curves each containing a Fold catastrophe, see Figure 3b.

3.3.2 D_4^- Elliptic Umbilic catastrophes

The scale space elliptic Umbilic catastrophe germ with perturbation is given by

$$L(x, y; t) = x^3 - xy^2 + 4xt + \lambda_3(y^2 + 2t) + \lambda_2 y . \quad (3)$$

Again, the critical curves and the catastrophe points follow from

$$\begin{cases} L_x &= 6x^2 + 4t - y^2 \\ L_y &= -2xy + 2\lambda_3 y + \lambda_2 \\ \det(H) &= 12x(2\lambda_3 - 2x) - 4y^2. \end{cases}$$

The unperturbed equation gives two critical points for all $t \neq 0$. At the origin a so-called scatter event occurs: the critical curve changes from y -axis to x -axis with




Figure 4: Critical paths of the D_4^- -catastrophe. a) Unperturbed. b) Small perturbation. c) Large perturbation.

increasing t , see Figure 4. Just as in the hyperbolic case, in fact two Fold catastrophes take place; in this case both an annihilation and a creation. The morsification is shown in Figure 4b. The critical curve on the right does not contain catastrophe points. The critical curve on the left, however, contains two Fold catastrophe points: a creation and an annihilation. Both are studied in detail in [44]. A brief sketch was given in section 2, where we could set $\lambda_2 = 0$, since only the creation at the origin was investigated.

So while increasing scale one will find two critical points, suddenly two extra critical points appear, of which one annihilates with one of the already existing ones. Finally, one ends up with again two critical points. Clearly, if the samples in scale are taken too large, one could completely miss the subsequent catastrophes, see *e.g.* Figure 4c. The properties of the creations will be discussed in the next section.

3.3.3 Creations

As we showed, a creation event occurs in case of a morsified elliptic Umbilic catastrophe. In most applications, however, creations are rarely found, giving rise to the (false) opinion that creations are caused by numerical errors and should be disregarded. The reason for their rare appearance lies in the specific requirements for the parameters in the (morsified) Umbilic catastrophe germ. Its general formulation is given by

$$L(x, y; t) = \frac{1}{6}L_{xxx}x^3 + \frac{1}{2}L_{xyy}xy^2 + L_{xt}xt + \frac{1}{2}L_{yy}(y^2 + 2t) + L_yy \quad (4)$$

In general, the spatial coefficients do *not* equal the derivatives evaluated in the coordinate system of the image. They follow from the alignment of the catastrophe in the plane defined by $y = 0$ and can have arbitrary value. Furthermore, the diffusion equation implies $L_{xt} \stackrel{\text{def}}{=} L_{xxx} + L_{xyy}$. Then the scale space evolution of the critical curves follow from

$$\begin{cases} \partial_x L = \frac{1}{2}L_{xxx}x^2 + L_{xt}t + \frac{1}{2}L_{xyy}y^2 \\ \partial_y L = L_{xyy}xy + L_{yy}y + L_y \\ \det(H) = L_{xxx}x(L_{xyy}x + L_{yy}) - L_{xyy}^2y^2. \end{cases}$$

Firstly we consider the case $L_y = 0$. Then Eq. (4) describes a Fold catastrophe (either annihilation or creation) at the origin, where the critical curve is positioned in the (x, t) -plane. A creation necessarily requires the constraint $L_{xxx}L_{xt} < 0$ at the catastrophe point. This constraint is sufficient.




Figure 5: a) The fraction of the space of the third order derivatives in which creations can occur as a function of the dimension according Theorem 1. b) Intersections of the curves $\det(H) = 0$ and $\partial_y L = 0$ with different values for L_y . For the value given by Theorem 2 the curves touch. c) Difference in intensity between the creation and the annihilation event for L_y increasing from 0 to its critical value.

Theorem 1 *At a catastrophe point in two spatial dimensions, if the third order derivatives of the general local form as given by Eq. (4) with $L_y = 0$, are uncorrelated, the number of creations has an a priori likelihood of $1/4$ relative to the total number of catastrophes. In n dimensions it is $\frac{1}{\pi} \arccos \frac{1}{\sqrt{n}}$.*

Proof. The requirement $L_{xxx}L_{xt} < 0$ can be rewritten to $L_{xxx}(L_{xxx} + L_{xyy}) < 0$. In the (L_{xxx}, L_{xyy}) -space this constraint is satisfied by all point sets in the area spanned by the lines through the origin with direction vectors $(1, 0)$ and $(1, -1)$, which is a quarter of the plane. For $n - D$ this extends to the area $L_{xxx}(L_{xxx} + L_{xy_1y_1} + \dots + L_{xy_{n-1}y_{n-1}}) < 0$ in $(L_{xxx}, L_{xy_iy_i})$ -space, with $\dim(\mathbf{y}) = n - 1$. This representing two intersecting planes with normal vectors $(1, 0, \dots, 0)$ and $(1, -1, \dots, -1)$. They make an angle of ϕ radians, given by

$$\cos \phi = \frac{(1, 0, \dots, 0) \cdot (1, -1, \dots, -1)}{|(1, 0, \dots, 0)| \cdot |(1, -1, \dots, -1)|} = \frac{1}{\sqrt{n}}$$

Then the fraction of the space follows by taking twice this angle and dividing by the complete angle of 2π , i.e. $\frac{1}{\pi} \arccos \frac{1}{\sqrt{n}}$. \square

Note that if $n = 1$, the fraction of the space where creations can occur is zero, for $n = 2$ it is a quarter. The also interesting case $n = 3$ yields a fraction that is slightly more than a quarter, whereas for $n \rightarrow \infty$ the fraction converges to a half, see Figure 5a. That is: the higher the dimensions, the easier critical points can be created.

The reason that in practice in two dimensional images the number of creations observed is (much) smaller than a quarter, is caused by the role of the perturbation parameters. It is possible to give a tight bound to the perturbation of Equation (4) in terms of L_y :

Theorem 2 *A creation and subsequent annihilation event occur in Equation (4) if and only if*

$$|L_y| \leq \frac{3}{16} L_{yy}^2 \sqrt{\frac{-3L_{xxx}}{L_{xyy}^3}} \quad (5)$$

Proof. The catastrophes satisfy $\partial_x L = \partial_y L = \det H = 0$. Since the solution of the system

$$\begin{aligned}\partial_y L &= L_y + y(L_{yy} + L_{xyy}x) = 0 \\ \det H &= L_{xxx}x(L_{yy} + L_{xyy}x) - L_{xyy}^2 y^2 = 0\end{aligned}\quad (6)$$

only contains spatial coordinates, their intersections define the spatial coordinates of the catastrophes. The catastrophe points form the local extrema of the critical curve in $(x, y; t)$ -space, *i.e.* at these points the tangent vector has no scale component. If the curves given by Eq. (6) touch, there is only a point of inflection in the critical curve, *i.e.* the critical curve in $(x, y; t)$ -space has a (Fold) catastrophe point. At this point of inflection, the spatial tangent vectors of the curves defined by Eq. (6) are equal.

Solving the system Eq. (6) with respect to y results in

$$y = -\frac{L_y}{L_{yy} + L_{xyy}x} = \pm \frac{1}{L_{xyy}} \sqrt{L_{xxx}x(L_{yy} + L_{xyy}x)}.$$

The equality of the tangent vectors at the point of inflection x_i, y_i yields

$$\frac{\partial}{\partial x} \left(-\frac{L_y}{L_{yy} + L_{xyy}x} \right) \Big|_{x_i, y_i} = \frac{\partial}{\partial x} \left(\pm \frac{1}{L_{xyy}} \sqrt{L_{xxx}x(L_{yy} + L_{xyy}x)} \right) \Big|_{x_i, y_i}$$

Solving both equalities results in

$$(x_i, y_i, L_y) = \left(-\frac{L_{yy}}{4L_{xyy}}, \pm \sqrt{\frac{-3L_{xxx}L_{yy}^2}{16L_{xyy}^3}}, \mp \frac{3L_{yy}^2}{16L_{xyy}} \sqrt{\frac{-3L_{xxx}}{L_{xyy}}} \right),$$

which gives the boundary values for L_y . \square

Note that Eq. (5) has only real solutions if $L_{xxx}L_{xyy} < 0$, *i.e.* at the D_4^- (morsified) catastrophe. As a consequence of Theorem 2, creations only occur if the perturbation is small enough. Again, this perturbation occurs in the coordinate system, obtained by the alignment of the catastrophe in the plane defined by $y = 0$.

Example. 1 Taking $L_{xxx} = 6, L_{xyy} = -12, L_{yy} = 2$ yielding $L = x^3 - 6xy^2 - 6xt + y^2 + 2t + L_y$, we obtain the “generic creation example” as given in section 2 with perturbation. Then Theorem 2 gives $|L_y| \leq \frac{1}{32}\sqrt{6}$ as a – relatively small compared to the other derivative values– bound for the occurrence of a creation – annihilation couple. In Figure 5b the ellipse $\det(H) = 0$ is plotted, together with the curves $\partial_y L = 0$ for $L_y = 0$ (resulting in two straight lines at $y = 0$ and $x = \frac{1}{6}$, intersecting at $(x, y) = (\frac{1}{6}, 0)$), and $L_y = 2^{-i}\sqrt{6}, i = 4, \dots, 7$. For $i > 5$, the perturbation is small enough and the intersection of $\partial_y L = 0$ and $\det H = 0$ contains two points. Thus a creation-annihilation is observed. If $i = 5$, L_y has its critical value and the curves touch. For larger values the curves do not intersect each other.

Obviously the perturbation L_y can be larger if L_{yy} increases. If so, the structure becomes more elongated. It is known by various examples of creations given in literature that elongated structures play an important role. In fact, the quintessential property is scale anisotropy.

Another reason that creations are rarely found is that their lifetime is rather limited: with increasing t the created critical points annihilate. If the scale steps are taken too large, one simply misses the creation – annihilation couple. This may be regarded as a dual expression for the previous explanation. In the chosen coordinate system this can be calculated explicitly.

Theorem 3 *The maximum lifetime of a creation given by Equation (4) is*

$$t_{lifetime} = \frac{-L_{xxx}L_{yy}^2}{2L_{xyy}^2(L_{xxx} + L_{xyy})}.$$

The difference in intensity of the critical point that is created and subsequently annihilated is

$$\frac{L_{xxx}(2L_{xxx} - L_{xyy})L_{yy}^3}{6L_{xyy}^3(L_{xxx} + L_{xyy})}.$$

Proof. Observe that the lifetime is bounded by the two intersections of $\partial_y L = 0$ and $\det(H) = 0$, see Figure 5b. As $|L_y|$ increases from zero, the two points move towards each other over the arch $\det(H) = 0$ until they reach the value given by theorem 2 with lifetime equal to zero. The largest arch length is obtained for $L_y = 0$. Then the spatial coordinates are found by

$$\partial_y L(x, y; t) = y(L_{xyy}x + L_{yy}) = 0$$

and

$$\det H = L_{xxx}x(L_{xyy}x + L_{yy}) - L_{xyy}^2y^2 = 0,$$

i.e. $(x, y) = (0, 0)$ and $(x, y) = (-\frac{L_{yy}}{L_{xyy}}, 0)$ The location in scale space is given by

$$\partial_x L(x, y; t) = \frac{1}{2}L_{xxx}x^2 - \frac{1}{2}L_{xyy}y^2 + L_{xt}t = 0.$$

Consequently, the first catastrophe takes place at the origin - since also $t = 0$ - with zero intensity. The second one is located at

$$(x, y; t) = \left(-\frac{L_{yy}}{L_{xyy}}, 0; \frac{-L_{xxx}L_{yy}^2}{2L_{xyy}^2(L_{xxx} + L_{xyy})}\right)$$

with intensity

$$L_{cat} = \frac{L_{xxx}(2L_{xxx} - L_{xyy})L_{yy}^3}{6L_{xyy}^3(L_{xxx} + L_{xyy})}.$$

Then the latter is also the maximum difference in intensity. \square

Example. 2 *To show the effect of the movement along the arch $\det(H) = 0$, see Figure 5c. Without loss of generality we took again $L_{xxx} = 6, L_{xyy} = -12, L_{yy} = 2$. Firstly, the two solutions to $\nabla L = 0 \wedge \det(H) = 0$ were calculated as function of L_y . Secondly, the difference of the intensity of the solutions was calculated for 766 subsequent values of $L_y, L_y \in [0, \dots, \frac{1}{32}\sqrt{6}]$. It is clearly visible that the intensity decreases monotonously with an increase of L_y . For this example we find that the lifetime is $\frac{1}{72}$, the difference in intensity $\frac{1}{18}$.*

From the proof of Theorem 3 it is again apparent that L_{yy} plays an important role in enabling a (long)lasting creation. To observe this in more detail, note that the curve

$$\det H = 0 \Leftrightarrow L_{xxx}(L_{xyy}x + L_{yy}) = L_{xyy}^2 y^2$$

is an ellipse (see also Figure 5b). Replacing x by $x - \frac{L_{yy}}{2L_{xyy}}$, it is centred at the origin:

$$L_{xxx}\left(x - \frac{L_{yy}}{2L_{xyy}}\right)L_{xyy}\left(x + \frac{L_{yy}}{2L_{xyy}}\right) = L_{xyy}^2 y^2 .$$

And consequently

$$L_{xxx}L_{xyy}\left(x^2 - \frac{L_{yy}^2}{4L_{xyy}^2}\right) = L_{xyy}^2 y^2 .$$

Setting $L_{xyy} = \frac{1}{b}$ and $L_{xxx}L_{xyy} = -\frac{1}{a^2}$, we find

$$\det H = 0 \Leftrightarrow x^2 + \frac{a^2}{b^2}y^2 = L_{yy}^2 \frac{b^2}{4} .$$

Assuming that we have a creation, $a^2 > 0$. The ellipse is enlarged with an increase of L_{yy}^2 . Obviously, at the annihilations of the Hyperbolic Umbilic catastrophe $a^2 < 0$, so $\det H = 0$ then describes a hyperbola.

3.4 D_5^\pm Parabolic Umbilic catastrophes

In the previous section we saw that the geometry significantly changed by taking either the term $-xy^2$, or the term $+xy^2$. Let us therefore, ignoring the perturbation terms λ_1 , λ_2 , and λ_3 , define the scale space Parabolic Umbilic catastrophe germ by

$$L(x, y; t) = \frac{1}{4!}x^4 + \frac{1}{2!}x^2t + \frac{1}{2!}t^2 + \delta\left(\frac{1}{2}xy^2 + xt\right) \quad (7)$$

where $\delta = \pm 1$ and t takes the role of λ_4 . Its critical curves and catastrophes follow from

$$\left\{ \begin{array}{l} L_x = \frac{1}{6}x^3 + xt + \delta\left(t + \frac{1}{2}y^2\right) \\ L_y = \delta xy \\ \det(H) = \delta x\left(\frac{1}{2}x^2 + t\right) - y^2 \end{array} \right.$$

So the catastrophe points are located at the origin (a double point) and at $(x, y; t) = \left(-\frac{3}{2}\delta, 0; -\frac{9}{8}\delta^2\right)$. The latter is a simple annihilation (a fold catastrophe), the former is a cusp catastrophe (three critical point change into one) for both values of δ , see Figure 6a-b. Indeed the geometry of the D_5 and its dual are not significantly different. Adding small perturbations by choosing the parameters λ_1 , λ_2 , and λ_3 , the morsified Cusp catastrophe remains, see Figure 6c-d. The critical curves at the Cusp breaks up into two curves, one with a Fold catastrophe, one without a catastrophe.




Figure 6: Critical paths of the D_5^\pm -catastrophe in the $(x, y; t)$ -space. a) Unperturbed, $\delta = -1$. b) Unperturbed, $\delta = 1$. c) Perturbed, $\delta = -1$. d) Perturbed, $\delta = 1$.

3.5 D_6^\pm Second Umbilic Catastrophes

Ignoring the perturbation terms $\lambda_1, \dots, \lambda_4$ for the moment, the scale space expression of the D_6^\pm -catastrophes are given by

$$L(x, y; t) = \frac{1}{5!}x^5 + \frac{1}{3!}x^3t + \frac{1}{2!}xt^2 + \delta\left(\frac{1}{2}xy^2 + xt\right), \quad (8)$$

where t takes the role of λ_5 and $\delta = \pm 1$. Its critical curves and catastrophes follow from

$$\begin{cases} L_x &= \frac{1}{4!}x^4 + \frac{1}{2}x^2t + \frac{1}{2}t^2 + \delta(t + \frac{1}{2}y^2) \\ L_y &= \delta xy \\ \det(H) &= \frac{1}{6}\delta x^2(x^2 + 6t) - y^2 \end{cases}$$

Setting $y = 0$, several catastrophes occur: At $(x, y; t) = (\pm\sqrt{-6\delta}, 0; \delta)$ two Fold annihilations if $\delta = -1$, at the origin a creation and at $(x, y; t) = (0, 0; -2\delta)$ again an annihilation, see Figure 7a for $\delta = 1$ and Figure 7b for $\delta = -1$.

It is clear that the morsification by t of the D_6^+ yields a D_4^- scatter followed (while increasing scale) by a D_4^+ double annihilation at the origin. The D_6^- shows a D_4^- scatter at the origin, followed by again a D_4^- scatter at some higher scale. Both images show that a part of the critical curve forms a loop: The created critical points annihilate with each other.

So if the perturbations are small (or if the measurement contains some uncertainty), one might not be able to distinguish between the involved Fold catastrophes. However, the scale space representation causes a separation into two non-generic catastrophes already mentioned. Further morsification gives more insight in the way *critical curves* can behave.

By taking $\lambda_1, \dots, \lambda_4 \neq 0$, the generic critical curves shown in Figure 7c-d are obtained. The morsification of the D_6^+ shows two critical curves behaving in an aesthetic




Figure 7: Critical paths of the D_6^\pm -catastrophe. a) Unperturbed, $\delta = 1$. b) Unperturbed, $\delta = -1$. c) Perturbed, $\delta = 1$. d) Perturbed, $\delta = -1$.

way, combining the morsifications of the D_4^\pm catastrophes, *i.e.* containing Fold annihilations and creations. Both created critical points on the right critical curve in Figure 7c annihilate at some larger scale.

The morsification of the D_6^- , on the other hand, still shows the loop close to the origin. Consequently, in contrast to the elliptic Umbilic catastrophe, now *both created branches annihilate with each other*: the critical curve in the centre of Figure 7d is a closed loop in scale space.

3.5.1 On generic loops

The perturbation term of the second Umbilic catastrophes in scale space is given by

$$\lambda_1 x + \lambda_2 y + \lambda_3(x^2 + 2t) + \lambda_4(y^2 + 2t).$$

If we assume that the image contains the catastrophe at the origin, we may set $\lambda_1 = \lambda_2 = \lambda_3 = 0$. Furthermore, if it contains an elongated structure at the creation, $|\lambda_4| > 0$. This perturbation causes the non-generic scatter event, visible in Figure 7b, to break up. While increasing $|\lambda_4|$ the two parabolae in the $x = 0$ -plane move along the critical loop, until they meet each other and change into two critical curves without a catastrophe, as shown in Figure 7d.

Setting $\lambda_4 = 2L_{yy}$ we get

$$L(x, y; t) = L_{xtt}\left(\frac{1}{5!}x^5 + \frac{1}{6}x^3t + \frac{1}{2}xt^2\right) + L_{xt}(xt + \frac{1}{2}xy^2) + L_{yy}\left(\frac{1}{2}x^2 + t\right),$$

where $L_{xtt} = L_{xxxx}$ and $L_{xt} = L_{xyy}$ in absence of other third and fifth order derivatives. Consequently,

$$\begin{cases} \partial_x L &= L_{xtt}\left(\frac{1}{4!}x^4 + \frac{1}{2}x^2t + \frac{1}{2}t^2\right) + L_{xt}\left(t + \frac{1}{2}y^2\right) \\ \partial_y L &= L_{xt}xy + L_{yy}y \\ \det(H) &= L_{xtt}\left(\frac{1}{6}x^3 + xt\right)(L_{xt}x + L_{yy}) - L_{xt}y^2 \end{cases}$$

One can verify that the solution $x = -\frac{L_{yy}}{L_{xt}}$ of $\partial_y L = 0$ describes the non-genericity on the loop if $L_{yy}^2 \leq (-3 + \sqrt{3})\frac{L_{xt}^3}{L_{xtt}}$ by solving $\partial_x L = 0$, so L_{yy} needs to be large enough and $L_{xtt}L_{xt} < 0$, *i.e.* only the morsified D_6^- pertains its loop. Now at the origin the creation takes place. Note that there $\partial_t L \neq 0$. The successive annihilation follows from $x = y = 0$ and $\partial_x L = L_{xtt}\frac{1}{2}t^2 + L_{xt}t = 0$ and takes place at $(x, y; t) = (0, 0; -\frac{2L_{xt}}{L_{xtt}})$. So the lifetime of the scale space loop is $t_l = -\frac{2L_{xt}}{L_{xtt}}$. Furthermore, the difference in intensity between creation and annihilation yields $L_{yy}t_l$. The loop can therefore persist over a certain range of scales. We will come back to these parameter setting and coordinate choice in section 4.

3.6 Morsification summary

All non-Fold catastrophes morsify to Fold catastrophes and Morse critical points. The morsification gives insight in the structure around the catastrophe point regarding the critical curves.

The morsification of the Umbilic catastrophes (the D_k) show that the trajectories in scale space of the created critical points fall into several classes.

The morsified D_4^+ -catastrophes describes two Fold annihilations. The morsified D_4^- catastrophe describes the creation of a pair of critical points and the annihilation of one of them with another critical point. So while tracing a critical branch of a critical curve both an annihilation and a creation event are traversed.

The morsified D_6^+ catastrophe describes the creation of a pair of critical points and the annihilation of both of them with two other critical points. So while tracing a critical branch of a critical curve successively an annihilation, a creation and again an annihilation event are traversed.

The morsified D_6^- -catastrophe describes an *isolated* closed critical curve, appearing *ex nihilo* with two critical branches that disappear at some larger scale.

So the morsified D_4^- (and its extension, the D_6^+) and D_6^- -catastrophes describe essentially different creation events. An important result lays on the area of tracing critical points. If one traces *only* critical points starting from the initial image, one will find the “ D_4^- ” creations, since they emerge as the starting point of a part of a critical curve that annihilates with one of the initial critical points. However, one will miss the “ D_6^- ” loops that occur somewhere in scale space, since they have no relation whatsoever to the critical points in the initial image. So fine-to-coarse tracing of critical points will not always yield the right result.

Note that the full morsification of the non-generic catastrophes always yields the generic Fold annihilations and creations and Morse critical points.

3.6.1 Other non-generic catastrophes

The catastrophes, their scale space formulation, and their morsifications, as treated in the previous sections, are only the beginning of an infinite set of possible descriptions on the behaviour of critical curves. Due to extreme local symmetries and inaccuracies – think of a large checkerboard pattern – one might encounter non-generic events to be described with higher other catastrophe models. These models follow straightforward from the route we described.

From the point of view of distinct appearances of critical curves, this expose suffices. Since the only possible generic catastrophes on critical curves are creations and annihilations of pairs of critical points, the possibilities are limited. Given an critical curve, it either originated from the initial image and the curve can contain several protuberances, i.e. sequences of successive creations and annihilations, or it didn't originate from the initial image and it forms a loop, perhaps also with protuberances. Both possibilities are described by the aforementioned catastrophe models. The scale space formulation of the “three exceptional catastrophes” don't yield any extra information, as one can verify.

4 Applications

In this section we give some examples to illustrate the theory presented in the previous sections. We will focus on the critical curves emerging from a creation event. An exam-




Figure 8: a) 181 x 217 artificial MR image. b) Image on scale 8.37

ple of desired symmetry in an image and consequently modelling by a Cusp catastrophe has been presented elsewhere [44]. Firstly, results on the artificial MR image of Figure 8a are presented. This image is taken from the web site <http://www.bic.mni.mcgill.ca/brainweb>. Secondly, an example of creation *ex nihilo*, the D_6^- -catastrophe, is shown by means of the classic “bridge”-image of Figure 11a.

4.1 D_4^- -catastrophe

The artificial MR image of Figure 8a was used as initial image for the scale space image. For visualisation purposes, we restricted to the scale range 8.37 – 33.1. The image at scale 8.37 (with only the large structures remaining) is shown in Figure 8b. This image contains 7 extrema.

The scale space image in this scale range contains 161 logarithmically sampled scales. At all scales the spatial critical points were calculated and connected, forming the critical paths. Figure 9 shows these critical paths in the $(x, y; t)$ -space. The bright curves represent the extrema, the dark ones the saddles. At the (approximate) catastrophe locations the curves are connected. Globally, the image shows annihilating pairs of critical points. Locally, however, the presence of extra branches of critical curves is visible.

A close-up of one of the critical paths is shown in Figure 10a. It clearly shows a critical curve containing two subsequent Fold annihilation – creation events. The critical curve evidently shows the appearance of an annihilation-creation-pair described by the D_3^- morsification. Note that the creation events would have been missed if the sampling was taking coarser, yielding one critical curve without protuberances in scale direction. Sampling without connecting critical paths yields the observation of temporarily created extrema (and saddles), *cf.* Simmons et al. [65].

4.2 D_6^- -catastrophe

Figure 11a shows the classical “bridge”-image: two mountains of different height (blobs with different intensity) connected by a small ramp and a deep valley between



Figure 9: Critical paths of the MR image in scale range 8.37 – 33.1.




Figure 10: a) Close-up of one of the critical paths of the MR image, showing a subsequent annihilation – creation event. b) Close-up, showing subsequent annihilation – creation events and loop events.

created (the latter if the dimension of the image is 2 or higher), it is important to describe also the local structure of the image around the non-generic events. These events might be encountered in practical usage of scale spaces and the non-generic catastrophes can be used to model these occurrences. We therefore embedded catastrophes in scale space. Scale acts as one of the perturbation parameters. The morsification of the catastrophes yields generic Fold annihilations and creations of pairs of critical points.

The A_k series can be used to model (almost) simultaneous annihilations of pairs of critical points at a location (or indistinguishable region) in scale space. If k is even, it models the annihilation of k critical points, if k is odd, it models the collision of k critical points where $k - 1$ annihilate and one remains.

For creations the D_k series can be used. Creations occur in different types. Critical paths in scale space can have protuberances, a subsequent occurrence of an annihilation and a creation. In scale space images this is visible by the creation of an extremum-saddle pair, of which one critical point annihilates at some higher scale with an already present critical point, while the other remains unaffected. It is also possible that critical paths form loops: the created pair annihilates at some higher scale. The possibility for both types to occur in practice was shown in the artificial MR image. This phenomena is known from physics, where it is used to describe the creation and successive annihilations of “virtual” elementary particles (and even the universe), see *e.g.* [1, 53, 54].

Furthermore we showed that the protuberances in the critical paths, expressed in canonical coordinates, occur only in case of a small local perturbation. In addition, creations are less likely to happen due to a special constraint on the combination of third order derivatives and local perturbation. We gave a dimension dependent expectation of this event and an upper bound for the perturbation in canonical coordinates.

The lifetime of a created pair is enlarged if the local structure is elongated. This was derived from the canonical formulation and visualised by the example of the bridge image in section 4.

Since the number of possible catastrophes is infinite, there is an infinite number of possible non-generic constellations in which (“infinite”) critical points are annihilated and created. We restricted ourselves to the situations in which at most 6 critical points annihilate and in which critical points are created, the latter divided into models representing protuberances and loops.

Finally, the calculations were based on the canonical coordinates. In general, it is not trivial to transform the local coordinate system to these nice formulated catastrophe germs. In that sense, the numerical values have no direct meaning. They do describe, however, the qualitative behaviour of the critical curves close to the location of the catastrophes and can therefore be used to model the type of behaviour encountered in practical usage of a scale space. We gave examples of the appearances of this behaviour in section 4 based on an artificial MR image.

The theory described in this paper extends the knowledge of the deep structure of Gaussian scale space, especially with respect to the behaviour of critical curves in the vicinity of creation events and the scale space lifetime of the created critical points. It emphasises the relevance of investigating the complete scale space image, instead of a series of images at different scales.

Acknowledgements

James Damon and Steve Pizer of the University of North Carolina are gratefully acknowledged for their comments on a preliminary version of this paper.

References

- [1] J. Argyris, C. Ciubotariu, and H.-G. Matuttis. Fractal space, cosmic strings and spontaneous symmetry breaking. *Chaos, Solitons and Fractals*, 12(1):1–48, January 2001.
- [2] V. I. Arnold, editor. *Geometrical Methods in the Theory of Ordinary Differential Equations*, volume 250 of *Grundlehren der mathematischen Wissenschaften: A Series of Comprehensive Studies in Mathematics*. Springer-Verlag, Berlin, 1983.
- [3] V. I. Arnold. *Catastrophe Theory*. Springer, Berlin, 1984.
- [4] V. I. Arnold, editor. *Ordinary Differential Equations*. Springer-Verlag, Berlin, 1992.
- [5] V. I. Arnold, editor. *Dynamical Systems VI: Singularity Theory I*, volume 6 of *Encyclopaedia of Mathematical Sciences*. Springer-Verlag, Berlin, 1993.
- [6] V. I. Arnold, editor. *Dynamical Systems VIII: Singularity Theory II & Applications*, volume 39 of *Encyclopaedia of Mathematical Sciences*. Springer-Verlag, Berlin, 1993.
- [7] V. I. Arnold, editor. *Dynamical Systems V: Bifurcation Theory and Catastrophe Theory*, volume 5 of *Encyclopaedia of Mathematical Sciences*. Springer-Verlag, Berlin, 1994.
- [8] K. Berkner. Resolution of singularities of convolutions with the gaussian kernel. *Proc. Amer. Math. Soc.*, 127(2):409–415, 1999.
- [9] J. W. Bruce and P. J. Giblin. *Curves and Singularities*. Cambridge University Press, 1984.
- [10] J. J. Clark. Singularity theory and phantom edges in scale space, 1988.
- [11] J. Damon. Local Morse theory for solutions to the heat equation and Gaussian blurring. *Journal of Differential Equations*, 115(2):386–401, 1995.
- [12] J. Damon. Generic properties of solutions to partial differential equations. *Arch. Rat. Mech. Anal.*, pages 353–403, 1997.
- [13] J. Damon. Local Morse theory for Gaussian blurred functions. In *Sporring et al. [66]*, pages 147–162, 1997.
- [14] J. Duncan and N. Ayache. Medical image analysis: Progress over two decades and the challenges ahead. *IEEE Transactions on Pattern Analysis and Machine Intelligence*, 22(1):85–105, 2000.
- [15] D. Eberly, R. Gardner, B. Morse, S. Pizer, and C. Scharlach. Ridges for image analysis. *Journal of Mathematical Imaging and Vision*, 4(4):353–373, 1994.
- [16] L. M. J. Florack. *Image Structure*, volume 10 of *Computational Imaging and Vision Series*. Kluwer Academic Publishers, Dordrecht, The Netherlands, 1997.
- [17] L. M. J. Florack and A. Kuijper. The topological structure of scale-space images. *Journal of Mathematical Imaging and Vision*, 12(1):65–80, February 2000.
- [18] L. M. J. Florack, B. M. ter Haar Romeny, J. J. Koenderink, and M. A. Viergeever. Scale and the differential structure of images. *Image and Vision Computing*, 10(6):376–388, July/August 1992.

- [19] L. M. J. Florack, B. M. ter Haar Romeny, J. J. Koenderink, and M. A. Viergever. Cartesian differential invariants in scale-space. *Journal of Mathematical Imaging and Vision*, 3(4):327–348, 1993.
- [20] L. M. J. Florack, B. M. ter Haar Romeny, J. J. Koenderink, and M. A. Viergever. General intensity transformations and differential invariants. *Journal of Mathematical Imaging and Vision*, 4(2):171–187, May 1994.
- [21] L. M. J. Florack, B. M. ter Haar Romeny, J. J. Koenderink, and M. A. Viergever. Linear scale-space. *Journal of Mathematical Imaging and Vision*, 4(4):325–351, 1994.
- [22] L. M. J. Florack, B. M. ter Haar Romeny, J. J. Koenderink, and M. A. Viergever. The Gaussian scale-space paradigm and the multiscale local jet. *International Journal of Computer Vision*, 18(1):61–75, April 1996.
- [23] A. T. Fomenko and T. L. Kunii. *Topological Modeling for Visualization*. Springer-Verlag, Tokyo, 1997.
- [24] R. Gilmore. *Catastrophe Theory for Scientists and Engineers*. Dover, 1993. Originally published by John Wiley & Sons, New York, 1981.
- [25] L. D. Griffin. Critical point events in affine scale-space. In *Sporring et al. [66]*, pages 165–180, 1997.
- [26] L. D. Griffin and A. Colchester. Superficial and deep structure in linear diffusion scale space: Isophotes, critical points and separatrices. *Image and Vision Computing*, 13(7):543–557, September 1995.
- [27] L. D. Griffin, A. Colchester, and G. Robinson. Scale and segmentation of grey-level images using maximum gradient paths. *Image and Vision Computing*, 10(5):389–402, 1992.
- [28] B. M. ter Haar Romeny, L. M. J. Florack, J. J. Koenderink, and M. A. Viergever, editors. *Scale-Space Theory in Computer Vision: Proceedings of the First International Conference, Scale-Space'97, Utrecht, The Netherlands*, volume 1252 of *Lecture Notes in Computer Science*. Springer-Verlag, Berlin, July 1997.
- [29] R. D. Henkel. Segmentation in scale space. In *Computer Analysis of Images and Patterns. Lecture Notes in Computer Science, vol. 970*. Springer-Verlag, pages 41–48, 1995.
- [30] T. Iijima. Basic theory of pattern normalization (for the case of a typical onedimensional pattern). *Bulletin of the Electrotechnical Laboratory*, 26:368–388, 1962. (in Japanese).
- [31] P. Johansen. On the classification of toppoints in scale space. *Mathematical Imaging and Vision*, 4(1):57–67, 1994.
- [32] P. Johansen, editor. *Proceedings of the Copenhagen Workshop on Gaussian Scale-Space Theory*, May 10–13 1996. Technical Rapport 96/19 ISSN 01078283.
- [33] P. Johansen. Local analysis of image scale space. In *Sporring et al. [66]*, pages 139–146, 1997.
- [34] P. Johansen. Local analysis of scale space. Technical Report 97/2, Department of Computer Science, University of Copenhagen, 1997.
- [35] P. Johansen, M. Nielsen, and O.F. Olsen. Branch points in one-dimensional Gaussian scale space. *Journal of Mathematical Imaging and Vision*, 13:193–203, 2000.
- [36] P. Johansen, S. Skelboe, K. Grue, and J. D. Andersen. Representing signals by their top-points in scale space. In *Proceedings of the International Conference on Image Analysis and Pattern Recognition (Paris, France, October 1986)*, pages 215–217. IEEE Computer Society Press, 1986.

- [37] S. Kalitzin. Topological numbers and singularities. In *Sporring et al. [66]*, pages 181–190, 1997.
- [38] S. N. Kalitzin, B. M. ter Haar Romeny, A. H. Salden, P. F. M. Nacken, and M. A. Viergever. Topological numbers and singularities in scalar images. Scale-space evolution properties. *Journal of Mathematical Imaging and Vision*, 9(3):253–296, November 1998.
- [39] S. N. Kalitzin, J. Staal, B. M. ter Haar Romeny, and M. A. Viergever. A computational method for segmenting topological point-sets and application to image analysis. *IEEE Transactions on Pattern Analysis and Machine Intelligence*, 23(5):447–459, 2001.
- [40] R. S. Keller. *Generic Transitions of Relative Critical Sets in Parametrized Families with Applications to Image Analysis*. PhD thesis, University of North Carolina, 1999.
- [41] M. Kerckhove, editor. *Scale-Space and Morphology in Computer Vision*, volume 2106 of *Lecture Notes in Computer Science*. Springer -Verlag, Berlin Heidelberg, 2001.
- [42] J. J. Koenderink. The structure of images. *Biological Cybernetics*, 50:363–370, 1984.
- [43] A. Kuijper and L.M.J. Florack. Calculations on critical points under gaussian blurring. In *Nielsen et al. [55]*, pages 318–329, 1999.
- [44] A. Kuijper and L.M.J. Florack. The application of catastrophe theory to image analysis, 2001. submitted.
- [45] A. Kuijper and L.M.J. Florack. Hierarchical pre-segmentation without prior knowledge. In *Proceedings of the 8th International Conference on Computer Vision (Vancouver, Canada, July 9–12, 2001)*, pages 487–493, 2001.
- [46] A. Kuijper, L.M.J. Florack, and M.A. Viergever. Scale space hierarchy. Technical Report UU-CS-2001-19, Department of Computer Science, Utrecht University, 2001.
- [47] A. Kuijper, L.M.J. Florack, and M.A. Viergever. Scale space hierarchy, 2001. submitted.
- [48] L. M. Lifshitz and S. M. Pizer. A multiresolution hierarchical approach to image segmentation based on intensity extrema. *IEEE Transactions on Pattern Analysis and Machine Intelligence*, 12(6):529–540, 1990.
- [49] T. Lindeberg. Scale–space behaviour of local extrema and blobs. *Journal of Mathematical Imaging and Vision*, 1(1):65–99, 1992.
- [50] T. Lindeberg. Detecting salient blob-like image structures and their scales with a scale-space primal sketch: a method for focus-of-attention. *International Journal of Computer Vision*, 11(3):283–318, 1993.
- [51] T. Lindeberg. *Scale-Space Theory in Computer Vision*. The Kluwer International Series in Engineering and Computer Science. Kluwer Academic Publishers, 1994.
- [52] Y.-C. Lu. *Singularity Theory and an Introduction to Catastrophe Theory*. Springer-Verlag, Berlin, second corrected printing edition, 1976.
- [53] E. M. Maslov and A. G. Shagalov. Dynamics of first-order phase transitions in the $\phi^4 - \phi^6$ model caused by the parametric instability of the metastable vacuum. *Physica D: Nonlinear Phenomena*, 152-153:769–778, 2001.
- [54] I. G. Moss. Heat kernel expansions for distributional backgrounds. *Physics Letters B*, 491(1-2):203–206, October 2000.
- [55] M. Nielsen, P. Johansen, O. Fogh Olsen, and J. Weickert, editors. *Scale-Space Theories in Computer Vision*, volume 1682 of *Lecture Notes in Computer Science*. Springer -Verlag, Berlin Heidelberg, 1999.

- [56] O. Fogh Olsen. Multi-scale watershed segmentation. In *Sporring et al. [66]*, pages 191–200, 1997.
- [57] O. Fogh Olsen. *Generic Image Structure*. PhD thesis, University of Copenhagen, Denmark, 2000.
- [58] O. Fogh Olsen and M. Nielsen. Generic events for the gradient squared with application to multi-scale segmentation. In *Ter Haar Romeny et al. [28]*, pages 101–112, 1997.
- [59] N. Otsu. *Mathematical Studies on Feature Extraction in Pattern Recognition*. PhD thesis, Electronical Laboratory, Ibaraki, Japan, 1981. (in Japanese).
- [60] T. Poston and I. N. Stewart. *Catastrophe Theory and its Applications*. Pitman, London, 1978.
- [61] J. H. Rieger. Generic properties of edges and corners on smooth greyvalue surfaces. *Biological Cybernetics*, 66:497–502, 1992.
- [62] J. H. Rieger. Generic evolutions of edges on families of diffused greyvalue surfaces. *Journal of Mathematical Imaging and Vision*, 5:207–217, 1995.
- [63] J. H. Rieger. Topographical properties of generic images. *International Journal of Computer Vision*, 23(1):79–92, 1997.
- [64] A. H. Salden. *Dynamic Scale-Space Paradigms*. PhD thesis, Utrecht University, 1996.
- [65] A. Simmons, S. R. Arridge, P. S. Tofts, and G. J. Barker. Application of the extremum stack to neurological MRI. *IEEE Transactions on Medical Imaging*, 17(3):371–382, June 1998.
- [66] J. Sporring, M. Nielsen, L. M. J. Florack, and P. Johansen, editors. *Gaussian Scale-Space Theory*, volume 8 of *Computational Imaging and Vision Series*. Kluwer Academic Publishers, Dordrecht, second edition, 1997.
- [67] R. Thom. *Stabilité Structurale et Morphogénèse*. Benjamin, New York, 1972.
- [68] R. Thom. *Structural Stability and Morphogenesis*. Benjamin-Addison Wesley, 1975. translated by D. H. Fowler.
- [69] T. Wada and M. Sato. Scale-space tree and its hierarchy. In *ICPR90*, volume II, pages 103–108, 1990.
- [70] J. A. Weickert, S. Ishikawa, and A. Imiya. On the history of Gaussian scale-space axiomatics. In *Sporring et al. [66]*, pages 45–59, 1997.
- [71] J. A. Weickert, S. Ishikawa, and A. Imiya. Linear scale-space has first been proposed in Japan. *Journal of Mathematical Imaging and Vision*, 10(3):237–252, 1999.
- [72] A. P. Witkin. Scale-space filtering. In *Proceedings of the Eighth International Joint Conference on Artificial Intelligence*, pages 1019–1022, 1983.
- [73] A. E. R. Woodcock and M. Davis, editors. *Catastrophe Theory*. Clarke, Irwin & Company Limited, 1978.
- [74] E. C. Zeeman. *Catastrophe Theory: Selected Papers, 1972-1977*. Addison-Wesley Publishing Company, 1977.
- [75] N. Zhao. *Feature Extraction of images by stable gaze tree*. PhD thesis, Dept. of Computer Science, Tokyo Inst. of Technology, Japan, 1985. (in Japanese).
- [76] N. Zhao and T. Iijima. A theory of feature extraction by the tree of stable view-points. *IECE Japan, Trans. D*, J68-D:1125–1132, 1985. (in Japanese).
- [77] N. Zhao and T. Iijima. Theory on the method of determination of view-point and field of vision during observation and measurement of figure. *IECE Japan, Trans. D*, J68-D:508–514, 1985. (in Japanese).