

The complex geometry of the spherical pendulum

Frits Beukers¹ and Richard Cushman¹

1 Introduction

In this paper we describe the geometry of the energy momentum mapping of the complexified spherical pendulum. For background on the classical spherical pendulum we refer the reader to [4, chpt. IV]. We show that this complex Hamiltonian system is Mumford-Jacobi completely integrable [9, p.3.51–3.53] and admits an analogue of the period lattice [5] and action angle coordinates [4, Apend. D]. Following the variation of the period lattice around a closed loop in the complement of the discriminant locus of a family of elliptic curves (see equation (9) below), we show that the complexified spherical pendulum has monodromy [4, p.383].

The monodromy of the complexified spherical pendulum also has been described in recent papers by Audin [2] (unpublished) and Gavrilov and Vivlo [6]. Audin starts with a Lax pair related to the complexified spherical pendulum. She then studies the geometry of a singular normalization \widetilde{C}_c of the family of affine elliptic curves C_c :

$$\mu^2 + \lambda^4 + 2c_2\lambda^3 + 2c_1\lambda^2 + 1 = 0,$$

which is part of the spectral curve of the Lax pair. Using Picard-Lefschetz techniques, she then computes the variation of the period map $H_1(\widetilde{C}_c, \mathbb{Z}) \rightarrow \mathbb{C}^2$ as $c = (c_1, c_2)$ traces out a curve in the complement of the discriminant locus of C_c .

The paper of Gavrilov and Vivolo starts with the discriminant locus of C_c and then constructs a set of action-angle coordinates, which gives a simple proof of Horozov's theorem for the spherical pendulum. Using these action-angle coordinates and Picard-Lefschetz theory, they show that the complex spherical pendulum has monodromy.

There are several difficulties with the approaches to monodromy used in the above papers. The most serious one is that the exact relation of the family C_c

¹Mathematics Institute, Univeristy of Utrecht, 3508TA Utrecht, the Netherlands

²version: 21 November 2001

with the dynamics of the complexified spherical pendulum is nowhere explained. Audin's choice of the singular normalization \widetilde{C}_c of the family C_c is not motivated from the dynamics. Also she does not explain what the generalized Jacobian for \widetilde{C}_c is. (The published literature on generalized Jacobians is inaccessible to dynamicists).

On the other hand, we think that our explanation of monodromy is direct. The key observation is that a level set of a regular value of the energy momentum map is a \mathbb{C}^* -bundle over an elliptic curve. Moreover, it is a quotient of \mathbb{C}^2 by a rank 3 lattice. The inverse of the quotient map is simply given by the action angle variables on the level set.

We note that the monodromy we find is not the same as the classic Picard-Fuchs monodromy described in [3, p.657–693], because it involves a differential form of the *third* kind, nor it is the same as the monodromy given in [1]. We have just given an analytic proof of a geometric explanation of monodromy suggested by Knörrer [8].

2 The complexified spherical pendulum

In this section we describe the complex Hamiltonian system given by the complexified spherical pendulum.

Let \langle , \rangle be the standard symmetric bilinear form on \mathbb{C}^3 , namely, $\langle x, y \rangle = \sum_{i=1}^3 x_i y_i$ for $x, y \in \mathbb{C}^3$. Set $Q = \{x \in \mathbb{C}^3 \mid \langle x, x \rangle = 1\}$. Then Q is the complexification of the 2-sphere S^2 in \mathbb{R}^3 . Let

$$M = TQ = \{(x, v) \in Q \times \mathbb{C}^3 \mid \langle x, v \rangle = 0\}$$

be the tangent bundle of Q with bundle projection $pr : M \rightarrow Q : (x, v) \rightarrow x$. On M there is a holomorphic nondegenerate closed 2-form ω which is obtained by pulling back the canonical symplectic form on the cotangent bundle of Q by the flat map associated to \langle , \rangle . Note that M is a smooth subvariety of $\mathbb{C}^3 \times \mathbb{C}^3$.

Recall [4, p.149] that as a constrained Hamiltonian system (H, M, ω) , the complexified spherical pendulum has Hamiltonian

$$H : M \rightarrow \mathbb{C} : (x, v) \rightarrow \frac{1}{2} \langle v, v \rangle + x_3. \quad (1)$$

Moreover, its motion is governed by the Hamiltonian vector field X_H , whose integral curves satisfy

$$\begin{aligned}\dot{x} &= v \\ \dot{v} &= -e_3 + (\langle x, e_3 \rangle - \langle v, v \rangle)x.\end{aligned}\tag{2}$$

The group $SO(2, \mathbb{C}) = \{A = \begin{pmatrix} a & -b \\ b & a \end{pmatrix} \in Gl(2, \mathbb{C}) \mid a^2 + b^2 = 1\}$ acts on M by

$$\tilde{\Phi} : SO(2, \mathbb{C}) \times M \rightarrow M : (A, (x, v)) \rightarrow \left(\begin{pmatrix} A & 0 \\ 0 & 1 \end{pmatrix} x, \begin{pmatrix} A & 0 \\ 0 & 1 \end{pmatrix} v \right).\tag{3}$$

This action is Hamiltonian, has momentum mapping

$$L : M \rightarrow \mathbb{C} : (x, v) \rightarrow x_1 v_2 - x_2 v_1,\tag{4}$$

and the associated Hamiltonian vector field X_L has integral curves which satisfy

$$\begin{aligned}\dot{x} &= e_3 \times x \\ \dot{v} &= e_3 \times v.\end{aligned}\tag{5}$$

Since $\tilde{\Phi}$ leaves H invariant, L is an integral of X_H . Let

$$\widetilde{\mathcal{E}M} : M \rightarrow \mathbb{C}^2 : (x, v) \rightarrow (H(x, v), L(x, v))$$

be the energy momentum mapping.

For later purposes it is convenient to introduce new coordinates $(\xi_1, \xi_2, z, \eta_1, \eta_2, y)$ on $\mathbb{C}^3 \times \mathbb{C}^3$ where

$$\xi_1 = x_1 + ix_2, \quad \xi_2 = x_1 - ix_2, \quad \eta_1 = v_1 + iv_2, \quad \eta_2 = v_1 - iv_2.$$

In these coordinates the defining equations of M are

$$\xi_1 \xi_2 + 1 = 1 - z^2 \quad \text{and} \quad \xi_1 \eta_2 + \xi_2 \eta_1 + 2zy = 0.$$

The energy momentum mapping $\widetilde{\mathcal{E}M}$ becomes

$$\mathcal{E}M : M \rightarrow \mathbb{C}^2 : (\xi, z, \eta, y) \rightarrow (c_1, c_2),\tag{6}$$

where

$$c_1 = \frac{1}{2} (\eta_1 \eta_2 + y^2) + z \quad \text{and} \quad c_2 = \frac{1}{i} (\xi_2 \eta_1 - \xi_1 \eta_2).$$

In addition, the $SO(2, \mathbb{C})$ -action $\widetilde{\Phi}$ becomes the \mathbb{C}^* -action

$$\Phi : \mathbb{C}^* \times M \rightarrow M : (\lambda, (\xi_1, \xi_2, z, \eta_1, \eta_2, y)) \rightarrow (\lambda\xi_1, \lambda^{-1}\xi_2, z, \lambda\eta_1, \lambda^{-1}\eta_2, y).$$

Proposition 2.1. Let M° be the subset of M consisting of all points $(\xi_1, \xi_2, z, \eta_1, \eta_2, y)$ where at least one of the monomials $\xi_1\xi_2, \xi_1\eta_2, \xi_2\eta_1, \eta_1\eta_2$ is nonzero. Then the following statements hold.

1. The \mathbb{C}^* -action Φ on M° is free and proper.
2. The map

$$\pi : M^\circ \rightarrow \mathbb{C}^4 : (\xi_1, \xi_2, z, \eta_1, \eta_2, y) \rightarrow (z, y, c_1, c_2)$$

is the quotient map $M^\circ \rightarrow M^\circ/\mathbb{C}^*$ of the \mathbb{C}^* -action Φ .

3. The image of the map π is the subvariety E° of \mathbb{C}^4 defined by

$$\{(z, y, c_1, c_2) \in \mathbb{C}^4 \mid y^2 = f_c(z)\} \setminus (\pm 1, 0, \pm 1, 0),$$

where

$$f_c(z) = 2(z^2 - 1)(z - c_1) - c_2^2. \quad (7)$$

4. The map $\pi : M^\circ \rightarrow E^\circ$ is the projection map of a \mathbb{C}^* -bundle over E° .

Proof.

1. First of all note that the map Φ_λ with $\lambda \neq 1$ has a fixed point if and only if $\xi_1 = \xi_2 = \eta_1 = \eta_2 = 0$. Hence \mathbb{C}^* acts freely on M° . To verify that the action Φ is proper, it suffices to show that for any two compact subsets K_1 and K_2 of M° the set $L = \{\lambda \in \mathbb{C}^* \mid K_1 \cap \Phi_\lambda(K_2) \neq \emptyset\}$ is compact. Since $K_2 \subseteq M^\circ$ is compact, at least one of the products $|\xi_1\xi_2|, |\xi_1\eta_2|, |\xi_2\eta_1|, |\eta_1\eta_2|$ restricted to K_2 is bounded below by a positive number. Suppose for example that $|\xi_1\xi_2|$ is bounded below by $\rho > 0$. In addition, the modulus of coordinates of points in K_1 are bounded above by some number R . Hence for every $\lambda \in L$ there is a point in K_2 whose ξ_1 and ξ_2 coordinates satisfy $|\lambda|\xi_1| < R$ and $|\lambda|^{-1}|\xi_2| < R$. Using the upper and lower bounds for $|\xi_1|$ and $|\xi_2|$ we find that $\rho/R^2 < |\lambda| < R^2/\rho$. Thus $L \subseteq \mathbb{C}$ is contained in a bounded set which is bounded away from zero. A standard argument using the continuity of Φ shows that L is closed. Consequently L is compact.

2. Given the equation of M and the definition of c_1, c_2 , the following equalities hold for points of M°

$$\begin{aligned}\xi_1\xi_2 &= 1 - z^2 \\ \xi_1\eta_2 &= -zy - ic_2 \\ \xi_2\eta_1 &= -zy + ic_2 \\ \eta_1\eta_2 &= 2(c_1 - z) - y^2.\end{aligned}\tag{8}$$

Note that the points of M° have the property that at least one of the right hand sides of the equations (8) is nonzero. Conversely, if we choose $p = (z, y, c_1, c_2)$ so that at least one of the right hand sides of (8) is nonzero, we see that the inverse image of p under the map π is a single \mathbb{C}^* -orbit. Hence π is the quotient map of the action Φ .

3. Note that the identity $(\xi_1\xi_2)(\eta_1\eta_2) = (\xi_1\eta_2)(\xi_2\eta_1)$ gives rise to

$$(1 - z^2)(2(c_1 - z) - y^2) = -(zy + ic_2)(-zy + ic_2),$$

which can be simplified to

$$y^2 = f_c(z) = 2(z^2 - 1)(z - c_1) - c_2^2.$$

4. Finally, if \mathbb{C}^* acts freely and properly on a complex manifold V , then the quotient map $V \rightarrow V/\mathbb{C}^*$ is the projection map of a \mathbb{C}^* -bundle over V/\mathbb{C}^* . In particular, this holds when $V = M^\circ$. \square

For any value of $c = (c_1, c_2)$ consider the affine elliptic curve

$$E_c : y^2 = f_c(z).$$

It is easy to check that E_c is smooth if and only if f_c has no multiple roots, that is, if and only if c does not lie on the discriminant locus Δ defined by

$$0 = \text{g.c.d.}(f_c, f'_c) = \frac{27}{4}c_2^4 + 2c_1(c_1^2 - 9)c_2^2 - 4(c_1^2 - 1)^2.\dagger\tag{9}$$

The quotient variety $E^\circ = M^\circ/\mathbb{C}^*$ can be considered as a family of affine elliptic curves E_c parametrized by $c = (c_1, c_2)$.

[†]Since $c_1 = \frac{3}{2}s - \frac{1}{2}\frac{1}{s}$, $c_2^2 = -\frac{1}{s}(1 - s^2)$, $s \in \mathbb{C}^*$ is a parametrization of Δ by rational functions, it follows that the genus its projectivization $\bar{\Delta}$ is 0. Because $\bar{\Delta}$ has three ordinary double points, it is an immersion of a 2-sphere in complex projective 2-space with three normal crossings.

Let

$$E^{ns} = \{(c_1, c_2, z, y) \in (\mathbb{C}^2 \setminus \Delta) \times \mathbb{C}^2 \mid y^2 = f_c(z)\}.$$

Then E^{ns} is the union of all smooth affine elliptic curves E_c . Since the points $(\pm 1, 0)$ lie in Δ , it follows that $E^{ns} \subseteq E^\circ$. Let M^{ns} be the inverse image of E^{ns} under the quotient map π and let M_c be the inverse image of E_c under π . Then M_c is the c -level set of the energy momentum map \mathcal{EM} . The geometrical structure of these level sets will become a major issue later on in this paper. By proposition 2.1 we know that $\pi : M^{ns} \rightarrow E^{ns}$ is a \mathbb{C}^* -bundle. If we restrict this bundle to the nonsingular curve E_c , we see that M_c is a \mathbb{C}^* -bundle over E_c .

That the discriminant locus Δ plays an important role can be seen from the following proposition.

Proposition 2.2. The derivative of the map $\mathcal{EM}|_{M^{ns}} : M^{ns} \rightarrow \mathbb{C}^2 \setminus \Delta$ has maximal rank at every point of M^{ns} .

Proof. The map $\mathcal{EM}|_{M^{ns}}$ is the composition of the projection map $\pi : M^{ns} \rightarrow E^{ns}$ followed by the map

$$\rho : E^{ns} \rightarrow \mathbb{C}^2 \setminus \Delta : (z, y, c_1, c_2) \rightarrow (c_1, c_2).$$

Since π is a quotient map of a free and proper action, its derivative has maximal rank. Thus it suffices to show that the derivative of ρ has maximal rank at every point of E^{ns} . Suppose not. Then there is a point p of E^{ns} where all the partial derivatives of $F(z, y, c_1, c_2) = -y^2 + f_c(z)$ vanish and $y^2 = f_c(z)$. Hence $y(p) = 0$ and $\frac{\partial f_c(z)}{\partial z}(p) = 0$. Moreover, $f_c(z(p)) = y^2(p) = 0$. Thus $z(p)$ is a double zero of f_c which is contrary to our assumption that $c \notin \Delta$. \square

3 A partial compactification of M_c and E_c

In this section we find a partial compactification of the affine elliptic curve E_c and the c -level set M_c of the energy momentum mapping.

Since the curves E_c in E^{ns} are affine, E_c is not complete. We would like to complete each E_c by adding the point at infinity. We do this as follows. The variety E^{ns} is an affine subvariety of $(\mathbb{C}^2 \setminus \Delta) \times \mathbb{C}^2$ with coordinates $((c_1, c_2), (z, y))$ where $y^2 = f_c(z)$ (7). We now embed the second factor \mathbb{C}^2 into the complex projective plane \mathbb{P}^2 by sending (z, y) to the projective point $(z : y : 1)$. Denote

the projective coordinates of \mathbb{P}^2 by $(z : y : u)$. Then E^{ns} can be completed to the variety \overline{E}^{ns} defined by

$$\left\{ (c_1, c_2), (z : y : u) \in (\mathbb{C}^2 \setminus \Delta) \times \mathbb{P}^2 \mid y^2 u = 2(z^2 - u^2)(z - c_1 u) - c_2^2 u^3 \right\}.$$

For each value of $c = (c_1, c_2) \notin \Delta$ the curve E_c has been completed by adding the point $((c_1, c_2), (0 : 1 : 0))$. We denote this completion by \overline{E}_c . Since $c \notin \Delta$ the curve \overline{E}_c is a smooth compact 1-dimensional complex manifold and hence is an elliptic curve.

We now complete the \mathbb{C}^* -bundle $\pi : M^{ns} \rightarrow E^{ns}$. Towards this goal, we introduce the variety $\overline{M}^{ns} \subseteq (\mathbb{C}^2 \setminus \Delta) \times \mathbb{P}^6$ with coordinates $((c_1, c_2), (\xi_1 : \xi_2 : z : \eta_1 : \eta_2 : y : u))$ defined by the equations

$$\begin{aligned} \xi_1 \xi_2 &= u^2 - z^2 \\ \xi_1 \eta_2 &= -zy - ic_2 u^2 \\ \xi_2 \eta_1 &= -zy + ic_2 u^2 \\ \eta_1 \eta_2 &= -y^2 - 2(c_1 u - z)u \end{aligned}$$

and the condition that at least one of the monomials $\xi_1 \xi_2, \xi_1 \eta_2, \xi_2 \eta_1, \eta_1 \eta_2$ is nonzero. On \overline{M}^{ns} we have a \mathbb{C}^* -action defined by

$$\begin{aligned} \Psi : \mathbb{C}^* \times \overline{M}^{ns} &\rightarrow \overline{M}^{ns} : \\ \left(\lambda, \left((c_1, c_2), (\xi_1 : \xi_2 : z : \eta_1 : \eta_2 : y : u) \right) \right) &\rightarrow \left((c_1, c_2), (\lambda \xi_1 : \lambda^{-1} \xi_2 : z : \lambda \eta_1 : \lambda^{-1} \eta_2 : y : u) \right). \end{aligned}$$

In a way similar to the proof of proposition 2.1 we can prove

Proposition 3.1. The map $\overline{\pi} : \overline{M}^{ns} \rightarrow \overline{E}^{ns}$ defined by

$$\overline{\pi} \left((c_1, c_2), (\xi_1 : \xi_2 : z : \eta_1 : \eta_2 : y : u) \right) = \left((c_1, c_2), (z : y : u) \right)$$

is the quotient map of the \mathbb{C}^* -action Ψ on \overline{M}^{ns} . Since Ψ is a free and proper \mathbb{C}^* -action, $\overline{\pi}$ is the projection map of a \mathbb{C}^* -bundle over \overline{E}^{ns} .

We easily check that M^{ns} embeds in \overline{M}^{ns} via

$$\left((c_1, c_2), (\xi_1, \xi_2, z, \eta_1, \eta_2, y) \right) \rightarrow \left((c_1, c_2), (\xi_1 : \xi_2 : z : \eta_1 : \eta_2 : y : 1) \right).$$

The restriction of $\overline{\pi}$ to M^{ns} is the projection map π of proposition 2.1. The inverse image of \overline{E}_c under $\overline{\pi}$ is denoted by \overline{M}_c . Thus \overline{M}_c is a \mathbb{C}^* -bundle over the elliptic curve \overline{E}_c .

4 Action angle coordinates

In this section we obtain a complex analogue of the period lattice and action angle coordinates for the complexified spherical pendulum.

We begin by choosing good coordinates on M^{ns} in which to express the symplectic form ω . Since ω is holomorphic on M it is holomorphic on M^{ns} . Let V be the Zariski open subset of M^{ns} given by

$$M^{ns} \cap \{(\xi_1, \xi_2, z, \eta_1, \eta_2, y) \in \mathbb{C}^6 \mid \xi_1 \xi_2 \neq 0, y \neq 0\}.$$

At every point p of E^{ns} with $y \neq 0$ the variables (ξ_1, z, c_1, c_2) can be used as local coordinates at p . At any point p of V we have $\xi_1 \neq 0$. Supposing that the values of (ξ_1, z, c_1, c_2) are known, then the values of ξ_2, η_1, η_2 follow from

$$\xi_2 = \frac{(1 - z^2)}{\xi_1}, \quad \eta_1 = \left(\frac{-zy + ic_2}{1 - z^2} \right) \xi_1, \quad \eta_2 = \frac{-zy - ic_2}{\xi_1}$$

by virtue of the defining equations of M^{ns} . Thus at every point of V the variables (ξ_1, z, c_1, c_2) can be used as local coordinates.

We now would like to write the symplectic form ω in terms of these coordinates.

Proposition 4.1. The symplectic form ω on M restricted to V is

$$\frac{dz}{y} \wedge dc_1 + i \left(\frac{zy - ic_2}{z^2 - 1} \frac{dz}{y} - \frac{d\xi_1}{\xi_1} \right) \wedge dc_2. \quad (10)$$

Proof. To see this, note that the canonical symplectic form on \mathbb{C}^6 can be written as $\frac{1}{2} (d\xi_1 \wedge d\eta_2 + d\xi_2 \wedge d\eta_1) + dz \wedge dy$. Since $\eta_1 = \frac{-zy + ic_2}{\xi_2}$ and $\eta_2 = -\frac{zy + ic_2}{\xi_1}$ on V , we obtain

$$d\eta_1 = \frac{-zy + ic_2}{\xi_2^2} d\xi_2 - \frac{1}{\xi_2} d(zy) + i \frac{1}{\xi_2} dc_2$$

and

$$d\eta_2 = \frac{zy + ic_2}{\xi_1^2} d\xi_1 - \frac{1}{\xi_1} d(zy) - i \frac{1}{\xi_1} dc_2.$$

Hence

$$\omega|_V = dz \wedge dy - \frac{1}{2} \left(\frac{d\xi_1}{\xi_1} + \frac{d\xi_2}{\xi_2} \right) \wedge d(zy) + \frac{1}{2i} \left(\frac{d\xi_1}{\xi_1} - \frac{d\xi_2}{\xi_2} \right) \wedge dc_2. \quad (11)$$

Since $\xi_1 \xi_2 = 1 - z^2$ on V , which differentiating gives

$$\frac{d\xi_1}{\xi_1} + \frac{d\xi_2}{\xi_2} = -\frac{2z}{1-z^2} dz.$$

Whence

$$\omega|_V = \frac{1}{1-z^2} dz \wedge dy + i \left(\frac{z}{z^2-1} dz - \frac{d\xi_1}{\xi_1} \right) \wedge dc_2. \quad (12)$$

Differentiating $y^2 = 2(c_1 - z)(1 - z^2) - c_2^2$ gives

$$dy = \frac{(3z^2 - 2c_1z - 1)}{y} dz + \frac{(1 - z^2)}{y} dc_1 - \frac{c_2}{y} dc_2,$$

which substituted into (12) yields (10). \square

We now prove

Proposition 4.2. The symplectic form ω can be extended holomorphically to \overline{M}^{ns} .

Proof. We will use equation (10) of proposition 4.1. We must extend ω to points of $\overline{M}^{ns} \setminus M^{ns}$. Consider the rational functions $s = z/y$ and $t = 1/y$ on E^{ns} . These functions can be extended holomorphically to the point at infinity of each E_c . At infinity they take the values $s = 0, t = 0$, where s is the local parameter at infinity. Also define new variables $\sigma_i = \xi_i/y$ and $\tau_i = \eta_i/y$ for $i = 1, 2$. In terms of these variables the defining equations of an affine part of \overline{M}^{ns} read

$$\begin{aligned} \sigma_1 \sigma_2 &= t^2 - s^2 \\ \sigma_1 \tau_2 &= -s - ic_2 t^2 \\ \sigma_2 \tau_1 &= -s + ic_2 t^2 \\ \tau_1 \tau_2 &= -1 + 2(c_1 t^2 - st). \end{aligned}$$

This affine part includes the points of $\overline{M}^{ns} \setminus M^{ns}$. Moreover, the defining equation for E^{ns} becomes

$$t = 2(s^2 - t^2)(s - c_1 t) - c_2^2 t^3. \quad (13)$$

From the above equations we see that $\tau_1 \tau_2 = -1$ when $s = t = 0$. In particular, τ_1 and τ_2 are nonzero above the points at infinity of M^{ns} . Hence $\sigma_1 = \sigma_2 = 0$ there. Let us rewrite ω in terms of s, t, τ_1 , and τ_2 . A straightforward calculation gives

$$\frac{dz}{y} = ds - \frac{s}{t} dt.$$

Differentiating (13) we obtain

$$dt = \frac{(6s^2 - 2c_1st - 2t^2) ds - 2t(s^2 - t^2) dc_1 - 2c_2t^3 dc_2}{1 + 4t(s - c_1t) + 2c_1(s^2 - t^2) + 3c_2^2t^2}.$$

Thus $\frac{s}{t} dt$ is holomorphic at $s = t = 0$ except possibly for the term $6 \frac{s^3}{t} dt$. However, using (13) we see that s^3 equals t times a polynomial in s and t . Thus s^3/t is holomorphic at infinity and hence the 1-form $\frac{dz}{y}$ can be holomorphically extended to the points above $s = t = 0$. A similar argument holds for the 2-form $\frac{dz}{y} \wedge dc_1$.

Now let us turn to the part of ω with dc_2 . First note that $\frac{c_2}{z^2-1} \frac{dz}{y}$ is holomorphic at infinity. Hence we can ignore that term. Next we rewrite the 1-form $\frac{zy}{z^2-1} \frac{dz}{y} - \frac{d\xi_1}{\xi_1}$. Using the relation $\xi_1 = -(zy + ic_2)/y\tau_2$ we obtain

$$\frac{d\tau_2}{\tau_2} + \frac{z dz}{z^2 - 1} + \frac{dy}{y} - \frac{d(zy + ic_2)}{zy + ic_2}.$$

The 1-form $\frac{d\tau_2}{\tau_2}$ is holomorphic at infinity since $\tau_2 \neq 0$ there. The last three terms are logarithmic derivatives which together equal

$$\frac{1}{2} \frac{d[(1 - 1/z^2)/(1 + ic_2/zy)]}{(1 - 1/z^2)/(1 + ic_2/zy)^2}.$$

Since $(1 - 1/z^2)/(1 + ic_2/zy)^2$ is holomorphic at infinity, its logarithmic derivative is also. Hence ω is holomorphic at infinity. \square

Corollary 4.3. Let $c = (c_1, c_2) \notin \Delta$. The differential forms

$$\omega_1 = \frac{dz}{y}, \quad \omega_2 = i \left(\frac{zy - ic_2}{z^2 - 1} \right) \frac{dz}{y} - \frac{d\xi_1}{\xi_1}$$

restricted to the Zariski open subset of M_c where $y(z^2 - 1) \neq 0$ can be extended holomorphically to all of \overline{M}_c . The extension of the 1-forms ω_1, ω_2 to \overline{M}_c will again be denoted by ω_1, ω_2 .

Proof. This follows from the observation that the 3-forms $\frac{dz}{y} \wedge dc_1 \wedge dc_2 = \omega \wedge dc_2$ and $\left(i \frac{zy - ic_2}{z^2 - 1} \frac{dz}{y} - \frac{d\xi_1}{\xi_1} \right) \wedge dc_2 \wedge dc_1 = \omega \wedge dc_1$ are holomorphic on \overline{M}^{ns} . \square

Motivated by the form of ω in proposition 4.1 we construct the complex analogue of the period lattice on \overline{M}_c as follows. Until further notice we assume that $c \notin \Delta$. Fix a base point $p_0 \in \overline{M}_c$ and consider the *Abel-Jacobi map*

$$\mathcal{AJ} : \overline{M}_c \rightarrow \mathbb{C}^2 : p \rightarrow (u_1(p), u_2(p)) = \left(\int_{p_0}^p \omega_1, \int_{p_0}^p \omega_2 \right), \quad (14)$$

where the integrals are taken along a path γ on \overline{M}_c which joins p_0 to p . Since the 1-forms ω_1 and ω_2 are closed, the functions u_1 and u_2 depend only on the homology class of the path on \overline{M}_c joining p_0 to p . The map \mathcal{AJ} is *multivalued* due to the fact that p_0 and p can be joined by nonhomologous paths, say γ and $\tilde{\gamma}$. The difference in the corresponding values of \mathcal{AJ} is given by a vector

$$\left(\int_{\Gamma} \omega_1, \int_{\Gamma} \omega_2 \right), \quad (15)$$

where Γ is the closed path beginning and ending at p_0 which is formed by first tracing out γ and then tracing out $\tilde{\gamma}$ backwards. Thus the multivaluedness of \mathcal{AJ} is given by vectors of the form (15) where Γ ranges over $H_1(\overline{M}_c, \mathbb{Z})$. Let

$$L_c = \left\{ \left(\int_{\Gamma} \omega_1, \int_{\Gamma} \omega_2 \right) \in \mathbb{C}^2 \mid \Gamma \in H_1(\overline{M}_c, \mathbb{Z}) \right\}.$$

Theorem 4.4. Let γ_1, γ_2 be a \mathbb{Z} -basis of $H_1(\overline{E}_c, \mathbb{Z})$. Suppose that the curves γ_i avoid the points of \overline{E}_c where $z^2 = 1$ or the point at infinity. For $j = 1, 2$ define

$$\lambda_j = \int_{\gamma_j} \frac{dz}{y}, \quad \mu_j = i \int_{\gamma_j} \frac{zy - ic_2}{z^2 - 1} \frac{dz}{y}. \quad (16)$$

Then L_c is a lattice in \mathbb{C}^2 called the *period lattice*. L_c has \mathbb{Z} -rank three and is generated by the vectors

$$(\lambda_1, \mu_1), \quad (\lambda_2, \mu_2), \quad (0, 2\pi).$$

Proof. Let $U_c = \overline{E}_c \setminus \{z^2 = 1, \infty\}$. Note that the three points $(1, -ic_2)$, $(-1, ic_2)$ and infinity are precisely the poles of the differential form $\Omega = i \frac{zy - ic_2}{z^2 - 1} \frac{dz}{y}$ on \overline{E}_c which are all of first order. Since $\Omega = i \left(\frac{zy - ic_2}{y(z+1)} \right) \frac{dz}{z-1}$, the residue of the pole at $(1, -ic_2)$ is $-i$. Similarly the residue at $(-1, ic_2)$ is $-i$. Hence the residue at infinity is $2i$. Consequently, the value of $\int_{\gamma} \Omega$, where γ is a closed loop in U_c , is determined by the homology class of $\gamma \in H_1(\overline{E}_c, \mathbb{Z})$ up to an integer multiple of 2π .

Let Γ be any closed path in \overline{M}_c . Without loss of generality we can assume that Γ lies in the open subset $V_c = \overline{\pi}^{-1}(U_c)$. Since $z^2 \neq 1, \infty$ in V_c , it follows that $\xi_1 \neq 0$. Moreover, the map

$$V_c \rightarrow U_c \times \mathbb{C}^* : (\xi_1, \xi_2, z, \eta_1, \eta_2, y) \rightarrow (z, y, \xi_1) \quad (17)$$

is an diffeomorphism. The image in U_c of the curve Γ under the projection $\bar{\pi}$ is homologous to an integral linear combination of γ_1 and γ_2 , say $\bar{\pi}(\Gamma) = m_1\gamma_1 + m_2\gamma_2$. Hence

$$\int_{\Gamma} \omega_1 = \int_{\bar{\pi}(\Gamma)} \frac{dz}{y} = m_1\lambda_1 + m_2\lambda_2$$

and

$$\int_{\Gamma} \omega_2 = i \int_{\Gamma} \frac{d\xi_1}{\xi_1} + i \int_{\bar{\pi}(\Gamma)} \frac{zy - ic_2}{z^2 - 1} \frac{dz}{y} = i \int_{\Gamma} \frac{d\xi_1}{\xi_1} + (m_1\mu_1 + m_2\mu_2) \bmod 2\pi.$$

Since we can let Γ encircle $\xi_1 = 0$ as many times as we want, the integral $i \int_{\Gamma} \frac{d\xi_1}{\xi_1}$ can be any multiple of 2π . Consequently, the set of values of $\int_{\Gamma} \omega_2$, where $\bar{\pi}(\Gamma)$ is homologous to $m_1\gamma_1 + m_2\gamma_2$, is $m_1\mu_1 + m_2\mu_2 + 2k\pi$ where $k \in \mathbb{Z}$ is arbitrary. This proves the theorem. \square

Another way of stating the result of theorem 4.4 is that the vectors

$$\begin{pmatrix} 0 \\ 2\pi \end{pmatrix}, \begin{pmatrix} \int_{\bar{\pi}^{-1}(\gamma_1)} \omega_1 \\ \int_{\bar{\pi}^{-1}(\gamma_1)} \omega_2 \end{pmatrix}, \begin{pmatrix} \int_{\bar{\pi}^{-1}(\gamma_2)} \omega_1 \\ \int_{\bar{\pi}^{-1}(\gamma_2)} \omega_2 \end{pmatrix} \quad (18)$$

generate the period lattice L_c .

Because $\omega_1 \wedge \omega_2 \neq 0$, the Abel-Jacobi map \mathcal{AJ} is locally invertible. It will turn out that this local inverse can be extended to all of \mathbb{C}^2 , thus defining a global holomorphic inverse. Although we could prove this at this point, we prefer to give an explicit construction of the inverse to \mathcal{AJ} in the next section.

Having discussed the period lattice we now prove the following complex analogue of action angle coordinates.

Proposition 4.5. For each fixed $c \in \mathbb{C}^2 \setminus \Delta$ there is an open neighborhood U of \bar{M}_c in \bar{M}^{ns} and local coordinates $(c_1, c_2, \tilde{u}_1, \tilde{u}_2)$ on U such that the symplectic form ω is $d\tilde{u}_1 \wedge dc_1 + d\tilde{u}_2 \wedge dc_2$. The coordinates $(c_1, c_2, \tilde{u}_1, \tilde{u}_2)$ are called *complex action-angle coordinates*. In action-angle coordinates the vector fields X_H and X_L on \bar{M}_c are $\frac{\partial}{\partial \tilde{u}_1}$ and $\frac{\partial}{\partial \tilde{u}_2}$, respectively.

Before proving the proposition we construct some preliminary local coordinates. To construct them we need a local section of the surjective submersion

$$\bar{\sigma} : \bar{M}^{ns} \rightarrow \mathbb{C}^2 \setminus \Delta : ((\xi_1 : \xi_2 : z : \eta_1 : \eta_2 : y : u), (c_1, c_2)) \rightarrow (c_1, c_2).$$

Note that the map $\bar{\sigma}$ is the composition the surjective submersion

$$\bar{\pi} : \bar{M}^{ns} \rightarrow \bar{E}^{ns}$$

with the surjective submersion

$$\bar{\rho} : \bar{E}^{ns} \rightarrow \mathbb{C}^2 \setminus \Delta : ((z : y : u), (c_1, c_2)) \rightarrow (c_1, c_2).$$

Since the fiber $\bar{\rho}^{-1}(c_1, c_2)$ is \bar{E}_c , which is compact, $\bar{\rho}$ is a locally trivial bundle. By construction the bundle $\bar{\pi}$ is locally trivial. Consequently, $\bar{\sigma}$ is a locally trivial bundle. Fix $c \in \mathbb{C}^2 \setminus \Delta$. Then $\bar{\sigma}^{-1}(c) = \bar{M}_c$. There is an open neighborhood W of c in $\mathbb{C}^2 \setminus \Delta$ such that the bundle $\bar{\sigma}|_U : U = \bar{\sigma}^{-1}(W) \rightarrow W$ is trivial. Let

$$\tau : W \subseteq \mathbb{C}^2 \setminus \Delta \rightarrow U = \bar{\sigma}^{-1}(W) \subseteq \bar{M}^{ns}$$

be a section of $\bar{\sigma}|_U$. We now prove

Lemma 4.6. The map

$$\begin{aligned} \psi : U = \bar{\sigma}^{-1}(W) \subseteq \bar{M}^{ns} &\rightarrow W \times \mathbb{C}^2 \subseteq W \times (\mathbb{C}^2 \setminus \Delta) : \\ p &\rightarrow \left(\bar{\sigma}(p), \int_{\tau(\bar{\sigma}(p))}^p \omega_1, \int_{\tau(\bar{\sigma}(p))}^p \omega_2 \right) = (c_1, c_2, \widehat{u}_1, \widehat{u}_2) \end{aligned} \quad (19)$$

is a local diffeomorphism.

Proof. Since $\dim \bar{\sigma}^{-1}(W) = \dim(W \times \mathbb{C}^2)$, it suffices to show that ψ is a submersion. Let π_1, π_2, π_3 be the projections onto the first, second and third factors of $W \times (\mathbb{C} \times \mathbb{C})$. Clearly the map $\pi_1 \circ \psi$ is a submersion. Now

$$d(\pi_2 \circ \psi)(p) = d\widehat{u}_1(p) = \omega_1(p) - \tau^* \omega_1(\bar{\sigma}(p))$$

and

$$d(\pi_3 \circ \psi)(p) = d\widehat{u}_2(p) = \omega_2(p) - \tau^* \omega_2(\bar{\sigma}(p)).$$

But $d(\pi_2 \circ \psi)(p) \wedge d(\pi_3 \circ \psi)(p)$ is nonzero because $\omega_1 \wedge \omega_2 = e_1 dz \wedge d\xi_1$ is nonzero and

- a). $\omega_1 \wedge \tau^* \omega_2 = f_1 dz \wedge dc_1 + f_2 dz \wedge dc_2,$
- b). $\omega_2 \wedge \tau^* \omega_1 = g_1 dz \wedge dc_1 + g_2 d\xi_1 \wedge dc_1 + g_3 dz \wedge dc_2 + g_4 d\xi_1 \wedge dc_2,$
- c). $\tau^* \omega_1 \wedge \tau^* \omega_2 = h_1 dc_1 \wedge dc_2.$

Therefore the map $(\pi_2 \times \pi_3) \circ \psi$ is a submersion, which implies that ψ is a submersion. \square

Proof of proposition 4.5. From lemma 4.6 it follows that $(c_1, c_2, \widehat{u}_1, \widehat{u}_2)$ are local coordinates on U . We now compute the 2-form

$$\begin{aligned} \widehat{d}u_1 \wedge dc_1 + \widehat{d}u_2 \wedge dc_2 &= (\omega_1 - \tau^* \omega_1) \wedge dc_1 + (\omega_2 - \tau^* \omega_2) \wedge dc_2 \\ &= \omega_1 \wedge dc_1 - h_1(c_1, c_2) dc_1 \wedge dc_2 + \omega_2 \wedge dc_2 - h_2(c_1, c_2) dc_1 \wedge dc_2, \\ &\quad \text{since } \tau^* \omega_1 \text{ and } \tau^* \omega_2 \text{ are 1-forms on } W \times \mathbb{C}^2 \\ &= \omega_1 \wedge dc_1 + \omega_2 \wedge dc_2 - h(c_1, c_2) dc_1 \wedge dc_2. \end{aligned} \quad (20)$$

The 2-form $h(c_1, c_2) dc_1 \wedge dc_2$ is closed. Choosing the neighborhood W to be simply connected and star shaped, it follows that there is a 1-form $\theta(c_1, c_2) = \theta_1 dc_1 + \theta_2 dc_2$ such that

$$-h(c_1, c_2) dc_1 \wedge dc_2 = d\theta(c_1, c_2). \quad (21)$$

Adjust the local section τ of the trivial bundle $\overline{\sigma}|U$ so as to define new angle coordinates

$$\widetilde{u}_1 = \widehat{u}_1 - \theta_1 \quad \widetilde{u}_2 = \widehat{u}_2 - \theta_2.$$

In action-angle coordinates $(c_1, c_2, \widetilde{u}_1, \widetilde{u}_2)$ we have

$$\begin{aligned} \widetilde{d}u_1 \wedge dc_1 + \widetilde{d}u_2 \wedge dc_2 &= (\widehat{d}u_1 - d\theta_1) \wedge dc_1 + (\widehat{d}u_2 - d\theta_2) \wedge dc_2 \\ &= \widehat{d}u_1 \wedge dc_1 + \widehat{d}u_2 \wedge dc_2 - d\theta(c_1, c_2) \\ &= \omega_1 \wedge dc_1 + \omega_2 \wedge dc_2, \end{aligned} \quad (22)$$

using (21) and (20). But the right hand side of (22) is the symplectic form ω on $U = \overline{\sigma}^{-1}(W)$.

In action-angle coordinates on U , the Hamiltonian function H (6) is c_1 . Thus

$$dc_1 = X_H \lrcorner (\widetilde{d}u_1 \wedge dc_1 + \widetilde{d}u_2 \wedge dc_2),$$

which implies that $X_H = \frac{\partial}{\partial \widetilde{u}_1}$. Similarly the momentum function L (6) is c_2 , which implies that $X_L = \frac{\partial}{\partial \widetilde{u}_2}$. This completes the proof of proposition 4.5. \square

An immediate consequence of proposition 4.5 is the fact that the complex spherical pendulum is Mumford-Jacobi completely integrable.

5 The inverse of the Abel-Jacobi map

In this section we show that the local inverse of the Abel-Jacobi map \mathcal{AJ} (14) constructed in preceding section extends to a global inverse defined on all of \mathbb{C}^2 .

We use the same notation as in theorem 4.4. We now recall the basic properties of the Weierstrass elliptic functions \wp , ζ , and σ . For more details see [10, chpt. XX, pp. 429–446]. Let Λ_c be the lattice in \mathbb{C} generated by the complex numbers λ_1 and λ_2 (16). Corresponding to Λ_c there is the Weierstrass \wp -function. There is also the ζ -function which is characterized by the properties: 1). $\zeta'(u) = -\wp(u)$ and 2). $\zeta(u) = \frac{1}{u} + O(u)$ about $u = 0$. The ζ function is not periodic. In fact there are two complex numbers η_1 and η_2 , called quasi-periods of \bar{E}_c , such that

$$\zeta(u + \lambda_1) = \zeta(u) + \eta_1, \quad \zeta(u + \lambda_2) = \zeta(u) + \eta_2.$$

More generally, corresponding to every $\lambda \in \Lambda_c$ there is a unique quasi-period $\eta(\lambda)$ such that $\zeta(u + \lambda) = \zeta(u) + \eta(\lambda)$. When $\lambda = m_1\lambda_1 + m_2\lambda_2$, we see that $\eta(\lambda) = m_1\eta_1 + m_2\eta_2$. Finally, the Weierstrass σ -function is defined by

$$\sigma(u) = u \prod_{\lambda \in \Lambda_c - \{0\}} \left(1 - \frac{u}{\lambda}\right) e^{u/\lambda + (u/\lambda)^2/2},$$

which satisfies $\sigma'(u)/\sigma(u) = \zeta(u)$. Furthermore, σ has the property that

$$\sigma(u + \lambda) = e^{\lambda/2 + \eta(\lambda)u} \sigma(u)$$

for every $\lambda \in \Lambda_c$.

To construct the inverse of the Abel-Jacobi map \mathcal{AJ} (14) we will need a product of σ -functions. Why the choices are made will become clear in the proof of theorem 5.1 below. Choose $a, b \in \mathbb{C}$ so that

$$2\wp(a) + \frac{1}{3}c_1 = 1, \quad 2\wp'(a) = -ic_2, \quad 2\wp(b) + \frac{1}{3}c_1 = -1, \quad 2\wp'(b) = ic_2.$$

Then define the function

$$\Sigma(u) = e^{(\zeta(a) + \zeta(b))u} \frac{\sigma(u-a)\sigma(u-b)}{\sigma(a)\sigma(b)\sigma(u)^2}.$$

As a consequence of the properties of the σ -function we obtain

$$\Sigma(u + \lambda) = \exp\left(- (a + b)\eta(\lambda) + (\zeta(a) + \zeta(b))\lambda\right)\Sigma(u).$$

Moreover, the principal part of Σ at $u = 0$ is $\frac{1}{u^2}$.

Theorem 5.1. For the point $p_0 \in \overline{M}_c$ in the definition of the Abel-Jacobi map \mathcal{AJ} (14) we take $(s, t, \sigma_1, \sigma_2, \tau_1, \tau_2) = (0, 0, 0, 0, 1, -1)$. Then the map

$$\mathcal{W} : \mathbb{C}^2 \rightarrow \overline{M}_c : (u_1, u_2) \rightarrow (z, y, \xi_1) = \left(2\wp(u_1) + \frac{1}{3}c_1, 2\wp'(u_1), e^{iu_2}\Sigma(u_1) \right) \quad (23)$$

is a global inverse of the Abel-Jacobi map. Since every fiber of \mathcal{W} is the lattice L_c , it follows that $\overline{M}_c = \mathbb{C}^2/L_c$.

Proof. First we note that the inverse of the map

$$\kappa : \overline{E}_c \rightarrow \mathbb{C} : \overline{\pi}(p) \rightarrow u_1(\overline{\pi}(p)) = \int_{\overline{\pi}(p_0)}^{\overline{\pi}(p)} \omega_1 = \int_{\overline{\pi}(p_0)}^{\overline{\pi}(p)} \frac{dz}{y}$$

is given by

$$\rho : \mathbb{C} \rightarrow \overline{E}_c : u_1 \rightarrow (2\wp(u_1) + \frac{1}{3}c_1, 2\wp'(u_1)) = (z, y). \quad (24)$$

To see this replace the variables z and y in ω_1 by their Weierstrass parametrizations. Then $du_1 = \frac{dz}{y}$ and $\rho(0) = \infty \in \overline{E}_c$.

We now consider $\int_{p_0}^p \omega_2$. First we replace the variables z, y in ω_2 by their Weierstrass parametrizations. Then $\frac{zy-ic_2}{z^2-1}$ becomes an elliptic function on \overline{E}_c with poles only at $a + \Lambda_c, b + \Lambda_c$ and $0 + \Lambda_c$, which are of first order with residues 1, 1, and -2 , respectively. Observe that the elliptic function

$$Z(u_1) = \zeta(u_1 - a) + \zeta(u_1 - b) - 2\zeta(u_1)$$

on \overline{E}_c has the same poles and residues as $\frac{zy-ic_2}{z^2-1}$. Thus the difference of $\frac{zy-ic_2}{z^2-1} - Z(u_1)$ is a holomorphic elliptic function and hence is a bounded entire function on \mathbb{C} , which is constant by Liouville's theorem. To compute this constant we evaluate the difference at $u_1 = 0$. We note that $\frac{zy-ic_2}{z^2-1} = -\frac{2}{u_1} + O(u_1)$ and $Z(u_1) = -\frac{2}{u_1} - \zeta(a) - \zeta(b) + O(u_1)$. Thus the constant is $\zeta(a) + \zeta(b)$, that is,

$$\frac{zy-ic_2}{z^2-1} = \zeta(u_1 - a) + \zeta(u_1 - b) - 2\zeta(u_1) + \zeta(a) + \zeta(b) = \frac{\Sigma'(u_1)}{\Sigma(u_1)}.$$

The second equality in the equation above explains why the function Σ was constructed. For the moment let p_1 be a point in \overline{M}_c . Then

$$u_2 = \int_{p_1}^p \omega_2 = i \int_{p_1}^p \frac{z(u_1)y(u_1) - ic_2}{z^2(u_1) - 1} du_1 - i \int_{p_1}^p \frac{d\xi_1}{\xi_1}$$

$$\begin{aligned}
&= i \int_{p_1}^p \frac{\Sigma'(u_1)}{\Sigma(u_1)} du_1 - i \int_{p_1}^p \frac{d\xi_1}{\xi_1} \\
&= i \log\left(\frac{\Sigma(u_1)}{\Sigma(u_1(p_1))}\right) - i \log\left(\frac{\xi_1}{\xi(p_1)}\right) \\
&= i \log(\Sigma(u_1)) - i \log \xi_1 + i \log\left(\frac{\xi(p_1)}{\Sigma(u_1(p_1))}\right).
\end{aligned}$$

We now let p_1 converge to the point p_0 given in the statement of the theorem. Recall that $\Sigma(u_1)$ has principal part $\frac{1}{u_1^2}$ at $u = 0$. Observe that

$$\xi_1 = -\frac{zy + ic_2}{y\tau_2} = -\frac{1}{\tau_2} \left(\frac{1}{u_1^2} + \dots \right),$$

using (8) and the fact that $y\tau_2 = \eta_2$. Since τ_2 converges to -1 as p_1 converges to p_0 , we see that $\lim_{p_1 \rightarrow p_0} \left(\frac{\xi(p_1)}{\Sigma(u_1(p_1))} \right) = 1$. Hence

$$u_2 = i \log \Sigma(u_1) - i \log \xi_1,$$

which implies that $\xi_1 = e^{i u_2} \Sigma(u_1)$.

Since (ξ_1, z, y) are coordinates for points p in a Zariski open subset V_c of \overline{M}_c (see (17)) and $(\mathcal{W} \circ \mathcal{AJ})(p) = p$, it follows that this equality holds for all points in \overline{M}_c . Thus \mathcal{W} (23) is a global inverse of the Abel-Jacobi map \mathcal{AJ} .

In order to show that each fiber of \mathcal{W} is the lattice L_c , we first compute an explicit expression for the period

$$\mu_j = i \int_{\gamma_j} \frac{zy - ic_2}{z^2 - 1} \frac{dz}{y},$$

see (16). Using the Weierstrass parametrization of \overline{E}_c and choosing the curve

$$[0, 1] \rightarrow \mathbb{C} : t \rightarrow t\lambda_j + (1-t)u_1^0, \quad u_1^0 \neq 0, a, b + \Lambda_c,$$

whose image under the map ρ (24) is γ_j , we find that

$$\begin{aligned}
\mu_j &= i \int_{u_1^0}^{u_1^0 + \lambda_j} \frac{z(u_1)y(u_1) - ic_2}{z^2(u_1) - 1} du_1 = i \int_{u_1^0}^{u_1^0 + \lambda_j} d \log \Sigma(u_1) \\
&= i \log\left(\frac{\Sigma(u_1^0 + \lambda_j)}{\Sigma(u_1^0)}\right) = -i \left[(a+b)\eta_j - (\zeta(a) + \zeta(b))\lambda_j \right].
\end{aligned}$$

In other words,

$$\mu(\lambda) = -i[(a+b)\eta(\lambda) - (\zeta(a) + \zeta(b))\lambda].$$

Now suppose that $(u_1, u_2), (u'_1, u'_2) \in \mathcal{W}^{-1}(z, y, \xi_1)$. Then u_1 and u'_1 correspond to the point $(z, y) = (2\wp(u_1) + \frac{1}{3}c_1, 2\wp'(u_1))$ on \overline{E}_c . Hence there is a $\lambda \in \Lambda_c$ such that $u'_1 = u_1 + \lambda$. Since u_2 and u'_2 correspond to

$$\xi_1 = e^{iu_2}\Sigma(u_1) = e^{iu'_2}\Sigma(u'_1) = e^{iu'_2}\Sigma(u_1 + \lambda),$$

it follows that

$$\begin{aligned} u_2 - u'_2 &= -i \log \left(\frac{\Sigma(u_1 + \lambda)}{\Sigma(u_1)} \right) \\ &= i[(a+b)\eta(\lambda) - (\zeta(a) + \zeta(b))\lambda] + 2\pi i k, \quad \text{for some } k \in \mathbb{Z} \\ &= -\mu(\lambda) + 2\pi i k. \end{aligned}$$

Thus (u_1, u_2) and (u'_1, u'_2) differ by some element of L_c . Consequently, each fiber of \mathcal{W} is the lattice L_c , which shows that \overline{M}_c is holomorphically diffeomorphic to \mathbb{C}^2/L_c . \square

6 Monodromy

In this section we show that the complexified spherical pendulum has monodromy. More precisely, we show that there is a noncontractible loop Γ in $\mathbb{C}^2 \setminus \Delta$ such that the bundle of period lattices

$$\nu : \bigsqcup_{c \in \gamma} L_c \rightarrow \Gamma$$

over Γ has classifying map given by $\begin{pmatrix} 1 & -1 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}$.

For the moment, consider the classical spherical pendulum. Let c lie in the set of regular values \mathcal{R} of its energy momentum mapping em . Then the motion of the spherical pendulum takes place on a smooth 2-dimensional torus T_c^2 in phase space TS^2 . The image of T_c^2 under the tangent bundle projection map is an annular region \mathcal{A} on S^2 bounded by two small circles lying in planes parallel to the equatorial plane. There are two important quantities associated to the motion on T_c^2 : one is the time $T(c)$ it takes starting on the upper boundary of \mathcal{A} to the motion to return to this boundary for the first time and the other is the angle $\Theta(c)$

measured along the upper boundary from starting point to first return point. As is well known

$$T(c) = 2 \int_{z_0}^{z_1} \frac{dz}{y} \quad \text{and} \quad \Theta(c) = 2 \int_{z_0}^{z_1} \frac{c_2}{1-z^2} \frac{dz}{y}, \quad (25)$$

where $y^2 = f_c(z) = 2(c_1 - z)(1 - z^2) - c_2^2$ and the discriminant of f_c is positive. Here $-1 < z_0 < z_1 < 1 < z_2$ are the three distinct real roots of f_c . Observe that $T(c)$ is the period of a holomorphic 1-form on \overline{E}_c and $\Theta(c)$ is a period of a differential form of the *third kind* on \overline{E}_c . In [5], [4, chpt. IV] it was shown that the real analytic function Θ on \mathcal{R} is multivalued – decreasing by 2π along any positively oriented closed curve Γ in \mathcal{R} which encircles the point $(1, 0)$. Another way to say this is to use the real period lattice \mathcal{L}_c defined by the 2-torus T_c^2 . The lattice \mathcal{L}_c is spanned by the vectors

$$X_L(p) \quad \text{and} \quad T(em(p)) \cdot X_H(p) + \frac{\Theta(em(p))}{2\pi} \cdot X_L(p),$$

where $p \in T^2(c)$ and X_H and X_L are the Hamiltonian vector fields corresponding to the energy and angular momentum of the spherical pendulum. Then the bundle of real period lattices

$$\tilde{\nu} : \coprod_{c \in \Gamma} \mathcal{L}_c \rightarrow \Gamma$$

over Γ has classifying map $\begin{pmatrix} 1 & -1 \\ 0 & 1 \end{pmatrix}$.

The following lemma describes the relation between the differential form Θ of the third kind on \overline{E}_c with the holomorphic 1-form ω_2 on \overline{M}_c .

Lemma 6.1. The period $\int_\gamma \omega_2$ modulo π is minus the period $\int_{\pi \circ \gamma} \Theta$ for any closed loop γ on \overline{M}_c which avoids the points with $z^2 = 1$.

Proof. First note that

$$\int_\gamma \Theta = \int_\gamma \frac{c_2}{1-z^2} \frac{dz}{y} = -i \int_\gamma \frac{(zy - ic_2)}{z^2 - 1} \frac{dz}{y} + i \int_\gamma \frac{zy}{z^2 - 1} \frac{dz}{y}.$$

By theorem 4.4 the periods of $i \frac{zy - ic_2}{(z^2 - 1)y} \frac{dz}{y}$, taken over \overline{E}_c , are, modulo 2π , equal to the periods of the form ω_2 . The periods of $i \frac{zy}{z^2 - 1} \frac{dz}{y}$ are multiples of π , since this 1-form is equal to $\frac{i}{2}$ times the logarithmic derivative $\frac{d(z^2 - 1)}{z^2 - 1}$. \square

The main goal of this section is to prove the following

Theorem 6.2. Consider the periods of the differential forms $\omega_1 = \frac{dz}{y}$ and $\Theta = \frac{c_2}{1-z^2} \frac{dz}{y}$ on \overline{E}_c as analytic functions of $c = (c_1, c_2)$. Then there is a closed path Γ in the c -space $\mathbb{C}^2 \setminus \Delta$ such that after analytic continuation along one circuit of Γ , the periods of ω_1 do not change and at least one period of Θ changes by 2π .

The idea of the proof is that we take a point c_0 in c -space close to the locus Δ . Through c_0 we consider the complex curve C of all parameter values c such that \overline{E}_c is isomorphic to \overline{E}_{c_0} . For every $c \in C$ we normalise the curve E_c to a single curve E in the Legendre form $v^2 = 2u(u-1)(u-t)$, where t is a fixed large number. However, the poles of the 1-form Θ still depend on the choice of $c \in C$. We let one of these poles, called s , run over a path which encloses the points $0, 1$ in \mathbb{C} . Then one of the periods of Θ changes by 2π .

To carry out the proof we need a few lemmata.

Lemma 6.3. Suppose $s, t \in \mathbb{C}$ satisfy $|s| < 2$ and $|t| > R$, where $R > 0$ is sufficiently large ($R = 100$ will do).

1. Define $a \in C$ by

$$a = \frac{1}{4} \left(-1 - t + 3s - \sqrt{t^2 + 2st - 2t - 3s^2 + 2s + 1} \right), \quad (26)$$

where the square root is chosen so that it lies close to t . Then a lies in $-\frac{t}{2} + \mathbb{C}[[s, \frac{1}{t}]]$ and the power series converges when $|s| < 2, |t| > R$.

2. The substitution

$$z = 1 + (u - s)/a, \quad y = v/a^{3/2} \quad (27)$$

transforms the defining equation $y^2 = f_c(z)$ of E_c into the defining equation $v^2 = 2u(u-1)(u-t)$ of E .

3.

$$c_1 = (t - 3s + 3a + 1)/a, \quad -c_2^2 = 2s(s-1)(s-t)/a^3.$$

4. The points on E_c with $z = 1$ correspond to points on E with $u = s$, while the points with $z = -1$ correspond to those on E with $u = -2a + s$.

Proof of lemma 6.3.

1. The statement about the analytic nature and the region of convergence of a as function of s, t is an elementary exercise.

2. We apply the inverse substitution $u = a(z-1) + s, v = a^{3/2}y$ to the curve $v^2 = 2u(u-1)(u-t)$. We get

$$y^2 = 2(z^3 + b_2z^2 + b_1z + b_0), \quad (28)$$

where

$$b_0 = \frac{1}{a^3}(s-a)((s-a)^2 - (1+t)(s-a) + t) \quad (29)$$

$$b_1 = (3s^2 + 2at + t - 2s + 3a^2 + 2a - 6as - 2st)/a^2, \quad (30)$$

$$b_2 = (-t + 3s - 3a - 1)/a. \quad (31)$$

Choose a in equation (26) so that $b_1 = -1$. Then the transformed equation (28) reads

$$y^2 = 2(z^2 - 1)(z + b_2) + 2(1 + b_2 + b_1 + b_0).$$

3. We can now identify c_1 with $-b_2$ and $-c_2^2$ with $2(b_0 + b_2)$. From the fact that $-\frac{1}{2}c_2^2a^3 = a^3(b_0 + b_2)$ and the definition of a a calculation using (30) shows that $-c_2^2 = 2s(s-1)(s-t)/a^3$.

4. From $u = a(z-1) + s$ we easily see that $z = \pm 1$ implies that $u = s$ and $u = -2a + s$, respectively. \square

Lemma 6.4. Using the transformation (30), the 1-forms $\frac{dz}{y}$ and Θ can be rewritten as

$$\frac{dz}{y} = a^{1/2} \frac{du}{v}, \quad \Theta = \frac{v(s)}{2i} \left(\frac{1}{u-s} - \frac{1}{u+2a-s} \right) \frac{du}{v}. \quad (32)$$

Here $v(s)$ denotes a suitable choice of branch of the function $\sqrt{2s(s-1)(s-t)}$.

Proof. The transformation of $\frac{dz}{y}$ is immediate. For Θ we note that

$$\frac{c_2}{1-z^2} \frac{dz}{y} = \frac{y(s)}{2i} \left(\frac{1}{z+1} - \frac{1}{z-1} \right) \frac{dz}{y},$$

where $y(s) = \sqrt{f_c(s)}$. Carrying out the transformation (30) is now straightforward. \square

From the lemma 6.3 and 6.4 we see that it suffices to study the variation of the periods of the 1-forms in (32).

We now assume that t has a fixed value and we let s run over a closed path Γ which encircles the points $0, 1$ exactly once. Since a (26) is close to $-\frac{1}{2}$ and is analytic in s , the value of $a^{1/2}$ returns to its original value after letting s trace out Γ once. Therefore the periods of $a^{1/2} \frac{du}{v}$ return to their original values. It remains to study the variation of the periods of Θ . For this we need the additional

Lemma 6.5. Consider the elliptic curve E defined by

$$v^2 = g(u) = 2u(u-1)(u-t).$$

Let s be a complex number $\neq 0, 1, t$ and let γ be a closed path on E avoiding the points where $u = s$. Define

$$p(s) = \int_{\gamma} \frac{v(s)}{u-s} \frac{du}{v}.$$

Then

$$p'(s) = \frac{s}{v(s)} \int_{\gamma} \frac{du}{v} - \frac{1}{v(s)} \int_{\gamma} u \frac{du}{v}, \quad (33)$$

where $'$ denotes differentiation with respect to s .

Proof. Differentiating p under the integral sign gives

$$p'(s) = \int_{\gamma} \frac{d}{ds} \left(\frac{v(s)}{u-s} \right) \frac{du}{v}.$$

But

$$\frac{d}{ds} \left(\frac{v(s)}{u-s} \right) = \frac{v'(s)}{u-s} + \frac{v(s)}{(u-s)^2} = \left(\frac{1}{2} \frac{g'(s)}{u-s} + \frac{g(s)}{(u-s)^2} \right) \frac{1}{v(s)},$$

using $v(s)^2 = g(s)$ and $2v(s)v'(s) = g'(s)$. Hence

$$p'(s) = \frac{1}{v(s)} \int_{\gamma} \left(\frac{1}{2} \frac{g'(s)}{u-s} + \frac{g(s)}{(u-s)^2} \right) \frac{du}{v}.$$

A straightforward computation, using the fact that $g(u)$ is a cubic polynomial with leading coefficient 2, gives

$$d\left(\frac{v}{u-s}\right) = \frac{1}{2} \frac{g'(s)}{u-s} \frac{du}{v} + \frac{g(s)}{(u-s)^2} \frac{du}{v} + (u-s) \frac{du}{v}.$$

Hence,

$$p'(s) = \frac{1}{v(s)} \left[\int_{\gamma} d\left(\frac{v}{u-s}\right) - \int_{\gamma} (u-s) \frac{du}{v} \right].$$

Since a period of an exact form is zero we obtain (33). \square

Proof of theorem 6.2. We use the same notation as in the discussion above. The variation $\Delta_{\Gamma} p$ of $p(s)$ along the path Γ is given by $\int_{\Gamma} p'(s) ds$. Hence, by lemma 6.5

$$\Delta_{\Gamma} p = \int_{\Gamma} \frac{s ds}{v(s)} \int_{\gamma} \frac{du}{v} - \int_{\Gamma} \frac{ds}{v(s)} \int_{\gamma} \frac{u du}{v}.$$

The Weierstrass parametrisation of the curve E is given by

$$u = 2\wp(\tau) - \frac{1}{3}(t+1), \quad v = 2\wp'(\tau).$$

Let τ_γ, τ_Γ be the elements in the period lattice L of \wp such that $\overline{E} = \mathbb{C}/L$ which correspond to the curves γ and Γ . Let η_γ and η_Γ be the quasi-periods of the Weierstrass ζ -function for L . Then

$$\Delta_\Gamma p = -2(\eta_\Gamma \tau_\gamma - \eta_\gamma \tau_\Gamma).$$

From the Legendre relations for elliptic functions it follows that

$$\Delta_\Gamma p = 4n\pi i,$$

where n is the negative of the intersection number of Γ and γ . By choosing γ such that it intersects Γ exactly once with intersection number 1, we can see to it that $p(s)$ changes by $4\pi i$ as s runs over Γ once.

To prove the theorem we show that $p(-2a+s)$ does not change value if we let s run over Γ . To that end we write $s' = -2a+s$ and note that it depends on s, t via $s' = t + \frac{s-s^2}{t} + O(\frac{1}{t^2})$. Since $s - s^2$ encircles the origin twice if we let s run over Γ , we see that s' encircles the point t twice. Denote this path by Γ' . Since Γ' lifts to a closed path in the τ -plane, we get $\Delta_{\Gamma'} p = 0$.

As a consequence of the preceding argument we see that

$$\frac{v(s)}{v(-2a+s)} p(-2a+s) = \int_\gamma \frac{v(s)}{u+2a-s} \frac{du}{v}$$

does not change value.

In lemma 6.4 we saw that

$$\Theta = \frac{1}{2i} \left(\frac{v(s)}{u-s} \frac{du}{v} - \frac{v(s)}{u+2a-s} \right) \frac{du}{v}.$$

Thus we conclude that there is a least one period of Θ which change by $\frac{1}{2i} 4\pi i = 2\pi$ when we make a circuit of Γ . □

We are now in a position to prove

Corollary 6.6 (to theorem 6.2). Running over the curve Γ once, the period of ω_1 does not change, whereas one of the periods of ω_2 decreases by 2π .

Proof. This follows immediately from lemma 6.1. \square

Another way to state the result of corollary 6.6 is that after making a circuit of Γ , the basis

$$\begin{pmatrix} 0 \\ 2\pi \end{pmatrix}, \begin{pmatrix} \int_{\bar{\pi}^{-1}(\gamma_1)} \omega_1 \\ \int_{\bar{\pi}^{-1}(\gamma_1)} \omega_2 \end{pmatrix}, \begin{pmatrix} \int_{\bar{\pi}^{-1}(\gamma_2)} \omega_1 \\ \int_{\bar{\pi}^{-1}(\gamma_2)} \omega_2 \end{pmatrix}$$

of the period lattice L_{c_0} for $c_0 \in \Gamma$ becomes either the basis

$$\begin{pmatrix} 0 \\ 2\pi \end{pmatrix}, \begin{pmatrix} \int_{\bar{\pi}^{-1}(\gamma_1)} \omega_1 \\ \int_{\bar{\pi}^{-1}(\gamma_1)} \omega_2 - 2\pi \end{pmatrix}, \begin{pmatrix} \int_{\bar{\pi}^{-1}(\gamma_2)} \omega_1 \\ \int_{\bar{\pi}^{-1}(\gamma_2)} \omega_2 \end{pmatrix},$$

in which case the bundle of period lattices over Γ has classifying map $M_1 = \begin{pmatrix} 1 & -1 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}$ or the basis

$$\begin{pmatrix} 0 \\ 2\pi \end{pmatrix}, \begin{pmatrix} \int_{\bar{\pi}^{-1}(\gamma_1)} \omega_1 \\ \int_{\bar{\pi}^{-1}(\gamma_1)} \omega_2 \end{pmatrix}, \begin{pmatrix} \int_{\bar{\pi}^{-1}(\gamma_2)} \omega_1 \\ \int_{\bar{\pi}^{-1}(\gamma_2)} \omega_2 - 2\pi \end{pmatrix}$$

for which the classifying map is $M_2 = \begin{pmatrix} 1 & 0 & -1 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}$. Since

$$\begin{pmatrix} -1 & 0 & 0 \\ 0 & 0 & -1 \\ 0 & -1 & 0 \end{pmatrix} M_2 \begin{pmatrix} -1 & 0 & 0 \\ 0 & 0 & -1 \\ 0 & -1 & 0 \end{pmatrix} = M_1,$$

the period lattice bundles over Γ with classifying maps M_1 and M_2 are isomorphic. This completes our treatment of monodromy in the complex spherical pendulum.

References

- [1] Alber, S. and Marsden, J.E., *Semiclassical monodromy and the spherical pendulum as a complex Hamiltonian system*, Fields Institute Communications, **8** (1996), 1–18.

- [2] M. Audin, *Monodromy in the spherical pendulum*, preprint September 2000, IRMA, Strasbourg.
- [3] Brieskorn, E. and Knörrer, H., “Plane Algebraic Curves”, Birkhäuser, Boston, 1986.
- [4] Cushman, R.H. and Bates, L.M., “Global Aspects of Classical Integrable Systems”, Birkhäuser, Basel, 1997.
- [5] Duistermaat, J.J., *On global action angle coordinates*, Commun. Pure Appl. Math. **33** (1980) 687–706.
- [6] L. Gavrilov and O. Vivolo, *The real period function of the A_3 singularity and perturbations of the spherical pendulum*, Compositio Mathematica, **123** (2000) 167–184.
- [7] Griffiths, P. and Harris, J., “Principles of Algebraic Geometry”, Wiley, N.Y., 1978.
- [8] Knörrer, H., Letter to R. Cushman dated 20 July 1983.
- [9] Mumford, D., “Tata Lectures on Theta II”, Birkhäuser, Boston, 1984.
- [10] Whittaker, E.T. and Watson, G.N., “A Course on Modern Analysis”, 4th ed., Cambridge University Press, Cambridge, UK., 1980.