

# Propositional Logics of Closed and Open Substitutions over Heyting's Arithmetic

Albert Visser

*Department of Philosophy, Utrecht University  
Heidelberglaan 8, 3584 CS Utrecht, The Netherlands  
email: Albert.Visser@phil.uu.nl*

October 24, 2005

## Abstract

In this note we compare propositional logics for closed substitutions and propositional logics for open substitutions in constructive arithmetical theories. We provide a strong example where these logics diverge in an essential way. We prove that for Markov's Arithmetic, i.e. Heyting's Arithmetic plus Markov's principle plus Extended Church's Thesis, the logic of closed and the logic of open substitutions are the same.

## 1 Introduction

In this paper, we study the propositional logics of constructive arithmetical theories. These logics contain precisely the propositional schemes such that all substitution instances of the given scheme are provable in the given theory. Prima facie, there is a difference between logics of *closed substitutions*, i.e., those substitutions where the range of the substitution consists entirely of *sentences*, and logics of *open substitutions*, where we allow *formulas* to be substituted. Of course, the logic of open substitutions for a given theory  $T$  will be a sublogic of the logic of closed substitutions for  $T$ , but the inclusion could be strict.

In Section 3, we will provide an 'essential' example to illustrate that, for some theories, the logic of closed substitutions and the logic of open substitutions are different. The example, say  $U$ , will be 'essential' in the sense that every consistent extension  $V$  of  $U$  is also an example. In fact, we will have the following. For every consistent extension  $V$  of  $U$ , the propositional logic of open substitutions is Intuitionistic Propositional Logic (IPC), and the propositional logic of closed substitutions is Classical Propositional Logic (CPC).

In Section 4, we will provide a sufficient condition to guarantee that the logic of open substitutions and the logic of closed substitutions are the same.

In Section 5, we study Markov's Arithmetic (MA), i.e., Heyting's Arithmetic, HA, plus Markov's Principle (MP), plus Extended Church's Thesis (ECT<sub>0</sub>). It

is well-known that the propositional logics of MA are not equal to IPC. See Section 2. We will show that MA satisfies the sufficient condition given in Section 4. Hence, the logic for open and the logic for closed substitutions in MA are the same.

Finally, in Section 6, we briefly discuss an ordering of open substitutions. This section is merely intended to draw attention to the existence of this further structure, not to present any definite results concerning it.

## Prerequisites

We will suppose that the reader is familiar with the basics of constructivism and arithmetical theories. A good two volume text book is [TvD88a], [TvD88b].

## 2 Logics and Theories

This section provides some basic definitions and contains a brief, admittedly incomplete, overview of what is known of propositional logics of arithmetical theories.

We will only consider arithmetical theories. We will treat theories as sets of sentences closed under the axioms and rules of constructive predicate logic. Here are a few of the theories that are of central interest.

- Heyting's Arithmetic HA.
- Markov's Arithmetic MA := HA + MP + ECT<sub>0</sub>.
- Peano's Arithmetic PA.

We will use s-ET for the scheme of sentential excluded third.

Closed substitutions will be assignments of sentences of the arithmetical language to propositional variables. Open substitutions will be assignments of formulas to propositional variables. We will call the sets of closed substitutions cSub and we will call the set of open substitutions oSub. As a default we will assume that the value of a substitution is almost everywhere ( $0 = 0$ ). This allows us to treat substitutions as finite objects. In case we do not adhere to this convention we will speak about *infinite substitutions*. Let  $U$  be any theory. We define:

- $\Lambda_U := \{\phi \mid \forall \sigma \in \text{cSub } U \vdash \phi^\sigma\}$ ,
- $\Lambda_U^\circ := \{\phi \mid \forall \sigma \in \text{oSub } U \vdash \phi^\sigma\}$ .<sup>1</sup>

Both  $\Lambda$  and  $\Lambda^\circ$  are monotonic in  $U$ . Clearly,  $\Lambda_U^\circ \subseteq \Lambda_U$ .<sup>2</sup> Note that, if  $\Lambda_U = \text{IPL}$ ,

<sup>1</sup>As usual, we take a formula to be provable iff its universal closure is provable.

<sup>2</sup>Alternativey, we can define the logics via the associated Lindenbaum Heyting algebras. Let  $\mathcal{H}$  be any Heyting algebra. We define  $\Lambda_{\mathcal{H}} := \{\phi \mid \forall \sigma : \text{PROP} \rightarrow \mathcal{H} \ \phi^\sigma = \top\}$ . Note that if there is an embedding of  $\mathcal{H}$  into  $\mathcal{G}$ , then  $\Lambda_{\mathcal{G}} \subseteq \Lambda_{\mathcal{H}}$ .

Let  $\mathcal{L}(U)$  be the sentential Lindenbaum algebra of  $U$  and let  $\mathcal{L}^\circ(U)$  be the formula Lindenbaum algebra of  $U$ . We have  $\Lambda_U = \Lambda_{\mathcal{L}(U)}$  and  $\Lambda_U^\circ = \Lambda_{\mathcal{L}^\circ(U)}$ .

then  $\Lambda_U^\circ = \text{IPL}$ . We give a quick example to show that  $\Lambda_U$  and  $\Lambda_U^\circ$  need not always be the same.

**Example 2.1** Consider  $\text{HA} + \text{s-ET}$ , i.e.,  $\text{HA}$  plus sentential Excluded Third. It is easily seen that  $\Lambda_{\text{HA}+\text{s-ET}} = \text{CPC}$ . On the other hand  $p \vee \neg p$  is not in  $\Lambda_{\text{HA}+\text{s-ET}}^\circ$ .

Suppose it was. Then,  $\text{HA} + \text{s-ET} \vdash \forall x (\exists y Txy \vee \neg \exists Txy)$ . It follows that:

$$\text{HA} \vdash \neg \neg \forall x (\exists y Txy \vee \neg \exists Txy).$$

But this contradicts the fact that:

$$\text{HA} + \text{ECT}_0 \vdash \neg \forall x (\exists y Txy \vee \neg \exists Txy),$$

where  $\text{HA} + \text{ECT}_0$  is consistent. In Section 3, we will produce a different and, in a sense, stronger example. The result of that section immediately implies that  $\Lambda_{\text{HA}+\text{s-ET}}^\circ$  is  $\text{IPC}$ .  $\square$

For many salient theories the sentential propositional logic is  $\text{IPC}$ . A result of the form ‘the sentential propositional logic of  $U$  is  $\text{IPC}$ ’ is often called: *de Jongh’s Theorem for  $U$* . We give a quick overview of a number of de Jongh’s Theorems. In all known cases we already have de Jongh’s Theorem for a set of substitutions of sentences of restricted complexity. In our table we also give these classes. We do not give the class for de Jongh’s original result, just because it is complicated. The class  $D\Pi_1^0$  consists of disjunctions of  $\Pi_1^0$ -sentences.

date	author	paper	theory	substitutions
1969	de Jongh	[dJ70]	$\text{HA}$	—
1973	Friedman	[Fri73]	$\text{HA}$	$\Pi_2^0$
1973	Smoryński	[Smo73]	$\text{HA} + \text{RFN}_{\text{HA}}$ $\text{HA} + \text{RFN}_{\text{HA}}$ $\text{HA} + \text{RFN}_{\text{HA}} + \text{MP}$	$\Sigma_1^0$ $D\Pi_1^0$ $\text{bool}(\Pi_1^0)$
1981	Gavrilenko	[Gav81]	$\text{HA} + \text{ECT}_0$	$\Sigma_1^0$
1981	Visser	[Vis81]	$\text{HA} + \text{DNS}$	$\Sigma_1^0$

Note that de Jongh’s Theorem is downwards persistent. In all cases we can obtain uniform versions of de Jongh’s Theorem: we can restrict the class of substitutions to a single infinite substitution. This means that the free Heyting algebra on countably many generators can be embedded in the Lindenbaum algebra of the given theory.

The above table shows that  $\Lambda_{\text{HA}+\text{MP}} = \Lambda_{\text{HA}+\text{ECT}_0} = \text{IPC}$ . Remarkably,  $\Lambda_{\text{MA}}$  turns out to be a proper extension of  $\text{IPC}$ . Consider the formulas  $\chi$  and  $\rho$ , which are defined as follows.

- $\chi := (\neg p \vee \neg q)$ ,
- $\rho := [(\neg \neg \chi \rightarrow \chi) \rightarrow (\neg \neg \chi \vee \neg \chi)] \rightarrow (\neg \neg \chi \vee \neg \chi)$

Clearly,  $\rho$  is IPC-invalid. We use  $\mathbf{r}$  for *Kleene realizability*. In his classical paper [Ros53], G.F. Rose showed that  $\rho$  is uniformly realizable, i.e.,

$$\exists e \forall \sigma \in \text{cSub } \mathbb{N} \models e \mathbf{r} \rho^\sigma.$$

Thus, Rose refuted a conjecture of Kleene that a propositional formula is IPC-provable if all its arithmetical instances are (truly and classically) realizable. Note the amazing fact that one and the same realizer realizes all instances! Inspecting the proof one sees that only a small part of classical logic is involved in the verification of realizability: Markov's Principle. See David McCarthy's paper [McC91] for a detailed analysis. Thus we obtain:

$$\exists e \forall \sigma \in \text{cSub } \text{HA} + \text{MP} \vdash e \mathbf{r} \rho^\sigma.$$

Since provable realizability in  $\text{HA} + \text{MP}$  implies provability in  $\text{MA}$ , we find that  $\rho \in \Lambda_{\text{MA}}$ .

The questions we have been asking for propositional logic can also be asked for predicate logic. Dick de Jongh, in an unpublished manuscript of 1969, proved completeness for Intuitionistic predicate logic IQC for interpretations in HA. For an abstract see [dJ70]. Daniel Leivant in his thesis [Lei75] proves uniform completeness of IQL for  $\Pi_2^0$ -substitutions w.r.t. HA. Finally, Jaap van Oosten gave a full semantical proof of de Jongh's Completeness Theorem in his paper [Oos91].

Returning to the propositional case, we end this section with a theorem of Georgi Gargov that in rather general circumstances the disjunction property for  $U$  implies the disjunction property for the propositional logic  $\Lambda_U$ . See [Gar84].<sup>3</sup>

Before stating the theorem, we introduce some notations. First we explain Guaspari's witness comparison notation. Suppose  $A$  is of the form  $\exists x A_0x$  and  $B$  is of the form  $\exists y B_0y$ . Suppose further that  $x$  is not free in  $B$  and  $y$  is not free in  $A$ .<sup>4</sup> We will write:

- $A \leq B \text{ :} \leftrightarrow \exists x (A_0x \wedge \forall y < x \neg B_0y)$ .
- $B < A \text{ :} \leftrightarrow \exists y (B_0y \wedge \forall x \leq y \neg A_0x)$ .

We will use ' $\Box_U$ ' for the arithmetization of provability in  $U$ . We will write ' $x \cdot y$ ' for Kleene application. Specifically, ' $x \cdot y = z$ ' means:  $\exists u (Txyu \wedge Uu = z)$ .

**Theorem 2.2** *Suppose that  $U$  is an RE arithmetical theory strong enough to have the following property:*

---

<sup>3</sup>In fact, Gargov's theorem is stated for the provability logics of extensions of HA. We give it for propositional logics of a wider class of theories. Gargov gives credit to M.M. Kipnis for one of the main ideas of his proof. This idea is contained in Kipnis' proof that the propositional logic of effective realizability has the disjunction property. See [Kip67].

<sup>4</sup>If  $A$  and  $B$  do not satisfy the variable conditions, we take suitable  $\alpha$ -variants that do.

- if  $e \cdot n = m$  and  $m \neq k$ , then  $U \vdash e \cdot n = m \wedge e \cdot n \neq k$ .<sup>5</sup>

Suppose further that  $U$  has the disjunction property. Then  $\Lambda_U$  has the disjunction property.

## Proof

Suppose  $U$  satisfies the conditions of the theorem. Suppose  $\Lambda_U \vdash \phi \vee \psi$ . Consider any two closed substitutions  $\sigma$  and  $\tau$ . For any natural number  $n$ , we define:

- $\rho_n(p) := (\text{sg}(n \cdot n) = 0 \wedge \sigma(p)) \vee (\text{sg}(n \cdot n) = 1 \wedge \tau(p))$ .

We define a recursive function  $E$  as follows.

- $E(n) := \begin{cases} 0 & \text{if } \Box_U \phi^{\rho_n} \leq \Box_U \psi^{\rho_n} \\ 1 & \text{if } \Box_U \psi^{\rho_n} < \Box_U \phi^{\rho_n} \end{cases}$ .

Clearly,  $E$  is total. Let  $e$  be an index of  $E$ . Suppose  $E(e) = 0$ . Then,  $e \cdot e = 0$ , and, hence,  $U \vdash e \cdot e = 0$ . By the definition of  $E$ , we have  $U \vdash \phi^{\rho_e}$ . On the other hand,  $U \vdash e \cdot e = 0 \rightarrow (p^{\rho_e} \leftrightarrow p^\sigma)$ . So  $U \vdash \phi^\sigma$ . Similarly, if  $E(e) = 1$ , we find that  $U \vdash \psi^\tau$ . Thus, for all  $\sigma$  and  $\tau$ , either  $U \vdash \phi^\sigma$  or  $U \vdash \psi^\tau$ .

Up to this point our proof was constructive. By classical reasoning, we conclude that  $\phi \in \Lambda_U$  or  $\psi \in \Lambda_U$ .  $\square$

Gargov's Theorem is paradigmatic for the fact that it is possible to prove properties of propositional logics of theories without having a characterization of those logics.

**Open Question 2.3** 1. Can we prove an analogue of Theorem 2.2 for  $\Lambda_U^\circ$ ?

2. Can we make the proof of Theorem 2.2 constructive? I conjecture *no*.

3. Does Theorem 2.2 work for extensions of  $i\mathbb{Q}$ , the constructive version of Robinson's Arithmetic? I think the answer must be *yes*. Specifically, I think that the classical proof that one can interpret  $I\Delta_0 + \Omega_1$  in  $\mathbb{Q}$ , should be transferable to the constructive case. If this is true, we can use this interpretation to obtain the desired result.

$\square$

## 3 An Essential Example

In this section we produce an example to the effect that there is a theory  $T$  that separates  $\Lambda$  and  $\Lambda^\circ$  in an essential way. The following theorem is the central result that immediately yields the desired example.

<sup>5</sup>Certainly, any theory extending  $i\mathcal{S}_2^1$ , the constructive version of Buss' theory  $\mathcal{S}_2^1$  has the desired property, but, clearly, a much weaker theory will suffice. Note that we may adapt the definition of Kleene's  $T$ -predicate to make the property easy to obtain. Also we could use a relative interpretation to enable us to work in a weaker theory.

**Theorem 3.1** *There is a consistent extension  $U$  of HA and an infinite substitution  $\sigma$ , such that  $\sigma$  witnesses the uniform completeness of IPC for open substitutions for every consistent extension  $V$  of  $U$ .*

We obtain the desired example by taking  $T := U + \mathbf{s}\text{-ET}$ . Clearly, every consistent extension  $W$  of  $T$  will satisfy  $\Lambda_W = \text{CPC}$  and  $\Lambda_W^\circ = \text{IPC}$ .

To prove the theorem, we need a lemma about the relationship between IPC and IQC.

**Lemma 3.2** *Suppose  $\text{IPC} \not\vdash \phi$ , where  $\phi$  is a formula in  $\vec{p} = p_0, \dots, p_{n-1}$ . Then IQC is consistent with  $\neg \forall x_0, \dots, x_{n-1} \phi(P(x_0), \dots, P(x_{n-1}))$ .*

### Proof

Suppose  $\text{IPC} \not\vdash \phi$ . Let  $\mathcal{K}$  be a finite Kripke model for propositional logic with root  $\mathfrak{b}$  such that  $\mathfrak{b} \not\Vdash \phi$ . We transform  $\mathcal{K}$  into a model  $\mathcal{K}^*$  for predicate logic with root  $\mathfrak{b}^*$  such that

$$\mathfrak{b}^* \Vdash \neg \forall x_0, \dots, x_{n-1} \phi(P(x_0), \dots, P(x_{n-1})).$$

Roughly,  $\mathcal{K}^*$  is the result of putting lots of copies of  $\mathcal{K}$  together.

- The nodes of  $\mathcal{K}^*$  are the finite, non-empty, sequences  $\langle k_0, \dots, k_{m-1} \rangle$  of nodes of  $\mathcal{K}$ . We write  $\text{lth}(\langle k_0, \dots, k_{m-1} \rangle) := m$ .
- $\mathfrak{b}^* := \langle \mathfrak{b} \rangle$ .
- $\sigma \preceq \tau \Leftrightarrow \text{lth}(\sigma) \leq \text{lth}(\tau)$  and  $\forall i < \text{lth}(\sigma) \sigma(i) \preceq \tau(i)$ .
- $D(\sigma) := \text{lth}(\sigma) \times n$ . (Remember that we took  $\vec{p} = p_0, \dots, p_{n-1}$ .) (Alternatively we could have taken  $D(\sigma) := \omega \times n$ , thus obtaining a model with constant domains.)
- $\sigma \Vdash P(\langle i, j \rangle) \Leftrightarrow \sigma_i \Vdash p_j$ . Here we assume that  $\langle i, j \rangle \in D(\sigma)$ .

It is easy to see that  $\mathcal{K}^*$  is a Kripke model for IQC. We have to show that, for every  $\sigma$ ,

$$\sigma \not\Vdash \forall x_0, \dots, x_{n-1} \phi(P(x_0), \dots, P(x_{n-1})).$$

Consider any  $\sigma$  and let  $\ell := \text{lth}(\sigma)$ . It is clearly sufficient to show:

$$\sigma * \langle \mathfrak{b} \rangle \not\Vdash \phi(P(\langle \ell, 0 \rangle), \dots, P(\langle \ell, n-1 \rangle)).$$

Consider the submodel  $\mathcal{K}^*[\sigma * \langle \mathfrak{b} \rangle]$  generated by  $\sigma * \langle \mathfrak{b} \rangle$ . We view this model as a propositional model where we identify the propositional variables  $p_0, \dots, p_{n-1}$  with  $P(\langle \ell, 0 \rangle), \dots, P(\langle \ell, n-1 \rangle)$ . It is easy to see that projection  $(\cdot)_\ell$  is a p-morphism from the model so conceived to  $\mathcal{K}$ . This immediately gives us the desired fact.  $\square$

Now we may prove the theorem.

## Proof

Daniel Leivant in his thesis [Lei75] proves uniform completeness of IQL for  $\Pi_2^0$ -substitutions w.r.t. HA. Combining Leivant's result with Lemma 3.2, we find a  $\Pi_2^0$ -predicate  $Q$  such that, for any  $\phi(p_0, \dots, p_{n-1})$  such that  $\text{IPC} \not\vdash \phi$ , we have:

$$\text{HA} \not\vdash \neg \neg \forall x_0, \dots, x_{n-1} \phi(Q(x_0), \dots, Q(x_{n-1})).$$

We can now take as our substitution  $\sigma(p_i) := Q((x)_i)$ . Here,  $(\cdot)_i$  is the projection function for an appropriate sequence coding, setting the value 0 in case  $i$  is bigger than or equal to the length of the sequence coded by  $x$ . Consider the theory  $U := \text{HA} + \{\neg \forall x \phi^\sigma \mid \text{IPC} \not\vdash \phi\}$ . If  $U$  were inconsistent, then there would be formulas  $\phi_0, \dots, \phi_{k-1}$  such that, for each  $j < k$ ,  $\text{IPC} \not\vdash \phi_j$  and  $\text{HA} \vdash \neg \bigwedge_j \neg \forall x \phi_j^\sigma$ . In other words:  $\text{HA} \vdash \neg \neg \bigvee_j \forall x \phi_j^\sigma$ . It follows that  $\text{HA} \vdash \neg \neg \forall x \bigvee_j \phi_j^\sigma$ . Hence, by the properties of  $\sigma$ ,  $\text{IPC} \vdash \bigvee_j \phi_j$  and, so, for some  $j < k$ ,  $\text{IPC} \vdash \phi_j$ . Quod non.

It is clear that every consistent extension  $V$  of  $U$  satisfies de Jongh's Theorem for open substitutions.  $\square$

Since HA proves the decidability of IPC, our theory  $U$  is in fact the same as HA plus all sentences of the form  $\forall \vec{p} \in \Pi_2^0 (\phi^\vec{p} \rightarrow \Box_{\text{IPC}} \phi)$ , where  $\phi$  is a propositional formula. Here the propositional quantifier is justified using a  $\Pi_2^0$ -truth predicate. Thus, we have established that HA plus a version of the completeness of propositional logic is consistent.

## 4 The Method of Attempted Counterexamples

We will explore an argument to show that  $\Lambda_U^\circ = \Lambda_U$ . The form of the argument is quite general, but, regrettably, the non-trivial applications are, until now, quite limited. In fact application to MA is the only non-trivial example I have.

The idea is as follows. Consider an open substitution  $\tau$ . We want to replace it by a closed one that behaves in ways that are 'sufficiently similar'. To realize this, we replace the free variable in  $\tau$  by (the paraphrase of) an 'attempted counterexample' (AC), say  $c$ . The partial constant is such that if  $T$  proves  $(\phi^\tau)(c)$ , then  $T$  proves  $\forall x \phi^\tau$ .

Here is a more precise presentation of the strategy. Consider a theory  $T$ . Suppose  $T$  is an arithmetical theory extending HA. Let  $Bx$  be a formula with only  $x$  free. We say that  $Ax$ , with only  $x$  free, is  $B$ -AC iff

- $T \vdash \forall x, y ((Ax \wedge Ay) \rightarrow x = y)$ ;
- whenever  $T \vdash \forall x (Ax \rightarrow Bx)$ , then  $T \vdash \forall x Bx$ .

**Example 4.1** Suppose  $T \not\vdash Bn$ . Then, it is easy to see that  $x = n$  is  $B$ -AC. So, the  $B$ -AC property is mainly interesting when, for all  $n$ ,  $T \vdash Bn$ .  $\square$

We say that  $T$  has the AC-property, if, for every  $Bx$ , there is a  $B$ -AC  $Ax$ . If we only have AC-formulas for the elements of a class  $\Gamma$ , we will speak about the  $\Gamma$ -AC property. Here is the desired application of the AC-property.

**Theorem 4.2** *Suppose  $T$  has the AC-property, then  $\Lambda_T^\circ = \Lambda_T$ .*

Before giving the proof, we formulate a convention. In this section, we will assume that open substitutions involve at most one free variable. This assumption does not restrict the generality of our results, since we are looking at theories that have coding of sequences.

### Proof

We already know that  $\Lambda_T^\circ \subseteq \Lambda_T$ . For the converse, suppose  $\phi \in \Lambda_T$ . Consider an open substitution  $\sigma$ . To show:  $T \vdash \phi^\sigma$ .

Suppose  $Ax$  is AC for  $\phi^\sigma x$ . Define  $\tilde{\sigma}$  by  $p^{\tilde{\sigma}} := \forall x (Ax \rightarrow p^\sigma x)$ . Note that  $\tilde{\sigma}$  is closed.

We claim that  $T \vdash Ay \rightarrow (p^\sigma y \leftrightarrow p^{\tilde{\sigma}})$ . Reason in  $T$ . Suppose  $Ay$ . First suppose  $p^{\tilde{\sigma}}$ . It is immediate that we obtain  $p^\sigma y$ . Conversely, suppose  $p^\sigma y$ . Consider any  $x$  and suppose  $Ax$ . It follows that  $x = y$ , and, hence,  $p^\sigma x$ . We may conclude that  $\forall x (Ax \rightarrow p^\sigma x)$ , i.e.  $p^{\tilde{\sigma}}$ .

It follows, by induction on  $\psi$ , that  $T \vdash Ay \rightarrow (\psi^\sigma y \leftrightarrow \psi^{\tilde{\sigma}})$ . Since  $\phi \in \Lambda_T$  and since  $\tilde{\sigma}$  is closed, we have:  $T \vdash \phi^{\tilde{\sigma}}$ . Ergo,  $T \vdash \forall y (Ay \rightarrow \phi^\sigma y)$ . Since  $A$  is an AC for  $\phi^\sigma$ , we find:  $T \vdash \forall y \phi^\sigma y$ . We conclude that  $\phi \in \Lambda_T^\circ$ .  $\square$

A formula  $B$  is  $T$ -stable if  $T \vdash \neg\neg B \rightarrow B$ . So  $B$  is  $T$ -stable iff  $B$  is  $T$ -equivalent to a negation. The next theorem articulates a basic insight.

**Theorem 4.3** *Every  $T$ -stable  $Bx$  has a  $B$ -AC formula  $Ax$  in  $T$ .*

### Proof

We define  $Ax := (\neg Bx \wedge \forall y < x By)$ . The uniqueness clause is clear. Suppose  $T \vdash \forall x (Ax \rightarrow Bx)$ . It follows that  $T \vdash \forall x (\forall y < x By \rightarrow \neg\neg Bx)$ . Hence, by stability and well-founded induction,  $T \vdash \forall x Bx$ .  $\square$

Note that Theorem 4.3 seems pretty useless. In the case of PA, where all formulas are stable, we already know that  $\Lambda_{\text{PA}}^\circ = \Lambda_{\text{PA}}$ . Fortunately, an adaptation of the proof of Theorem 4.3 will show that MA has the full AC-property, and so,  $\Lambda_{\text{MA}}^\circ = \Lambda_{\text{MA}}$ .



**Example 4.4** Let  $U$  be HA plus s-ET, i.e. sentential excluded third. Consider  $Cx := \exists y Txy \vee \neg \exists y Txy$ . Suppose we have a  $C$ -AC  $Ax$ .

Reason in  $U$ . Suppose  $Ax$ . By the uniqueness clause from the definition of  $C$ -AC, it follows that  $\neg \forall y (Ay \rightarrow Cy)$  implies  $\neg (Ax \rightarrow Cx)$ . However, together with  $Ax$  this implies  $\neg C(x)$ , which is a contradiction. Hence, by s-ET we find  $\forall y (Ay \rightarrow Cy)$ . Using  $Ax$  again we get  $Cx$ . Thus, we may conclude that  $(Ax \rightarrow Cx)$  without any assumptions.

So,  $U \vdash \forall x (Ax \rightarrow Cx)$ . Since  $A$  is  $C$ -AC,  $U \vdash \forall x Cx$ . This, however, is false (consider realizability). So  $U$  does not have the AC-property.  $\square$

## 5 Markov's Arithmetic

In the present section we study Markov's Arithmetic. We prove that MA has the AC-property.

### 5.1 General Facts

In this subsection, we provide some general facts about MA. First we provide two characterizations of MA.

**Theorem 5.1** *We have:*

1.  $MA \vdash A$  iff, for some  $n$ ,  $HA + MP_{PR} \vdash n \mathbf{r} A$ ;
2.  $MA \vdash A$  iff, for some  $n$ ,  $PA \vdash n \mathbf{r} A$ .

The proof is just a minor adaptation of the proof of [Tro73], Theorem 3.2.25.

### Proof

Suppose  $MA \vdash A$ . Since, all principles of MA are realizable over  $HA + MP_{PR}$ , we find  $HA + MP_{PR} \vdash n \mathbf{r} A$ .

Suppose  $HA + MP_{PR} \vdash n \mathbf{r} A$ . Since,  $HA + MP_{PR}$  is included in MA, we find  $MA \vdash n \mathbf{r} A$ . Hence,  $MA \vdash A$ .

Suppose  $HA + MP_{PR} \vdash n \mathbf{r} A$ . Then, clearly,  $PA \vdash n \mathbf{r} A$ .

Suppose  $PA \vdash n \mathbf{r} A$ . Over  $HA + MP_{PR}$ , the formula  $n \mathbf{r} A$  is equivalent to a negative formula, say  $B$ . So  $PA \vdash B$ . Since, by the double negation translation, PA is conservative over HA w.r.t. negative formulas, we find  $HA \vdash B$ . Thus,  $HA + MP_{PR} \vdash B$  and, hence,  $HA + MP_{PR} \vdash n \mathbf{r} A$ .  $\square$

A propositional formula  $\phi$  is *effectively realizable* if there is a recursive function  $F$ , such that, for all closed substitutions  $\sigma$ ,  $\mathbb{N} \models F(\sigma) \mathbf{r} \phi^\sigma$ .

**Theorem 5.2** *All  $\phi$  in  $\Lambda_{MA}$  are effectively realizable.*

## Proof

We apply Theorem 5.1(2). Take  $F(\sigma)$  to be the number  $n$  provided by the smallest PA-proof of a sentence of the form  $n \mathbf{r} \phi^\sigma$ .  $\square$

Our next theorem is an immediate consequence of Theorem 2.2, using the fact that MA has the disjunction property.

**Theorem 5.3**  $\Lambda_{\text{MA}}$  has the disjunction property.

## 5.2 MA has the AC-property

In this subsection, we show that MA has the AC-property. Consider any  $Cx$ . We define:

$$\text{D1 } Bxe := (\exists u (e \cdot x) \cdot 0 = u \wedge \forall v ((e \cdot x) \cdot 0 = v \rightarrow v \mathbf{r} Cx)).$$

We find, using the Gödel Fixed Point Lemma a formula  $Ax$  such that:

$$\text{D2 } \text{MA} \vdash Ax \leftrightarrow \neg \neg [\exists p \exists e < p \{ \text{proof}_{\text{MA}}(p, e \mathbf{r} \forall z (Az \rightarrow Cz)) \wedge \\ \forall q < p \forall f < q \neg \text{proof}_{\text{MA}}(q, f \mathbf{r} \forall z (Az \rightarrow Cz)) \wedge \\ \neg Bxe \wedge \forall y < x Bye \}].$$

Here we assume that  $z$  can be uniquely extracted from a formula of the form  $z \mathbf{r} E$ . We can easily arrange this to be the case.

Remember that any formula of the form  $v \mathbf{r} Cx$  is almost negative. Thus, in the presence of  $\text{MP}_{\text{PR}}$  it becomes negative (modulo provable equivalence). It follows that  $Bxe$  is a MA-stable formula. We will show that  $A$  is  $C$ -AC, i.e.,

1.  $\text{MA} \vdash \forall x, y ((Ax \wedge Ay) \rightarrow x = y)$ ,
2.  $\text{MA} \vdash \forall x (Ax \rightarrow Cx) \Rightarrow \text{MA} \vdash \forall x Cx$ .

The uniqueness (1) is easy to see. We prove (2). Suppose  $\text{MA} \vdash \forall x (Ax \rightarrow Cx)$ . Then, we can find an proof  $p$  and a number  $e$ , such that  $p$  witnesses that  $\text{MA} \vdash e \mathbf{r} \forall z (Az \rightarrow Cz)$ . We pick  $p^*$  and  $e^*$  such that  $p^*$  is smallest with this property and  $e^*$  is the witness produced by  $p^*$ . By  $\Sigma$ -completeness, we find:

$$\text{MA} \vdash \text{proof}_{\text{MA}}(p^*, e^* \mathbf{r} \forall z (Az \rightarrow Cz)) \wedge \\ \forall q < p^* \forall f < q \neg \text{proof}_{\text{MA}}(q, f \mathbf{r} \forall z (Az \rightarrow Cz)) \quad (1)$$

Hence, from D2, we have:

$$\text{MA} \vdash Ax \leftrightarrow \neg \neg (\neg Bxe^* \wedge \forall y < x Bye^*). \quad (2)$$

On the other hand, we have  $\text{MA} \vdash e^* \mathbf{r} \forall z (Az \rightarrow Cz)$ . Spelling this out we find that:

$$\text{MA} \vdash \forall x \forall y (y \mathbf{r} Ax \rightarrow (\exists u (e^* \cdot x) \cdot y = u \wedge \forall v ((e^* \cdot x) \cdot y = v \rightarrow v \mathbf{r} Cx))). \quad (3)$$

Since,  $Ax$  is a negation, we have  $\text{MA} \vdash Ax \leftrightarrow 0 \text{ r } Ax$ . Hence:

$$\text{MA} \vdash \forall x (Ax \rightarrow (\exists u (e^* \cdot x) \cdot 0 = u \wedge \forall v ((e^* \cdot x) \cdot 0 = v \rightarrow v \text{ r } Cx))). \quad (4)$$

In other words, by D1, we have:

$$\text{MA} \vdash \forall x (Ax \rightarrow Bxe^*). \quad (5)$$

Comparing Equations (2) and (5), we find  $\text{MA} \vdash \forall x \neg Ax$ . Hence, by Equation (2),

$$\text{MA} \vdash \forall x (\forall y < xBye^* \rightarrow \neg\neg Bxe^*). \quad (6)$$

Hence, using the stability of  $B$ , we obtain  $\text{MA} \vdash \forall x Bxe^*$ . Thus, putting  $f^* := \Lambda w.((e^* \cdot w) \cdot 0)$ , we find:  $\text{MA} \vdash f^* \text{ r } \forall x Cx$ . Ergo, applying  $\text{ECT}_0$  in  $\text{MA}$ , we have:  $\text{MA} \vdash \forall x Cx$ .

**Remark 5.4** Note that we use many special features of  $\text{MA}$ . E.g., the proof does not seem to go through if we replace  $\text{ECT}_0$  by  $\text{CT}_0$ . The proof does generalize to any RE extension  $U$  of  $\text{MA}$  that is closed under the rule: if  $U \vdash E$ , then, for some  $n$ ,  $U \vdash n \text{ r } E$ .  $\square$

## 6 Structure on Open Substitutions

To simplify inessentially, let's suppose that all open substitutions only involve one arithmetical variable. We consider substitutions of a fixed finite  $\vec{p}$ . We fix some theory  $T$ . We define:

- $\sigma \sqsubseteq_T \tau$  iff, for some primitive recursive  $f$ ,  $T \vdash \forall x \bigwedge_{p \in \vec{p}} (p^\sigma(x) \leftrightarrow p^\tau(fx))$ .
- $(\sigma \sqcup_T \tau)(p)(x) := (\exists z ((x = 2z \wedge p^\sigma(z)) \vee (x = 2z + 1 \wedge p^\tau(z)))$ .

We will suppress the subscript  $T$  as long as  $T$  is clear from the context. Clearly, if  $\sigma \sqsubseteq_T \tau$ , then  $T \vdash \forall x \phi^\tau(x) \rightarrow \forall x \phi^\sigma(x)$ . We also have:

$$T \vdash \forall x \phi^{\sigma \sqcup \tau} \leftrightarrow (\forall x \phi^\sigma \wedge \forall x \phi^\tau).$$

It is easy to see that  $\sqcup$  is the supremum of  $\sqsubseteq$ . We define further:

- $\phi$  is o-exact for  $T$  iff there is an open  $\vec{p}$ -substitution  $\sigma$  such that, for all  $\vec{p}$ -formulas  $\psi$ ,  $T \vdash \psi^\sigma$  iff  $\Lambda_T^\circ(\vec{p}) \vdash \phi \rightarrow \psi$ .
- We will say that the pair  $\langle \sigma, \phi \rangle$  is an o-exact pair for  $T$  if  $\sigma$  witnesses the o-exactness for  $T$  of  $\phi$ .

**Theorem 6.1** 1. Let  $\langle \sigma, \phi \rangle$  and  $\langle \tau, \psi \rangle$  be o-exact for  $T$ . Suppose  $\sigma \sqsubseteq_T \tau$ . Then,  $\Lambda_T^\circ \vdash \phi \rightarrow \psi$ .

2. Suppose  $\langle \sigma, \phi \rangle$  and  $\langle \tau, \psi \rangle$  are o-exact. Then  $\langle \sigma, \phi \rangle + \langle \tau, \psi \rangle := \langle \sigma \sqcup \tau, \phi \vee \psi \rangle$  is o-exact. Thus, o-exact formulas are closed under disjunction.

## Proof

Ad (1). We have:

$$\begin{aligned}\Lambda_T^\circ \vdash \psi \rightarrow \chi &\Rightarrow T \vdash \forall x \chi^\tau \\ &\Rightarrow T \vdash \forall x \chi^\sigma \\ &\Rightarrow \Lambda_T^\circ \vdash \phi \rightarrow \chi\end{aligned}$$

Ad (2). We have:

$$\begin{aligned}\Lambda_T^\circ \vdash (\phi \vee \psi) \rightarrow \chi &\Leftrightarrow \Lambda_T^\circ \vdash \phi \rightarrow \chi \text{ and } \Lambda_T^\circ \vdash \psi \rightarrow \chi \\ &\Leftrightarrow T \vdash \forall x \chi^\sigma \text{ and } T \vdash \forall x \chi^\tau \\ &\Leftrightarrow T \vdash \forall x \chi^\sigma \wedge \forall x \chi^\tau \\ &\Leftrightarrow T \vdash \forall x \chi^{\sigma \sqcup \tau}\end{aligned}$$

□

## References

- [dJ70] D.H.J. de Jongh. The maximality of the intuitionistic predicate calculus with respect to Heyting's Arithmetic. *The Journal of Symbolic Logic*, 36:606, 1970.
- [Fri73] H. Friedman. Some applications of Kleene's methods for intuitionistic systems. In A.R.D. Mathias and H. Rogers, editors, *Cambridge Summerschool in Mathematical Logic*, pages 113–170. Springer Verlag, Berlin, Heidelberg, New York, 1973.
- [Gar84] G. K. Gargov. A note on the provability logics of certain extensions of Heyting's Arithmetic. In *Mathematical Logic, proceedings of the conference on Mathematical Logic, dedicated to the memory of A. A. Markov (1903–1979), Sofia, September 22–23, 1980*, pages 20–26. Bulgarian Academy of Sciences, 1984.
- [Gav81] Yu. V. Gavrilenko. Recursive realizability from the intuitionistic point of view. *Soviet Mathematical Dokl.*, 23:9–14, 1981.
- [Kip67] M.M. Kipnis. A property of propositional formulas. *Soviet Math. Dokl.*, 8:620–621, 1967.
- [Lei75] D. Leivant. *Absoluteness in intuitionistic logic*, volume 73. Mathematical Centre Tract, Amsterdam, 1975.
- [McC91] D.C. McCarty. Incompleteness in intuitionistic metamathematics. *The Notre Dame Journal of Formal Logic*, 32:323–358, 1991.

- [Oos91] J. van Oosten. A semantical proof of De Jongh's Theorem. *Archive for Mathematical Logic*, 23:105–114, 1991.
- [Ros53] G.F. Rose. Propositional calculus and realizability. *Transactions of the American Mathematical Society*, 61:1–19, 1953.
- [Smo73] C. Smoryński. Applications of Kripke Models. In A.S. Troelstra, editor, *Metamathematical Investigations of Intuitionistic Arithmetic and Analysis*, Springer Lecture Notes 344, pages 324–391. Springer, Berlin, 1973.
- [Tro73] A.S. Troelstra. *Metamathematical investigations of intuitionistic arithmetic and analysis*. Springer Lecture Notes 344. Springer Verlag, Berlin, 1973.
- [TvD88a] A.S. Troelstra and D. van Dalen. *Constructivism in Mathematics, vol 1*. Studies in Logic and the Foundations of Mathematics, vol. 121. North Holland, Amsterdam, 1988.
- [TvD88b] A.S. Troelstra and D. van Dalen. *Constructivism in Mathematics, vol 2*. Studies in Logic and the Foundations of Mathematics, vol. 123. North Holland, Amsterdam, 1988.
- [Vis81] A. Visser. *Aspects of diagonalization and provability*. Ph.D. Thesis, Department of Philosophy, Utrecht University, 1981.