

from each element and is therefore affected by the Fe:O ratio.

Aziz, Chergui and collaborators observed a decrease in the magnitude of the sub-background portion of the FeCl<sub>3</sub> TFY for more concentrated solutions, which led them to conclude that there was improved charge-transfer efficiency from the Fe to the water (reduced fluorescence efficiency). However, they neglect to consider the effect of changing the Fe:O ratio on the magnitude of the PFY from each element. For low-concentration samples, the magnitude of the decrease in the O PFY will be greater than the magnitude of the increase in the Fe PFY, resulting in more sub-background intensity. As the Fe concentration in the sample is increased, the Fe PFY peak intensity will increase more than the O PFY dip will decrease, resulting in a net reduction in the sub-background intensity, consistent with the concentration-dependent measurements of FeCl<sub>3</sub> by Aziz *et al.* This can be shown by considering the expression for the PFY<sup>4</sup> from each element with tabulated mass attenuation coefficients from the NIST

database<sup>5</sup>. A similar argument can be used to explain why there is no sub-background for the 1M FeCl<sub>3</sub> ethanol solution, where the Fe:O ratio is relatively high.

Solute–solvent charge transfer is an important process in many biological and electrochemical systems and deserves significant attention using all possible methods. Although TFY may be sensitive to changes in the radiative decay rates of the elements in the sample at some level, the structure and intensity of TFY spectra is predominantly determined by the ratio of the linear attenuation coefficients of these elements. For this reason, the TFY spectra presented by Aziz, Chergui and collaborators cannot be used as proof of solute–solvent electron transfer. □

#### References

1. Aziz, E. F., Rittmann-Frank, M. H., Lange, K. M., Bonhommeau, S. & Chergui, M. *Nature Chem.* **2**, 853–857 (2010).
2. Nilsson, A. *Nature Chem.* **2**, 800–802 (2010).
3. Achkar, A. J. *et al. Phys. Rev. B* **83**, 081106(R) (2011).
4. Stöhr, J. *NEXAFS Spectroscopy* (Springer, 1992).
5. Chantler, C. J. *Phys. Chem. Ref. Data* **24**, 71 (1995).

T. Z. Regier<sup>1,2\*</sup>, A. J. Achkar<sup>3</sup>, D. Peak<sup>4</sup>, J. S. Tse<sup>2</sup> and D. G. Hawthorn<sup>3</sup>

<sup>1</sup>Canadian Light Source, University of Saskatchewan, 101 Perimeter Road, Saskatoon S7N 0X4, Canada. <sup>2</sup>Department of Physics and Engineering Physics, University of Saskatchewan, 116 Science Place, Saskatoon S7N 5E2, Canada. <sup>3</sup>Department of Physics and Astronomy, University of Waterloo, 200 University Ave. West, Waterloo N2L 3G1, Canada. <sup>4</sup>College of Agriculture and Bioresources, University of Saskatchewan, 51 Campus Drive, Saskatoon S7N 5A8, Canada. \*e-mail: tom.regier@lightsources.ca

#### Acknowledgements

This research is supported by NSERC. The research described in this Correspondence was performed at the Canadian Light Source, which is supported by NSERC, NRC, CIHR and the University of Saskatchewan.

#### Author contributions

T.Z.R. wrote the correspondence and performed the data analysis. D.P. performed the measurement. All authors contributed to the development of the concepts.

Published online 12 August 2012

# Dips and peaks in fluorescence yield X-ray absorption are due to state-dependent decay

**To the Editor** — In a 2010 paper Aziz, Chergui and colleagues observe fluorescence yield (FY) detected X-ray absorption spectra that are concentration-dependent and show both dips and peaks<sup>1</sup>. In this comment I will show that all observed spectral features are a consequence of the relative ratio of background and edge emission, combined with energy-dependent X-ray emission decay channels.

Fluorescence yield detection can be used as the basis for an X-ray absorption measurement. The FY of an energy-dependent edge absorption ( $\mu_x(\omega)$ ) on a constant background ( $\mu_B$ ) is derived in detail in the Supplementary Information (section A). It is given as:

$$I_{\text{FY}} = \frac{\mu_B \alpha_B}{\mu_x(\omega) + \mu_B + \mu_x(\omega_B') + \mu_B(\omega_B')} + \frac{\mu_x(\omega) \alpha_x}{\mu_x(\omega) + \mu_B + \mu_x(\omega_x') + \mu_B(\omega_x')} \quad (1)$$

The effective absorption coefficient  $\mu$  includes the concentration of the species

and  $\alpha$  is the efficiency of the X-ray emission decay, an intrinsic property of the core hole. Note that the two denominators are different as they contain the self-absorption at the X-ray emission energies of the edge ( $\omega_x'$ ) and the background ( $\omega_B'$ ). For a dilute species the effective absorption coefficient  $\mu_x$  is very small, implying that the  $\mu_x$ -dependent terms in the denominator can be neglected. The consequence is that the FY is proportional to the X-ray absorption coefficient of the edge absorption plus a constant background (Supplementary Information, section B). In partial fluorescence yield (PFY) an X-ray emission detector is used to select only part of the X-ray emission spectrum. In the dilute limit, the PFY intensity is directly proportional to the X-ray absorption coefficient without any background signal. Increasing the concentration does result in saturation effects in PFY experiments of concentrated samples<sup>2,3</sup>.

If the X-ray emission of the background dominates, a negative signal for the total fluorescence yield is likely, as in the case of a dilute iron species dissolved in water

as measured in ref. 1. The reason for the domination of the background X-ray emission is that the main water X-ray emission energy  $\omega_B'$  has an energy below the oxygen K edge, implying that it has small self-absorption. The main iron emission energies  $\omega_x'$  have an energy above the oxygen K edge and the emitted X-rays are strongly re-adsorbed by the water. Because the iron X-rays are strongly self-absorbed, the first term dominates equation (1). It can be shown (Supplementary Information, section C) that this term can be rewritten as a signal that is negatively proportional to the absorption constant, exactly as observed in ref. 1.


The negative signal from background emission and the positive signal from edge emission occur simultaneously and their relative magnitude determines the sign of the TFY signal. Increasing the edge-element concentration influences the relative values of background emission and edge emission. Because the self-absorption at the edge emission energy  $\mu(\omega_x')$  is much larger than  $\mu(\omega_B')$ , increasing the concentration will increase the edge signal more than it will

decrease the background signal. This will slowly turn a negative spectrum (dips) into a positive spectrum (peaks). In other words, an increase in the concentration will decrease the X-ray penetration depth, lowering the chances for self-absorption, which mainly affects the strongly self-absorbed iron edge signal.

The last issue to explain is the occurrence of dips and peaks in the same spectrum. Above we have implicitly assumed that the decay rate ( $\alpha_i$ ) is, over a short energy range, independent of the X-ray energy. However, it is known that in case of  $3d$  metal L edges, the fluorescence decay can show strong variations over the edge. The fluorescence decay for states at the high-energy side of the  $L_3$  and  $L_2$  edges is increased by a factor of five<sup>4</sup>. This effect is correlated with the total moment  $J$  value of the core-hole state. Low- $J$  core-hole states have stronger fluorescence decay than high- $J$  states. The low- $J$  states occur at the high-end side of the  $L_3$  edge implying that the high-end side of the  $L_3$  edge will have stronger edge fluorescence, which implies a tendency to more positive intensity at the high end of the  $L_3$  edge. Because of its  $2p_{1/2}$  core hole one expects the  $L_2$  edge to have on overall a stronger fluorescence than the  $L_3$  edge.

In the case of iron ions in water, part of the spectrum appears as dips and other parts as peaks<sup>1</sup>. Using the state-dependent fluorescence decay, the simultaneous occurrence of dips and peaks appears if one assumes that the fluorescence decay is high for  $J = 3/2$  states, intermediate for  $J = 5/2$  states and low for  $J = 7/2$  states. This behaviour is simulated in Fig. 1 and is in agreement with the experimental observations made in ref. 1. A more detailed theoretical treatment would include crystal field and charge-transfer effects and explicitly calculate the integrated FY XAS signals over the different fluorescence channels, similar to the  $Ni^{2+}$  calculations in ref. 4. Such treatment is not expected to modify the main observations made in Fig. 1.

In conclusion, it is shown that the behaviour of fluorescence yield L-edge X-ray absorption spectra is a consequence



**Figure 1** | Simulated state-dependent decay rates yielding dips and peaks in a FY XAS spectrum. From top to bottom (i) Atomic multiplet calculation of the iron  $L_{2,3}$  X-ray absorption spectrum of  $Fe^{3+}$  ions; (ii) final states with  $J = 5/2$ ; (iii) final states with  $J = 7/2$  (solid line) and  $J = 3/2$  (dashed line); (iv) Difference spectrum of the  $J = 7/2$  minus  $J = 3/2$  spectrum, simulating the effects of the state dependent FY decay rates.

of the relative ratio of background and edge X-ray emission, combined with energy-dependent X-ray emission decay channels. The combination of these effects explains the dips and peaks of the L edge of iron in water, including their concentration dependence. Because all observations as observed in ref. 1 are naturally explained from the fundamental behaviour of fluorescence yield detection, the introduction of an alternative explanation described only in qualitative terms does not contribute to a better understanding. To study alternative phenomena, it is necessary that different predictions arise and dedicated experiments are performed to distinguish the interpretation. In all cases where FY is used the intrinsic effects of concentration dependence, combined background and edge emission and energy-

dependent X-ray emission strengths must be included in such analysis. □

#### References

1. Aziz, E. F., Rittmann-Frank, M. H., Lange, K., Bonhommeau S. & Chergui, M. *Nature Chem.* **2**, 853–857 (2010).
2. De Groot, F. M. F. & Kotani, A. *Core Level Spectroscopy of Solids* (Taylor & Francis/CRC, 2008).
3. Achkar, A. J. *et al. Phys. Rev. B* **83**, 081106 (2011).
4. De Groot, F. M. F., Arrio, M. A., Sainctavit, P., Cartier, C. & Chen, C. T. *Solid State Comm.* **92**, 991–995 (1994).

#### Frank M. F. de Groot

Debye Institute for Nanomaterials Science,  
Utrecht University, Universiteitsweg 99,  
3584 CG, Utrecht, Netherlands.  
e-mail: f.m.f.degroot@uu.nl

Additional information  
Supplementary information accompanies this paper at  
[www.nature.com/naturechemistry](http://www.nature.com/naturechemistry).

Published online 12 August 2012

## Reply to 'Dark channel fluorescence...' and 'Dips and peaks...'

**Aziz *et al.* reply to Regier *et al.* and de Groot** — Dips appearing in the total fluorescence yield (TFY) at the L-edge spectra of  $3d$ -transition metal ions in aqueous solution were recently argued to be useful

for characterizing ultrafast charge-transfer processes to the solvent<sup>1,2</sup>. Yet, to prove experimentally such charge delocalization, additional studies would be required that are capable of explicitly tracking electronic

relaxation channels. Indeed, the existence of the charge-delocalization channel (leading to a quenched fluorescence) has been, in our opinion, unequivocally demonstrated in a later comparative study of the transmission,