

## ORIGINAL ARTICLE

# Phthalates, perfluoroalkyl acids, metals and organochlorines and reproductive function: a multipollutant assessment in Greenlandic, Polish and Ukrainian men

Virissa Lenters,<sup>1</sup> Lützen Portengen,<sup>1</sup> Lidwien A M Smit,<sup>1</sup> Bo A G Jönsson,<sup>2</sup> Aleksander Giwercman,<sup>3</sup> Lars Rylander,<sup>2</sup> Christian H Lindh,<sup>2</sup> Marcello Spanò,<sup>4</sup> Henning S Pedersen,<sup>5</sup> Jan K Ludwicki,<sup>6</sup> Lyubov Chumak,<sup>7</sup> Aldert H Piersma,<sup>8</sup> Gunnar Toft,<sup>9</sup> Jens Peter Bonde,<sup>10</sup> Dick Heederik,<sup>1</sup> Roel Vermeulen<sup>1</sup>

► Additional material is published online only. To view please visit the journal online (<http://dx.doi.org/10.1136/oemed-2014-102264>).

For numbered affiliations see end of article.

## Correspondence to

Dr Roel Vermeulen, Institute for Risk Assessment Sciences, Utrecht University, P.O. Box 80.178, Utrecht 3508 TD, The Netherlands; R.C.H.Vermeulen@uu.nl

Received 8 April 2014

Revised 2 August 2014

Accepted 20 August 2014

## ABSTRACT

**Objectives** Numerous environmental contaminants have been linked to adverse reproductive health outcomes. However, the complex correlation structure of exposures and multiple testing issues limit the interpretation of existing evidence. Our objective was to identify, from a large set of contaminant exposures, exposure profiles associated with biomarkers of male reproductive function.

**Methods** In this cross-sectional study (n=602), male partners of pregnant women were enrolled between 2002 and 2004 during antenatal care visits in Greenland, Poland and Ukraine. Fifteen contaminants were detected in more than 70% of blood samples, including metabolites of di(2-ethylhexyl) and diisononyl phthalates (DEHP, DiNP), perfluoroalkyl acids, metals and organochlorines. Twenty-two reproductive biomarkers were assessed, including serum levels of reproductive hormones, markers of semen quality, sperm chromatin integrity, epididymal and accessory sex gland function, and Y:X chromosome ratio. We evaluated multipollutant models with sparse partial least squares (sPLS) regression, a simultaneous dimension reduction and variable selection approach which accommodates joint modelling of correlated exposures.

**Results** Of the over 300 exposure–outcome associations tested in sPLS models, we detected 10 associations encompassing 8 outcomes. Several associations were notably consistent in direction across the three study populations: positive associations between mercury and inhibin B, and between cadmium and testosterone; and inverse associations between DiNP metabolites and testosterone, between polychlorinated biphenyl-153 and progressive sperm motility, and between a DEHP metabolite and neutral  $\alpha$ -glucosidase, a marker of epididymal function.

**Conclusions** This global assessment of a mixture of environmental contaminants provides further indications that some organochlorines and phthalates adversely affect some parameters of male reproductive health.

## INTRODUCTION

There has been considerable scientific and public concern, particularly over the past two decades,

## What this paper adds

- Some ubiquitous environmental contaminants have been linked to changes in male reproductive health; however, interpretation of the mostly single-pollutant assessments is difficult given the complex mixture of daily exposures.
- In a study of around 600 fertile men from Greenland, Poland and Ukraine, we assessed associations between 15 contaminants measured in serum and 22 biomarkers of male reproductive function using sparse partial least squares regression for variable selection.
- We identified 8 perturbed biomarker outcomes, including a decrease in serum testosterone levels with increasing diisononyl phthalate levels, and confirmed a previously reported decrease in progressive sperm motility with increasing polychlorinated biphenyl-153 levels.

that exposure to environmental contaminants may impede development in utero, perturb hormonal homeostasis and regulation, and contribute to subfertility in humans. This concern stems in part from evidence of geographic variability in semen quality,<sup>1</sup> secular trends of decreasing testosterone<sup>2</sup> and reports, although inconsistent and contested, of secular declines in sperm concentrations and semen quality.<sup>3–4</sup>

Despite the surge in studies on environmental contaminants and male reproductive health, interpretation of the evidence base is hampered by the piecemeal approach to many investigations. In observational epidemiology, often one class of compounds is correlated with a subset of outcomes, disregarding correlations among background low-level exposures. We chose to simultaneously evaluate a large set of environmental contaminant exposures and biomarkers of male reproductive function in a pregnancy-based study of men from Greenland, Poland and Ukraine.<sup>5–6</sup> The set of contaminants comprises high priority<sup>7</sup> legacy and emerging

**To cite:** Lenters V, Portengen L, Smit LAM, et al. *Occup Environ Med* Published Online First: [please include Day Month Year] doi:10.1136/oemed-2014-102264

contaminants of concern, suspected to interfere with the endocrine system: high-molecular-weight phthalates, plasticisers used in a wide variety of consumer products, including polyvinyl chloride; perfluoroalkyl acids (PFAAs), surfactants used in many applications, such as water and stain-repellent coatings; three non-essential metals; one polychlorinated biphenyl and two organochlorine pesticides. These leach from products or are deposited directly into the environment, leading to near-ubiquitous human exposure,<sup>8–9</sup> and all, except phthalates, are persistent and bioaccumulate.

We sought an approach to identify exposure–outcome associations between the 19 exposures, representing four classes of xenobiotic compounds, and 22 biomarker outcomes, reflecting various aspects of male reproductive function. Given the relatively weak prior information for the many possible associations, we decided to take a more agnostic, data-driven approach. That the exposures are highly correlated represented a methodological challenge. For this reason, we used sparse partial least squares (sPLS) regression modelling, which relates linear combinations of a subset of the most predictive exposures to an outcome via linear regression.<sup>10</sup> Unlike conventional ordinary least squares (OLS) regression, this dimension reduction approach does not suffer from instability in estimates and failure to converge due to multicollinearity, and enabled a joint assessment of the full set of measured biomarkers of low-level environmental contaminant exposures and reproductive function.

## MATERIALS AND METHODS

### Study populations

Study design and data collection procedures have been described previously.<sup>6</sup> Briefly, pregnant women and their male partners were recruited during routine antenatal care visits from 2002 through 2004 inclusive at a large central hospital in Warsaw, Poland, at three hospitals and eight antenatal clinics in Kharkiv, Ukraine, and at local hospitals in 19 municipalities and settlements across Greenland. Of the 1710 couples enrolled (45% participation rate), male partners were consecutively invited to participate in a semen study until approximately 200 men at each of the three locations had agreed (38% participation rate). Participants provided a semen sample, a venous blood sample and information on lifestyle factors (further details provided in the online supplementary methods).

### Exposure and outcome assessment

Phthalate metabolites and PFAAs were analysed in serum samples by liquid chromatography-tandem mass spectrometry<sup>9</sup> (expanded in the online supplementary methods). Phthalates included secondary oxidised metabolites of di(2-ethylhexyl) phthalate (DEHP) [mono(2-ethyl-5-hydroxyhexyl) phthalate (MEHHP, alternatively 5OH-MEHP), mono(2-ethyl-5-oxohexyl) phthalate (MEOHP, alternatively 5oxo-MEHP) and mono(2-ethyl-5-carboxypentyl) phthalate (MECPP, alternatively 5cx-MEPP)], and of diisononyl phthalate (DiNP) [mono(4-methyl-7-hydroxyoctyl) phthalate (MHiNP, alternatively 7OH-MMeOP), mono(4-methyl-7-oxooctyl) phthalate (MOiNP, alternatively 7oxo-MMeOP) and mono(4-methyl-7-carboxyheptyl) phthalate (MOiCP, alternatively 7cx-MMeHP)]. Analysed PFAAs [a subset of per- and polyfluoroalkyl substances (PFASs), also referred to as perfluorinated compounds (PFCs)] included perfluorooctane sulfonic acid (PFOS), perfluorooctanoic acid (PFOA), perfluorohexane sulfonic acid (PFHxS), perfluorononanoic acid (PFNA), perfluorodecanoic acid (PFDA), perfluoroundecanoic acid (PFUnDA) and perfluorododecanoic acid (PFDoDA). Metals were measured in whole

blood samples; cadmium (Cd) and lead (Pb) by inductively coupled plasma-mass spectrometry (Thermo X7, Thermo Elemental, Winsford, UK) and mercury (Hg) by cold vapour atom fluorescence spectrophotometry. 2,2',4,4',5,5'-hexachlorobiphenyl (PCB-153) and the dichlorodiphenyltrichloroethane metabolite, 1,1-dichloro-2,2-bis(*p*-chlorophenyl)-ethylene (*p,p'*-DDE), were analysed by gas chromatography-mass spectrometry,<sup>11</sup> as was hexachlorobenzene (HCB) according to a modified method by Otero *et al.*<sup>12</sup>

Twenty-two markers of reproductive function (listed in table 1) were assessed following standardised protocols<sup>13–17</sup>: reproductive hormones, conventional semen characteristics, markers of sperm chromatin integrity and apoptosis, markers of epididymal and accessory sex gland function, and the proportion of Y:X chromosome-bearing sperm. Analytical details are provided in the online supplementary methods and table S1.

### Statistical analysis

We used sPLS regression modelling to assess associations between the exposure profiles and each outcome. In PLS regression, an X-matrix of exposures and Y outcome vector are simultaneously decomposed into latent variables and regressed in a way that maximises the covariance between X and Y.<sup>18–19</sup> These latent variables, called components, are orthogonal, linear combinations of the full set of input variables. Chun and Keleş<sup>10</sup> proposed sPLS, embedding variable selection into the PLS algorithm by eliminating 'uninformative' X-variables via penalisation. A penalty term ( $L_1$ ) is applied during dimension reduction, and regression coefficients are obtained via PLS for the reduced set of X-variables (see online supplementary methods). Model complexity is a function of the number of components used to construct the model (K) and the degree of sparsity ( $\eta$ ). To determine the optimal model (avoid overfitting), we performed Monte Carlo cross-validation,<sup>20</sup> and added a null (intercept only) model in the cross-validation loop to assess significance. We tested models with K values 1–5 and  $\eta$  values between 0.01 and 0.99 (more sparse) in steps of 0.01, using a fivefold cross-validation, repeated (with different random partitions of the data) 500 times. We selected the most parsimonious model within one SE of the overall minimum mean squared error of prediction (MSEP).<sup>19</sup>

We took several data pretreatment and exploration steps before undertaking sPLS analyses. We imputed exposure data below the limit of detection (LOD) (0–18%) and values missing-at-random in the matrix of exposures (2–16%). Values were imputed from a log-normal probability distribution via single conditional imputation, dependent on the population, and detected values for the other exposures. In the primary analyses, HCB, PCB-153 and *p,p'*-DDE were lipid adjusted (see online supplementary methods), and four contaminants for which <70%<sup>21</sup> of the samples had concentrations above the LOD were excluded: MEOHP, MOiNP, PFUnDA and PFDoDA. To approximate normality, we used natural logarithm (ln)-transformed values for all exposure variables and for 14 of the 22 outcome variables.

To address potential confounding, we employed a two-stage regression approach<sup>22</sup>: first, each outcome and each exposure were separately regressed on potential confounders, and second, sPLS regression models were fit inputting the residuals. We a priori selected the set of potential confounders,<sup>9</sup> including study population and serum cotinine level, and variably age, body mass index (BMI), abstinence period and time of blood sampling, depending on the outcome (as specified in table 1). We imputed missing covariate data (0.8–16%; as elaborated in the online supplementary methods). To assess the potential for

**Table 1** Summary statistics for biomarkers of reproductive function and potential confounders among 602 male partners of pregnant women

Outcomes and covariates	n*	Greenland (n=199) GM (5, 95 P)	Warsaw, Poland (n=197) GM (5, 95 P)	Kharkiv, Ukraine (n=206) GM (5, 95 P)	p Value†
Male reproductive hormones in serum§¶					
Follicle-stimulating hormone (IU/L)	456	4.38 (1.70, 9.20)	3.46 (1.40, 7.40)	3.67 (1.60, 9.40)	0.001
Luteinising hormone (IU/L)	456	4.05 (1.80, 8.50)	3.81 (1.90, 7.10)	3.79 (1.90, 7.60)	0.34
Inhibin B (ng/L)**	456	184.21 (83.00, 314.00)	157.94 (71.00, 275.00)	194.74 (98.00, 321.00)	<0.001
Sex hormone-binding globulin (nmol/L)	455	26.90 (15.60, 46.70)	21.61 (9.45, 40.90)	26.19 (15.50, 43.20)	<0.001
Total testosterone (nmol/L)**	456	14.81 (7.27, 24.41)	13.08 (6.83, 20.76)	18.01 (10.26, 26.89)	<0.001
Free testosterone (nmol/L)**	455	0.31 (0.18, 0.48)	0.29 (0.16, 0.42)	0.39 (0.22, 0.58)	<0.001
Estradiol (pmol/L)	454	63.44 (40.50, 98.70)	71.50 (45.80, 112.60)	80.73 (52.50, 137.50)	<0.001
Conventional semen characteristics					
Semen volume (mL)††,‡‡	535	3.21 (1.20, 7.20)	3.37 (1.25, 7.19)	3.29 (1.59, 7.57)	0.68
Sperm concentration (10 <sup>6</sup> /mL)††	600	51.12 (10.75, 199.00)	55.95 (7.15, 257.50)	53.07 (10.15, 185.80)	0.65
Total sperm count (10 <sup>6</sup> /ejaculate)††,‡‡	533	160.15 (31.50, 692.30)	186.68 (18.82, 1061.96)	179.46 (34.20, 804.80)	0.39
Morphologically normal (%)**	598	6.97 (2.00, 13.00)	6.66 (2.00, 14.00)	7.37 (1.00, 15.00)	0.20
Progressive motility (% A+B)**,††,§§	565	55.29 (22.00, 81.00)	60.98 (19.00, 87.00)	55.08 (14.50, 87.00)	0.007
Sperm chromatin integrity§,††					
SCSA DNA fragmentation index (%)	546	7.66 (3.32, 20.28)	10.07 (4.16, 31.50)	10.83 (3.54, 31.56)	<0.001
High DNA stainability (%)	546	11.37 (4.37, 29.88)	8.66 (3.60, 25.14)	10.29 (4.35, 26.55)	<0.001
TUNEL DNA fragmentation index (%)	462	2.95 (0.79, 13.37)	11.64 (2.57, 38.28)	6.46 (1.90, 30.18)	<0.001
Apoptotic markers§,††					
Fas positivity (%)**	454	25.00 (2.22, 70.60)	48.57 (3.13, 96.42)	28.24 (0.00, 90.60)	<0.001
Bcl-xL positivity (%)**	286	26.54 (0.44, 90.65)	18.05 (0.81, 69.64)	66.01 (9.54, 99.17)	<0.001
Epididymal and accessory sex gland function§,††,‡‡					
Neutral $\alpha$ -glucosidase (mU/ejaculate)	448	16.32 (4.80, 52.00)	21.81 (6.55, 71.52)	17.51 (4.86, 52.56)	0.001
Prostate-specific antigen ( $\mu$ g/ejaculate)	507	3187.74 (1078.00, 8330.85)	3708.59 (1312.37, 11127.89)	2497.94 (784.08, 7707.70)	<0.001
Zinc ( $\mu$ mol/ejaculate)	497	4.68 (1.06, 12.63)	6.31 (1.50, 25.23)	4.22 (1.11, 16.56)	<0.001
Fructose ( $\mu$ mol/ejaculate)	506	47.00 (12.67, 153.85)	42.24 (6.10, 161.65)	39.36 (10.69, 117.35)	0.18
Y chromosome sperm cells (%)**	411	51.26 (48.52, 55.49)	50.30 (48.79, 52.11)	50.76 (48.72, 53.61)	<0.001
Age (years)**	597	31.34 (20.83, 43.15)	30.34 (25.36, 36.80)	27.88 (20.67, 38.75)	<0.001
BMI (kg/m <sup>2</sup> )**	592	25.99 (20.56, 31.77)	25.81 (20.43, 31.93)	24.20 (19.71, 29.41)	<0.001
Abstinence period (days)	557	2.86 (0.50, 7.10)	4.27 (1.00, 30.00)	3.41 (1.50, 7.00)	<0.001
Current smoker, n (%)	597	144 (72.7%)	53 (27.2%)	136 (66.7%)	<0.001
Current smokers: cotinine (ng/mL)	589	113.73 (0.96, 465.23)	23.95 (<LOD, 358.42)	33.59 (<LOD, 447.36)	<0.001
Time of blood sampling, <12:00, n (%)	504	17 (16.8%)	187 (94.9%)	183 (88.8%)	<0.001

\*Number available out of 602.

†Test for difference in levels between the three study populations: ANOVA for means and  $\chi^2$  test for proportions.

‡In adjusted analyses, adjusted for time of blood sampling.

§In adjusted analyses, adjusted for age.

¶In adjusted analyses, adjusted for BMI.

\*\*Outcome was not ln-transformed; arithmetic mean presented.

††In adjusted analyses, adjusted for ln-abstinence period.

‡‡Samples with spillage were excluded (n=67 for semen volume and sperm count; n=41–52 for the 4 epididymal and accessory sex gland function markers).

§§Samples with a delay of &gt;1 h from collection to semen analysis were excluded (n=28).


BMI, body mass index; GM, geometric mean; LOD, limit of detection; P, percentile; SCSA, sperm chromatin structure assay; TUNEL, terminal deoxynucleotidyl transferase dUTP nick end-labelling.

overadjustment, in a secondary analysis we evaluated sPLS models with exposure and outcome variables only prestandardised (centred) for study population. This was considered the minimal adjustment, necessary due to the large differences in central tendencies between the populations for the exposures (figure 1), and to a lesser degree but still conspicuous, for the outcomes. Mean centring also removed baseline offsets, while scaling by log-transformation reduced the variance differences between X-variables, and thus the relative importance during dimension reduction.<sup>23</sup>

As an additional screening approach, we tested 330 single-pollutant, covariate-adjusted OLS linear regression models for associations between the 15 exposures (with >70% detection frequency) and 22 outcomes. To adjust for multiple comparisons, we computed the false discovery rate (FDR),<sup>24</sup> and set the

significance threshold at an FDR <10%. In sensitivity analyses, we tested the four contaminant exposures with <70% detection frequency dichotomised as detect versus non-detect variables, and tested models excluding participants with imputed confounder and (missing-at-random) exposure data. We also assessed models with lipid-unadjusted organochlorines, including total lipids as a covariate. Finally, we tested for heterogeneity in effect estimates between study populations with an interaction term, and by assessing plots of the population-stratified regressions.

sPLS and OLS regression coefficients are presented per ln-unit increase in exposure, and for interpretability, converted into the per cent change in outcome per IQR increase in exposure; for ln-transformed outcomes, the proportional change, and for untransformed outcomes, the absolute change relative to the



**Figure 1** Box plots of the distributions of exposure biomarker concentrations per study population. MEHHP, mono(2-ethyl-5-hydroxyhexyl) phthalate; MECPP, mono(2-ethyl-5-carboxypentyl) phthalate; MHiNP, mono(4-methyl-7-hydroxyoctyl) phthalate; MOiCP, mono(4-methyl-7-carboxyheptyl) phthalate; Hg, mercury; PFHxS, perfluorhexane sulfonic acid; PFOS, perfluorooctane sulfonic acid; PFOA, perfluorooctanoic acid; PFNA, perfluorononanoic acid; PFDA, perfluorodecanoic acid; HCB, hexachlorobenzene; PCB, polychlorinated biphenyl; DDE, dichlorodiphenyldichloroethylene.

mean outcome level. Statistical analyses were performed using R V.3.0.1 (R Foundation for Statistical Computing, Vienna, Austria), and specifically the *spls* package for sPLS modelling.<sup>25</sup>

## RESULTS

Descriptive characteristics of the study populations are shown in table 1. Participants from Kharkiv were slightly younger than their counterparts from Greenland and Warsaw, and fewer participants from Warsaw smoked. In exploring the structure of the exposure data, we observed large variations within and between the three study populations (figure 1). Levels differed across study populations for all contaminant exposures (analysis of variance  $p$  value  $<0.05$ , online supplementary table S2), except for MOiCP. Levels were higher in samples from Greenland compared to Warsaw and Kharkiv for many exposures, especially for Hg and PCB-153. Correlations within the X-matrix ranged from  $r_p = -0.48$  to  $0.91$  (figure 2). We also observed distributional differences for many outcomes (table 1), although distributions for the three populations were generally more similar for the outcomes than for the exposures.

Of the 22 sPLS regression models tested in the primary analysis ( $n=286$ – $600$  complete cases), the X-matrix significantly improved the prediction error beyond the null model for 8 outcomes: luteinising hormone (LH), inhibin B, total testosterone, free testosterone, semen volume, progressive sperm motility, terminal deoxynucleotidyl transferase dUTP nick end-labelling assay (TUNEL) DNA fragmentation index (DFI), and neutral  $\alpha$ -glucosidase (NAG). Within each of these significant sPLS regression models, 1 to 3 exposures within the X-matrix contributed to a total of 10 significant exposure–outcome associations (table 2). The optimal models were all  $K=1$  component models and quite sparse ( $\eta$  between  $0.64$  and  $0.99$ ).

In the secondary analysis with exposure and outcome variables only prestandardised for study population but not for other

potential confounders, sex hormone-binding globulin (SHBG) was additionally selected (the inclusion of the X-matrix improved the MSEF beyond the null model), and the LH model was no longer selected (see online supplementary table S3). Extra exposures were also selected, for a total of 24 associations.

In the covariate-adjusted OLS regression analyses, 8 overlapping exposure–outcome associations, comprising the same 8 outcomes but fewer exposures (8 of 10) compared to the primary sPLS analyses, were identified as significant ( $FDR <10\%$ ). Two associations, between MHiNP and total testosterone and TUNEL DFI, were significant at an  $FDR <5\%$  significance threshold (see online supplementary figure S2). An additional association between MEHHP and TUNEL DFI, not selected in the sPLS model, was significant in OLS analyses (see online supplementary table S3). Consistent with the sPLS modelling results, a greater number of associations was significant in the secondary analysis with OLS models only adjusted for study population compared to the primary analysis with further adjusted models (15 compared to 9; 14 of which overlapped with those selected in sPLS secondary analysis only prestandardised for study population). No additional associations were detected in the sensitivity analysis testing contaminants with  $<70\%$  detection frequency inputted as binary detect versus non-detect variables. In a reanalysis using only the complete (non-imputed) exposure and covariate data, 2 associations— $p,p'$ -DDE and LH, and MHiNP and total testosterone—were selected in both the sPLS and OLS modelling (with  $n=321$ – $480$  and  $328$ – $586$  complete cases, respectively, except for Bcl-xL positivity models which had  $192$ – $252$  complete cases). This analysis had reduced power, and while CIs were wider, coefficients for the complete case analysis were remarkably similar to the primary analysis with imputed data (data not shown). Coefficients for lipid unadjusted organochlorines, adjusted for total lipids, were only slightly different in magnitude (see footnotes of table 2).

**Figure 2** Pairwise Pearson correlation coefficients, also represented as a heat map, of the exposure biomarkers. MEHHP, mono(2-ethyl-5-hydroxyhexyl) phthalate; MECPP, mono(2-ethyl-5-carboxypentyl) phthalate; MHiNP, mono(4-methyl-7-hydroxyoctyl) phthalate; MOiCP, mono(4-methyl-7-carboxyheptyl) phthalate; Hg, mercury; PFHxS, perfluorhexane sulfonic acid; PFOS, perfluorooctane sulfonic acid; PFOA, perfluorooctanoic acid; PFNA, perfluorononanoic acid; PFDA, perfluorodecanoic acid; HCB, hexachlorobenzene; PCB, polychlorinated biphenyl; DDE, dichlorodiphenyldichloroethylene.

	MEHHP	MECPP	MHiNP	MOiCP	Hg	Cd	Pb	PFHxS	PFOS	PFOA	PFNA	PFDA	HCB	PCB-153	<i>p,p'</i> -DDE
MEHHP	1														
MECPP	0.34	1													
MHiNP	0.40	0.34	1												
MOiCP	0.21	0.46	0.61	1											
Hg	0.28	-0.26	0.21	-0.02	1										
Cd	0.12	-0.07	0.05	0.03	0.20	1									
Pb	0.03	-0.01	0.01	0.01	0.18	0.28	1								
PFHxS	0.24	-0.32	0.20	-0.06	0.69	0.07	0.04	1							
PFOS	0.28	-0.32	0.24	-0.03	0.74	0.08	0.08	0.91	1						
PFOA	0.11	-0.23	0.15	0.02	0.36	-0.14	-0.17	0.71	0.7	1					
PFNA	0.21	-0.16	0.16	-0.02	0.63	0.08	0.20	0.63	0.65	0.42	1				
PFDA	0.28	-0.27	0.21	-0.03	0.78	0.09	0.14	0.81	0.87	0.61	0.80	1			
HCB	0.06	0.05	-0.02	-0.02	0.21	0.18	0.30	-0.24	-0.15	-0.48	0.18	-0.01	1		
PCB-153	0.25	-0.19	0.17	-0.06	0.76	0.27	0.29	0.46	0.55	0.09	0.54	0.60	0.53	1	
<i>p,p'</i> -DDE	0.00	0.07	-0.12	-0.10	0.12	0.02	0.20	-0.15	-0.15	-0.33	0.22	0.02	0.55	0.34	1

For 5 of the 10 sPLS-selected associations, there was a significant interaction between exposure and study population in the OLS sensitivity analysis (table 2). It was apparent that 8 associations were consistent in slope direction across study populations in regression models stratified by population (see online supplementary figure S3). Poland exhibited a slope in the opposite direction for LH and *p,p'*-DDE, and for the oxidative phthalates and TUNEL DFI. Associations with consistent slope directions, and their  $\beta_{\text{sPLS}}$  corresponding overall per cent change per IQR increase in exposure, included a 11% increase in

inhibin B with Hg; a 6% decline in total testosterone with MHiNP; a 3% and 2% decline in free testosterone with DiNP metabolites, and a 5% increase with Cd; a 10% decline in semen volume with MEHHP; a 11% decline in the fraction of progressively motile sperm with PCB-153; and a 16% decline in NAG with MEHHP.

## DISCUSSION

Using a multipollutant modelling approach, we identified 10 associations between biomarkers of environmental contaminant

**Table 2** Exposure–outcome associations and regression coefficients selected from multipollutant sPLS regression modelling and corresponding single-pollutant OLS regression coefficients

Outcome	Exposure*	sPLS		OLS		
		K, $\eta$ †	$\beta_{\text{sPLS}}$	$\beta_{\text{OLS}}$	95% CI	p Value
LH* (IU/L)	<i>p,p'</i> -DDE (ng/g)	1, 0.99	0.083	0.083‡,§	(0.031 to 0.135)	0.002¶
Inhibin B (ng/L)	Hg (ng/mL)	1, 0.99	10.79	10.82	(3.90 to 17.73)	0.002¶
Total testosterone (nmol/L)	MHiNP (ng/mL)	1, 0.90	-1.14	-1.15‡	(-1.74 to -0.57)	0.0001¶
Free testosterone (nmol/L)	MHiNP (ng/mL)	1, 0.64	-0.0113	-0.019‡	(-0.032 to -0.007)	0.002¶
	MOiCP (ng/mL)		-0.0091	-0.010‡	(-0.020 to 0.000)	0.043
	Cd (ng/mL)		0.0091	0.012	(0.001 to 0.023)	0.029
Semen volume* (mL)	MEHHP (ng/mL)	1, 0.99	-0.106	-0.106	(-0.167 to -0.045)	0.0008¶
Progressive sperm (%)	PCB-153 (ng/g)	1, 0.99	-3.37	-3.37§	(-5.48 to -1.25)	0.002¶
TUNEL DFI* (%)	MHiNP (ng/mL)	1, 0.99	-0.218	-0.218‡	(-0.332 to -0.104)	0.0002¶
NAG* (mU/ejaculate)	MEHHP (ng/mL)	1, 0.77	-0.163	-0.164	(-0.255 to -0.073)	0.0005¶

Regression coefficients ( $\beta$ ) are expressed per ln-unit change in exposure. Refer to online supplementary table S3 for per cent changes and coefficients from models only adjusted for study population. sPLS and OLS models were adjusted for study population, cotinine and for additional potential confounders as specified in table 1; sPLS models by inputting exposures and outcomes 'prestandardised' for covariates (residuals).

\*Variable ln-transformed in statistical analyses.

†sPLS tuning parameters: K, the number of components used and  $\eta$ , the degree of sparsity.

‡Interaction p value <0.10 for the cross-product term between exposure and study population (see online supplementary figure S3 for population-stratified regression plots).

§Sensitivity analysis:  $\beta_{\text{OLS}}$  (95% CI) for models with organochlorines unadjusted for lipids (ng/mL), and with total lipids (g/L) included as an additional covariate: LH and *p,p'*-DDE, 0.070 (0.017, 0.123); progressive sperm and PCB-153, -3.592 (-5.871 to -1.313).

¶Significant after adjustment for multiple comparisons in single-pollutant OLS regression modelling (FDR <10%).

DDE, dichlorodiphenyldichloroethylene; DFI, DNA fragmentation index; FDR, false discovery rate; Hg, mercury; LH, luteinising hormone; MHiNP, mono(4-methyl-7-hydroxyoctyl) phthalate; MOiCP, mono(4-methyl-7-carboxyheptyl) phthalate; MEHHP, mono(2-ethyl-5-hydroxyhexyl) phthalate; NAG, neutral  $\alpha$ -glucosidase; OLS, ordinary least squares; PCB, polychlorinated biphenyl; sPLS, sparse partial least squares; TUNEL, terminal deoxynucleotidyl transferase dUTP nick end-labelling.

exposure and biomarkers of male reproductive function from several hundred associations tested in (sub)fertile men from Greenland, Warsaw and Kharkiv. This is the first time several of the relationships have been assessed in an epidemiological study, or simultaneously.

One of the most statistically robust and consistent associations was between DiNP metabolites and decreasing free (bioactive) testosterone and total testosterone. DEHP metabolites did not exceed the significance threshold in confounder-adjusted analyses but exhibited inverse relationships. DiNP exposure is outpacing DEHP exposure in Europe, where DiNP use is less restricted than that of DEHP. These inverse associations are consistent with an antiandrogenic effect, which has more often been observed compared to pro-androgenic effects in experimental studies.<sup>26</sup> Phthalates are generally considered to act by inhibiting Leydig cell synthesis of testosterone. While most experimental studies have focused on in utero exposure, DEHP and its primary metabolite MEHP were recently shown to suppress steroidogenesis of adult human testis explants in vitro, without affecting production of the hormone insulin-like factor 3 (INSL3).<sup>27</sup> Several epidemiological studies have found inverse associations between (urinary) MEHP and testosterone in men; in 74 occupationally exposed workers and 63 controls in China,<sup>28</sup> and in a pooled analysis of 425 men of infertile couples and 425 men recruited through a US infertility clinic.<sup>29</sup> No association was found in 234 young Swedish men sampled during routine medical conscript examinations.<sup>30</sup> Joensen *et al*<sup>31</sup> considered another exposure metric, the proportion of the summed urinary metabolites excreted as the primary metabolite, as a marker of DEHP and DiNP metabolism; %MEHP and %MiNP were associated (significant at  $p < 0.05$  or near-significant) with total testosterone and free androgen index in 881 healthy young men from Denmark around 19 years of age. However, no significant associations were reported for the sum of primary and secondary metabolites. Primary monoester phthalate metabolites were not measured in the current study—only secondary oxidative metabolites—due to the lipase activity of serum and risk of contamination from the diesters from the sampling devices.

We found a clear dose-dependent decrease in serum testosterone levels with increasing levels of MHiNP. We recently discovered, however, that the measurement of MHiNP was possibly confounded by an unknown co-eluting agent. Nevertheless, in an additional analysis using the molar sum of the other two DiNP metabolites versus all three oxidative metabolites tested in relation to total testosterone, the magnitude of the regression coefficients was similar:  $\beta_{OLS} -0.713$  ( $p$  value=0.0037) for  $\Sigma DiNP_{MOiNP+MOiCP}$  versus  $-0.929$  (0.0007) for  $\Sigma DiNP_{om}$ . Moreover, that other phthalate metabolites were significantly inversely associated with testosterone lends weight to the evidence that one or more phthalates is indeed related to decreasing testosterone levels.

We observed a positive association between cadmium and free testosterone in the sPLS analyses, although this association was not significant in the OLS analyses adjusted for potential confounders, notably cotinine. Disentangling the effect of smoking and cadmium is difficult as smoking is an important source of cadmium exposure. A cadmium–testosterone association has previously been reported in 219 men recruited from two US infertility clinics,<sup>32</sup> and in a representative sample of 1262 US men,<sup>33</sup> although this latter association did not sustain statistical significance after adjustment for smoking. An experimental study of chronic low-dose oral exposure in rats reported

cadmium-related elevated testosterone<sup>34</sup>; however, the underlying mechanism remains unknown.

We observed decreasing NAG levels with increasing MEHHP. Sperm undergo maturation in the epididymis, and NAG is a marker of epididymal function. A smaller study ( $n=234$ ) of urinary phthalates and NAG in Swedish conscripts was null.<sup>30</sup> In a study of 40 rats administered di-*n*-butyl phthalate, epididymal weight and NAG activity decreased dose dependently, in line with our finding.<sup>35</sup>

We found decreasing sperm DNA damage with increasing levels of the secondary hydroxylated metabolite of DiNP, counter to the hypothesis of a deleterious effect of exposure. However, the exposure–outcome relationship was inconsistent in direction, exhibiting a positive slope in the Polish study population. We detected associations for DNA fragmentation assessed with TUNEL, but not the sperm chromatin structure assay (SCSA). TUNEL represents DNA strand breaks measured directly in the ejaculated spermatozoa, whereas SCSA is a measure of susceptibility to acid denaturation.<sup>36</sup> Sperm DNA damage might be mediated by testicular oxidative stress. In rats, DEHP decreased antioxidant enzymes and concomitantly increased reactive oxygen species.<sup>37</sup> In 379 US men seeking infertility treatment, urinary monoethyl phthalate and MEHP were positively associated with neutral Comet assay parameters.<sup>38</sup> MEHP was more strongly associated when adjusted for oxidative metabolites (MEHHP and MEOHP), and similar to our finding for MEHHP, the metabolites were inversely associated with DNA damage. Hauser *et al*<sup>38</sup> suggested that an increased ratio of oxidative metabolites to MEHP could represent interindividual differences in rates of DEHP metabolism to ‘less toxic’ metabolites, or it may reflect inhibition of phase 1 enzymes involved in xenobiotic metabolism.

Some of the exposure–outcome associations tested in the current analysis have previously been tested in the same or overlapping study populations in analyses considering a single-pollutant or a single class of chemicals (refer to refs. <sup>39–41</sup>; previous investigations of PCB-153 and *p,p'*-DDE were reviewed by Bonde *et al*<sup>5</sup>). In addition to identifying several novel associations, we confirmed, using a more conservative significance threshold and joint modelling approach, several previously reported associations, including an unfavourable inverse association between PCB-153 and the proportion of progressively motile sperm.<sup>5 17</sup> Other smaller studies have reported comparable findings, for instance, for PCB-153 in 305 Swedish men,<sup>42</sup> and for PCB-138 but not PCB-153 and the odds of <50% progressive motility in 212 US men recruited at an infertility clinic (uncorrected for multiple comparisons).<sup>43</sup>

We also confirm a previously reported strong positive association between mercury and inhibin B.<sup>40</sup> As inhibin B, predominantly excreted by Sertoli cells, is correlated with sperm concentration and counts and serves as a marker of spermatogenesis,<sup>44</sup> this positive association is contrary to a deleterious effect of exposure. The authors<sup>40</sup> suggest that this association may be driven by a higher consumption of seafood—particularly relevant for Greenlandic Inuit—and the resultant concomitant exposure to mercury and  $\omega$ -3 polyunsaturated fatty acids. There are indications that the latter may beneficially impact certain semen quality parameters.<sup>45</sup> This mercury–inhibin B association was not observed in 219 US men recruited from infertility clinics.<sup>32</sup>

We observed an association between MEHHP and a decline in semen volume, as previously reported for this study population.<sup>39</sup> A null finding for MEHP was reported for a study of

234 young Swedish men,<sup>30</sup> and a statistically significant increase in semen volume for the highest versus lowest %MiNP quartile in 881 Danish men was interpreted by the authors as a chance finding.<sup>31</sup>

We also observed a positive, yet inconsistent-across-populations, association between *p,p'*-DDE and increasing LH, as previously reported.<sup>17</sup> As LH regulates testosterone production by Leydig cells, this is in line with an antiandrogenic effect of *p,p'*-DDE. A null finding was reported for 341 men from a US infertility clinic with lower *p,p'*-DDE levels (geometric mean 236 ng/g lipids).<sup>46</sup>

We found indications of associations between phthalates, metals and organochlorines and SHBG, which did not sustain significance on adjustment for confounders, specifically BMI. These non-significant perturbations in SHBG, together with the other associations with hormones, might be indicative of effects of exposure profiles on hypothalamo–pituitary–gonadal axis regulation. Similar to the potential (over)adjustment for cotinine, adjustment for BMI may introduce overadjustment bias rather than (only) prevent confounding bias, given that some exposures may causally influence BMI.

A limitation of this assessment is the cross-sectional nature of this study, although, besides phthalates, all measured contaminants are relatively stable over time and as such were most likely reasonable proxies of recent past exposure as related to the measured outcomes. However, while blood and semen samples were collected on the same day for nearly all of the participants from Poland and Ukraine, for around 60% of the participants from Greenland, samples were collected more than 3 days apart (elaborated in the online supplementary methods), contributing to misclassification of exposure, especially for the phthalates. Analyses excluding these participants yielded generally consistent results.

Recent advances in molecular epidemiology (high-throughput screening techniques) and the introduction of the exposome concept<sup>47</sup> imply that epidemiological researchers will be confronted more and more with the challenging task of analysing data with high numbers of measured analytes with often complex correlation structures. Single-pollutant modelling may result in a high potential for false positives (type I errors), although multiple testing corrections and the recently proposed environment-wide association study<sup>48</sup> design addresses this. Using multipollutant OLS regression modelling may lead to inflated variances and flipped signs of parameter estimates, thus making it difficult to disentangle the independent effects of these variables. Methods which project data onto a lower dimensional space, like PLS regression, can better cope with collinear variables and, as a corollary, reduce the number of models tested.<sup>10 18</sup> While PLS-based methods are well established in chemometrics and bioinformatics, they are only recently gaining traction in genomics, metabolomics and epidemiology.<sup>49</sup>

In the current analysis, the associations selected using sPLS rather closely matched those we would have selected using single-pollutant OLS models. However, if we had screened for associations using multipollutant OLS models, accounting for correlations between exposures, these estimates would have suffered from multicollinearity and been unreliable (variance inflation factors exceeded 3 for 8 to 9 of the 15 exposure terms across multipollutant OLS models).

As sPLS captures the complementary contribution of X-variables, it has the power to detect multiple pollutants which might not be detected in single-pollutant modelling. This can be explained by additivity, and/or consistency at large, in that noise from measurement error and biological variation can be

averaged out when a larger number of X-variables inform the latent variable. This was evident for the free testosterone model, and for several models in the sensitivity analysis with inputs only prestandardised for study population, in which up to 8 exposures were selected. Further, in constructing latent variables, sPLS regression delineates groupings in the X-matrix of exposures, which is a step forward in tackling the long-standing challenge of assessing mixture effects of contaminants.

Some aspects of (s)PLS modelling implementation are underdeveloped, such as quantifying uncertainty (stability of the selection), and explicitly correcting for multiple testing, beyond the implicit sparseness step. We estimated an empirical null distribution using a Monte Carlo approach to establish a significance threshold. A caveat of sPLS is that penalisation methods optimise prediction and will tend to select one of two highly correlated exposures with nearly equal coefficient sizes.<sup>19</sup> Competing multipollutant modelling approaches exist<sup>22</sup> (eg, tree-based models, elastic net penalised regression); however, simulation and validation assessments, specifically for data structures relevant for environmental epidemiology (often low dimensional), are lacking.

There is no straightforward way to adjust for potential confounders in (s)PLS analyses. Adjusting for confounders in sPLS models by inputting residuals in a two-step approach should, as with OLS modelling,<sup>50</sup> yield unbiased regression coefficients, although with a slight loss of efficiency under certain (sample size, correlation) scenarios. We observed large contrasts in exposures across the three study populations, and some differences in outcomes such as TUNEL DFI. To reduce the risk of ecological bias due to potential between-population confounding, in all analyses we prestandardised (mean centred) the exposure and outcome data by population (sPLS models) or adjusted by population (OLS models). This may have artificially reduced true contrast in exposure and outcome data between study populations, and negatively affected our power to detect some associations.

While (s)PLS modelling has some power to detect non-linear relationships (weighted over components), the PLS and OLS regression coefficients reflect linear relationships. In an exploratory analysis with generalised additive modelling, this assumption of linearity was met for most exposure–outcome relationships (<2% deviated from linearity, with an estimated degree of freedom >1.0). Further research is needed to compare the efficacy of sPLS regression with other multipollutant modelling and variable selection approaches, including non-parametric techniques.

## CONCLUSIONS

We used a systematic, multipollutant modelling approach to perform a global assessment of biomarkers of contaminant exposure and male reproduction function in one of the largest studies to date. We identified several environmentally perturbed outcomes, including a robust inverse association between DiNP phthalates and circulating testosterone, and between PCB-153 and sperm motility.

### Author affiliations

<sup>1</sup>Division of Environmental Epidemiology, Institute for Risk Assessment Sciences, Utrecht University, Utrecht, The Netherlands

<sup>2</sup>Division of Occupational and Environmental Medicine, Lund University, Lund, Sweden

<sup>3</sup>Reproductive Medicine Centre, Skåne University Hospital Malmö, Lund University, Malmö, Sweden

<sup>4</sup>Laboratory of Toxicology, Unit of Radiation Biology and Human Health, ENEA Casaccia Research Center, Rome, Italy

<sup>5</sup>Centre for Arctic Environmental Medicine, Nuuk, Greenland

<sup>6</sup>Department of Toxicology and Risk Assessment, National Institute of Public Health-National Institute of Hygiene, Warsaw, Poland

<sup>7</sup>Department of Social Medicine and Organization of Public Health, Kharkiv National Medical University, Kharkiv, Ukraine

<sup>8</sup>Laboratory for Health Protection Research, National Institute for Public Health and the Environment (RIVM), Bilthoven, The Netherlands

<sup>9</sup>Department of Occupational Medicine, Danish Ramazzini Center, Aarhus University Hospital, Aarhus, Denmark

<sup>10</sup>Department of Occupational and Environmental Medicine, Bispebjerg Hospital, University of Copenhagen, Copenhagen, Denmark

**Contributors** VL, LP, DH and RV conceived and designed the current data analyses. JPB, GT, AG, LR, BAGJ, MS, HSP, JKL and LC designed and collected data for the cohort study. BAGJ and CHL performed chemical analyses. VL analysed the data and drafted the manuscript, with supervision from LP and RV. All authors, particularly LP, RV, DH, LAMS, AHP, JPB, GT, AG and MS, participated in the interpretation of results and revision of the manuscript. All authors approved the final version of the manuscript.

**Funding** This work was supported by the European Commission's 7th and 5th Framework Programmes (grants FP7-ENV-2008-1-226217 and QLK4-CT-2001-00202).

**Competing interests** None.

**Patient consent** Obtained.

**Ethics approval** Polish Bioethical Committee, Ethical Committee for Human Research in Greenland, Commission on Ethics and Bioethics of Kharkiv National Medical University.

**Provenance and peer review** Not commissioned; externally peer reviewed.

## REFERENCES

- Jørgensen N, Asklund C, Carlsen E, *et al.* Coordinated European investigations of semen quality: results from studies of Scandinavian young men is a matter of concern. *Int J Androl* 2006;29:54–61.
- Travis TG, Araujo AB, O'Donnell AB, *et al.* A population-level decline in serum testosterone levels in American men. *J Clin Endocrinol Metab* 2007;92:196–202.
- Jørgensen N, Joensen UN, Jensen TK, *et al.* Human semen quality in the new millennium: a prospective cross-sectional population-based study of 4867 men. *BMJ Open* 2012;2.
- Carlsen E, Giwercman A, Keiding N, *et al.* Evidence for decreasing quality of semen during past 50 years. *BMJ* 1992;305:609–13.
- Bonde JP, Toft G, Rylander L, *et al.* Fertility and markers of male reproductive function in Inuit and European populations spanning large contrasts in blood levels of persistent organochlorines. *Environ Health Perspect* 2008;116:269–77.
- Toft G, Axmon A, Giwercman A, *et al.* Fertility in four regions spanning large contrasts in serum levels of widespread persistent organochlorines: a cross-sectional study. *Environ Health* 2005;4:26.
- Groshart C, Okkerman PC. *Towards the establishment of a priority list of substances for further evaluation of their role in endocrine disruption—preparation of a candidate list of substances as a basis for priority setting.* European Commission DG ENV, 2000. [http://ec.europa.eu/environment/archives/docum/pdf/bkh\\_main.pdf](http://ec.europa.eu/environment/archives/docum/pdf/bkh_main.pdf) (accessed Apr 2014).
- CDC (Centers for Disease Control and Prevention). Fourth National Report on Human Exposure to Environmental Chemicals. Atlanta, GA. 2009. <http://www.cdc.gov/exposurereport/pdf/fourthreport.pdf> (accessed Apr 2014).
- Lindh CH, Rylander L, Toft G, *et al.* Blood serum concentrations of perfluorinated compounds in men from Greenlandic Inuit and European populations. *Chemosphere* 2012;88:1269–75.
- Chun H, Keles S. Sparse partial least squares regression for simultaneous dimension reduction and variable selection. *J R Stat Soc Series B Stat Methodol* 2010;72:3–25.
- Jönsson BAG, Rylander L, Lindh C, *et al.* Inter-population variations in concentrations, determinants of and correlations between 2,2',4,4',5,5'-hexachlorobiphenyl (CB-153) and 1,1-dichloro-2,2-bis(p-chlorophenyl)-ethylene (p,p'-DDE): a cross-sectional study of 3161 men and women from Inuit and European populations. *Environ Health* 2005;4:27.
- Otero R, Santiago-Silva M, Grimalt JO. Hexachlorocyclohexanes in human blood serum. *J Chromatogr A* 1997;778:87–94.
- Elzanaty S, Rignell-Hydbom A, Jönsson BAG, *et al.* Association between exposure to persistent organohalogen pollutants and epididymal and accessory sex gland function: multicentre study in Inuit and European populations. *Reprod Toxicol* 2006;22:765–73.
- Stronati A, Manicardi GC, Cecati M, *et al.* Relationships between sperm DNA fragmentation, sperm apoptotic markers and serum levels of CB-153 and p,p'-DDE in European and Inuit populations. *Reproduction* 2006;132:949–58.
- Spanò M, Toft G, Hagmar L, *et al.* Exposure to PCB and p,p'-DDE in European and Inuit populations: impact on human sperm chromatin integrity. *Hum Reprod* 2005;20:3488–99.
- Toft G, Rignell-Hydbom A, Tyrkiel E, *et al.* Semen quality and exposure to persistent organochlorine pollutants. *Epidemiology* 2006;17:450–8.
- Giwercman AH, Rignell-Hydbom A, Toft G, *et al.* Reproductive hormone levels in men exposed to persistent organohalogen pollutants: a study of Inuit and three European cohorts. *Environ Health Perspect* 2006;114:1348–53.
- Wold S, Sjöström M, Eriksson L. PLS-regression: a basic tool of chemometrics. *Chemom Intell Lab* 2001;58:109–30.
- Hastie TJ, Tibshirani R, Friedman J. *The elements of statistical learning: data mining, inference, and prediction.* 2nd ed. New York, NY: Springer, 2009.
- Xu Q, Liang Y, Du Y. Monte Carlo cross-validation for selecting a model and estimating the prediction error in multivariate calibration. *J Chemom* 2004;18:112–20.
- Lubin JH, Colt JS, Camann D, *et al.* Epidemiologic evaluation of measurement data in the presence of detection limits. *Environ Health Perspect* 2004;112:1691–6.
- Chadeau-Hyam M, Campanella G, Jombart T, *et al.* Deciphering the complex: methodological overview of statistical models to derive OMICS-based biomarkers. *Environ Mol Mutagen* 2013;54:542–57.
- Van den Berg RA, Hoefsloot HCJ, Westerhuis JA, *et al.* Centering, scaling, and transformations: improving the biological information content of metabolomics data. *BMC Genomics* 2006;7:142.
- Benjamini Y, Hochberg Y. Controlling the false discovery rate: a practical and powerful approach to multiple testing. *J R Stat Soc Ser B* 1995;57:289–300.
- Chung D, Chun H, Keles S. spls: Sparse Partial Least Squares (SPLS) Regression and Classification. R Packag version 22-1. 2013. <http://cran.r-project.org/package=spls> (accessed Jan 2014).
- Albert O, Jégou B. A critical assessment of the endocrine susceptibility of the human testis to phthalates from fetal life to adulthood. *Hum Reprod Update* 2014;20:231–49.
- Desdoits-Lethimonier C, Albert O, Le Bizec B, *et al.* Human testis steroidogenesis is inhibited by phthalates. *Hum Reprod* 2012;27:1451–9.
- Pan G, Hanaoka T, Yoshimura M, *et al.* Decreased serum free testosterone in workers exposed to high levels of di-n-butyl phthalate (DBP) and di-2-ethylhexyl phthalate (DEHP): a cross-sectional study in China. *Environ Health Perspect* 2006;114:1643–8.
- Mendiola J, Meeker JD, Jørgensen N, *et al.* Urinary concentrations of di (2-ethylhexyl) phthalate metabolites and serum reproductive hormones: pooled analysis of fertile and infertile men. *J Androl* 2012;33:488–98.
- Jönsson BAG, Richthoff J, Rylander L, *et al.* Urinary phthalate metabolites and biomarkers of reproductive function in young men. *Epidemiology* 2005;16:487–93.
- Joensen UN, Frederiksen H, Jensen MB, *et al.* Phthalate excretion pattern and testicular function: a study of 881 healthy Danish men. *Environ Health Perspect* 2012;120:1397–403.
- Meeker JD, Rossano MG, Protas B, *et al.* Environmental exposure to metals and male reproductive hormones: circulating testosterone is inversely associated with blood molybdenum. *Fertil Steril* 2010;93:130–40.
- Menke A, Guallar E, Shiels MS, *et al.* The association of urinary cadmium with sex steroid hormone concentrations in a general population sample of US adult men. *BMC Public Health* 2008;8:72.
- Zeng X, Jin T, Zhou Y, *et al.* Changes of serum sex hormone levels and MT mRNA expression in rats orally exposed to cadmium. *Toxicology* 2003;186:109–18.
- Zhou D, Wang H, Zhang J. Di-n-butyl phthalate (DBP) exposure induces oxidative stress in epididymis of adult rats. *Toxicol Ind Health* 2011;27:65–71.
- Evenson DP, Larson KL, Jost LK. Sperm chromatin structure assay: its clinical use for detecting sperm DNA fragmentation in male infertility and comparisons with other techniques. *J Androl* 2002;23:25–43.
- Kasahara E, Sato EF, Miyoshi M, *et al.* Role of oxidative stress in germ cell apoptosis induced by di(2-ethylhexyl)phthalate. *Biochem J* 2002;365:849–56.
- Hauser R, Meeker JD, Singh NP, *et al.* DNA damage in human sperm is related to urinary levels of phthalate monoester and oxidative metabolites. *Hum Reprod* 2007;22:688–95.
- Specht IO, Toft G, Hougaard KS, *et al.* Associations between serum phthalates and biomarkers of reproductive function in 589 adult men. *Environ Int* 2014;66:146–56.
- Mocevic E, Specht IO, Marott JL, *et al.* Environmental mercury exposure, semen quality and reproductive hormones in Greenlandic Inuit and European men: a cross-sectional study. *Asian J Androl* 2013;15:97–104.
- Toft G, Jönsson BAG, Lindh CH, *et al.* Exposure to perfluorinated compounds and human semen quality in arctic and European populations. *Hum Reprod* 2012;27:2532–40.
- Richthoff J, Rylander L, Jönsson BAG, *et al.* Serum levels of 2,2',4,4',5,5'-hexachlorobiphenyl (CB-153) in relation to markers of reproductive function in young males from the general Swedish population. *Environ Health Perspect* 2003;111:409–13.
- Hauser R, Chen Z, Pothier L, *et al.* The relationship between human semen parameters and environmental exposure to polychlorinated biphenyls and p,p'-DDE. *Environ Health Perspect* 2003;111:1505–11.



- 44 Jørgensen N, Liu F, Andersson A-M, *et al.* Serum inhibin-b in fertile men is strongly correlated with low but not high sperm counts: a coordinated study of 1,797 European and US men. *Fertil Steril* 2010;94:2128–34.
- 45 Safarinejad MR. Effect of omega-3 polyunsaturated fatty acid supplementation on semen profile and enzymatic anti-oxidant capacity of seminal plasma in infertile men with idiopathic oligoasthenoteratospermia: a double-blind, placebo-controlled, randomised study. *Andrologia* 2011;43:38–47.
- 46 Ferguson KK, Hauser R, Altshul L, *et al.* Serum concentrations of p, p'-DDE, HCB, PCBs and reproductive hormones among men of reproductive age. *Reprod Toxicol* 2012;34:429–35.
- 47 Wild CP. Complementing the genome with an "exposome": the outstanding challenge of environmental exposure measurement in molecular epidemiology. *Cancer Epidemiol Biomarkers Prev* 2005;14:1847–50.
- 48 Patel CJ, Bhattacharya J, Butte AJ. An Environment-Wide Association Study (EWAS) on type 2 diabetes mellitus. *PLoS ONE* 2010;5:e10746.
- 49 Holmes E, Loo RL, Stamler J, *et al.* Human metabolic phenotype diversity and its association with diet and blood pressure. *Nature* 2008;453:396–400.
- 50 Xing G, Lin C-Y, Xing C. A comparison of approaches to control for confounding factors by regression models. *Hum Hered* 2011;72:194–205.



## Phthalates, perfluoroalkyl acids, metals and organochlorines and reproductive function: a multipollutant assessment in Greenlandic, Polish and Ukrainian men

Virissa Lenters, Lützen Portengen, Lidwien A M Smit, et al.

*Occup Environ Med* published online September 10, 2014  
doi: 10.1136/oemed-2014-102264

---

Updated information and services can be found at:  
<http://oem.bmj.com/content/early/2014/09/10/oemed-2014-102264.full.html>

---

	<i>These include:</i>
<b>Data Supplement</b>	"Supplementary Data" <a href="http://oem.bmj.com/content/suppl/2014/09/10/oemed-2014-102264.DC1.html">http://oem.bmj.com/content/suppl/2014/09/10/oemed-2014-102264.DC1.html</a>
<b>References</b>	This article cites 45 articles, 9 of which can be accessed free at: <a href="http://oem.bmj.com/content/early/2014/09/10/oemed-2014-102264.full.html#ref-list-1">http://oem.bmj.com/content/early/2014/09/10/oemed-2014-102264.full.html#ref-list-1</a>
<b>P&lt;P</b>	Published online September 10, 2014 in advance of the print journal.
<b>Email alerting service</b>	Receive free email alerts when new articles cite this article. Sign up in the box at the top right corner of the online article.

---

### Notes

---

Advance online articles have been peer reviewed, accepted for publication, edited and typeset, but have not yet appeared in the paper journal. Advance online articles are citable and establish publication priority; they are indexed by PubMed from initial publication. Citations to Advance online articles must include the digital object identifier (DOIs) and date of initial publication.

---

To request permissions go to:  
<http://group.bmj.com/group/rights-licensing/permissions>

To order reprints go to:  
<http://journals.bmj.com/cgi/reprintform>

To subscribe to BMJ go to:  
<http://group.bmj.com/subscribe/>