

# Hyper-resolution global hydrological modelling: what is next? “Everywhere and locally relevant”

Marc F. P. Bierkens,<sup>1,17\*</sup>

Victoria A. Bell,<sup>2</sup>

Peter Burek,<sup>3</sup>

Nathaniel Chaney,<sup>4</sup>

Laura E. Condon,<sup>5</sup>

Cédric H. David,<sup>6,7</sup>

Ad de Roo,<sup>3</sup> Petra Döll,<sup>8</sup>

Niels Drost,<sup>9</sup>

James S. Famiglietti,<sup>6,7</sup>

Martina Flörke,<sup>10</sup>

David J. Gochis,<sup>11</sup>

Paul Houser,<sup>12</sup>

Rolf Hut,<sup>13</sup>

Jessica Keune,<sup>14</sup>

Stefan Kollet,<sup>14,18</sup>

Reed M. Maxwell,<sup>5</sup>

John T. Reager,<sup>6,7</sup>

Luis Samaniego,<sup>15</sup>

Edward Sudicky,<sup>16</sup>

Edwin H. Sutanudjaja,<sup>1</sup>

Nick van de Giesen,<sup>13</sup>

Hessel Winsemius<sup>13,17</sup>

and Eric F. Wood<sup>4</sup>

<sup>1</sup> Utrecht University, Utrecht, The Netherlands

<sup>2</sup> Centre for Ecology and Hydrology, Wallingford, UK

<sup>3</sup> Joint Research Centre, Ispra, Italy

<sup>4</sup> Princeton University, Princeton, NJ, USA

<sup>5</sup> Integrated Ground Water Modeling Center, Colorado School of Mines, Golden, CO, USA

<sup>6</sup> University of California Center for Hydrologic Modeling, Irvine, CA, USA

<sup>7</sup> Jet Propulsion Laboratory, California Institute of Technology, Pasadena, CA, USA

<sup>8</sup> Goethe University of Frankfurt, Frankfurt, Germany

<sup>9</sup> Netherlands eScience Center, Amsterdam, The Netherlands

<sup>10</sup> University of Kassel, Kassel, Germany

<sup>11</sup> NCAR HR Regional Modelling, Boulder, CO, USA

<sup>12</sup> George Mason University, Fairfax, VA, USA

<sup>13</sup> Delft University of Technology, Delft, The Netherlands

<sup>14</sup> Agrosphere (IBG-3), Forschungszentrum Jülich GmbH, Jülich, Germany

<sup>15</sup> UFZ Helmholtz Centre for Environmental Research, Leipzig, Germany

<sup>16</sup> University of Waterloo, Waterloo, Canada

<sup>17</sup> Deltares, Delft, The Netherlands

<sup>18</sup> Centre for High-Performance Scientific Computing in Terrestrial Systems, Geoverbund ABC/J, Germany

\*Correspondence to:

Marc F. P. Bierkens, Utrecht University, Utrecht, The Netherlands.

E-mail: m.f.p.bierkens@uu.nl

## Background

Between 15 and 17 March 2010, a workshop was held at Princeton University entitled ‘Meeting a Grand Challenge to Hydrology: The Global Monitoring of Earth’s Terrestrial Water’. The goal of this workshop was to assess the need for developing hyper-resolution (0.1–1 km) global hydrology and land surface models and to make an inventory on what obstacles need to be overcome to make hyper-resolution models a reality. The primary output from this workshop was a position paper formulating a number of science questions that would benefit from hyper-resolution modelling and key challenges to overcome to make this possible (see Wood *et al.*, 2011).

Since the Princeton workshop and the paper, several groups have been working on making high-resolution ‘Locally Relevant Hydrological Models Everywhere’ a reality. For instance, WaterGAP (Döll *et al.*, 2003) now runs at 5 min globally (Flörke *et al.*, 2013) as does PRC-GLOBWB (Van Beek *et al.*, 2011), whereas LISFLOOD (De Roo *et al.*, 2000; Van Der Knijff *et al.*, 2010) runs at 6 min globally. Using a modular python-based framework, NOAA-MP (Niu *et al.*, 2011) is being coupled to Dynamic TOPMODEL (Beven and Freer, 2001) for 30-m continental simulations. At the same time, the Land Information System software has been developed to support high-performance land surface modelling and data assimilation with horizontal resolutions down to 1 km (Peters-Lidard *et al.*, 2007). Parallel to these efforts is the community of models configured to be as physically based as possible (e.g. three-dimensional variably saturated coupled to shallow water equations for surface runoff) using high-performance computing technology to scale up from catchments to basins to continents (Kollet and Maxwell, 2006; Camporese *et al.*, 2010;

Received 29 October 2014

Accepted 31 October 2014

Brunner and Simmons, 2012; Maxwell, 2013). With global hydrological models ever increasing their resolution and physically based catchment models their domains, the two approaches are bound to meet in the middle in the near future. Aside from hydrology, but relevant to the present initiative, are ongoing efforts to improve the hydrology in land surface models (Milly and Shmakin, 2002; Balsamo *et al.*, 2009; Milly *et al.*, 2014) and dynamic vegetation models (Rost *et al.*, 2008; Biemans *et al.*, 2011), some of which act as land components in earth system models (Oleson *et al.*, 2004; Best *et al.*, 2011; Guimberteau *et al.*, 2012). Table I provides a (non-exhaustive) overview of the various large-scale models from the different communities and their properties.

Undoubtedly, all these efforts face similar problems: which processes are best explicitly modelled and which are better parameterized? How to cope with the computer processing costs that increase exponentially with resolution; not only from finer resolution grids but also because many processes previously parameterized at the lower resolution now have to be addressed in a spatially explicit manner? How to best utilize increasingly powerful computing resources including parallel computers? How to obtain and process the data required to feed the huge parameterization needs of these models? How, if at all, to calibrate such models, validate their simulation results and perform uncertainty analyses on them?

Because many of the groups are at the moment trying to find solutions to these problems independently, we organized a follow-up workshop in Utrecht to exchange experiences and learn from each other. The main outcome of the workshop was the foundation of the open HyperHydro network. Here, we present the objectives of and the rationale behind this initiative, the scientific questions that it will attempt to answer and the major obstacles we need to work on.

## The Objective, the Rationale and the Science

The HyperHydro initiative is an open network of scientists with the objective of simulating large-scale terrestrial hydrology and water resources at hyper-resolution with acceptable accuracy. By using the term ‘open network’, we intend that scientists are free to join and co-operate and that participation in its activities is voluntary: Participants use their own time and resources to participate in the network’s activities. There are a number of reasons that motivate this initiative:

### Science

There are science questions related to global environmental change that require hyper-resolution descriptions of terrestrial water resources (see also Wood *et al.*, 2011). We will mention four of these: (1) In

global biogeochemical cycling, a proper description of the fate and export of, e.g. N, P, C, and Si requires a very detailed representation of streams, ponds, lakes, reservoirs, floodplains and wetlands because much of the retention on land can be concentrated in the smallest of water bodies (Bouwman *et al.*, 2013). (2) Related to this, reconciling rates of soil erosion to observed riverine sediment export rates is impossible at the present time because of an insufficient description of along-river sediment transport, which in turn is largely related to not resolving important small scale processes such as sediment trapping, river bank erosion and bedload transport (Walling, 2009; Wisser *et al.*, 2013); (3) Recently, several groups have been working on the assessment of global flood hazard and risk (Pappenberger *et al.*, 2012; Hirabayashi *et al.*, 2013; Winsemius *et al.*, 2013). These approaches thus far only consider the probability of flooding based on simplified hydrodynamics or downscaled volumes of overbank discharge. However, flood risk also needs flow velocities, time to flooding and flood duration times to convert hazard into risk. This in turn requires very detailed (<100 m) resolution hydrodynamic modelling including the effects of flood defence structures. (4) The non-uniqueness of parameters and process descriptions in hydrological models (Beven and Binley, 1992) and the fact that different processes are active at different time scales in different catchments making each catchment unique (Beven, 2000), calls for generally applicable hydrological models with better process understanding that provide acceptable predictions in ungauged catchments (PUB, Hrachowitz *et al.*, 2014). Many of the approaches suggested in PUB, such as regionalization methods and similarity frameworks, and its successor Panta Rhei (Montanari *et al.*, 2013), e.g. optimality and catchment co-evolution, would be greatly helped if we had ‘hydrological and water resources models of everywhere’ at a locally relevant resolution. These models and their output could then be explored for scale dependent first order controls on runoff, evaporation, groundwater recharge and others.

### Opportunity

Not always popular with hydrologists but no less an important reason to pursue hyper-resolution models is the phrase ‘Because we can!’. On the one hand, there have been enormous advances in global data availability, such as (a) low-resolution hydrologic states from remote sensing: soil moisture (de Jeu *et al.*, 2008), evaporation (Sheffield *et al.*, 2006; Miralles *et al.*, 2011), total terrestrial water storage (Swenson *et al.*, 2006; Famiglietti and Rodell, 2013), lake and reservoir levels (Swenson and Wahr, 2009; Gao *et al.*, 2012); (b) high-

Table I. A (non-exhaustive) overview of the various large-scale models from the different communities and their properties

Model name	Class of model (six types) <sup>a</sup> :	Domain	Resolution (space and time)	Discretization type (grid, polygon or sub-catchment)	Soil and groundwater dynamics	Routing	Human water use/Reservoirs	Surface energy balance	Vegetation	Carbon cycling	reference	Institutes	url
Mac-PDM	GHM	Globe	0.5°; daily	Grid	Vertical soil	Basin aggregation of runoff	No/No	No	Fixed	No	Gosling and Arnell (2011)	University of Nottingham (UK)	
WBMPPlus	GHM	Globe	0.5°; daily	Grid	Vertical soil, groundwater reservoir	1 D Muskingum–Cunge over channel network	Yes/Yes	No	Fixed, climatology of phenology, irrigated area change	No	Wisser <i>et al.</i> (2010)	University of New Hampshire (USA) City University of New York (USA)	www.wsag.unh.edu/wbm.html
WaterGAP	GHM	Globe	0.5°; 5' daily	Grid	Vertical soil, groundwater reservoir	1D Kinematic wave + Manning channel network	Yes/Yes	No	Fixed, climatology of phenology, irrigated area change	No	Döll <i>et al.</i> (2003), Müller Schmied <i>et al.</i> (2014)	Kassel University (Germany) Goethe University Frankfurt (Germany)	www.watergap.de
H08	GHM	Globe	1.0°; daily (3 h for energy balance)	Grid	Vertical soil	Accumulation along 1.0° channel network	Yes/Yes	Yes	Simple crop growth model	No	Hamasaki <i>et al.</i> (2008)	Institute for Environmental Studies (Japan)	http://h08.nies.go.jp/h08/index.html
PCR-GLOBWB	GHM	Globe	0.5°; 5' daily	Grid	Vertical soil, groundwater reservoir or lateral groundwater (optional)	Kinematic wave + Manning channel network, approximate floodplain inundation	Yes/Yes	No	Fixed, climatology of phenology, irrigated area change	No	Van Beek <i>et al.</i> (2011), Wada <i>et al.</i> (2014)	Utrecht University (Netherlands) Delfares (Netherlands)	www.globalhydrology.nl
VIC	LSM/GHM	Globe	0.5°; daily (3 h for energy balance)	Grid	Vertical soil, groundwater-like reservoir	Unit Hydrograph per cell + linearized St. Venant over channel network.	No/No	Yes	Fixed, climatology of phenology	No	Gao <i>et al.</i> (2009)	Princeton University (USA) University of Washington (USA)	http://www.hydro.washington.edu/Lettenmaier/Models/VIC/index.shtml
MATSURU	LSM/GHM	Globe	1.0°; daily (6 h for energy balance)	Grid	Vertical soil, groundwater reservoir	CAMA Flood: 1D Diffusive wave + Manning for	Yes/Yes	Yes	Simple crop growth model	No	Koirala <i>et al.</i> (2014)	IIS, University of Tokyo (Japan)	

M. F. P. BIERKENS *ET AL.*

INVITED COMMENTARY

Model	Model Type	Resolution	Time Step	Grid	Soil/Reservoir	Channel/Network	Routing	Phenology	Optional	Author	Institution	Website		
LaD	LSM	Globe	1.0°; daily	Grid	Vertical soil, groundwater reservoir	friction slope channel network, floodplain inundation.	Basin aggregation of runoff	No/No	Yes	Constant	No	Milly and Shumakin (2002)	NOAH Geophysical Fluids Dynamics Laboratory (USA)	<a href="http://www.gfdl.noaa.gov/land-model">http://www.gfdl.noaa.gov/land-model</a>
LM3 (part of ESM2)	LESM	Globe	daily (sub-daily for energy balance) 1.0°; daily (sub-daily for energy balance)	Grid	Vertical soil, groundwater reservoir and topography-based groundwater dynamics	Non-linear storage – outflow relationship per cell along the channel network	No/No	Yes	Dynamic	Yes	Milly <i>et al.</i> (2014)			
HTESSEL	LSM/LESM	Globe	Variable, 0.25–1.0° globally; 3 hourly time steps	Grid	Vertical soil	Basin aggregation of runoff	No/No	Yes	Constant	Optional: coupling with LPJG-euss is possible	Balsamo <i>et al.</i> (2009)	European Centre for Medium Range Weather Forecasting (UK)		
LPJml	DVM	Globe	0.5°; daily	Grid	Vertical soil	Linear reservoir per cell along the channel network	Yes/Yes*	No	Dynamic + phenology + plant physiology	Yes	Rost <i>et al.</i> (2008), Biemans <i>et al.</i> (2011)	PIK- Potsdam Institute for Climate Research (Germany)	<a href="http://www.pik-potsdam.de/research/projects/activities/biosphere-water-modelling/lpjml">www.pik-potsdam.de/research/projects/activities/biosphere-water-modelling/lpjml</a>	
JULES	LSM/LESM	Globe	0.5°; 3 hourly time steps	Grid	Vertical soil	Routing over 1° grid using an advection scheme.	Yes (irrigation only)/Yes	Yes	Dynamic + phenology + plant physiology	Yes	Best <i>et al.</i> (2011)	Centre for Ecology and Hydrology (UK)	<a href="https://jules.jchmr.org">https://jules.jchmr.org</a>	
ORCHIDEE	LSM/LESM	Globe	0.5°; half hourly time steps	Grid	Vertical soil	No built-in routing; but has been used with routing schemes.	Yes (irrigation only)/No	Yes	Dynamic + phenology + plant physiology	Yes	Guimberteau <i>et al.</i> (2012)	Institut Pierre Simon Laplace (France)	<a href="http://labex.ipsl.fr/orchidee/">http://labex.ipsl.fr/orchidee/</a>	
CLM	LESM	Globe	Variable, standard 0.9° by 1.25° LAT-LON; 6 hourly	Grid	Vertical soil, groundwater reservoir.	Linear reservoir per cell along the channel network	No/No	Yes	Dynamic + phenology + plant physiology	Yes	Lawrence <i>et al.</i> (2011).	National Center for Atmospheric Research (USA)	<a href="http://www.cesm.ucar.edu/models/cm/">http://www.cesm.ucar.edu/models/cm/</a>	

(Continues)

Table I. (Continued)

Model name	Class of model (six types) <sup>a</sup> :	Domain	Resolution (space and time)	Discretization type (grid, polygon or sub-catchment)	Soil and groundwater dynamics	Routing	Human water use/Reservoirs	Surface energy balance	Vegetation	Carbon cycling	reference	Institutes	url
WRF-Hydro	LES/DVM	Variable	Variable, (sub-) hourly time steps	Finite difference	Vertical soil, groundwater reservoir or lateral groundwater (optional)	2D-diffusion wave over topography	No/Yes	Yes	Dynamic phenology depending on LSM selection	No	Gochis <i>et al.</i> (2013)	National Center for Atmospheric Research (USA)	http://www.ral.ucar.edu/projects/wrf_hydro/
Lisflood	RHM	Europe, Africa	Europe (5 km), Global (0.1 degrees)	grid	Vertical soil, groundwater reservoir	Kinematic wave + Manning over channel network	Yes/Yes	No	LAI observed / LAI climatology for scenarios	No	Van der Knijff <i>et al.</i> (2010)	EC-JRC at Ispra (Italy)	http://floods.jrc.ec.europa.eu/lisflood-model.html
E-Hype	RHM	Europe	Median catchment size 215 km <sup>2</sup> (15 × 15 km); daily time step	Sub-catchments	Vertical soil, groundwater reservoir	Variable velocity (translation) + linear reservoir (attenuation) along a channel network	Yes/no	No	Partly Fixed, partly with seasonal climatology of crop cover.	No	Lindström <i>et al.</i> (2010)	SMHI – Swedish Meteorological and Hydrological Institute (Sweden)	http://hype.sourceforge.net/
mHM	RHM	Europe	1-24 km (Europe) 1/8° (USA) 1° (Africa, Asia, Lat-Ame), hourly-daily	Grid	Vertical soil, Groundwater reservoir.	ID Muskingum–Cunge over channel network.	No/No	No	LAI-observed. Dynamic phenology (option)	No	Samaniego <i>et al.</i> (2010), Kumar <i>et al.</i> (2013)	Helmholtz Centre for Environmental Research – UFZ	http://www.ufz.de/index.php?en=31389
RAPID	RHM (fast routing module)	Basins	Median catchment size 3 km <sup>2</sup> ; 3-hourly time steps (outputs, routing at 15 min)	Sub-catchments	No	ID Muskingum–Cunge over channel network	No/No	No	No	No	David <i>et al.</i> (2011)	University of Texas at Austin	http://rapid-hub.org.

(Continues)

Table I. (Continued)

Model name	Class of model (six types) <sup>a</sup> : Domain	Resolution (space and time)	Discretization type (grid, polygon or sub-catchment)	Soil and groundwater dynamics	Routing	Human water use/Reservoirs	Surface energy balance	Vegetation	Carbon cycling	Institutes	url
HydroGeoSphere	FL-PDE Sub-watershed to Basin to Continental (3D)	Dynamically variable, fully implicit	Control-Volume Finite Element, unstructured mesh or structured finite difference	3D-Richards Equation	2D-diffusion wave over topography	Yes/Yes	Yes	Variable	No	University of Waterloo (Canada) (2012); Hwang <i>et al.</i> (2014) (Canada) Aquanty (Canada)	www.aquanty.com
ParFlow	FL-PDE US, Basins Euro-Cordex	Variable, (sub-) hourly time steps	Finite difference	3D-Richards Equation	2D-kinematic wave over topography	No/No	Yes, when coupled to Community Land Model	Fixed	No	University of Lawrence Livermore National lab (USA) Shrestha <i>et al.</i> (2014), Colorado School of Mines (USA) Centre for High-Performance Scientific Computing in Terrestrial Systems, Geoverbund ABC/J	http://computation.llnl.gov/casc/parflow/parflow_home.html
CATHY	FL-PDE Basins	Variable, (sub-) hourly time steps	Finite elements (subsurface flow), finite difference (surface water flow)	3D-Richards Equation	2D-diffusion wave over topography	No/No	No	Fixed	No	Campopese <i>et al.</i> (2010) University of Padua (Italy) University of Quebec (Canada)	

<sup>a</sup> GHM, global hydrological model (conceptual bucket type; often human impacts); RHM, regional hydrological model (conceptual bucket type; often human impacts); LSM, land surface model (used in climate or numerical weather prediction models; energy balance); DVM, dynamic vegetation model (water, vegetation, carbon); LESM, Land models used in earth system models (water, energy, carbon); FI-PDE, fully implicit partial differential equations



resolution parameter fields: global land cover at 1 km (GLCC), global soils and soil properties at 1 km (<http://soilgrids.org>), global surface lithology at 1 km (Hartmann and Moosdorf, 2012), surface elevation and global river networks at 15 arc-second (500 m) (Lehner *et al.*, 2008) and 3 arc-second (100 m). (c) Global online databases of field observations: discharge (GRDC, MOPEX, EWA-FRIEND), evaporation and carbon fluxes (FLUXNET), soil moisture (ISMN (Dorigo *et al.*, 2011)); (d) low-resolution forcing datasets of meteorological forcing (WFD (Weedon *et al.*, 2011); CRU TS3.21 (Harris *et al.*, 2014); ERA-INTERIM (Dee *et al.*, 2011), MERRA (Rienecker *et al.*, 2011)). On the other hand, advances in computational capabilities and data storage have exceeded petaflop and petabyte capabilities, making it possible to model large domains at very high resolution. At the same time, new solvers and simulation platforms are being developed that take advantage of these new computational possibilities and can efficiently handle very large problems. The earth system modelling and climate modelling community will certainly grasp the opportunity. It is imperative that hydrologists are central to this effort in order to safeguard the sound incorporation of hydrologic principles at hyper-resolutions.

### *Society*

Most developed countries have dedicated hydrological models to support water resources management and planning under conditions of environmental change, or for operational forecasting and early warning. However, in many developing countries, such models are not yet available or are poorly constrained because of the lack of local data. For these developing countries, access to information about water resources that is timely and locally relevant can be a great asset (GEOSS, 2009). The issue here is an ethical argument on the equity of information (Lievrouw and Farb, 2003). One could discuss whether detailed information on water resources status is really of use to local water managers, or the local population in general, given that one needs to be a specialist to be able to interpret these data and because it arises from technological push rather than stakeholder pull. However, we believe that adoption of new technology is always likely to be slow unless it is made freely available and presented in the right manner. Similarly, the potential future applications of new technology are often unpredictable, and as a result, the development of new capabilities to model the global water cycle may produce additional, currently unrecognized benefits. In any case, river basin managers often consider 1–5 km resolution models more useful than 0.5° models. Obviously, how and in what form to make this information available needs further thought.

## **Obstacles and Challenges**

The goal to provide locally relevant hydrological information globally is far from trivial and poses a series of significant challenges (see also Wood *et al.*, 2011). To name a few:

### *Scale separation and scale-related breakdown of concepts and assumptions*

When moving from the common resolution of 0.5° (order of 50 km at the equator) down to 1 km and smaller, many concepts that have been designed to resolve small-scale processes at the sub-grid scale break down. One could even talk about the ‘loss of the sub-grid paradigm’. First, simple cell fractionation has to be replaced by explicit dynamics. For instance, surface runoff at 0.5° can be conceptualized as the fraction of saturated soil assuming a univariate spatial frequency distribution of sub-grid surface elevation or associated with a distribution of soil storage within a cell (Moore, 1985; Blyth *et al.*, 2004). However, when moving to higher resolutions, the explicit spatial juxtaposing of saturated soils with time has to be accounted for, e.g. by using concepts related to the topographic index (Beven and Freer, 2001). Second, very much related to the reasons above, is the breaking up of connections between compartments of the hydrological system. In 0.5° models, water stress assessments are based on the assumption that water demand is satisfied by available surface water and groundwater within the same grid cell. This assumption works well because most regional water redistribution works fit within a 50 × 50 km area around the location where water is consumed. However, at resolutions of 10 km and finer, inter-cell redistribution of water from abstraction points to hotspots of water consumption, or from lateral groundwater flow, need to be taken into account (Krakauer *et al.*, 2014). Third, assumptions about dominant physics break down. For example, consider the way we model heat and water fluxes between the land and atmosphere. It is generally assumed that sensible and latent heat fluxes are proportional to the vertical gradient of potential temperature and specific humidity. This only holds true if vertical gradients are much larger than horizontal gradients (Holtslag and Ek, 2008), for example, when using average fluxes over large horizontal domains. However, for spatial resolutions approaching 100 m and below, horizontal advection becomes important, and new theories to correctly model land-atmosphere heat and moisture exchange are needed. Finally, and this is indeed a great challenge, in an ideal world we would have sub-grid concepts and parameterizations that change consistently with scale as described in, e.g. Blöschl and Sivapalan (1995) and Bierkens *et al.* (2000).

### *Lack of information and epistemic uncertainty*

In a comment on the paper of Wood *et al.* (2011), Beven and Cloke (2012) rightly point out that moving to higher resolutions will pose huge challenge as epistemic uncertainties will become very large because of lack of process and parameter knowledge at such high resolutions. Indeed, there is much we do not know about the global land surface properties at these resolutions that need to be added. For instance, relating local water demand and water abstraction at high resolution requires knowledge of local water redistribution systems that is not available globally. Although the number of high-resolution datasets on topography, land use, geology, soil properties and others, is increasing, these data are also the products of remote sensing, land surface models, statistical downscaling techniques or combinations thereof and thus full of inaccuracies. Finally, several studies (e.g. Döll and Fiedler, 2008; Biemans *et al.*, 2009) have shown that precipitation that is used to force global hydrological models contributes to a large part of the uncertainty in model output. High-resolution meteorological datasets of sufficient accuracy are thus greatly needed. This does not mean that current lack of information and related uncertainty should deter us from attempting hyper-resolution modelling. First, the number and quality of high-resolution datasets are increasing, and they are expected to increase even further as new missions with better sensors are launched. Second, our globalized digital world makes local datasets universally accessible, enabling better ground-truthing of these datasets. Third, because many of the envisioned hyper-resolution models will have global coverage, multi-model predictions of parameters and variables are available for all locations in the world. A comparison of multiple model results and their fit to observations will reveal pathways for improvement of both process description and their parameterizations. Moreover, multi-model ensemble simulations would provide an approximation of predictive uncertainty, arising from both model structure and forcing datasets.

### *Computational challenges*

Obviously, hyper-resolution global modelling results in large computational demands in terms of CPU time and storage requirements (Kollet *et al.*, 2010; Maxwell, 2013). Just increasing the spatial resolution by a factor of 10 increases calculation and storage requirements by a factor of 100. Moreover, as stated before, when increasing resolution, many processes that use simple sub-grid parameterizations may need to be replaced by explicit process dynamics further increasing computational efforts. Finally, running hyper resolution models real-time in an operational setting, and incorporating

incoming observations using data-assimilation schemes, will further increase the computational demand. Fortunately rapid developments in both hardware and software architecture support these increased computational challenges. Powerful PCs with multi-core and multi-thread CPUs or small linux clusters are commonplace with many hydrology groups. Fibre-optic cable networks are present on many universities as well as large quantities of idle desktops making grid computing a possibility. Moreover, massively parallel computing facilities involving many hundreds of thousands of cores can be found in the larger research centres, with the largest having >10 petaflop performance and expecting to reach exaflop ability in 2020. This tremendous computational power can only be exploited if the computations behind a model can be performed in parallel. Here, domain decomposition methods can be used (e.g. using a hierarchical structure based on continents, basins, catchments, sub-catchments, etc.). If the code can be adequately restructured, parallel processing software such as Open MPI allows information to travel between different computers (nodes) and OpenMP to allow further breakup of operations to different cores (processors). Alternatively, smart robust numerical methods need to be developed that solve the equations relying on only a limited number of cores. Note, however, that efficient numerical methods are also needed in the case of massive parallel computing if one requires a good scaling to a large numbers of cores.<sup>1</sup>

### **The Hyperhydro Network ([www.hyperhydro.org](http://www.hyperhydro.org))**

The HyperHydro network is open to the broader scientific community for anyone who wants to cooperate in activities furthering the development of hyper-resolution models. The governance structure is quite informal, with a post box and a website hosted at Utrecht University and three working groups (WG) loosely organized around the sets of challenges laid out above. These groups are:

#### *WGI Test beds*

The objective of this working group is to set up a test bed for comparing different large-scale models at different resolutions. Data-sets will be provided for different domains resolutions: Global at 5 min; CONUS and CORDEX Europe at 1 km; the Rhine-Meuse, Illinois and California at 100 m. Starting with the smallest domains, the WG will run a collection of different models on the same datasets (forcing, soil,

<sup>1</sup>In parallel computing, 'good scaling' means that the computation times keep decreasing significantly when more processors are used.



geology, etc., output data) for decreasing resolution. This enables the assessment of model uncertainty (by model inter comparison) and model bias (by model-data comparison) but also the effects of resolution on model uncertainty and bias. The WG will set up the data and simulation protocols for participating groups to perform their analyses. The ultimate goal is to improve model conceptualizations and parameterizations and make these robust under change of model resolution.

### WG2 Common framework

Members of this group together work on computational challenges, including parallel computing and model component coupling. This will take the form of collecting information on and developing tools for solving computational issues (e.g. parallelization), developing a common framework for coupling modules based on existing couplers and standards and setting up a common platform for generating comparison statistics (as related to the work by WG1).

### WG3 Parameter and concepts

The goal of WG3 is to develop concepts and possible solutions on how to deliver the information needed to achieve hyper-resolution (<1 km) globally. Their work will consist of compiling an overview of useful high-resolution global datasets that can be used in parameterizing global hyper-resolution models, creating new high-resolution datasets using auxiliary information, devising new model concepts to replace sub-grid parameterizations with computationally frugal, spatially explicit physics and to create new hyper-resolution global forcing data of sufficient accuracy.

HyperHydro will set up regular meetings at international conferences and symposia to discuss progress and set targets. Working group leads will be responsible for its impetus and progress, and we expect that results of working group activities will be reported through peer-reviewed publications. We invite all those who are interested to look on the website [www.hyperhydro.org](http://www.hyperhydro.org) for current activities and past results and urge those who want to be on the mailing list or actively participate in one of the working groups to send an email to the lead author.

## ACKNOWLEDGEMENTS

The contents of this paper reflect the discussions that were held among the authors during and after the Workshop ‘Hyper-resolution global hydrological modelling: the next step’ held on 13–14 February 2014 in Utrecht, the Netherlands. The authors took part in these discussions. Cédric H. David, James S. Famiglietti, and John T. Reager were partially supported by the Jet Propulsion Laboratory, California Institute of Technology, under a

contract with the National Aeronautics and Space Administration. Niels Drost, Rolf Hut, and Edwin Sutanudjaja were supported by the Netherlands eScienceCenter under the project eWaterCycle.

## REFERENCES

- Balsamo G, Viterbo P, Beljaars A, van den Hurk B, Hirschi M, Betts AK, Scipal K. 2009. A revised hydrology for the ECMWF model: verification from field site to terrestrial water storage and impact in the integrated forecast system. *Journal of Hydrometeorology* 10: 623–643.
- Beven KJ, Binley AM. 1992. The future of distributed models: model calibration and uncertainty prediction. *Hydrological Processes* 6: 279–298.
- Beven KJ. 2000. On the future of distributed modelling in hydrology. *Hydrological Processes* 14: 16–17.
- Beven KJ, Freer JE. 2001. A dynamic TOPMODEL. *Journal of Hydrology* 15: 1993–2011.
- Beven KJ, Cloke HL. 2012. Comment on ‘‘Hyperresolution global land surface modeling: Meeting a grand challenge for monitoring Earth’s terrestrial water’’ by Eric F. Wood *et al.* *Water Resources Research* 48: W01801.
- Best MJ, Pryor M, Clark DB, Rooney GG, Essery RLH, M’enanad CB, Edwards JM, Hendry MA, Porson A, Gedney N, Mercado LM, Sitch S, Blyth E, Boucher O, Cox PM, Grimmond CSB, Harding RJ. 2011. The Joint UK Land Environment Simulator (JULES), model description – Part 1: Energy and water fluxes. *Geoscientific Model Development* 4: 677–699.
- Biemans H, Haddeland I, Kabat P, Ludwig F, Hutjes RWA, Heinke J, Von Bloh W, Gerten D. 2011. Impact of reservoirs on river discharge and irrigation water supply during the 20th century. *Water Resources Research* 47: W03509.
- Biemans H, Hutjes RWA, Kabat P, Gerten D, Rost S. 2009. Effects of precipitation uncertainty on discharge calculations for main river basins. *Journal of Hydrometeorology* 10: 1011–1025.
- Bierkens MFP, Finke PA, de Willigen P. 2000. *Upscaling and Downscaling Methods for Environmental Research*. Kluwer Academic Publishers: Dordrecht; 190.
- Blyth EM, Finch J, Robinson M, Rosier P. 2004. Can soil moisture be mapped onto the terrain? *Hydrology and Earth System Science* 8: 923–930.
- Blöschl G, Sivapalan M. 1995. Scale issues in hydrological modelling: a review. *Hydrological Processes* 9: 251–290.
- Bouwman AF, Bierkens MFP, Griffioen J, Hefting MM, Middelburg JJ, Middelkoop H, Slomp CP. 2013. Nutrient dynamics, transfer and retention along the aquatic continuum from land to ocean: towards integration of ecological and biogeochemical models. *Biogeosciences* 10: 1–22.
- Brunner P, Simmons CT. 2012. HydroGeoSphere: a fully integrated, physically based hydrological model. *Ground Water* 50: 170–176.
- Camporese M, Paniconi C, Putti M, Orlandini S. 2010. Surface-subsurface flow modeling with path-based runoff routing, boundary condition-based coupling, and assimilation of multisource observation data. *Water Resources Research* 46: W02512.
- David CH, Maidment DR, Niu G-Y, Yang Z-L, Habets F, Eijkhout V. 2011. River network routing on the NHDPlus dataset. *Journal of Hydrometeorology* 12: 913–934.
- Dee DP, Uppala SM, Simmons AJ, Berrisford P, Poli P, Kobayashi S, Andrae U, Balmaseda MA, Balsamo G, Bauer P, Bechtold P, Beljaars ACM, van de Berg L, Bidlot J, Bormann N, Delsol C, Dragani R,

- Fuentes M, Geer AJ, Haimberger L, Healy SB, Hersbach H, Hólm EV, Isaksen L, Kållberg P, Köhler M, Matricardi M, McNally AP, Monge-Sanz BM, Morcrette J-J, Park B-K, Peubey C, de Rosnay P, Tavolato C, Thépaut J-N, Vitart F. 2011. The ERA-Interim reanalysis: configuration and performance of the data assimilation system. *Quarterly Journal of the Royal Meteorological Society* 137: 553–597.
- De Jeu RAM, Wagner W, Holmes TRH, Dolman AJ, van de Giesen NC, Friesen J. 2008. Global soil moisture patterns observed by space borne microwave radiometers and scatterometers. *Surveys in Geophysics* 29: 399–420.
- De Roo APJ, Wesseling CG, Van Deursen WPA. 2000. Physically-based river basin modelling within a GIS: the LISFLOOD model. *Hydrological Processes* 14: 1981–1992.
- Döll P, Kaspar F, Lehner B. 2003. A global hydrological model for deriving water availability indicators: model tuning and validation. *Journal of Hydrology* 270: 105–134.
- Döll P, Fiedler K. 2008. Global-scale modeling of groundwater recharge. *Hydrology and Earth System Science* 12: 863–885.
- Dorigo WA, Wagner W, Hohensinn R, Hahn S, Paulik C, Xaver A, Gruber A, Drusch M, Mecklenburg S, van Oevelen P, Robock A, Jackson T. 2011. The International Soil Moisture Network: a data hosting facility for global in situ soil moisture measurements. *Hydrology and Earth System Sciences* 15: 1675–1698.
- Famiglietti JS, Rodell M. 2013. Water in the balance. *Science* 340: 1300.
- Flörke M, Kynast E, Bärlund I, Eisner S, Wimmer F, Alcamo J. 2013. Domestic and industrial water uses of the past 60 years as a mirror of socio-economic development: a global simulation study. *Global Environmental Change* 23: 144–156.
- Gao H, Tang Q, Shi X, Zhu C, Bohn TJ, Su F, Sheffield J, Pan M, Lettenmaier DP, Wood EF. 2009. Water budget record from variable infiltration capacity (VIC) model. In Algorithm Theoretical Basis Document for Terrestrial Water Cycle Data Records (in review)
- Gao H, Birkett C, Lettenmaier DP. 2012. Global monitoring of large reservoir storage from satellite remote sensing. *Water Resources Research* 48: W09504.
- GEOS. 2009. The Global Earth Observation System of Systems Strategic Targets. Document 12(Rev1) Group on Earth Observations (<https://www.earthobservations.org/documents.php>) [Accessed on August 28 2014].
- Gochis DJ, Yu W, Yates DN. 2013. The WRF-Hydro model technical description and user's guide, version 1.0. NCAR Technical Document. 120 pages. Available online at: [http://www.ral.ucar.edu/projects/wrf\\_hydro/](http://www.ral.ucar.edu/projects/wrf_hydro/) [Accessed on August 28 2014].
- Gosling SN, Arnell NW. 2011. Simulating current global river runoff with a global hydrological model: model revisions, validation, and sensitivity analysis. *Hydrological Processes* 25: 1129–1145.
- Guimberteau M, Drapeau G, Ronchail J, Sultan B, Polcher J, Martinez J-M, Prigent C, Guyot J-L, Cochonneau G, Espinoza JC, Filizola N, Fraizy P, Lavado W, De Oliveira E, Pombosa R, Noriega L, Vauchel P. 2012. Discharge simulation in the sub-basins of the Amazon using ORCHIDEE forced by new datasets. *Hydrology and Earth System Science* 16: 911–935.
- Harris I, Jones PD, Osborn TJ, Lister DH. 2014. Updated high-resolution grids of monthly climatic observations – the CRU TS3.10 Dataset. *International Journal of Climatology* 34: 623–642.
- Hanasaki N, Kanae S, Oki T, Masuda K, Motoya K, Shirakawa N, Shen Y, Tanaka K. 2008. An integrated model for the assessment of global water resources – part 1: model description and input meteorological forcing. *Hydrology and Earth System Science* 12: 1007–1025.
- Hartmann J, Moosdorf N. 2012. The new global lithological map database GLiM: a representation of rock properties at the Earth surface. *Geochemistry, Geophysics, Geosystems* 13: Q12004.
- Holtlag AAM, Ek M. 2008. Physics of evaporation and atmospheric boundary layers over land. In *Climate and the Hydrological Cycle*, Bierkens MFP, Dolman AJ, Troch PA (eds). IAHS Special Publication 8: Wallingford UK; 39–56.
- Hrachowitz M, Savenije HHG, Blöschl G, McDonnell JJ, Sivapalan M, Pomeroy JW, Arheimer B, Blume T, Clark MP, Ehret U, Fenicia F, Freer JE, Gelfan A, Gupta HV, Hughes DA, Hut RW, Montanari A, Pande S, Tetzlaff D, Troch PA, Uhlenbrook S, Wagener T, Winsemius HC, Woods RA, Zehe E, Cudennec C. 2014. A decade of predictions in ungauged basins (PUB)—a review. *Hydrological Sciences Journal* 58: 1198–1255.
- Hwang H-T, Park Y-J, Sudicky EA, Forsyth PA. 2014. A parallel computational framework to solve flow and transport in integrated surface-subsurface hydrologic systems. *Environmental Modelling & Software* 61: 39–58.
- Hirabayashi Y, Roobavannan M, Sujan K, Lisako K, Dai Y, Satoshi W, Hyungjun K, Shinjiro K. 2013. Global flood risk under climate change. *Nature Climate Change* 3: 816–821.
- Koirala S, Yeh P-J, Hirabayashi Y, Kanae S, Oki T. 2014. Global-scale land surface hydrologic modeling with the representation of water table dynamics. *Journal of Geophysical Research, [Atmospheres]* 119: 75–89.
- Kollet SJ, Maxwell RM. 2006. Integrated surface-groundwater flow modeling: a free-surface overland flow boundary condition in a parallel groundwater flow model. *Advances in Water Resources* 29: 945–958.
- Kollet SJ, Maxwell RM, Woodward CS, Smith SG, Vanderborght J, Vereecken H, Simmer C. 2010. Proof-of-concept of regional scale hydrologic simulations at hydrologic resolution utilizing massively parallel computer resources. *Water Resources Research* 46: W04201.
- Krakauer NY, Li LH, Fan Y. 2014. Groundwater flow across spatial scales: importance for climate modeling. *Environmental Research Letters* 9: 034003.
- Kumar R, Livneh B, Samaniego L. 2013. Toward computationally efficient large-scale hydrologic predictions with a multiscale regionalization scheme. *Water Resources Research* 49: 5700–5714.
- Lawrence DM, Oleson KW, Flanner MG, Thornton PE, Swenson SC, Lawrence PJ, Zeng X, Yang Z-L, Levis S, Sakaguchi K, Bonan GB, Slater AG. 2011. Parameterization improvements and functional and structural advances in version 4 of the Community Land Model. *Journal of Advances in Modeling Earth Systems* 3: 27. Art. M03001.
- Lehner B, Verdin K, Jarvis A. 2008. New global hydrography derived from spaceborne elevation data. *Eos, Transactions AGU* 89: 93–94.
- Lievrouw LA, Farb SE. 2003. Information and equity. *Annual Review of Information Science and Technology* 37: 499–540.
- Lindström G, Pers CP, Rosberg R, Strömqvist J, Arheimer B. 2010. Development and test of the HYPE (Hydrological Predictions for the Environment) model – A water quality model for different spatial scales. *Hydrology Research* 41: 295–319.
- Maxwell RM. 2013. A terrain-following grid transform and preconditioner for parallel, large-scale, integrated hydrologic modeling. *Advances in Water Resources* 53: 109–117.
- Milly PC, Shmakin AB. 2002. Global modeling of land water and energy balances. Part I: The land dynamics (LaD) model. *Journal of Hydrometeorology* 3: 283–299.
- Milly PC, Malyshev S, Shevliakova E, Dunne KA, Findell KL, Gleeson T, Liang Z, Phillips P, Stouffer RJ, Swenson SC. 2014. An enhanced model of land water and energy for global hydrologic and earth-system studies. *Journal of Hydrometeorology* 15: 1739–1761.
- Miralles DG, Holmes TRH, De Jeu RAM, Gash JH, Meesters AGCA, Dolman JA. 2011. Global land-surface evaporation estimated from satellite-based observations. *Hydrology and Earth System Science* 15: 453–469.

- Montanari A, Young G, Savenije HHG, Hughes D, Wagener T, Ren LL, Koutsoyiannis D, Cudennec C, Toth E, Grimaldi S, Blöschl G, Sivapalan M, Beven K, Gupta H, Hipsey M, Schaeffli B, Arheimer B, Boegh E, Schymanski SJ, Di Baldassarre G, Yu B, Hubert P, Huang Y, Schumann A, Post D, Srinivasan V, Harman C, Thompson S, Rogger M, Viglione A, McMillan H, Characklis G, Pang Z, Belyaev V. 2013. “Panta Rhei—Everything Flows”: Change in hydrology and society—The IAHS Scientific Decade 2013–2022. *Hydrological Sciences Journal* 58: 1256–1275.
- Moore RJ. 1985. The probability-distributed principle and runoff production at point and basin scales, *Hydrological Sciences Journal* 30: 273–297.
- Müller Schmied H, Eisner S, Franz D, Wattenbach M, Portmann FT, Flörke M, Döll P. 2014. Sensitivity of simulated global-scale freshwater fluxes and storages to input data, hydrological model structure, human water use and calibration. *Hydrology and Earth System Science* 18: 3511–3538.
- Niu G-Y, Tang Z-L, Mitchel KE, Chen F, Ek MB, Barlage M, Kumar A, Manning K, Niyogi D, Rosero E, Tewari M, Xia Y. 2011. The community Noah land surface model with multiparameterization option (Noah-MP): 1. Model description and evaluation with local measurements. *Journal of Geophysical Research* 116: D12109.
- Oleson K, Dai Y, Bonan G, Bosilovich M, Dickinson R, Dirmeyer P, Hoffman F, Houser PR, Levis S, Niu G, Thornton P, Vertenstein M, Yang Z, Zeng X. 2004. Technical Description of the Community Land Model (CLM), NCAR Technical Note, NCAR/TN-461 + STR.
- Pappenberger F, Dutra E, Wetterhall F, Cloke HL. 2012. Deriving global flood hazard maps of fluvial floods through a physical model cascade. *Hydrology and Earth System Science* 16: 4143–4156.
- Peters-Lidard CD, Houser PR, Tian Y, Kumar SV, Geiger J, Olden S, Lighty L, Doty B, Dirmeyer P, Adams J, Mitchell K, Wood EF, Sheffield J. 2007. High performance earth system modeling with NASA/GSFC’s land information system. *Innovations in Systems and Software Engineering* 3: 157–165.
- Rienecker MM, Suarez MJ, Gelaro R, Todling R, Bacmeister J, Liu E, Bosilovich MG, Schubert SD, Takacs L, Kim G-K, Bloom S, Chen J, Collins D, Conaty A, da Silva A, Gu W, Joiner J, Koster RD, Lucchesi R, Molod A, Owens T, Pawson S, Pegion PH, Redder CR, Reichle R, Robertson FR, Ruddick AG, Sienkiewicz M, Woollen J. 2011. MERRA: NASA’s Modern-Era Retrospective Analysis for Research and Applications. *Journal of Climate* 24: 3624–3648.
- Rost S, Gerten D, Bondeau A, Lucht W, Rohwer J, Schaphoff S. 2008. Agricultural green and blue water consumption and its influence on the global water system. *Water Resources Research* 44: W09405.
- Samaniego L, Kumar R, Attinger S. 2010. Multiscale parameter regionalization of a grid-based hydrologic model at the mesoscale. *Water Resources Research* 46: W05523.
- Sheffield J, Goteti G, Wood EF. 2006. Development of a 50-yr high-resolution global dataset of meteorological forcings for land surface modeling. *Journal of Climate* 19: 3088–3111.
- Shrestha P, Sulis M, Masbou M, Kollet S, Simmer C. 2014. A scale-consistent Terrestrial Systems Modeling Platform based on COSMO, CLM and ParFlow. *Monthly Weather Review* 142: 3466–3483.
- Swenson S, Yeh P-J-F, Wahr J, Famiglietti J. 2006. A comparison of terrestrial water storage variations from GRACE with in situ measurements from Illinois. *Geophysical Research Letters* 33: L16401.
- Swenson S, Wahr J. 2009. Monitoring the water balance of Lake Victoria, East Africa, from space. *Journal of Hydrology* 370: 163–176.
- Van Beek LPH, Wada Y, Bierkens MFP. 2011. Global monthly water stress: 1. Water balance and water availability. *Water Resources Research* 47: W07517.
- Van der Knijff JM, Younis J, De Roo APJ. 2010. LISFLOOD: a GIS-based distributed model for river-basin scale water balance and flood simulation. *International Journal of Geographical Information Science* 24: 189–212.
- Wada Y, Wisser D, Bierkens MFP. 2014. Global modeling of withdrawal, allocation and consumptive use of surface water and groundwater resources. *Earth System Dynamics* 5: 15–40.
- Walling DE. 2009. *The Impact of Global Change on Erosion and Sediment. Transport by Rivers*. UNESCO: Paris.
- Weedon GP, Gomes S, Viterbo P, Shuttleworth JW, Blyth E, Österle H, Adam JC, Bellouin N, Boucher O, Best M. 2011. Creation of the WATCH Forcing Data and its use to assess global and regional reference crop evaporation over land during the twentieth century. *Journal of Hydrometeorology* 12: 823–848.
- Winsemius HC, Van Beek LPH, Jongman B, Ward PJ, Bouwman A. 2013. A framework for global river flood risk assessments. *Hydrology and Earth System Science* 17: 1871–1892.
- Wisser D, Fekete BM, Vörösmarty CJ, Schumann AH. 2010. Reconstructing 20th century global hydrography: a contribution to the Global Terrestrial Network- Hydrology (GTN-H). *Hydrology and Earth System Science* 14: 1–24.
- Wisser D, Frohling S, Hagen S, Bierkens MFP. 2013. Beyond peak reservoir storage? A global estimate of declining water storage capacity in large reservoirs. *Water Resources Research* 49: 5732–5739.
- Wood EF, Roundy JK, Troy TJ, van Beek LPH, Bierkens MFP, Blyth E, de Roo A, Doell P, Ek M, Famiglietti J, Gochis D, van de Giesen N, Houser P, Jaffe P, Kollet S, Lehner B, Lettenmaier DP, Peters-Lidard C, Sivapalan M, Sheffield J, Wade A, Whitehead P. 2011. Hyperresolution global land surface modeling: meeting a grand challenge for monitoring Earth’s terrestrial water. *Water Resources Research* 47: W05301. DOI: 10.1029/2010WR010090.

## Data repositories referred to

- CRU – monthly meteorological forcing from observations. <https://climatedataguide.ucar.edu/climate-data/cru-ts321-gridded-precipitation-and-other-meteorological-variables-1901>
- ERA-Interim daily meteorological forcing from ECMRWF re-analysis. [http://data-portal.ecmwf.int/data/d/interim\\_daily/](http://data-portal.ecmwf.int/data/d/interim_daily/)
- EWA-Friend European catchment data. [http://www.bafg.de/GRDC/EN/04\\_spldtdbss/42\\_EWA/ewa\\_node.html](http://www.bafg.de/GRDC/EN/04_spldtdbss/42_EWA/ewa_node.html)
- FLUXNET: Water vapour, energy and CO<sub>2</sub> land-atmosphere fluxes from towers. <http://fluxnet.ornl.gov/obtain-data>
- GLCC land cover data. <http://landcover.usgs.gov/landcoverdata.php>
- GRDC global runoff data. [http://www.bafg.de/GRDC/EN/Home/homepage\\_node.html](http://www.bafg.de/GRDC/EN/Home/homepage_node.html)
- ISMN global network of soil moisture data. <https://ismn.geo.tuwien.ac.at/ismn/>
- MERRA daily meteorological forcing from the NASA Goddard Earth Observing System Data Assimilation System Version 5. <http://gmao.gsfc.nasa.gov/research/merra/>
- MOPEX US catchment data. [ftp://hydrology.nws.noaa.gov/pub/gcip/mopex/US\\_Data/](ftp://hydrology.nws.noaa.gov/pub/gcip/mopex/US_Data/)