

Complete Genome Sequence of *Campylobacter iguaniorum* Strain 1485E^T, Isolated from a Bearded Dragon (*Pogona vitticeps*)

Maarten J. Gilbert,^a William G. Miller,^b Emma Yee,^b Marja Kik,^c Jaap A. Wagenaar,^{a,d,e} Birgitta Duim^{a,e}

Department of Infectious Diseases and Immunology, Faculty of Veterinary Medicine, Utrecht University, Utrecht, The Netherlands^a; Produce Safety and Microbiology Research Unit, Agricultural Research Service, U.S. Department of Agriculture, Albany, California, USA^b; Department of Pathobiology, Faculty of Veterinary Medicine, Utrecht University, Utrecht, The Netherlands^c; Central Veterinary Institute of Wageningen UR, Lelystad, The Netherlands^d; WHO Collaborating Center for Campylobacter/OIE Reference Laboratory for Campylobacteriosis, Utrecht, The Netherlands^e

***Campylobacter iguaniorum* has been isolated from reptiles. This *Campylobacter* species is genetically related to *Campylobacter fetus* and *Campylobacter hyointestinalis*. Here we present the first whole-genome sequence for this species.**

Received 28 July 2014 Accepted 4 August 2014 Published 21 August 2014

Citation Gilbert MJ, Miller WG, Yee E, Kik M, Wagenaar JA, Duim B. 2014. Complete genome sequence of *Campylobacter iguaniorum* strain 1485E^T, isolated from a bearded dragon (*Pogona vitticeps*). *Genome Announc.* 2(4):e00844-14. doi:10.1128/genomeA.00844-14.

Copyright © 2014 Gilbert et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 3.0 Unported license](https://creativecommons.org/licenses/by/3.0/).

Address correspondence to Maarten J. Gilbert, m.j.gilbert@uu.nl, or William G. Miller, william.miller@ars.usda.gov.

Campylobacter iguaniorum is genetically related to, but distinct from, the species *Campylobacter fetus* and *Campylobacter hyointestinalis* and has recently been proposed as a novel *Campylobacter* species (Gilbert et al., submitted for publication). Reptiles, chelonians, and lizards in particular appear to be the primary reservoir of this *Campylobacter* species (1). The pathogenicity of this species is unknown; strains have been recovered from reptiles with and without clinical signs of disease (1, 2). Here, we report the first whole-genome sequence of the *C. iguaniorum* type strain 1485E^T (LMG 28143^T), isolated from a bearded dragon (*Pogona vitticeps*).

Sequencing was performed using shotgun and paired-end reads obtained on a Roche 454 FLX genome sequencer. Using the Newbler assembler (v. 2.6), we assembled a total of 220,914 454 reads into a single chromosomal scaffold of 12 contigs and a single megaplasmid contig, providing a draft genome sequence with a coverage of 68×. All 454 base calls were validated using 1,570,644 Illumina MiSeq reads, providing an additional 178× coverage. Scaffold gaps were filled as described (3). Sequences across the contig junctions were confirmed with Sanger sequencing. Homopolymeric GC tracts were characterized using the high-depth MiSeq reads.

The circular genome size of *C. iguaniorum* strain 1485E^T is 1,684,608 bp, with an average G+C content of 35.9%. A 70,030-bp megaplasmid is present. Protein-, rRNA-, and tRNA-encoding genes were identified as described (3). The genome was annotated based on *C. fetus* subsp. *testudinum* strain 03-427^T (accession number CP006833) (4, 5), with further annotation using Artemis (6), the identification of Pfam domains [v.26.0 (7)], and BLASTP comparisons to proteins in the NCBI nonredundant (nr) database. The chromosomal genome encodes 1663 putative protein-coding genes (including 18 probable pseudogenes), 43 tRNA genes, and 3 rRNA operons. The megaplasmid encodes an additional 111 putative protein-coding genes. A total of 34 homopolymeric GC tracts (≥8 bp) were identified, of which 23 were hypervariable. Many of these hypervariable GC tracts reside in surface-structure-related genes, as in other campylobacters; notably, however, eight are located within autotransporter domain-

containing genes, whose role in *C. iguaniorum* biology remains to be determined.

In contrast to *C. fetus*, strain 1485E^T does not encode an S-layer. Also, a defined lipooligosaccharide (LOS) region, bounded by *waa* genes and containing multiple glycosyltransferases, is absent from strain 1485E^T; instead, five separate predicted glycosylation regions were identified. One of these regions is >50 kb and has a deviant G+C content (30.0%), suggesting possible acquisition via lateral transfer. A clustered regularly interspaced short palindromic repeats (CRISPR)-Cas system is present. Multiple genes encoding respiratory enzymes not identified to date within the *C. fetus* group are also present. Interestingly, as in reptile-associated *C. fetus* subsp. *testudinum*, a putative tricarballoylate catabolism pathway was identified.

The whole-genome sequence of strain 1485E^T supports the proposal of *C. iguaniorum*. Further genome analysis and comparison can provide valuable insights into host adaptation, virulence, taxonomic structure, and evolution of this novel reptile-associated *Campylobacter* species.

Nucleotide sequence accession numbers. The complete genome sequence of *C. iguaniorum* strain 1485E^T has been deposited in GenBank under accession numbers [CP009043](https://www.ncbi.nlm.nih.gov/nuccore/CP009043) and [CP009044](https://www.ncbi.nlm.nih.gov/nuccore/CP009044).

ACKNOWLEDGMENTS

We thank Linda van der Graaf-van Bloois for useful advice and technical support. We thank Mary Chapman and Nathaniel Simon for the generation of Illumina MiSeq reads and Nathaniel Simon for the further assistance in the final assembly and genome closure.

This work was supported by USDA-ARS CRIS project 5325-42000-047-00D.

REFERENCES

- Gilbert MJ, Kik M, Timmerman AJ, Severs TT, Kusters JG, Duim B, Wagenaar JA. 2014. Occurrence, diversity, and host association of intestinal *Campylobacter*, *Arcobacter*, and *Helicobacter* in reptiles. *PLoS One* 9:e101599. <http://dx.doi.org/10.1371/journal.pone.0101599>.
- Benejat L, Gravet A, Sifré E, Ben Amor S, Quintard B, Mégraud F, Lehours P. 2014. Characterization of a *Campylobacter fetus*-like strain iso-

- lated from the faeces of a sick leopard tortoise (*Stigmochelys pardalis*) using matrix-assisted laser desorption/ionization time of flight as an alternative to bacterial 16S rDNA phylogeny. *Lett. Appl. Microbiol.* 58:338–343. <http://dx.doi.org/10.1111/lam.12194>.
3. Merga JY, Winstanley C, Williams NJ, Yee E, Miller WG. 2013. Complete genome sequence of the *Arcobacter butzleri* cattle isolate 7h1h. *Genome Announc.* 1(4):e00655-13. <http://dx.doi.org/10.1128/genomeA.00655-13>.
 4. Gilbert MJ, Miller WG, Yee E, Blaser MJ, Wagenaar JA, Duim B. 2013. Complete genome sequence of *Campylobacter fetus* subsp. *testudinum* strain 03–427^T. *Genome Announc.* 1(6):e01002-13. <http://dx.doi.org/10.1128/genomeA.01002-13>.
 5. Fitzgerald C, Tu ZC, Patrick M, Stiles T, Lawson AJ, Santovenia M, Gilbert MJ, van Bergen M, Joyce K, Pruckler J, Stroika S, Duim B, Miller WG, Loparev VL, Sinnige JC, Fields PI, Tauxe RV, Blaser MJ, Wagenaar JA. 4 June 2014. Description of *Campylobacter fetus* subsp. *testudinum* subsp. nov., isolated from humans and reptiles. *Int. J. Syst. Evol. Microbiol.* <http://dx.doi.org/10.1099/ijs.0.057778-0>.
 6. Rutherford K, Parkhill J, Crook J, Horsnell T, Rice P, Rajandream MA, Barrell B. 2000. Artemis: sequence visualization and annotation. *Bioinformatics* 16:944–945. <http://dx.doi.org/10.1093/bioinformatics/16.10.944>.
 7. Punta M, Coggill PC, Eberhardt RY, Mistry J, Tate J, Boursnell C, Pang N, Forslund K, Ceric G, Clements J, Heger A, Holm L, Sonnhammer EL, Eddy SR, Bateman A, Finn RD. 2012. The Pfam protein families database. *Nucleic Acids Res.* 40:D290–D301. <http://dx.doi.org/10.1093/nar/gkr1065>.