

Effects of neonicotinoids and fipronil on non-target invertebrates

L. W. Pisa · V. Amaral-Rogers · L. P. Belzunces · J. M. Bonmatin · C. A. Downs ·
D. Goulson · D. P. Kreutzweiser · C. Krupke · M. Liess · M. McField · C. A. Morrissey ·
D. A. Noome · J. Settele · N. Simon-Delso · J. D. Stark · J. P. Van der Sluijs · H. Van Dyck ·
M. Wiemers

Received: 8 May 2014 / Accepted: 15 August 2014 / Published online: 17 September 2014
© Springer-Verlag Berlin Heidelberg 2014

Abstract We assessed the state of knowledge regarding the effects of large-scale pollution with neonicotinoid insecticides and fipronil on non-target invertebrate species of terrestrial, freshwater and marine environments. A large section of the assessment is dedicated to the state of knowledge on sublethal effects on honeybees (*Apis mellifera*) because this important pollinator is the most studied non-target invertebrate species.

Lepidoptera (butterflies and moths), Lumbricidae (earthworms), Apoidae sensu lato (bumblebees, solitary bees) and the section “other invertebrates” review available studies on the other terrestrial species. The sections on freshwater and marine species are rather short as little is known so far about the impact of neonicotinoid insecticides and fipronil on the diverse invertebrate fauna of these widely exposed habitats.

Responsible editor: Philippe Garrigues

L. W. Pisa (✉) · N. Simon-Delso · J. P. Van der Sluijs
Environmental Sciences, Copernicus Institute, Utrecht University,
Heidelberglaan 2, 3584 CS Utrecht, The Netherlands
e-mail: l.w.pisa@uu.nl

V. Amaral-Rogers
Buglife, Bug House, Ham Lane, Orton Waterville, Peterborough PE2
5UU, UK

L. P. Belzunces
Laboratoire de Toxicologie Environnementale, INRA, UR 406
Abeilles & Environnement, Site Agroparc, 84000 Avignon, France

J. M. Bonmatin
Centre de Biophysique Moléculaire, UPR 4301 CNRS, affiliated to
Orléans University and to INSERM, 45071 Orléans cedex 02, France

C. A. Downs
Haereticus Environmental Laboratory, P.O. Box 92, Clifford,
VA 24533, USA

D. Goulson
School of Life Sciences, University of Sussex, Sussex BN1 9RH, UK

D. P. Kreutzweiser
Canadian Forest Service, Natural Resources Canada, 1219 Queen
Street East, Sault Ste Marie, ON P6A 2E5, Canada

C. Krupke
Department of Entomology, Purdue University, West Lafayette, IN,
USA

M. Liess
Department System-Ecotoxicology, Helmholtz Centre for
Environmental Research, UFZ, Permoserstrasse 15, 04318 Leipzig,
Germany

M. McField
Healthy Reefs for Healthy People Initiative, Smithsonian Institution,
Belize City, Belize

C. A. Morrissey
Department of Biology and School of Environment and
Sustainability, University of Saskatchewan, 112 Science Place,
Saskatoon, SK S7N 5E2, Canada

D. A. Noome
Task Force on Systemic Pesticides, 46, Pertuis-du-Sault,
2000 Neuchâtel, Switzerland

D. A. Noome
Kijani, Kasungu National Park, Private Bag 151, Lilongwe, Malawi

J. Settele · M. Wiemers
Department of Community Ecology, Helmholtz-Centre for
Environmental Research, UFZ, Theodor-Lieser-Str. 4, 06120 Halle,
Germany

J. Settele
German Centre for Integrative Biodiversity Research (iDiv),
Halle-Jena-Leipzig, Deutscher Platz 5e, 04103 Leipzig, Germany

For terrestrial and aquatic invertebrate species, the known effects of neonicotinoid pesticides and fipronil are described ranging from organismal toxicology and behavioural effects to population-level effects. For earthworms, freshwater and marine species, the relation of findings to regulatory risk assessment is described. Neonicotinoid insecticides exhibit very high toxicity to a wide range of invertebrates, particularly insects, and field-realistic exposure is likely to result in both lethal and a broad range of important sublethal impacts. There is a major knowledge gap regarding impacts on the grand majority of invertebrates, many of which perform essential roles enabling healthy ecosystem functioning. The data on the few non-target species on which field tests have been performed are limited by major flaws in the outdated test protocols. Despite large knowledge gaps and uncertainties, enough knowledge exists to conclude that existing levels of pollution with neonicotinoids and fipronil resulting from presently authorized uses frequently exceed the lowest observed adverse effect concentrations and are thus likely to have large-scale and wide ranging negative biological and ecological impacts on a wide range of non-target invertebrates in terrestrial, aquatic, marine and benthic habitats.

Keywords Pesticides · Neonicotinoids · Fipronil · Non-target species · Invertebrates · Honeybee · Earthworms · Butterflies · Freshwater habitat · Marine habitat

Introduction

Neonicotinoids and fipronil are relatively new, widely used, systemic compounds designed as plant protection products to kill insects which cause damage to crops. They are also used in veterinary medicine to control parasites such as fleas, ticks and worms on domesticated animals and as pesticides to control non-agricultural pests. Other papers in this special issue have shown that neonicotinoid insecticides and fipronil

are presently used on a very large scale (e.g. Simon-Delso et al. 2014, this issue), are highly persistent in soils, tend to accumulate in soils and sediments, have a high runoff and leaching potential to surface and groundwater and have been detected frequently in the global environment (Bonmatin et al. 2014, this issue). Effects of exposure to the large-scale pollution with these neurotoxic chemicals on non-target insects and possibly other invertebrates can be expected as identified for other insecticides. However, for the majority of insect and other invertebrate species that are likely to be exposed to neonicotinoids and fipronil in agricultural or (semi)natural ecosystems, no or very little information is available about the impact of these pesticides on their biology. Here we assess the present state of knowledge on effects on terrestrial and aquatic invertebrates.

Terrestrial invertebrates

Honeybees

Many studies have focused on investigating the effects of neonicotinoids and fipronil on honeybees (*Apis mellifera*). Apart from its cultural and honey production value, the honeybee is the most tractable pollinator species and critical for the production of many of the world's most important crops (Klein et al. 2007; Breeze et al. 2011). Losses of honeybees are generally measured as winter loss on national to regional level, and indications are that honeybee populations undergo high losses in many parts of the world (Oldroyd 2007; Stokstad 2007; van Engelsdorp and Meixner 2010; Van der Zee et al. 2012a, b).

No single cause for high losses has been identified, and high losses are associated with multiple factors including pesticides, habitat loss, pathogens, parasites and environmental factors (Decourtye et al. 2010; Mani et al. 2010; Neumann and Carreck 2010; Kluser et al. 2011). Apart from direct biotic and abiotic factors, changes in honeybee populations also depend on the economic value of honeybees and thus on human effort (Aizen and Harder 2009; Mani et al. 2010). Neonicotinoids are among the most used insecticides worldwide and are thus prime targets for investigating possible relationships with high honeybee losses.

Acute and chronic lethal toxicity to honeybees

Neonicotinoids and fipronil show high acute toxicity to honeybees (Table 1). The neonicotinoid family includes imidacloprid, clothianidin and thiamethoxam (the latter is metabolized to clothianidin in the plant and in the insect). Imidacloprid, clothianidin and thiamethoxam belong to the nitro-containing neonicotinoids, a group that is generally more toxic than the cyano-containing neonicotinoids, which includes acetamiprid and thiacloprid. Although neonicotinoids

N. Simon-Delso
Beekeeping Research and Information Centre (CARI), Place Croix
du Sud 4, 1348 Louvain-la-Neuve, Belgium

J. D. Stark
Puyallup Research and Extension Centre, Washington State
University, Puyallup, WA 98371, USA

J. P. Van der Sluijs
Centre for the Study of the Sciences and the Humanities, University
of Bergen, Postboks 7805, 5020 Bergen, Norway

H. Van Dyck
Behavioural Ecology and Conservation Group, Biodiversity
Research Centre, Earth and Life Institute, Université Catholique de
Louvain (UCL), Croix du Sud 4-5, bte L7.07.04,
1348 Louvain-la-Neuve, Belgium

Table 1 Toxicity of insecticides to honeybees, compared to DDT. Dose used is given in gram per hectare, median lethal dose (LD₅₀) is given in nanogram per bee. The final column expresses toxicity relative to DDT (DDT is 1). Source: Bonmatin (2011)

Pesticide	®Example	Main use	Typical dose (g/ha)	Acute LD ₅₀ (ng/bee)	Ratio of LD ₅₀ as compared to DDT
DDT	Dinocide	Insecticide	200–600	27,000	1
Thiacloprid	Proteus	Insecticide	62.5	12,600	2.1
Amitraz	Apivar	Acaricide	–	12,000	2.3
Acetamiprid	Supreme	Insecticide	30–150	7,100	3.8
Coumaphos	Perizin	Acaricide	–	3,000	9
Methiocarb	Mesurool	Insecticide	150–2,200	230	117
Tau-fluvalinate	Apistan	Acaricide	–	200	135
Carbofuran	Curater	Insecticide	600	160	169
Λ-cyhalotrin	Karate	Insecticide	150	38	711
Thiametoxam	Cruiser	Insecticide	69	5	5,400
Fipronil	Regent	Insecticide	50	4.2	6,475
Imidacloprid	Gaucho	Insecticide	75	3.7	7,297
Clothianidin	Poncho	Insecticide	50	2.5	10,800
Deltamethrin	Decis	Insecticide	7.5	2.5	10,800

are applied as foliar insecticides with possible direct exposure risks to honeybees, a large part of neonicotinoid use consists of seed coating or root drench application. Fipronil belongs to the phenylpyrazole family of pesticides and, like the neonicotinoids, has systemic properties (Simon-Delso et al. 2014).

Given that neonicotinoids and fipronil act systemically in plants, oral lethal doses for honeybees have been extensively studied for these compounds. Unlike many older classes of insecticides, neonicotinoids may be more toxic when ingested (Suchail et al. 2001; Iwasa et al. 2004). The level of neonicotinoids and fipronil that honeybees are exposed to in the nectar and pollen of treated plants varies greatly, although there are trends based upon application method. Generally, soil drenches and foliar application result in higher concentrations of the active compounds in plants than seed treatments, with the latter application used in large, annual cropping systems like grain crops, cotton and oilseed crops.

In practice, the honeybee lethal dose 50 (LD₅₀) for these pesticides varies for a wide range of biotic and abiotic conditions. The LD₅₀ of imidacloprid, for example, has shown values between 3.7 and 40.9, 40 and 60, 49 and 102 and 490 ng/bee (Nauen et al. 2001; Schmuck et al. 2001; Suchail et al. 2001; DEFRA 2007, 2009). This variation, of a factor 100 (5–500 ng/bee), has been observed not only between colonies but also among bees taken from a single colony. A major component of this observed variation likely stems from the discrepancy in the contact and oral toxicities of these compounds, with contact lethal doses generally being higher than oral lethal doses. However, contact with the floral parts is frequent when the bees visit flowers, and this

is different from the topical application used in laboratory conditions.

Other sources of variability may be attributed to differences in environmental conditions during testing as well as any inherent differences in the condition of the tested bees themselves. For example, data have shown that measured LD₅₀ values for bees vary with temperature (Medrzycki et al. 2011), the age of bees (Schmuck 2004; Medrzycki et al. 2011), the honeybee subspecies tested (Suchail et al. 2000), the pattern of exposure (Illarionov 1991; Belzunces 2006) and prior exposure of bees to pesticides (Belzunces 2006). Given the large variability of honeybee toxicity data, it has been suggested that LD₅₀ values should only be used to compare levels of toxicity among pesticides rather than drawing conclusions about the risk of mortality posed to honeybees via environmental exposure to pesticides (Belzunces 2006).

Oral subchronic exposure to imidacloprid and six of its metabolites induced a high toxicity at concentrations of 0.1, 1 and 10 ppb (part per billion) or ng/g, whereas the metabolites olefin-imidacloprid and 5-OH-imidacloprid were toxic in acute exposure. The main feature of subchronic toxicity is the absence of a clear dose–effect relationship that can account for a maximum effect of the lowest concentration due to the existence of multiple molecular targets, as has been demonstrated in the honeybee (Déglise et al. 2002; Thany et al. 2003; Thany and Gauthier 2005; Barbara et al. 2008; Gauthier 2010; Dupuis et al. 2011; Bordereau-Dubois et al. 2012). The absence of clear dose–effect relationships has also been observed in other studies, at higher concentrations (Schmuck 2004).

Existence of non-monotonic dose–response relations implies that some chemicals, including neonicotinoids,

have unexpected and potent effects at (very) low doses. These non-linear and often non-intuitive patterns are due to the complex interplay of receptor binding and gene reprogramming effects of such substances and can generate unexpected dose–response relationships, many of which are still being mapped out (Fagin 2012; Charpentier et al. 2014). This poses major challenges to risk assessment based on the classical log-probit model.

As previously reviewed by van der Sluijs et al. (2013), there are no standardised protocols for measuring chronic lethal effects. In traditional risk assessment of pesticides, they are usually expressed in three ways: LD₅₀: the dose at which 50 % of the exposed honeybees die (usually within a 10 day time span); no observed effect concentration (NOEC): the highest concentration of a pesticide producing no observed effect; and lowest observed effect concentration (LOEC): the lowest concentration of a pesticide producing an observed effect.

For imidacloprid, including its neurotoxic metabolites, lethal toxicity can increase up to 100,000 times compared to acute toxicity when the exposure is extended in time (Suchail et al. 2001). There has been some controversy on the findings of that study, which are discussed in detail by Maxim and Van der Sluijs (2007, 2013). However, the key finding that exposure time amplifies the toxicity of imidacloprid is consistent with later findings, implying that the standard 10 day chronic toxicity test for bees is far too short for testing neonicotinoids and fipronil, given their persistence and hence the likely chronic exposure of bees under field conditions. Indeed, honeybees fed with 10⁻¹ of the LC₅₀ of thiamethoxam showed a 41.2 % reduction of life span (Oliveira et al. 2013). Recent studies have shown that chronic toxicity of neonicotinoids can more adequately be expressed by time to 50 % mortality instead of by the 10 day LD₅₀ (Sánchez-Bayo 2009; Maus and Nauen 2010; Tennekes 2010; Tennekes 2011; Tennekes and Sánchez-Bayo 2012; Mason et al. 2013; Rondeau et al. 2014). There is a linear relation between the logarithm of the daily dose and the logarithm of the time to 50 % mortality (Tennekes 2010, 2011; Tennekes and Sánchez-Bayo 2012; Tennekes and Sánchez-Bayo 2013; Rondeau et al. 2014). Sanchez-Bayo and Goka (2014) demonstrated that field-realistic residues of neonicotinoid insecticides in pollen pose high risk to honeybees and bumblebees, whilst in the field synergisms with ergosterol inhibiting fungicides will further amplify these risks. They found that imidacloprid poses the highest risk to bumblebees (31.8–49 % probability to reach the median lethal cumulative dose after 2 days feeding on field-realistic dose in pollen) and thiamethoxam the highest risk to honeybees (3.7–29.6 % probability to reach median lethal cumulative dose). In experiments with honeybee colonies, similar, long-term chronic effects have been found with typical times of 80–120 days for 1 ppm dinotefuran and 400 ppb clothianidin (Yamada et al. 2012). Note that these

studies used concentrations that are on the uppermost limit of the currently reported ranges of concentrations found in pollen and nectar in the field. However, such data are sparse and limited to a few crops only, so it cannot yet be concluded whether such concentrations are rare or common in the field—the question of “field-relevant dose” is not yet fully resolved, and it is likely that there is a wide range in these values over space and time (Van der Sluijs et al. 2013).

Field and laboratory studies attempting to test field-realistic lethal doses have shown variable, often conflicting, results. In one study, chronic oral and contact exposure during 10–11 days to 1 µg/bee of acetamiprid and 1,000 µg/bee of thiamethoxam caused no significant worker mortality (Aliouane et al. 2009). Conversely, laboratory studies using imidacloprid showed high worker mortality when honeybees consumed contaminated pollen (40 ppb) (Decourtye et al. 2003, 2005) and contaminated sugar syrup (0.1, 1.0 and 10 ppb) (Suchail et al. 2001). These results were contrary to those of field studies performed by Schmuck et al. (2001), who reported no increased worker mortality when colonies were exposed to sunflower nectar contaminated with imidacloprid at rates from 2.0 to 20 µg/kg. Faucon et al. (2005) also found no worker mortality in a field study of honeybees fed imidacloprid in sugar syrup. A meta-analysis by Cresswell (2011) concluded that oral exposure to imidacloprid at realistic field concentrations did not result in worker mortality, although a subsequent study by Yamada et al. (2012) feeding a range of dinotefuran (1–10 ppm) and clothianidin (0.4–4 ppm) concentrations demonstrated colony failure within 104 days in each case, suggesting that detection of colony-level effects may require longer post-exposure observation.

Field studies to investigate the exposure of bees to pesticides face major difficulties. For the analysis of very low concentrations of compounds present in pollen, nectar, bees or other matrices, appropriate methods that meet validity criteria of quantitative analysis have to be developed. Pilling et al. (2013) exposed bees to thiamethoxam-treated maize and oilseed rape but were not able to quantify concentrations lower than 1 ppb, although this may be a result of the authors using a lower seed treatment application than is used in normal agricultural practice. Even though both treatment and control colonies experienced relatively high losses (mostly queens laying only drone brood) and the authors were unable to undertake any statistical analysis due to a lack of replication, they wrongly concluded that there is a low risk to honeybees from exposure to treated maize and oilseed rape.

Also, in terms of activity and feeding behaviour, bees might not be foraging on treated crops in (exactly) the same way as they would do on untreated crops (Colin et al. 2004). Furthermore, comparison of treated and control areas can be totally flawed because control fields might not be “clean” but treated with other pesticides,

including insecticides. The recent study of Pilling and co-workers on thiamethoxam (Pilling et al. 2013) is illustrative for this case as it did not provide information about the treatment status of the control plots.

For mass-dying of bees in spring near corn fields during sowing of neonicotinoid-treated seeds, there now is a one to one proven causal link. Acute intoxication occurs through exposure to the dust cloud around the pneumatic sowing machines during foraging flights to adjacent forests (providing honeydew) or nearby flowering fields (Apenet 2010; Girolami et al. 2012; Tapparo et al. 2012; Krupke et al. 2012; Pochi et al. 2012; Tapparo et al. 2012). In these cases, dead bees have typically been found to have high levels of seed treatment neonicotinoids on, or in, their bodies. Such mass colony losses during corn sowing have been documented in Italy, Germany, Austria, Slovenia, the USA and Canada (Gross 2008; Krupke et al. 2012; Sgolastra et al. 2012; Tapparo et al. 2012). In response to the incidents, the adherence of the seed coating has been improved owing to better regulations, and an improved sowing technique has recently become compulsory throughout Europe (European Commission 2010). However, despite the deployment of air deflectors in the drilling machines and improved seed coating techniques, emissions are still substantial and the dust cloud remains acutely toxic to bees (Biocca et al. 2011; Marzaro et al. 2011; Girolami et al. 2012; Tapparo et al. 2012; Sgolastra et al. 2012).

Acute lethal effects of neonicotinoids dispersed as particles in the air seem to be promoted by high environmental humidity (Girolami et al. 2012). Honeybees also transport toxic dust particles on their bodies into the hive (Girolami et al. 2012). Sunny and warm days also seem to favour the dispersal of active substances (Greatti et al. 2003).

Sublethal effects on honeybees

Effects on activity, locomotion, metabolism and ontogenetic development Imidacloprid, thiamethoxam and clothianidin have been shown to rapidly induce flight muscle paralysis in honeybees exposed to guttation drops containing these substances, resulting in the cessation of wing movements (Girolami et al. 2009). Imidacloprid further impairs the mobility of bees, as reflected by decreases in running and walking and increases in the time that exposed bees remain stationary (Medrzycki et al. 2003). However, when exposed to sub-chronic doses of neonicotinoids, decreases in locomotion were not observed in honeybees and bumblebees by Cresswell et al. (2012b).

Ontogenetic development is a crucial period that determines the physiological and functional integrity of adult individuals. Thus, in addition to the effects on adults, neonicotinoids may act on larval development with consequences for the adult stage. Adult honeybees exposed to

imidacloprid during the larval stage exhibit impairment of olfactory associative behaviour (Yang et al. 2012). This could be due to altered neural development. Impairments in mushroom body development in the bee brain and the walking behaviour of honeybee workers have been observed in individuals exposed to imidacloprid during the larval period (Tomé et al. 2012). Effects on adult bees exposed during the larval stage could also be attributed to the induction of cell death by imidacloprid in larvae (Gregorc and Ellis 2011). In the early stages of adult life, after emergence, imidacloprid can disrupt the development of hypopharyngeal glands by decreasing the size of the acini and by increasing the expression of hsp70 and hsp90 (Smodis Skerl et al. 2009; Hatjina et al. 2013). Derecka et al. (2013) provided beehives in the field for 15 days with syrup tainted with 2 µg/l imidacloprid. They found that these levels of imidacloprid, at the low end of the field-realistic range, significantly impact energy metabolism in worker bee larvae.

Impacts of pesticides on metabolism may affect the detoxifying, intermediary and energetic metabolism pathways. Imidacloprid impairs brain metabolism in the honeybee which results in an increase of cytochrome oxidase in mushroom bodies (Decourtye et al. 2004a, b).

Effects on behaviour, learning and memory Optimal function of the honeybee nervous system is critical to individual and colony functioning (Desneux et al. 2007; Thompson and Maus 2007). Increasing levels of research effort have been devoted to developing an improved understanding of how sublethal exposure to neonicotinoids and fipronil may affect the honeybee nervous system. There is evidence that sublethal exposure can affect learning, memory and orientation in honeybees.

Laboratory experiments administering a single dose of imidacloprid demonstrated that learning was altered (Guez et al. 2001; Lambin et al. 2001), and exposure to chronic sublethal doses has demonstrated that learning and foraging are impaired by imidacloprid and fipronil (Decourtye et al. 2003). Furthermore, thiamethoxam has been shown to decrease memory capacity (Aliouane et al. 2009). The methodologies and doses varied in these laboratory tests, but all used concentrations above 20 ppb; this is towards the upper end of concentrations found in most field situations. These concentrations would not be expected to be found in pollen or nectar following seed treatment applications, but have been found in cucurbit flowers following soil drench applications (Dively and Hooks 2010). Field experiments offer the potential for powerful tests; however, results have been mixed, and many studies focus on honeybee orientation to and from a feeding source. A study that trained honeybee foragers to a sugar syrup reward in a complex maze demonstrated that 38 % of bees found the food source following ingestion of 3 ng/bee of

thiamethoxam, compared with 61 % in an unexposed control group (Decourtye and Devillers 2010). A series of studies training foragers to orient to a sugar feeder found that foragers were unable to return to the hive after ingesting imidacloprid at concentrations ranging from 100 to 1,000 ppb (Bortolotti et al. 2003; Ramirez-Romero et al. 2005; Yang et al. 2008). In contrast, other semi-field studies have shown no effects upon foraging or survivorship following exposure to canola, maize and sunflowers grown from neonicotinoid-treated seeds (Schmuck et al. 2001; Cutler and Scott-Dupree 2007; Nguyen et al. 2009). Possible explanations for these conflicting results may be that when given a range of foraging opportunities, honeybees may reduce foraging visits to food sources containing pesticides (Mayer and Lunden 1997; Colin et al. 2004), or that neonicotinoids do not have effects on colonies in the exposure regimes tested here.

Recently, Henry et al. (2012a, b) described the results of innovative field experiments using radio frequency identification (RFID) tags to determine the colony-level effects of orientation impairment upon foragers fed a sublethal dose of imidacloprid (1.42 ng in 20 μ l of sucrose syrup). In two separate experiments, treated foragers failed to return to the colony at rates of 10.2 and 31.6 %, relative to untreated foragers feeding upon the same flowering plants. A higher risk of not returning was associated with the more difficult orientation tasks. Using these forager loss rates, the researchers modelled the colony-level effects and found significant, largely consistent deviations from normal colony growth rates, in some cases to levels that may put the colony at risk of collapse. A subsequent suggestion by Cresswell and Thompson (2012) to alter the simulation slightly to reflect the period when seed-treated crops are flowering demonstrated that the risk of collapse was no longer evident. However, a follow-up calculation by Henry et al. (2012a) using a larger dataset that incorporated a range of empirically derived colony growth estimates revealed even higher deviations from normal than the original work: a more serious negative outcome for colonies. The variable outcomes based upon model assumptions reflect uncertainties that have plagued honeybee researchers and further underscore the importance of ensuring that models are robust and represent a range of scenarios. The key contribution of this work was the demonstration that sublethal doses can impose a stressor (i.e. non-returning foragers) that can have significant negative outcomes on a colony level.

Learning and memory represent fundamental functions involved in the interaction of individuals with their environment and are critical in enabling bees to respond to the requirements of the colony throughout their life. Imidacloprid impairs learning and olfactory performance via both acute and chronic exposure pathways, and summer bees appear more sensitive than winter bees (Decourtye et al. 2003). These effects are observed not only in the laboratory but also in

semi-field conditions, and bees do not recover after exposure ceases. Results obtained with acetamiprid and thiamethoxam showed that the action of neonicotinoids depends on the level/degree of exposure and cannot be generalized to structurally related compounds. Unlike contact exposure, oral exposure of acetamiprid resulted in an impairing of long-term retention of olfactory learning (El Hassani et al. 2008). Conversely, for thiamethoxam, subchronic exposure, but not acute exposure, elicited a decrease of olfactory memory and an impairment of learning performance (El Hassani et al. 2008; Aliouane et al. 2009).

Neonicotinoids have specific routes of metabolism in insects, particularly in the honeybee, that lead to complex influences on learning and memory processes. Imidacloprid and thiamethoxam are metabolized into toxic metabolites that may potentially bind to different honeybee nicotinic acetylcholine receptors (Nauen et al. 2001; Suchail et al. 2001, 2004a; Nauen et al. 2003; Ford and Casida 2006; Benzidane et al. 2010; Casida 2011). The metabolism of acetamiprid results in the appearance of different metabolites in the honeybee, among which 6-chloronicotinic acid is toxic in chronic exposure but not in acute exposure and remains stable for at least 72 h, especially in the head and the thorax (Suchail et al. 2001, 2004a; Brunet et al. 2005). Considering the presence of multiple active metabolites over time, it is very difficult to ascertain what steps of the memory process (acquisition, consolidation or retrieval) are affected by imidacloprid, acetamiprid, thiamethoxam or their metabolites.

Habituation may be defined as “a form of learning that consists in the gradual and relatively prolonged decrease of the intensity or the frequency of a response following the repeated or prolonged stimulus responsible for eliciting such a response” (Braun and Bicker 1992; Epstein et al. 2011a, b; Belzunces et al. 2012). Habituation can be regarded as an important adaptive behaviour because it allows individuals to minimize their response and, therefore, their energy investment, towards unimportant stimuli. The neonicotinoid imidacloprid alters patterns of habituation in honeybees following contact exposure to a sublethal dose (Guez et al. 2001; Lambin et al. 2001). Imidacloprid-induced changes in habituation appear to vary depending on the age of bees and time after exposure. Furthermore, these changes in habituation may be due to factors such as differential sensitivity of different nicotinic acetylcholine receptors (nAChRs) to imidacloprid (Déglise et al. 2002; Thany et al. 2003; Thany and Gauthier 2005; Barbara et al. 2008; Gauthier 2010; Dupuis et al. 2011; Bordereau-Dubois et al. 2012; Farooqui 2013), or the accumulation of imidacloprid metabolites like olefin and 5-hydroxy-imidacloprid, which can delay or accelerate habituation, respectively (Guez et al. 2001, 2003).

Olfaction and taste are very important physiological senses for honeybees (Detzel and Wink 1993; Giurfa 1993; Balderrama et al. 1996; Goulson et al. 2001; Reinhard et al.

2004; Gawleta et al. 2005; Couvillon et al. 2010; Maisonnasse et al. 2010; Kather et al. 2011). The effects of neonicotinoids on gustation can be explored by studying the modulation of the gustatory threshold, which can be defined as the lowest concentration of a sucrose solution applied to the antenna that triggers a feeding response. Different active compounds have been shown to induce dissimilar effects on gustation in honeybees. For example, fipronil increases the gustatory threshold of bees subjected to contact exposure (El Hassani et al. 2005). Whilst similar results were found for imidacloprid, acetamiprid decreases the threshold of bees that are exposed orally, but not topically (El Hassani et al. 2009). Thiamethoxam elicits a decrease in honeybee responsiveness to sucrose, and exposure to acetamiprid increases the responsiveness of honeybees to water regardless of exposure route (El Hassani et al. 2008; Aliouane et al. 2009).

The discrepancy in the effects observed could be explained in part by neonicotinoid metabolism that induced the appearance of toxic metabolites (Suchail et al. 2004a, b; Brunet et al. 2005) and by the existence of different nAChRs that are either sensitive and resistant to particular neonicotinoids (Déglise et al. 2002; Thany et al. 2003; Thany and Gauthier 2005; Barbara et al. 2008; Gauthier 2010; Dupuis et al. 2011; Bordereau-Dubois et al. 2012). Although it has been demonstrated in pollinating flies and in beetles, the repellent effect of imidacloprid and other neonicotinoids has not been investigated in the honeybee (Easton and Goulson 2013).

Accurate navigation is essential for efficient foraging and, hence, for colony health and survival. Neonicotinoids and fipronil may impair navigation in different ways. Sublethal exposure of honeybees to clothianidin and imidacloprid elicits a decrease in foraging activity and induces longer foraging flights (Schneider et al. 2012). Thiamethoxam induces high mortality by causing failure in the homing behaviour of foraging bees, leading to large losses of foragers from the colony (Henry et al. 2012a, b). Although this effect has been demonstrated for the pyrethroid deltamethrin for almost 20 years (Vandame et al. 1995), impacts on the homing behaviour of foraging bees continue to be left out of the assessment process for pesticide registration.

Proper foraging behaviour is essential for both individual bees and the colony as a whole because it determines the availability of food (stores) and, consequently, the survival of the colony. Exposure to imidacloprid, clothianidin and fipronil can lead to reductions in the proportion of active bees in the hive and, furthermore, initiate behaviours that can reduce the efficiency of foraging flights. For example, exposed individuals may spend longer periods of time at a food source, decrease the frequency of visits, increase the time between foraging trips, engage in longer foraging flights, reduce foraging distances, exhibit problems revisiting the same feeding site or exhibit reductions in visual learning capacities (Nielsen et al. 2000; Morandin and Winston 2003;

Colin et al. 2004; Ramirez-Romero et al. 2005; Yang et al. 2008; Han et al. 2010; Schneider et al. 2012; Teeters et al. 2012). Fischer et al. (2014) exposed adult honeybees to sublethal doses of imidacloprid (7.5 and 11.25 ng/bee), clothianidin (2.5 ng/bee) and thiacloprid (1.25 µg/bee) and subsequently tracked the flight paths of individual bees with harmonic radar. The rate of successful return was significantly lower in treated bees, the probability of a correct turn at a salient landscape structure was reduced and less directed flights during homing flights were performed. These findings show that sublethal doses of these three neonicotinoids either block the retrieval of exploratory navigation memory or alter this form of navigation memory. Reproduction and colony development may be regarded as integrative endpoints for assessing the final impacts of pesticides on bees as both are a compulsory condition of social insect physiology.

Neonicotinoids such as thiacloprid, thiamethoxam and imidacloprid decrease brood production, larval eclosion, colony growth rate and the number of queens reared in bumblebees (Tasei et al. 2000; Mommaerts et al. 2010; Whitehorn et al. 2012). Studies suggest that the reduction in brood production may be associated with a reduction in pollen and sugar consumption by adult bees (Laycock et al. 2012a, b). The rearing of honeybees on brood comb containing high levels of pesticide residues results in delayed larval development and emergence and shortened adult longevity (Wu et al. 2011). Since the brood combs in the latter study contained five neonicotinoids at relatively high concentrations, it is difficult to ascribe the observed effects to any one pesticide, or pesticide class. An epidemiological study involving Hill's criteria (minimal conditions that prove evidence of a causal relationship) revealed conflicting results on the involvement of dietary traces of neonicotinoids in the decline of honeybee populations (Cresswell et al. 2012a) and could not establish a causal link between observations of bee decline and neonicotinoid use rates.

Interaction with pathogens

Detrimental effects of pesticides might be increased in combination with other environmental stress agents (Mason et al. 2013). Specific pathogens and parasites are ancestral companions of (some) honeybee populations, and accidental movement of parasites and pathogens by man has exposed both honeybees and wild bees to non-native enemies to which they may have reduced resistance (e.g. Goulson 2003; Graystock et al. 2013a, b). Imidacloprid can act synergistically with the pathogen *Nosema* spp. by increasing *Nosema*-induced mortality (Alaux et al. 2010). It affects social immunity and so increases the number of *Nosema* spores in the guts of bees from imidacloprid-exposed colonies exposed in cage studies (Pettis et al. 2012). Sequential exposure to *Nosema ceranae* can sensitize bees to thiacloprid by eliciting potentiation that

leads to high mortality rates, a feature shared with fipronil (Vidau et al. 2011; Aufauvre et al. 2012). Similarly, other experiments with fipronil and *N. ceranae* have demonstrated reciprocal sensitization (Aufauvre et al. 2012). Furthermore, exposure to pesticides during embryonic and post-embryonic development may alter the susceptibility of adult bees to pathogens. For example, adult honeybees reared in brood combs containing high levels of pesticide residues exhibit higher levels of infection by *N. ceranae* and higher levels of *Nosema* spores (Wu et al. 2012).

Di Prisco et al. (2013) demonstrated that clothianidin negatively modulates nuclear factor kappa-light-chain-enhancer of activated B cells (NF- κ B, a protein involved in DNA transcription) immune signaling in insects and adversely affects honeybee antiviral defences controlled by this transcription factor. They identified a negative modulator of NF- κ B activation specific for insects. Exposure to clothianidin, by enhancing the transcription of the gene encoding this inhibitor, reduces immune defences and promotes the replication of the deformed wing virus present in honeybees. Similar immunosuppression was found to be induced by imidacloprid. The occurrence of this insecticide-induced viral proliferation at sublethal doses that are well within field-realistic concentrations suggests that the studied neonicotinoids are likely to have a negative effect under field conditions.

Synergistic effects with other pesticides

In agricultural ecosystems, honeybees are seldom exposed to only a single pesticide. Combined exposures could be of high concern because they can elicit synergies and potentiations. For example, thiacloprid acts synergistically with ergosterol biosynthesis inhibitor (EBI) fungicides in bees exposed in laboratory conditions but not in tunnel conditions (Schmuck et al. 2003).

Analyses of honeybees and colony contents indicate that honeybees are indeed frequently exposed to multiple pesticides simultaneously (Mullin et al. 2010; Krupke et al. 2012; Paradis et al. 2013). However, the study of pesticide mixtures can be challenging (Lydy et al. 2004), and there is a paucity of information in the literature regarding the mixtures encountered by honeybees. Triazole fungicides have been found in pollen collected from colonies (Krupke et al. 2012) and have been shown to synergize toxicity of some neonicotinoids (thiacloprid and acetamiprid) up to 559-fold in the laboratory, although the same results have not been shown in semi-field studies (Schmuck et al. 2003). Piperonyl butoxide also has been found in pollen and has been shown to synergize toxicity of some neonicotinoids (thiacloprid and acetamiprid) up to 244-fold in the laboratory (Iwasa et al. 2004). Despite the challenges associated with this type of research, this is a clear research gap that should be addressed in the future, given that

honeybees rarely encounter only a single pesticide during foraging and/or in the hive.

Toxicity to bumblebees and solitary bees

Bumblebees (genus *Bombus*) are primitive social bees. Colonies start from overwintering queens, build up to a few hundred adult workers and break down when new queens and males are produced. A small number of bumblebee species are commercially reared for pollination, but many of the non-managed bumblebees also contribute substantially to crop pollination (Chagnon et al. 1993; Bosch and Kemp 2006; Greenleaf and Kremen 2006; Goulson 2010). Solitary bees that are also commonly managed in agricultural settings include the alfalfa leafcutter bee (*Megachile rotundata*), alkali bees (*Nomia melanderi*), blue orchard bees (*Osmia lignaria*) and Japanese horn-faced bees (*Osmia cornifrons*). *M. rotundata* is the major pollinator of alfalfa, which is grown as a high value livestock feed in North America. It is often considered a domesticated species, although populations frequently occur naturally. This species contributed US\$5.26 billion to the value of alfalfa hay in 2009 (Calderone 2012). In addition to managed bees, there are more than 20,000 species of wild bees in the world, many of which contribute to crop pollination, and all of them contribute to pollination of wild flowers.

There are few long-term population-level studies involving bumblebees and other bee species, and in many cases, the impacts of pesticide exposure and dosage are unclear. These species differ from honeybees in that they generally exhibit smaller foraging ranges and often prefer to nest in the ground. Therefore, populations located near agricultural operations and associated pesticide applications may have fewer alternative options for food and habitat resources. Furthermore, ground-nesting species may face additional exposure risks (i.e. pesticide-contaminated soil) that are not encountered by honeybees, but which remain to be evaluated. Finally, whilst bumblebees tend to be bigger, solitary bees are often smaller than honeybees; thus, these species likely receive a different dose relative to their body weight than honeybees do.

Likely levels of exposure of wild bee species are poorly understood. Whilst neonicotinoid levels have been quantified in the nectar and pollen of various crop plant species (Cresswell 2011; Anon 2012), the degree to which wild bees utilize these resources has not been measured, and furthermore, basic values of toxicity, such as LD₅₀ and LC₅₀, are completely lacking for the vast majority of these species. The few studies that do exist have employed a range of methods with conflicting results so that drawing general conclusions is difficult at this stage. Moreover, these studies are criticised for low sample size, which limits power to detect effects and/or highly unnatural laboratory conditions.

It is clear that neonicotinoids and fipronil are highly toxic to all bee species tested so far, which in addition to honeybees

includes various *Bombus* species, several social stingless bee species and the solitary species *O. lignaria* and *M. rotundata* (Scott-Dupree et al. 2009; Valdovinos-Núñez et al. 2009; Gradish et al. 2010; Mommaerts et al. 2010; Tomé et al. 2012). Cresswell et al. (2012a, b) demonstrated that bumblebees exhibit sublethal responses to imidacloprid at 10 ppb, whilst honeybees were unaffected at this concentration. Scott-Dupree et al. (2009) found that *O. lignaria* is more sensitive to both clothianidin and imidacloprid than *Bombus impatiens*, with *M. rotundata* more sensitive still. Stark et al. (1995) found no difference in the 24 h contact LD₅₀ of imidacloprid between honeybees and the solitary bee species *M. rotundata* and *N. melanderi*. Scott-Dupree et al. (2009) demonstrated that *B. impatiens* individuals were more tolerant of thiamethoxam and clothianidin than *O. lignaria* and *M. rotundata*. However, the orchard bee *O. lignaria* exhibits delayed hatching and development when fed imidacloprid at rates from 30 to 300 µg/kg (Abbott et al. 2008). Arena and Sgolastra (2014) compared the acute toxicity of numerous pesticides and found that *Scaptotrigona postica* and *M. rotundata* were more sensitive than honeybees to fipronil, whilst *N. melanderi* was more tolerant. Together, these results suggest that “other” bees may be at least as sensitive, if not more sensitive, to neonicotinoids than honeybees, although more work is clearly needed.

A number of studies have used queenless micro-colonies of bumblebees (containing only workers) to examine the sublethal effects of cumulative neonicotinoid exposure to low, field-realistic doses. Several have found no detectable effects; for example, Tasei et al. (2000) exposed *Bombus terrestris* micro-colonies to 6–25 ppb of imidacloprid and found no significant response. Similarly, Franklin et al. (2004) exposed *B. impatiens* to concentrations of up to 36 ppb of clothianidin without detecting an impact (see also Morandin and Winston 2003). Most recently, Laycock et al. (2012a, b) exposed micro-colonies of *B. terrestris* to a range of concentrations of imidacloprid (0–125 µg/l) and detected a 30 % reduction in fecundity at doses as low as 1 ppb. In the only comparable work on other bee species, Abbott et al. (2008) injected concentrations of up to 300 ppb of neonicotinoids into pollen stores of *O. lignaria* and *M. rotundata* with no measurable impact on larval development.

Interestingly, negative effects seem to be detected more frequently and at lower concentrations when bees have to forage at a distance, even when the distances are tiny. Mommaerts et al. (2010) found no impact of imidacloprid exposure on micro-colonies of *B. terrestris* at field-realistic concentrations when food was provided in the nest, but when workers had to walk just 20 cm down a tube to gather food they found significant sublethal effects on foraging activity, with a median sublethal effect concentration (EC₅₀) of just 3.7 ppb. The same researchers also studied queenright colonies foraging in a glasshouse where food was 3 m from their nest and found that ingestion of 20 ppb of imidacloprid caused

significant worker mortality, including bees dying at the feeder. Significant mortality was also observed at 10 ppb, but not at 2 ppb. This may explain why some lab studies have failed to find effects.

With impacts more pronounced when bees have to leave the colony, one might predict more marked effects when bees are foraging naturally, travelling kilometres across the landscape (Knight et al. 2005; Osborne et al. 2008). Only four studies have examined impacts of exposure to neonicotinoids on non-*Apis* bees when free-flying in the landscape. Tasei et al. (2001) placed *Bombus lucorum* colonies in the field for 9 days, either adjacent to an imidacloprid-treated field or a control field of sunflowers. During this time, 54 % more of the foragers from the ten imidacloprid-exposed colonies failed to return compared to the ten control colonies; however, this difference was not statistically significant as sample sizes were very small. After 9 days, the colonies were returned to the lab and fed ad libitum. Treated colonies grew more slowly but the difference was not significant. Gill et al. (2012) provided *B. terrestris* colonies with feeders containing 10 ppb of imidacloprid in sugared water whilst simultaneously allowing bees freedom to forage outside the nest. Bees exposed to imidacloprid brought back pollen less often and tended to bring back smaller loads, compared to control bees. Feltham et al. (2014) simulated exposure of queenright *B. terrestris* colonies to a crop of flowering oilseed rape, providing them with sugared water and pollen containing 0.7 and 6 ppb of imidacloprid, respectively, for 2 weeks. They found a 57 % reduction in the mass of pollen brought back to colonies, which persisted for at least 4 weeks after treatment ceased. Only one study to date has attempted to examine the effects of exposure to neonicotinoids on colony-level development of bumblebees under field conditions; Whitehorn et al. (2012) used the same field-realistic doses as Feltham et al. (2014) and then allowed colonies to develop naturally in the field. They recorded significantly reduced nest growth and an 85 % decrease in queen production in imidacloprid-exposed colonies compared to control colonies. This reduction in colony performance is likely due to a combination of factors such as reduced pollen input (as demonstrated by Gill et al. 2012 and Feltham et al. 2014) and perhaps impaired fecundity of queens (following Laycock et al. 2012a, b). In an 11 week greenhouse study, caged queenright colonies of *B. impatiens* were fed treatments of 0, 10, 20, 50 and 100 ppb of imidacloprid, respectively, and clothianidin in sugar syrup (50%) (Scholer and Krischik 2014). At 6 weeks, queen mortality was significantly higher in 50 and 100 ppb and by 11 weeks in 20–100 ppb neonicotinyl-treated colonies. Starting at 20 ppb, there is a statistically significant reduction in queen survival (37 % for imidacloprid, 56 %

for clothianidin), worker movement, colony consumption and colony weight compared to 0 ppb treatments. At 10 ppb imidacloprid and 50 ppb clothianidin, fewer males were produced (Scholer and Krischik 2014).

Bryden et al. (2013) conceived a model to simulate bumblebee colony development to assess the colony-level impacts of well-known sublethal effects on individuals. Their study shows that bumblebee colonies fail when exposed to sustained sublethal levels of pesticide. This is explained by impairment of colony function. Social bee colonies have a positive density dependence, and they are subject to an Allee effect. There is a critical stress level for the success of a colony such that a small increase in the level of stress can make the difference between failure and success.

It seems likely that intoxicated bees are fully able to gather food when it is presented to them within the nest, but when bees have to navigate over realistic distances to extract nectar and pollen from complex, patchily distributed flowers, the effects of intoxication become evident. Studies have focused mainly on behavioural effects in adult bees shortly after exposure to neonicotinoids, but there is evidence from both honeybees (Yang et al. 2012) and stingless bees (Tomé et al. 2012) that exposure during larval stages can impair development of the central nervous system and, hence, result in reduced adult performance several weeks after colony exposure. Therefore, the implications for risk assessment are clear; lab trials, and even trials where colonies are placed immediately adjacent to treated crops, are not appropriate for detecting these impacts. Similarly, experiments need to run for many weeks to examine the long-term effects of exposure on bee health.

The existing toxicological data suggests that impacts on diverse bee taxa are broadly similar at the level of the individual bee, with some evidence that bumblebees and solitary bees may be more susceptible than honeybees. It is clear that field-realistic doses of neonicotinoids can have a range of significant detrimental effects on larval development, adult fecundity, adult foraging behaviour and colony performance in social species. However, the effects of neonicotinoids on the vast majority of bee species have not been examined, and caution is necessary when extrapolating from social to solitary species. No studies have evaluated the impacts of neonicotinoids on solitary species under field conditions. It might plausibly be argued that the large colony size exhibited by honeybees and some stingless bees could buffer these species against reductions in foraging performance, as well as any navigational errors on the part of workers; however, this is unlikely to be the case for either bumblebee colonies, which have just a few hundred workers at most, or solitary bees, where a single female has sole responsibility for provisioning of offspring. Thus, impacts at the population level may be inversely related to levels of sociality. This possibility awaits experimental investigation.

Butterflies and moths (Lepidoptera)

Among agricultural practices, pesticide use is known to impact butterflies and moths; however, based on observational field data, it is difficult to distinguish the impacts of pesticides from other agricultural customs, such as fertilizer application or landscape simplification, e.g. by removal of hedgerows (Geiger et al. 2010). In the case of butterflies or moths that inhabit structures adjacent to areas where pesticides are applied via aerial spraying, indirect effects of drift from spraying may pose risks both during and after applications (Sinha et al. 1990). In the 1980s for example, helicopter application of pesticides in vineyards of the Mosel Valley in Germany nearly led to the extinction of an isolated population of the Apollo butterfly (*Parnassius apollo*) which was restricted to adjacent rocky slopes (Kinkler et al. 1987; Richarz et al. 1989; Schmidt 1997). In Northern Italy, butterfly communities in natural grasslands have suffered drastic declines downwind of intensively sprayed orchards, leading to the disappearance of all but the most generalist species (Tarmann 2009). Furthermore, spray applications of pesticides may alter soil quality (Freemark and Boutin 1995) and thereby indirectly affect the larvae and pupae of moth species residing in the upper layers of the soil surface during the spring.

Contrary to other non-target species (e.g. bees, birds, spiders, ground beetles), very few comparative pesticide sensitivity tests have been carried out for butterflies and moths. This is surprising given the significant role these insects play for conservation programs. One such study conducted by Brittain et al. (2010b) evaluated the impact of pesticides on various groups of pollinators. When comparing intensively managed systems (high pesticide application rates) with less intensively managed systems (fewer pesticide applications), the authors demonstrated that fewer bumblebee and butterfly species were observed in intensively managed habitat patches. The study also demonstrated that wild bees have higher pesticide-related risks than butterflies (Brittain et al. 2010b).

Moreover, studies by Feber et al. (1997) and Rundlöf et al. (2008) have demonstrated negative impacts of pesticides on butterflies. Both studies evaluated the impacts of organic versus conventional agriculture on butterfly populations. In each case, organic farms were found to host greater numbers and species of butterflies. This response was likely due in part to reduced applications of herbicides in organic systems, as herbicides reduce the abundance of host and nectar plants that are crucial for the survival of larvae as well as adults (Boggs 2003). In contrast, similar studies comparing Lepidopteran communities between organic and conventional agriculture systems found no differences in the number or species richness of butterflies (Weibull et al. 2000 and Brittain et al. 2010a). In the case of these studies, characteristics of the surrounding landscape such as the absence of specific vegetation

elements (e.g. hedgerows or floral nectar sources) at both the local and regional scales, or the broad scale application of pesticides, may have influenced the outcome of the findings.

In contrast to the few ecological and ecotoxicological studies on the direct and indirect impacts of pesticides on non-target Lepidoptera, numerous results are available on the impacts of pesticides on the butterfly and moth species that are regarded as agricultural pests during the larval stage (Haynes 1988; Davis et al. 1991a, b, 1993; Liang et al. 2003). The impacts of systemic pesticides on Lepidoptera have been investigated for some 32 pest species of moths in nine different families (Table 2). This represents a tiny fraction of the estimated 200,000 Lepidoptera species. The results demonstrate considerable variation in the impact of pesticides on different species of Lepidoptera. For example, Doffou et al. (2011a, b) noted that the susceptibility of two cotton pests, *Pectinophora gossypiella* (Gelechiidae) and *Cryptophlebia leucotreta* (Tortricidae), to acetamiprid differs almost 3-fold (LD_{50} =11,049 and 3,798 ppm, respectively). First instar *Cydia pomonella* caterpillars (Tortricidae) are more than 100 times as sensitive as final fifth instar caterpillars, with an LC_{50}/LC_{90} of 0.84/1.83 and 114.78/462.11 ppm, respectively (Stara and Kocourek 2007a, b).

Not surprisingly, different neonicotinoid compounds vary in toxicity. Thiacloprid and acetamiprid for example are recorded to have stronger effects on the survival of *Phyllonorycter ringoniella* than all other neonicotinoid substances (Funayama and Ohsumi 2007a, b). Acetamiprid has been shown to be more toxic than thiacloprid in several studies, but the degree of difference varies greatly. For example, a study by Cichon et al. (2013) found thiacloprid to be two times as toxic to *C. pomonella* as acetamiprid (LC_{99}/LC_{50} =1.55/0.17 vs 0.71/0.08 ppm, respectively), whilst Magalhaes and Walgenbach (2011) recorded a 60-fold difference in the sensitivity of the same species to these compounds (LC_{50} =1.06 and 65.63 ppm, respectively).

Many studies have documented systemic pesticide resistance in Lepidoptera; for example, *Phthorimaea operculella* has been found to be resistant to fipronil (Doğramacı and Tingey 2007), *Spodoptera litura* to both fipronil and imidacloprid (Huang et al. 2006a, b; Ahmad et al. 2008; Abbas et al. 2012), *C. pomonella* to acetamiprid and thiacloprid (Cichon et al. 2013; Knight 2010; Stara and Kocourek 2007a, b), and *Plutella xylostella* to acetamiprid (Ninsin et al. 2000a, b). In the latter field study from Japan, an almost 10-fold higher dosage was required to reach the same lethal concentration ($LC_{50/95}$ =315/2,020 compared to 35.1/137 ppm in susceptible laboratory colonies). Applications of such high concentrations may further increase negative impacts on non-target species of insects. Even low sublethal doses can have severe impacts on Lepidoptera populations. In a study on *Helicoverpa armigera* by Ahmad et al. (2013), a

16th of the LC_{50} of imidacloprid (5.38 ppm) increased the next generation survival rate by a factor of 4 (i.e. equivalent to LC_{10}) compared to a treatment with the LC_{50} dose. Sublethal effects included a significant reduction in the survival and fecundity as well as increased mortality in the first and subsequent generations. Asaro and Creighton (2011a, b) noted that loblolly pines appeared to be protected from the Nantucket pine tip moth (*Rhyacionia frustrana*) even 1 year after treatment, and the treatment effect apparently was not confined to the target pest species, but extended to further non-target insect species.

There is a clear need for studies on the impact of pesticides on butterflies and moths and in particular those species that are not agricultural pests, but which commonly inhabit managed landscapes. Extensive studies on the direct and indirect effects of pesticides on these non-target groups are urgently needed on different geographical scales and across long time periods (Aebischer 1990) and which include all developmental stages of butterflies and moths (i.e. egg, larva, pupa, adult). It is of paramount importance to include varying intensities of pesticide applications, their persistence and their interplay with biotic and abiotic factors (Longley and Sotherton 1997; Brittain et al. 2010b).

Other invertebrates

This section will review the studies on neonicotinoids and non-target organisms, in particular the predatory invertebrates of natural pest species. Biological pest control plays an important role in integrated pest management (Byrne and Toscano 2007; Peck and Olmstead 2010; Prabhaker et al. 2011; Khani et al. 2012) with studies suggesting that predators may contribute to the similarity in crop yields between non-treated and pesticide-treated fields (Albajes et al. 2003; Seagraves and Lundgren 2012).

Routes of exposure

Non-target organisms can be exposed to neonicotinoid pesticides in a variety of ways. Predatory invertebrates may become contaminated by consuming pests such as leafhoppers or aphids that feed on treated crops (Albajes et al. 2003; Papachristos and Milonas 2008; Moser and Obrycki 2009; Prabhaker et al. 2011; Khani et al. 2012). Direct contamination through the diet can also be a problem for other beneficial plant-feeding invertebrates (Dilling et al. 2009; Girolami et al. 2009; Moser and Obrycki 2009; Prabhaker et al. 2011; Khani et al. 2012). For example, several species of hoverfly and parasitoid wasps attack agricultural pests, but also subsidise their diet with nectar. Therefore, these insects can be affected by neonicotinoids, which are translocated into the nectar and pollen of treated crop plants (Stapel et al. 2000; Krischik et al. 2007).

Table 2 Studies on the effects of systemic pesticides in Lepidoptera

Family	Species	Host	Imidacloprid	Thiamethoxam	Clothianidin	Acetamiprid	Thiacloprid	Dinotefuran	Fipronil
Gelechiidae	<i>Pectinophora gossypiella</i>	Cotton				Doffou et al. (2011a, b)			
Gelechiidae	<i>Plithorinaca operculella</i>	Potato	Symington (2003)				Saour (2008)		Dogramaci and Tingey (2008)
Gracillariidae	<i>Camieria ohridella</i>	Horse chestnut tree	Stygar et al. (2013)						
Gracillariidae	<i>Phyllocnistis citrella</i>	Citrus	Villanueva-Jimenez and Hoy (1998), Setamou et al. (2010)						
Gracillariidae	<i>Phyllonorycter ringoniella</i>	Apple	Funayama and Ohsumi (2007a, b)	Funayama and Ohsumi (2007a, b)	Funayama and Ohsumi (2007a, b)	Funayama and Ohsumi (2007a, b)		Funayama and Ohsumi (2007a, b)	
Lyonetiidae	<i>Leucopiera coffeella</i>	Coffee		Diez-Rodriguez et al. (2006)					
Noctuidae	<i>Agrotis ipsilon</i>	Corn and various crops			Kullik et al. (2011a)				
Noctuidae	<i>Helicoverpa armigera</i>	Various crops	Ahmad et al. (2013)						
Noctuidae	<i>Helicoverpa zea</i>	Cotton	Kilpatrick et al. (2005)	Kilpatrick et al. (2005)		Kilpatrick et al. (2005)			Pedibhotla et al. (1999)
Noctuidae	<i>Heliothis virescens</i>	Tobacco							
Noctuidae	<i>Lacanobia subijuncta</i>	Apple and various fruits		Brunner et al. (2005)	Brunner et al. (2005)				Fang et al. (2008)
Noctuidae	<i>Sesamia inferens</i>	Rice							Ahmad et al. (2008), Huang et al. (2006a, b)
Noctuidae	<i>Spilarcia obliqua</i>	Polyphagous	Ansari et al. (2012)						
Noctuidae	<i>Spodoptera litura</i>	Polyphagous	Abbas et al. (2012)						
Psychidae	<i>Thyridopheryx ephemeriformis</i>	Thuja and other ornamental plants			Rhainds and Sadof (2009)			Rhainds and Sadof (2009)	
Pyrilidae	<i>Acrobasis vaccinii</i>	Blueberry							Mann et al. (2009)
Pyrilidae	<i>Cacablastis cactorum</i>	Opuntia	Bloem et al. (2005)			Wise et al. (2010)	Wise et al. (2010)		Fang et al. (2008), He et al. (2013), Chen and Klein (2012), Cheng et al. (2010), He et al. (2007, 2008), Li et al. (2007)
Pyrilidae	<i>Chilo infuscatellus</i>	Sugarcane							
Pyrilidae	<i>Chilo suppressalis</i>	Rice	Yu et al. (2007a, b)						
Pyrilidae	<i>Ostrinia nubilalis</i>	Stored grain	Yue et al. (2003)	Yu et al. (2007a, b)					Durham et al. (2001, 2002), Siegfried et al. (1999)
Pyrilidae	<i>Plodia interpunctella</i>	Stored grain	Yue et al. (2003)	Yue et al. (2003)					
Pyrilidae	<i>Tryporyza incertulas</i>	Rice	Wang et al. (2005)						
Sesiidae	<i>Pennisetia marginata</i>	Raspberry	McKern et al. (2007)						

Table 2 (continued)

Family	Species	Host	Imidacloprid	Thiamethoxam	Clothianidin	Acetamiprid	Thiacloprid	Dinotefuran	Fipronil
Tortricidae	<i>Choristoneura rosaceana</i>	Apple		Brunner et al. (2005)	Brunner et al. (2005)	Brunner et al. (2005), Dunley et al. (2006)			
Tortricidae	<i>Cryptophlebia leucolenta</i>	Cotton				Doffou et al. (2011a, b)			
Tortricidae	<i>Cydia pomonella</i>	Apple		Brunner et al. (2005)	Brunner et al. (2005)	Brunner et al. (2005), Cichon et al. (2013), Magalhaes and Walgenbach (2011), Knight (2010), Magalhaes and Walgenbach (2011), Mota-Sanchez et al. (2008)	Cichon et al. (2013), Magalhaes and Walgenbach (2011), Stara and Kocourek (2007), Voudouris et al. (2011), Reyes et al. (2007)		
Tortricidae	<i>Epiphyas postvittana</i>	Trees	Taverner et al. (2012)						
Tortricidae	<i>Grapholita lobarzewskii</i>	Apples	Charmillot et al. (2007)					Taverner et al. (2011, 2012)	
Tortricidae	<i>Grapholita molesta</i>	Apple		Jones et al. (2012)			Magalhaes and Walgenbach (2011)	Charmillot et al. (2007)	
Tortricidae	<i>Pandemis pyrusana</i>	Apple							
Tortricidae	<i>Rhyacionia frustrana</i>	Pine trees		Brunner et al. (2005)	Brunner et al. (2005)	Brunner et al. (2005), Dunley et al. (2006)			Asaro and Creighton (2011)
Yponomeutidae	<i>Plutella xylostella</i>	Cabbage	Asaro and Creighton (2011a, b) Hill and Foster (2000)			Ninsin et al. (2000a, b), Sayyed and Crickmore (2007), Ninsin and Tanaka (2005), Ninsin (2004a, b), Ninsin and Miyata (2003)			Li et al. (2006), Sayyed and Wright (2004), Shi et al. (2004), Zhou et al. (2011)

Other routes of exposure include contact with treated surfaces, exposure to sprays or consumption of guttation droplets (Papachristos and Milonas 2008; Prabhaker et al. 2011; Khani et al. 2012). For example, neonicotinoid soil drenches or injections have been found to adversely affect the development of Lepidoptera larvae pupating within the soil (Dilling et al. 2009), whilst soil drenches have been found to significantly lower the overall abundance of insect species and species richness. In one study, imidacloprid was used on eastern hemlock (*Tsuga canadensis*) to effectively control the hemlock woolly adelgid (*Adelges tsugae*); however, the abundance of non-target detritivorous, fungivorous and phytophagous invertebrates was significantly lower in soil drench and injection treatments, when compared to untreated plots (Dilling et al. 2009).

Parasitoid wasps such as *Gonatocerus ashmeadi* can come into contact with neonicotinoids when emerging from the eggs of its host. One such host, the glassy-winged sharpshooter (*Homalodisca itripennis*), a common agricultural pest of many different crops, lays its eggs on the underside of leaves, beneath the epidermal layer. If eggs are laid on neonicotinoid-treated plants, *G. ashmeadi* nymphs may be exposed to toxins when they emerge from the egg and chew through the leaf to get to the surface (Byrne and Toscano 2007).

A 3 year study by Peck (2009) found that when imidacloprid was used as a lawn treatment to target neonate white grubs (Coleoptera: Scarabaeidae), it exhibited cumulative detrimental effects on the abundance of Hexapods, Collembola, Thysanoptera and Coleoptera adults, which were suppressed by 54–62 % overall throughout the course of the study. Population numbers of non-target organisms can also be indirectly affected by a reduction in prey or host species (Byrne and Toscano 2007; Dilling et al. 2009).

Diptera

Of the Diptera, the genus *Drosophila* provides well-known and convenient model species for toxicity testing. Mechanisms of resistance to imidacloprid and its metabolism have been studied in *Drosophila melanogaster*. Particularly, cytochrome P450 monooxygenases (CYPs) are involved, as is the case in mosquitoes (Riaz et al. 2013). According to Kalajdzic et al. (2012), three P450 genes (Cyp4p2, Cyp6a2 and Cyp6g1)

located on the 2R chromosome were highly up-regulated in imidacloprid-resistant flies. However, the same authors did not find that imidacloprid induced expression of Cyp6g1 and Cyp6a2 (Kalajdzic et al. 2013). More recently, it has been shown that imidacloprid was metabolized to eight derivatives in *D. melanogaster*. In this process, only the P450 Cyp6g1 was involved in the enhanced metabolism in vivo (Hoi et al. 2014). Direct toxicity (LC₅₀) has been determined for various *D. melanogaster* strains. For instance, the toxicity of several neonicotinoids and butene-fipronil was evaluated (Arain et al. 2014) with neonicotinoids being less toxic than butene-fipronil. It was suggested that differences exist between adults and larvae. Acute LC₅₀ values can be compared to LC₅₀ measured after chronic exposure, within two studies. With a mutant strain, Frantzios et al. (2008) found a decrease by a factor of 2 for adult flies (acute vs chronic) and a factor of 3 for larvae. Very recently, Charpentier and co-workers have distinguished between male and female flies, from a field strain (Charpentier et al. 2014). Here, the chronic LC₅₀ was 29 times lower than the acute LC₅₀ for males; it was 172 times lower for females and 52 times lower for larvae. Additionally, this study demonstrated that a significant increase of mortality (27–28 %), with a V-shape, was occurring at concentrations 1,100 and 4,600 times lower than the chronic LC₅₀ for males and females, respectively. Other parameters that are crucial for reproduction were tested (mating and fecundity). The LOEC was determined at a concentration that is 3,300,000 and more than 7,900,000 times lower than the acute LC₅₀ for females and males, respectively. These data can be linked to data concerning mortalities observed by chronic exposure of bees at very low concentrations.

Hymenoptera (excluding bees)

A few studies have investigated the effect of neonicotinoid pesticides on parasitic wasps used as biological control agents. Stapel et al. (2000) found that the parasitoid wasp *Microplitis croceipes* had significantly reduced foraging ability and longevity after feeding on extrafloral nectar of cotton (*Gossypium hirsutum*) treated with imidacloprid. Prabhaker et al. (2007) give acute toxicity for two different exposure times for the parasitic wasp species *Eretmocerus eremicus*, *Encarsia formosa*, *Aphytis melinus* and *G. ashmeadi* (Table 3).

Table 3 Acute neonicotinoid toxicity for different Hymenoptera species (Prabhaker et al. 2007)

Species	48 h exposure time mg (AI)/ml		24 h exposure time mg (AI)/ml
	Acetemiprid	Thiamethoxam	Imidacloprid
<i>Eretmocerus eremicus</i>	108.27	1.01	1.93
<i>Encarsia formosa</i>	12.02	0.397	0.980
<i>Gonatocerus ashmeadi</i>	0.134	1.44	2.63
<i>Aphytis melinus</i>	0.005	0.105 (24 h exposure time)	0.246

In another study, *Anagyrus pseudococci* (a nectar-feeding wasp) was fed using buckwheat (*Fagopyrum esculentum*) flowers that had been exposed to imidacloprid as a soil treatment, applied at the label rate. Only 38 % of the wasps survived after 1 day, compared to 98 % fed on untreated flowers. This decreased to 0 % survivorship after 7 days for treated flowers, compared to 57 % on the untreated flowers (Krischik et al. 2007).

As stated in the section on exposure routes, exposure to imidacloprid did not affect mortality of *G. ashmeadi* (a parasitoid wasp) during development within its host, and the adults were sensitive during emergence from the host egg. When mortality was assessed within 48 h of emergence, the LC₅₀ for the parasitoid was 66 ng of imidacloprid per cm² leaf (Byrne and Toscano 2007).

Neonicotinoids are commonly used in household products as highly concentrated bait formulations to control ants (Rust et al. 2004; Jeschke et al. 2010); however, the use of agrochemical products at less concentrated doses is an issue for non-target ants. For the leafcutter ant *Acromyrmex subterraneus subterraneus*, Galvanho et al. (2013) found that sublethal doses of imidacloprid reduced grooming behaviour. Grooming behaviour in this ant is a defence against pathogenic fungi like *Beauveria* species. Barbieri et al. (2013) recently discovered that interactions between different ant species may be negatively affected using sublethal doses of neonicotinoids. In interspecific interactions, individuals of a native ant species (*Monomorium antarcticum*) lowered their aggression towards an invasive ant species (*Linepithema humile*) although survival was not affected. Exposed individuals of *L. humile* displayed an increase in aggression with the outcome that the probability of survival was reduced.

Hemiptera

Whilst many Hemiptera are acknowledged as being problematic agricultural pests, a number are important predators of these pests, although they do also feed on some plant tissues, which would be contaminated by neonicotinoids (Prabhaker

et al. 2011). Table 4 shows LC₅₀ rates for different Hemiptera species.

Neuroptera

It is not only the agricultural use of neonicotinoids that affects beneficial invertebrates. In one study, Marathon 1 % G, a product for amateur use on flowers containing imidacloprid, had been found to affect lacewings (*Chrysopa* spp.) when used at the label rate. Survival rates on untreated flowers were found to be 79 % for adults, compared to 14 % on treated flowers over a 10 day period (Rogers et al. 2007).

Coleoptera

A number of studies have looked into the effects of neonicotinoids on various taxa of Coleoptera such as Histeridae (Hister beetles) (Kunkel et al. 1999), Carabidae (ground beetles) (Kunkel et al. 2001; Mullin et al. 2010) and Coccinellidae (ladybird beetles) (Smith and Krischick 1999; Youn et al. 2003; Lucas et al. 2004; Papachristos and Milonas 2008; Moser and Obrycki 2009; Eisenback et al. 2010; Khani et al. 2012).

Some Coleoptera, notably in the carabid and staphyliniid families, are voracious predators and are a vital aspect of integrated pest management. For example, although the provision of beetle banks as nesting habitat takes land out of crop production, in wheat (*Triticum aestivum*) fields, any losses have been found to be more than offset by savings from a reduced need for aphid-controlling pesticides (Landis et al. 2000).

Many of these beetle groups are undergoing rapid declines. In the UK, three quarters of carabid species have reduced in numbers, half of which have been undergoing population declines of more than 30 %, although the reason for these considerable declines are unknown (Brooks et al. 2012). These groups have been particularly useful as bioindicators, due to their sensitivity to habitat changes especially in agricultural environments (Kromp 1999; Lee et al. 2001). In the EU Draft Assessment Report for imidacloprid, acute toxicity tests were

Table 4 LC₅₀ rates for different Hemiptera species

Species	Chemical	LC ₅₀ residual contact (mg AI/l)		Reference
		Nymphs	Adults	
<i>Orius Laevigatus</i>	Imidacloprid	0.04	0.3	Delbeke et al. (1997)
<i>Hyaliodes vitripennis</i>	Thiacloprid	1.5	0.3	Bostanian et al. (2005)
<i>Hyaliodes vitripennis</i>	Thiamethoxam	1.43	0.5	Bostanian et al. (2005)
<i>Geocoris punctipes</i>	Imidacloprid		5,180	Prabhaker et al. (2011)
	Thiamethoxam		2,170	
<i>Orius insidiosus</i>	Imidacloprid		2,780	
	Thiamethoxam		1,670	

undertaken on the carabid beetle *Poecilus cupreus*, finding the larvae to be highly sensitive. Despite the rapporteur Member State deeming that the concentrations tested were too high for it to conclude no risk to carabids for use on sugar beet, there was no indication of further research required (EFSA 2006).

When exposed to turf plots treated with imidacloprid, the carabid beetle *Harpalus pennsylvanicus* displayed a range of neurotoxic problems including paralysis, impaired walking and excessive grooming. These abnormal behaviours then rendered the individuals vulnerable to predation from ants (Kunkel et al. 2001). A study by Mullin et al. (2010) exposed 18 different carabid species to corn seedlings treated to field-relevant doses of either imidacloprid, thiamethoxam or clothianidin. Nearly 100 % mortality was observed for all species over 4 days.

Coccinellids predators are well known for their ability to control common pests, both in agricultural and domestic environments. In soil treatments of imidacloprid, reduced mobility and delayed reproduction have been found in pollen-feeding species such as *Coleomegilla maculata* (Smith and Krischick 1999), whilst egg production and oviposition periods of the Mealybug destroyer (*Cryptolaemus montrouzieri*) (Khani et al. 2012) and *Hippodamia undecimnotata* (Papachristos and Milonas 2008) were significantly reduced. Table 5 shows available acute toxicity for some coccinellid species.

Harmonia axyridis (harlequin ladybird) larvae were exposed to corn seedlings grown from seeds treated with the label recommended doses of either thiamethoxam or clothianidin. Seventy-two percent of the larvae exhibited neurotoxic symptoms such as trembling, paralysis and loss of coordination, with only 7 % recovery from the poisoning (Moser and Obrycki 2009).

Arachnida

In addition to crop protection, applications of neonicotinoid insecticides in veterinary medicine have expanded. Imidacloprid is applied to domestic pets as a

spot-on formulation against ear mites (*Otodectes cynotis*) (Jeschke et al. 2010). However, studies on mites have found a positive effect on population numbers. Zeng and Wang (2010) found that sublethal doses of imidacloprid (determined for the green peach aphid (*Myzus persicae*)) significantly increased the hatch rate of eggs and pre-adult survivorship of the carmine spider mite (*Tetranychus cinnabarinus*). James and Price (2002) also found that imidacloprid increased egg production by 23–26 % in two-spotted spider mites (*Tetranychus urticae*) in the laboratory. Another study found that fecundity of this species was slightly elevated when treated with thiamethoxam (Smith et al. 2013).

Szczepaniec et al. (2013) discovered that the application of neonicotinoids suppressed expression of plant defence genes when applied to cotton and tomato plants. These genes alter the levels of phytohormones and decrease the plant's resistance to spider mites (*T. urticae*). When mites were added to the crops, population growth increased from 30 to over 100 % on neonicotinoid-treated plants in the greenhouse and up to 200 % in the field experiment. This study was prompted after the same author had investigated an outbreak of *T. urticae* in New York City, USA. In an attempt to eradicate the emerald ash borer beetle (*Agrillus planipennis*) from Central Park, imidacloprid was applied to trees as a soil drench and trunk injections. This resulted in an outbreak of *T. urticae* on elms due to the natural predators being poisoned through ingestion of prey exposed to imidacloprid, combined with fecundity elevation in the mites themselves (Szczepaniec et al. 2011).

Another study found that thiamethoxam and imidacloprid treatments significantly increased two-spotted spider mite (*T. urticae*) densities on cotton plants when compared to the untreated controls (Smith et al. 2013). This study suggested that the increased usage of neonicotinoids could explain the recent infestation increases of two-spotted spider mite occurring in various crops across the mid-south of the USA.

Table 5 Acute neonicotinoid toxicity for different Coccinellid species

Species	Chemical	LD ₅₀ (ng AI per beetle)	LC ₅₀ (µg AI/ml)	Reference
<i>Sasajiscymnus tsugae</i>	Imidacloprid	0.71		Eisenback et al. (2010)
<i>Harmonia axyridis</i>	Imidacloprid		364	Youn et al. (2003)
<i>Harmonia variegata</i>	Thiamethoxam		788.55	Rahmani et al. (2013)
<i>Cryptolaemus montrouzieri</i>	Imidacloprid		17.25–23.9	Khani et al. (2012)
<i>Coccinella undecimpunctata</i>	Imidacloprid		34.2	Ahmad et al. (2011)
<i>Coccinella undecimpunctata</i>	Acetamiprid		93.5	Ahmad et al. (2011)
<i>Coleomegilla maculata</i> —adult	Imidacloprid	0.074		Lucas et al. (2004)
<i>Coleomegilla maculata</i> —larvae	Imidacloprid	0.034		Lucas et al. (2004)

Earthworms (Lumbricidae)

Earthworms are vitally important members of the soil fauna, especially in agricultural soils where they can constitute up to 80 % of total soil animal biomass (Luo et al. 1999). They play critical roles in the development and maintenance of soil physical, chemical and biological properties (Lee 1985). Their activities improve soil structure by increasing porosity and aeration, facilitating the formation of aggregates and reducing compaction (Edwards and Bohlen 1996; Mostert et al. 2000). Soil fertility is enhanced by earthworm effects on biogeochemical cycling (Coleman and Ingham 1988; Bartlett et al. 2010), the modification of microbial biomass and activity (Sheehan et al. 2008), breakdown of plant litter (Knollengberg et al. 1985) and the mixing of litter with soil (Wang et al. 2012a).

Neonicotinoid and other systemic insecticides can pose a risk of harm to earthworm survival and behaviour, potentially disrupting soil development and maintenance processes. The same neural pathways that allow neonicotinoids to act against invertebrate pests (Elbert et al. 1991) are also present in earthworms (Volkov et al. 2007). Thus, when neonicotinoids are applied for the protection of agricultural and horticultural crops, earthworms can be exposed by direct contact with the applied granules or seeds, or with contaminated soil or water. Moreover, their feeding activities may result in ingestion of contaminated soil and organic particles (e.g. Wang et al. 2012b). Foliar residues in plant litter after systemic uptake from soils or from direct plant injections also pose a risk to litter-feeding earthworms that consume the contaminated plant litter (e.g. Kreutzweiser et al. 2009).

Neonicotinoids can persist and move in soils thereby increasing the likelihood that earthworms will be exposed for extended periods of time. Laboratory and field trials with neonicotinoids have demonstrated that their half-life in soils varies depending on soil conditions but can range from several weeks to several years (Cox et al. 1997; Sarkar et al. 2001; Cox et al. 2004; Bonmatin et al. 2005; Fossen 2006; Gupta and Gajbhiye 2007; Goulson 2003). Imidacloprid is the most widely used neonicotinoid, and its adsorption to soils is increased by moisture and organic matter content (Broznic et al. 2012), resulting in increased imidacloprid concentrations in organic-rich soils compared to low-organic soils (Knoepp et al. 2012). Earthworms generally prefer moist, organic-rich soils. When soil organic carbon content is low, the high solubility of imidacloprid renders it mobile and it is readily moved through soils (Broznic et al. 2012; Knoepp et al. 2012; Kurwadkar et al. 2013), thereby increasing the likelihood that earthworms could be exposed to the pesticide in soils outside the direct area of application.

Effects on survival

Neonicotinoids can be highly toxic to earthworms. However, reported median lethal concentrations (LC₅₀) were variable depending on the particular insecticide, test conditions, route of exposure and duration (Table 6). In 13 separate studies, the reported LC₅₀ ranged from 1.5 to 25.5 ppm, with a mean of 5.8 and median of 3.7 ppm. In seven studies that reported lowest concentrations at which effects on survival were measureable, those lowest effective concentrations ranged from 0.7 to 25 ppm, with a mean of 4.7 and median of 1.0 ppm. *Eisenia fetida* was the most common test species in these survival studies and represented the range of reported lethal concentrations, giving little indication from among these studies that other species were more sensitive than *E. fetida*.

When compared to other common insecticides, neonicotinoids tend to be among the most toxic to earthworms. Wang et al. (2012a) tested the acute toxicities of 24 insecticides to *E. fetida* and found that the neonicotinoids were the most toxic in soil bioassays and that acetamiprid and imidacloprid in particular were the two most toxic insecticides overall. They also reported that a contact toxicity bioassay demonstrated that the neonicotinoids were extremely toxic by a contact route of exposure (LC₅₀ of 0.0088 to 0.45 µg cm⁻²), although the units of contact toxicity concentration were difficult to compare to standard lethal concentrations. Across a broader range of 45 pesticides, Wang et al. (2012b) found that in soil bioassays, the neonicotinoid insecticide, clothianidin, was the most toxic pesticide to *E. fetida*. Alves et al. (2013) compared three insecticides used for seed treatment and reported that imidacloprid was the most toxic to earthworms. In soil bioassays with five different insecticides, Mostert et al. (2002) found that imidacloprid was the second most toxic (behind carbaryl) to earthworms. We found only two studies that reported the toxicity of fipronil, another common, agricultural systemic insecticide, and both found it to be substantially (at least 100 times) less lethal to earthworms than the neonicotinoids (Mostert et al. 2002; Alves et al. 2013).

Effects on reproduction

Only a few studies tested sublethal effects of neonicotinoids on earthworm reproduction, but it is apparent that reductions in fecundity can occur at low concentrations (Table 6). Baylay et al. (2012) reported EC₅₀s for imidacloprid and thiacloprid against cocoon production by *Lumbricus rubellus* of 1.5 and 1.3 ppm, respectively, whilst Gomez-Eyles et al. (2009) found similar EC₅₀s for the same two insecticides at 1.4 and 0.9 ppm for *E. fetida*. The latter study also reported measurable reductions in cocoon production at 0.3 ppm of thiacloprid. Alves et al. (2013) reported an EC₅₀ for reproduction effects of imidacloprid on *Eisenia andrei* of 4 ppm with measureable

Table 6 Impacts of neonicotinoids and fipronil on earthworms. The impact rating scale is as follows: —, large decrease; —, moderate decrease; 0, little or no measurable effect (where little is either a small or a brief change); +, moderate increase; and ++, large increase. Endpoints are listed together, separated by a semi-colon, for studies that examined multiple measurement endpoints. Lowest effective concentration is the lowest concentration at which a significant effect was reported, not necessarily the mathematically modelled lowest effective concentration

Taxa	Insecticides	Location	Measurement endpoint	Impact	LC/EC ₅₀	Lowest effective concentration	Reference
<i>Eisenia fetida</i>	Imidacloprid	China	Contact toxicity survival; soil toxicity survival	—; —	LC ₅₀ =0.027 µg cm ⁻² ; LC ₅₀ =2.82 ppm		Wang et al. (2012a)
<i>Eisenia fetida</i>	Imidacloprid	France	Survival; biochemical (hsp70); avoidance	—; —; ++		0.66; 0.66; 0.2 ppm	Dittbrenner et al. (2012)
<i>Eisenia fetida</i>	Imidacloprid	France	Survival; body mass	—; —		0.66; 0.2 ppm	Dittbrenner et al. (2011a)
<i>Eisenia fetida</i>	Imidacloprid	UK	Cocoon production; weight change	—; —	EC ₅₀ =1.41; EC ₅₀ =2.77 ppm		Gomez-Eyles et al. (2009)
<i>Eisenia fetida</i>	Imidacloprid	China	Survival	—	LC ₅₀ =2.30 ppm		Zang et al. (2000)
<i>Eisenia fetida</i>	Imidacloprid	China	Survival	—	LC ₅₀ =2.30 ppm	1 ppm	Luo et al. (1999)
<i>Eisenia fetida</i>	Imidacloprid	Canada	Survival; weight loss	—; —		25; 14 ppm	Kreutzweiser et al. (2008b)
<i>Eisenia fetida</i>	Fipronil	Brazil	Survival; reproduction; avoidance	0; —; +		>1,000; 62; >10 ppm	Alves et al. (2013)
<i>Eisenia fetida</i>	Clothianidin	China	Contact toxicity survival; soil toxicity survival	—; —	LC ₅₀ =0.28 µg cm ⁻² ; LC ₅₀ =6.06 ppm		Wang et al. (2012b)
<i>Eisenia fetida</i>	Thiacloprid	China	Contact toxicity survival; soil toxicity survival	—; —	LC ₅₀ =0.45 µg cm ⁻² ; LC ₅₀ =10.96 ppm		Wang et al. (2012a)
<i>Eisenia fetida</i>	Thiacloprid	UK	Cocoon production; weight change	—; —	EC ₅₀ =0.968; EC ₅₀ =19.0 ppm	0.291; 1.91 ppm	Gomez-Eyles et al. (2009)
<i>Eisenia fetida</i>	Acetamiprid	China	Contact toxicity survival; soil toxicity survival	—; —	LC ₅₀ =0.0088 µg cm ⁻² ; LC ₅₀ =1.52 ppm		Wang et al. (2012a)
<i>Eisenia fetida</i>	Nitenpyram	China	Contact toxicity survival; soil toxicity survival	—; —	LC ₅₀ =0.22 µg cm ⁻² ; LC ₅₀ =3.91 ppm		Wang et al. (2012a)
<i>Lumbricus terrestris</i>	Imidacloprid	France	Survival; biochemical (hsp70); avoidance	0; +; 0		4 ppm	Dittbrenner et al. (2012)
<i>Lumbricus terrestris</i>	Imidacloprid	France	Survival; body mass	0; —		2 ppm	Dittbrenner et al. (2011b)
<i>Lumbricus terrestris</i>	Imidacloprid	USA	Feeding activity; abundance	—; —		43 mg m ⁻²	Tu et al. (2011)
<i>Lumbricus terrestris</i>	Imidacloprid	France	Burrowing	—		2 ppm	Dittbrenner et al. (2011b)
<i>Lumbricus terrestris</i>	Imidacloprid	France	Body mass change; cast production	—; —	NA; EC ₅₀ =0.84 ppm	0.66; 0.66 ppm	Dittbrenner et al. (2010)
<i>Lumbricus terrestris</i>	Imidacloprid	France	Cast production; body mass change	—; —	LC ₅₀ =10.7 ppm	1.89; 0.189 ppm	Capowiez et al. (2010)
<i>Lumbricus terrestris</i>	Imidacloprid and thiacloprid mixture	UK	Survival; weight change; cocoon production; metabolism	0; —; —; 0	EC ₅₀ imidacloprid=1.46 and EC ₅₀ thiacloprid=1.28 ppm		Baylay et al. (2012)
<i>Aporrectodea caliginosa</i>	Imidacloprid	France	Survival; biochemical (hsp70); avoidance	0; —; ++		2; 2 ppm	Dittbrenner et al. (2012)
<i>Aporrectodea caliginosa</i>	Imidacloprid	France	Survival; body mass	—; —		2; 0.66 ppm	Dittbrenner et al. (2011a)
<i>Aporrectodea caliginosa</i>	Imidacloprid	France	Burrowing	—		0.2 ppm	Dittbrenner et al. (2011b)
<i>Aporrectodea caliginosa</i>	Imidacloprid	France	Body mass change; cast production	—; —	NA; EC ₅₀ =0.76 ppm	0.66; 0.66 ppm	Dittbrenner et al. (2010)
<i>Aporrectodea nocturna</i>	Imidacloprid	France	Weight loss; avoidance; burrowing	—; +; —		0.5; 0.1; 0.05 ppm	Capowiez and Berard (2006)
<i>Aporrectodea nocturna</i>	Imidacloprid	France	Burrowing	—		0.1 ppm	Capowiez et al. (2006)
<i>Aporrectodea nocturna</i>	Imidacloprid	France	Survival, weight loss	—; —	LC ₅₀ =3.74 ppm	0.1 ppm	Capowiez et al. (2005)
<i>Aporrectodea nocturna</i>	Imidacloprid	France	Burrowing	—		0.01 ppm	Capowiez et al. (2003)
<i>Allobophora icterica</i>	Imidacloprid	France	Weight loss; avoidance; burrowing	—; +; —		0.5; 0.01; 0.05 ppm	Capowiez and Berard (2006)
<i>Allobophora icterica</i>	Imidacloprid	France	Burrowing	—		0.1 ppm	Capowiez et al. (2006)
<i>Allobophora icterica</i>	Imidacloprid	France	Survival, weight loss	—; —	LC ₅₀ =2.81 ppm	0.1 ppm	Capowiez et al. (2005)
<i>Allobophora icterica</i>	Imidacloprid	France	Burrowing	—		0.01 ppm	Capowiez et al. (2003)
<i>Dendrobaena octaedra</i>	Imidacloprid	Canada	Survival; leaf decomposition	0; —		31 ppm	Kreutzweiser et al. (2009)
<i>Dendrobaena octaedra</i>	Imidacloprid	Canada	Survival; weight loss; reproduction; leaf decomposition	—; —; —; —	LC ₅₀ =5.7 ppm	3; 3; 7; 7 ppm	Kreutzweiser et al. (2008b)

Table 6 (continued)

Taxa	Insecticides	Location	Measurement endpoint	Impact	LC/EC ₅₀	Lowest effective concentration	Reference
<i>Dendrobaena octaedra</i>	Imidacloprid	Canada	Survival; weight loss; reproduction; leaf decomposition	0; –; 0; –		11; 3.2 ppm	Kreutzweiser et al. (2008a)
<i>Eisenia andrei</i>	Imidacloprid	Brazil	Survival; reproduction; avoidance	–; –; ++	LC ₅₀ =25.53; EC ₅₀ =4.07; EC ₅₀ =0.11 mg/kg	25; 0.75; 0.13 ppm	Alves et al. (2013)
Pheretima group	Imidacloprid	South Africa	Survival	–	LC ₅₀ =3.0 ppm		Mostert et al. (2002)
Pheretima group	Fipronil	South Africa	Survival	0		>300 ppm	Mostert et al. (2002)
<i>Apporectodea</i> spp.	Clothianidin	USA	Abundance; biomass; cast production	–; –; –	NA, field applications		Larson et al. (2012)

adverse effects at 0.7 ppm. Kreutzweiser et al. (2008b) tested the effects of imidacloprid in forest litter on the litter-dwelling earthworm *Dendrobaena octaedra* and reported significant reductions in cocoon production among surviving earthworms at 7 ppm.

Effects on behaviour

A number of studies focused on behavioural endpoints under the premise that effects on behaviour are often ultimately linked to population or community effects (Little 1990; Dittbrenner et al. 2012). The behavioural attributes considered here are avoidance behaviour, burrowing, cast production and weight change (as an indicator of feeding behaviour). Among the 31 reported values for behavioural effects, weight change was the most common, followed by burrowing, avoidance behaviour and cast production (Table 6). Only a few studies gave median effective concentrations (EC₅₀), and they ranged from 0.1 (avoidance) to 19 (weight change) ppm, with a mean EC₅₀ of 3.7 and median of 1.3 ppm. These behavioural EC₅₀s were about 1.5 to 2.8 times lower than the mean and median lethal concentrations of 5.8 and 3.7 ppm.

However, many more studies reported lowest concentrations at which behavioural effects were detected, and those ranged from 0.01 to 14 ppm with a mean of 1.2 and median of 0.5 ppm. Thus, measurable behavioural effects were more sensitive endpoints than measurable survival effects. Measurable behavioural effects occurred at concentrations of about two to four times lower than the mean and median lowest effective concentrations on survival of 4.7 and 1.0 ppm. Burrowing (smaller, shorter, more narrow burrows) was the most sensitive behavioural endpoint with effects detected at mean and median concentrations of 0.3 and 0.07 ppm (range 0.01 to 2, *n*=8). Avoidance behaviour was the next most sensitive endpoint with effects detected at mean and median concentrations of 0.5 and 0.13 ppm (*n*=5), followed by cast production (mean 1.1, median 0.7 ppm, *n*=3) and weight change (mean 2.1, median 0.7 ppm, *n*=13). All of these indicate that measurable adverse effects on earthworm behaviour would be expected at neonicotinoid concentrations below 1 ppm in soil.

Risks to earthworms

The actual risk of harmful effects on earthworm populations posed by neonicotinoid insecticides will depend on exposure concentration, exposure duration, route of exposure, rate of uptake and inherent species sensitivity. From the toxicity studies reviewed here, it appears that individual earthworms across all common species are at risk of mortality if they consume soil or organic particles with neonicotinoid insecticide concentrations of about 1 ppm or higher for several days. Higher numbers (up to 50 %) of earthworms would be

expected to be at risk of mortality when concentrations reach about 3 ppm and higher. Although it was difficult to compare the exposure concentrations to standard bioassays, it appears that the risk of mortality from surface contact exposure can be ten times or more higher than the risk of mortality from consumption of contaminated soils (Wang et al. 2012a). On the other hand, the route of exposure can affect the likelihood of lethal effects on earthworms. When earthworms were exposed to foliar residues in leaf litter from imidacloprid-injected trees, a significant feeding inhibition effect was detected that reduced leaf consumption but did not cause earthworm mortality, even at concentrations of about 10 ppm (Kreutzweiser et al. 2008a).

The risk of sublethal effects on some important behavioural attributes is higher than the risk of mortality to individuals. Insecticide effects on burrowing and avoidance behaviours would be expected at concentrations of about 0.1 to 0.5 ppm and higher. Whilst alterations in burrowing behaviour, especially reductions in burrowing depths, have implications for the transfer properties of soils (Capowiez et al. 2006; Dittbrenner et al. 2011b), the consequences in real-world field conditions are not clear. Fewer, smaller and shorter burrows could reduce air, water and solute transport through soils affecting overall soil ecology, but none of the studies we found actually tested these implications in experimental or field settings.

The concentrations that pose risk of mortality (assuming high toxicity by contact exposure) and sublethal effects on earthworms fall within the range of reported field concentrations, albeit at the upper end of that range of concentrations. Dittbrenner et al. (2011b) indicate that predicted environmental concentrations for imidacloprid in agricultural soils would be about 0.3 to 0.7 ppm, suggesting risks of at least sublethal effects on earthworms could be quite high. Bonmatin et al. (2005) reported that imidacloprid in soils can reach several hundred parts per billion shortly after sowing of treated seeds. Soil samples from a tea plantation treated with clothianidin had average concentrations of up to 0.45 ppm shortly after application (Chowdhury et al. 2012). Donnarumma et al. (2011) found concentrations of imidacloprid in soils at about 0.6 to 0.8 ppm by 2 weeks after application of treated seeds. Ramasubramanian (2013) reported clothianidin concentrations in soils of 0.27 to 0.44 ppm up to 3 days after single applications and 0.51 to 0.88 ppm by 3 days after double applications of water-soluble granules. Collectively, these studies show that operational applications of neonicotinoids can result in soil concentrations that are likely to pose a high risk of sublethal effects and potential risk of lethal effects (especially by contact toxicity) to earthworms.

At least two issues related to the assessment of risk to earthworms from exposure to neonicotinoids have not been adequately addressed in the published literature. The first is the length of exposure periods in toxicity testing compared to

the length of exposure to persistent concentrations in natural soils. Most toxicity tests are short term, in the order of days to weeks. On the other hand, neonicotinoid residues can persist in soils for months to years (Bonmatin et al. 2014, this issue). For most pesticides, lethal or effective concentrations become lower as exposure periods increase, and this is likely the case for neonicotinoids (Tennekes 2010; Tennekes and Sánchez-Bayo 2012, 2013; Rondeau et al. 2014). It is plausible that long-term low-level concentrations of neonicotinoids in soils may pose higher risk to earthworms than what can be inferred from the published toxicity tests. The second issue pertains to the heterogeneous distribution of neonicotinoid residues in natural soils. When residues enter the soil at the surface from spray or granule deposition or from litter fall, concentrations in soils are likely to be higher on or near the surface than in deeper soils. Residues entering soils from planted seed or from contaminated water are likely to be higher at or near the source of contamination than elsewhere. Both situations would result in concentration “hot spots” near the points of entry. Conversely, most toxicity tests prepare test concentrations as parts per million (or equivalent) and assume complete mixing. Therefore, levels of exposure to earthworms at or near those hot spots in natural soils will consequently be higher than would be predicted from residue analyses of bulk samples from laboratory or field test systems.

Mortality or behavioural effects on individual earthworms do not necessarily translate to population effects with ecological consequences. Populations of organisms with short generation times (e.g. several generations per year as is the case for most earthworm species) and/or high dispersal capacity have a higher likelihood of recovery from pesticide-induced population declines than those with longer regeneration periods and limited dispersal capacity (Kreutzweiser and Sibley 2013). However, the tendency for neonicotinoids to persist in organic soils reduces the likelihood of this recovery pathway because subsequent generations may be exposed to concentrations similar to those to which the parent generation was exposed. Life history strategies and their influences on community responses and recovery from pesticide effects have been demonstrated by population modelling of other non-target organisms (Wang and Grimm 2010), and similar principles may apply to assessing risks to overall earthworm populations and communities. Population models that account for differential demographics and population growth rates within communities have been shown to provide more accurate assessments of potential pesticide impacts on populations and communities than conventional lethal concentration estimates can provide (Stark and Banks 2003). The use of ecological models to incorporate a suite of factors including seasonal variations, community assemblage mechanisms and lethal and sublethal insecticide effects and their influences on the risks to organisms, populations or communities can provide useful insights into receptor/pesticide interactions and

can thereby improve risk assessments (Bartlett et al. 2010). Ecological and population modelling combined with pesticide exposure modelling and case-based reasoning (drawing on past experience or information from similar chemical exposures) can provide further refinements and improve risk assessment for earthworm communities and their ecological function (van den Brink et al. 2002). Empirical field studies of earthworm population responses to realistic field concentrations of neonicotinoids are lacking and would greatly improve risk assessment efforts.

Aquatic invertebrates

Freshwater invertebrates

Aquatic invertebrates are extremely important components of aquatic ecosystems. They play roles as decomposers, grazers, sediment feeders, parasites and predators. They also provide much of the food that vertebrates associated with these systems feed upon. Pesticides, including neonicotinoids, reach surface waters through various routes, but in particular through atmospheric deposition (drift) after application by various sprayers, by surface runoff and by seepage of contaminated groundwater. Aquatic invertebrates are particularly susceptible to pesticides. Unlike terrestrial organisms, aquatic organisms generally cannot avoid exposure easily by moving to uncontaminated areas, particularly when pesticides are water soluble. Uptake of pesticides in aquatic invertebrates occurs through respiration (gills and trachea), feeding and through the epidermis, be it cuticle or skin.

Neonicotinoids have been used for a comparatively shorter period of time than other insecticides. However, they are found in freshwater systems more and more frequently. For example, surface water monitoring for pesticides in California has revealed that imidacloprid has frequently exceeded water quality guidelines of 1 ppb (Starner and Goh 2012). In the Washington State, USA, the State Department of Ecology and the State Department of Agriculture have been monitoring salmon-bearing rivers and streams for pesticides, including imidacloprid for a number of years and this insecticide is frequently found (<http://agr.wa.gov/PestFert/natresources/SWM/>).

However, even though imidacloprid and other neonicotinoids are present in freshwater systems, the question remains to what extent such concentrations affect aquatic organisms in the field. Here we discuss a number of studies dealing with neonicotinoid toxicity to aquatic invertebrates and make some observations about their potential impact on aquatic ecosystems.

Laboratory studies

Crustacea and Amphipoda Several laboratory studies have been published on the toxicity of the neonicotinoid imidacloprid on a range of aquatic invertebrates (Table 7). Stark and Banks (2003) developed acute toxicity data and population-level toxicity data for the water flea *Daphnia pulex* exposed to thiamethoxam (Actara). Thiamethoxam was the least toxic insecticide evaluated in this study of seven insecticides, and its LC₅₀ of 41 ppm was well above any anticipated concentration expected to be found in surface water systems.

Chen et al. (2010) estimated the acute toxicity of imidacloprid to the water flea, *Ceriodaphnia dubia* (LC₅₀=2.1 ppb), and the chronic toxicity to *C. dubia* populations. The effects of the adjuvant, R-11 alone and in combination with imidacloprid were also assessed. In the population study, exposure of *C. dubia* to imidacloprid concentrations of 0.3 ppb reduced population size to 19 % of the control population. This concentration is well below the U.S. EPA's expected environmental concentration of 17.4 ppb, indicating that imidacloprid may cause damage to aquatic invertebrates in the field.

The acute and chronic effects of imidacloprid on the amphipod *Gammarus pulex* were studied by Nyman et al. (2013). Feeding by *G. pulex* and body lipid content were significantly reduced after exposure to a constant imidacloprid concentration of 15 ppb. Furthermore, *G. pulex* individuals were unable to move and feed after 14 days of constant exposure resulting in a high level of mortality.

Interestingly, the standard test organism *Daphnia magna* is especially insensitive to neonicotinoids (Beketov and Liess 2008). An acute LC₅₀ of around 7,000 ppb is several orders of magnitude above effective concentrations found for several other invertebrates. This implies that *D. magna* cannot be used as a sensitive test organism protective for many species.

Insecta Acute toxicity estimates of neonicotinoids on aquatic insects have also been published. LC₅₀ estimates for aquatic insects range from 3 to 13 ppb. Imidacloprid LC₅₀ estimates for the mayfly *Baetis rhodani*, the black fly *Simulium latigonium* (Beketov and Liess 2008) and the mosquito *Aedes taeniorhynchus* (Song et al. 1997) are 8.5, 3.7 and 13 ppb, respectively. LC₅₀ estimates for *B. rhodani* and *S. latigonium* exposed to thiacloprid were 4.6 and 3.7 ppb, respectively (Beketov and Liess 2008). A chronic LC₅₀ of 0.91 ppb was reported for the midge *Chironomus tentans* for imidacloprid (Stoughton et al. 2008). A study on the effects of imidacloprid as a mixture with the organophosphate insecticides dimethoate and chlorpyrifos on the midge *Chironomus dilutus* found that imidacloprid acted synergistically with chlorpyrifos and antagonistically with dimethoate (LeBlanc et al. 2012).

Table 7 Selection of studies on the effects of imidacloprid on freshwater macrophuna

	Compound	Experimental design	Effect	LC ₅₀ /EC ₅₀	LOAEL	Reference
Aquatic taxa						
Oligochaeta	Imidacloprid	10 day exposure to contaminated sediment	Survival, growth, behaviour, avoidance		<0.05 mg/kg	Sardo and Soares (2010)
<i>Chironomus tentans</i> and <i>Hyalella azteca</i>	Imidacloprid	Standard toxicity test	Survival	0.91 µg/l (28 days)		Stoughton et al. (2008)
Mesocosm communities	Neonics and other insecticides		Drift response			Berghahn et al. (2012)
<i>Daphnia</i> , <i>Gammarus pulex</i>	Imidacloprid		Survival			Ashauer et al. (2011)
Mayflies	Imidacloprid		Nymph abundance emergence patterns and adult body size			Alexander et al. (2008)
<i>Ceriodaphnia dubia</i>	Imidacloprid	Lab toxicity tests	Mortality	2.1 ppb		Chen et al. (2010)
<i>D. magna</i>	Imidacloprid	Lab toxicity tests	Population growth rate	10.4 mg/l		Song et al. (1997)
<i>Aedes aegypti</i>	Imidacloprid	Lab toxicity tests	Mortality	44 ppb		Song et al. (1997)
<i>Aedes taeniorhynchus</i>	Imidacloprid	Lab toxicity tests	Mortality	13 ppb		Song et al. (1997)
Mayflies, Oligochaetes	Imidacloprid		Feeding inhibition			Alexander et al. (2008)
Odonata, Libellulidae	Imidacloprid, fipronil	Field	Larval and adult survival, emergence			Jinguji et al. (2013)
Macro-invertebrate community	Imidacloprid	Stream mesocosm	Community diversity, leaf litter breakdown			Pestana et al. (2009)
Crustacean: <i>Asellus aquaticus</i> , <i>Gammarus fossarum</i>	Imidacloprid and atrazine	Standard toxicity test	Survival, respiration, electron transport system			Lukancic et al. (2010)
Caddisfly: <i>Sericoostoma</i> , <i>Chironomus riparius</i>	Imidacloprid	Standard toxicity test	Burrowing behaviour; antipredator behaviour			Pestana et al. (2009)
Ostracoda, <i>Daphnia magna</i>	Imidacloprid	Lab toxicity test	Survival			Sánchez-Bayo (2006)
<i>Chironomus diutus</i>	Imidacloprid + mixtures (chlorpyrifos, dimethoate)	Lab toxicity test	Survival			LeBlanc et al. (2012)
Terrestrial taxa						
Aphidius ervi	Imidacloprid + cadmium	Lab toxicity tests	Population growth rate			Kramarz and Stark (2003)

Oligochaetes Sardo and Soares (2010) investigated the effects of imidacloprid on the aquatic oligochaete *Lumbriculus variegatus*. They exposed this worm species to imidacloprid concentrations ranging from 0.05 to 5.0 mg/kg sediment. Mortality was fairly low (35 % in the highest concentration), but *L. variegatus* avoided imidacloprid-contaminated sediment. Furthermore, individual growth (biomass) was inhibited at all concentrations tested compared to the control.

Mesocosm studies Alexander et al. (2008) examined the effect of imidacloprid as a 12 day pulse or 20 day continuous exposure on the mayflies *Epeorus* spp. and *Baetis* spp. Nymph densities were reduced after both types of exposures. Sublethal effects were observed as well. Adults were smaller and had smaller head and thorax size after exposure to imidacloprid concentrations as low as 0.1 ppb. However, these effects were only found in males.

Within community test systems, neonicotinoids had strong effects especially on insects (Hayasaka et al. 2012). However, to our knowledge, all experiments investigating a dose–response relationship observed effects at the lowest concentrations evaluated. Hence, it is difficult to establish a NOEC. Within outdoor mesocosm studies, a LOEC of 1.63 ppb was estimated for imidacloprid. Adverse effects on benthic communities with 5 % reductions in the abundance of invertebrates were observed by Pestana et al. (2009). For thiacloprid, strong effects on sensitive long living insects were observed at pulsed exposure to 0.1 ppb (Liess and Beketov 2011), the lowest effective concentration observed so far in communities.

Berghahn et al. (2012) conducted stream mesocosm studies whereby 12 h pulses of imidacloprid (12 ppb) were introduced three times at weekly intervals. Results showed that drift of insects and the amphipod *Gammarus roeseli* increased after exposure to pulses of imidacloprid. These results indicated that imidacloprid was having a negative effect on *G. roeseli*.

In another stream mesocosm study, Böttger et al. (2013) evaluated pulses of imidacloprid on *G. roeseli*. The number of brood carrying females was reduced in the imidacloprid treatments compared to the control groups in the last 3 weeks of the study.

The populations of an aquatic invertebrate, the common mosquito *Culex pipiens*, exposed over several generations to repeated pulses of low concentrations of the neonicotinoid thiacloprid, continuously declined and did not recover in the presence of a less sensitive competing species, the water flea *D. magna*. By contrast, in the absence of a competitor, insecticide effects on the more sensitive species were only observed at concentrations one order of magnitude higher, and the species recovered more rapidly after a contamination event. The authors conclude that repeated toxicant pulse of populations that are challenged with interspecific competition may result

in a multigenerational culmination of low-dose effects (Liess et al. 2013).

Risk to aquatic ecosystems A species sensitivity distribution (SSD) of acute toxicity data for a wider range of species, including ostracods, cladocerans and other aquatic organisms, predicts a hazardous concentration for 5 % of aquatic species (HC5) for imidacloprid in water in the range 1.04–2.54 ppb (Sanchez-Bayo and Kouchi 2012).

Van Dijk et al. (2013) developed a regression analysis for abundance of aquatic macro-invertebrate species and nearby imidacloprid concentrations in Dutch surface waters. Data from 8 years of nationwide monitoring covering 7,380 different locations of macro-invertebrate samples and 801 different locations of imidacloprid samples were pooled. Next, the biological samples (macro-invertebrate abundance counts) were combined with nearby (in space and time) chemical samples (imidacloprid concentrations), and next, a statistical analysis was done on the complete pooled combined dataset. They found that macro-invertebrate abundance consistently declines along the gradient of increasing median nearby imidacloprid concentration in the pooled dataset. This pattern turned out to be robust: it is independent of year and location. Overall, a significant negative relationship ($P < 0.001$) was found between abundance of all macro-invertebrate species pooled and nearby imidacloprid concentration. A significant negative relationship was also found for abundance of each of the pooled orders Amphipoda, Basommatophora, Diptera, Ephemeroptera and Isopoda, and for several species separately. The order Odonata had a negative relationship very close to the significance threshold of 0.05 ($P = 0.051$). In accordance with previous research, a positive relationship between abundance and nearby imidacloprid pollution was found for the order Actinedida. However, other pesticides were not included into the analyses by Van Dijk et al. (2013). Therefore, possible co-linearity or synergisms between neonicotinoids and other pollutants still need to be further explored (Vijver and Van den Brink 2014).

Pesticide exposure was identified to strongly reduce the amount and abundance of vulnerable invertebrate species in streams using the SPEAR approach (Liess and von der Ohe 2005). The approach was extended from German streams to Australian, Danish, French and Finnish streams revealing the same effects of pesticide exposure on vulnerable invertebrate species (Rasmussen et al. 2013; Liess et al. 2008; Schäfer et al. 2012). Beketov et al. (2013) analysed the effect of pesticide presence on invertebrate species richness in European (Germany and France) and Australian streams. They found an overall reduction of 42 % for Europe and 27 % for Australia in species richness between uncontaminated and heavily contaminated streams. The limitation of these studies in the context of assessment of neonicotinoid impact is that toxicity was mainly due to insecticides, other than neonicotinoids, as general usage of the latter only increased recently.

The results of laboratory and semi-field (mesocosm) studies indicate that aquatic invertebrates are very sensitive to the neonicotinoid insecticides. However, most of the studies we found in the literature were conducted with imidacloprid. For pesticide risk assessment, the published results to date indicate that it may be difficult to predict community-level effects using the tiered aquatic effect assessment scheme and acute and chronic toxicity data. When extrapolating from acute and chronic single species test systems, the assessment factors identified by the uniform principle of the relevant EU legislation (1107/2009) do not predict safe concentrations in multi-species outdoor mesocosms. For example, acute laboratory effects of thiacloprid on sensitive insect species show that effects occur after exposure to the range of 3–13 ppb. Accordingly, an assessment factor of 100 would indicate a safe concentration of 0.03 to 0.13 ppb for thiacloprid. However, outdoor mesocosm results employing a pulsed exposure show a LOEC below 0.1 ppb for thiacloprid (Liess and Beketov 2011). Lower concentrations were not investigated. Obviously, an assessment factor higher than 100 is needed to identify safe concentrations on the basis of acute test results. For the HC5 calculated on acute lethal concentrations, an assessment factor of larger than 10 is necessary (Liess and Beketov 2012). Additionally, in a laboratory study, chronic effects of sensitive insect species were exhibited after exposure to 0.91 ppb imidacloprid. Employing an assessment factor of 10 would indicate a safe concentration of approximately 0.1 ppb imidacloprid. However, this concentration is not safe according to the results obtained in complex community investigations. Unfortunately, to the best of our knowledge, no community-level investigation with imidacloprid evaluating a range of concentrations below 0.1 ppb has been published. This type of study would help with determining a NOEC for imidacloprid. Overall, the results of the published literature indicate that certain neonicotinoids have the potential to cause significant damage to aquatic ecosystems by causing negative effects in individuals and populations of aquatic invertebrates at very low concentrations. Protective concentrations for these products in aquatic systems still need to be determined.

Marine and coastal invertebrates

There is very limited information regarding the assessment of the environmental toxicology and contamination of neonicotinoids in marine ecosystems. Standardised environmental toxicological characterization focuses on only a few species models and rarely examines species that represent keystone organisms in marine or coastal ecosystems (CCME 2007). Monitoring and surveillance of neonicotinoid pollution in marine coastal habitats are non-existent.

Toxicology The earliest published marine ecotoxicological studies of neonicotinoids were with opossum shrimps

(*Mysidopsis bahia*) which are distributed in marine coastal waters (Ward 1990, 1991; Lintott 1992). Median LC₅₀ (96 h) for the technical grade of imidacloprid was 34.1 ppb with a mortality-NOEC of 13.3 ppb (Ward 1990). Exposure to a commercial formulation (ADMIRE) of imidacloprid resulted in a 96 h mortality-NOEC of 21 ppb. Maximum acceptable toxicant concentrations for *M. bahia* to imidacloprid were 23 parts per trillion (ppt) for growth effects and 643 ppt for reproductive effects (Ward 1991).

Toxicology for other marine arthropods includes *Artemia* spp. and a brackish water mosquito (*Aedes taeniorhynchus*). The 48 h LC₅₀ for *Artemia* was 361 ppm, whilst *Aedes* exhibited a 72 h LC₅₀ of 21 ppb, and a 48 h LC₅₀ of 13 ppb for an early instar stage of development (Song et al. 1997; Song and Brown 1998). Osterberg et al. (2012) demonstrated that in the blue crab (*Callinectes sapidus*), megalopae were an order of magnitude more sensitive than juveniles to lethal effects of imidacloprid (24 h-LC₅₀=10 ppb for megalopae vs 24 h-LC₅₀=1,1 ppb for juveniles).

There are no known published OECD/EPA parameter-based studies on non-arthropod marine invertebrates. For the marine mussel, *Mytilus galloprovincialis*, a transcriptomic and proteomic survey was conducted as a response to imidacloprid and thiacloprid exposures (Dondero et al. 2010). This study concluded that the two neonicotinoids induced distinct toxicodynamic responses and that caution should be heeded when conducting ecological risk assessments for chemical mixtures that target the same receptor. Rodrick (2008) demonstrated that imidacloprid had an effect on oyster hemocyte immunocompetence and that there was an additive effect when oysters were exposed to a compound stress of salinity and exposure to imidacloprid. Tomizawa et al. (2008) used the gastropod *Aplysia californica* as a model to characterize imidacloprid and thiacloprid as agonists of the acetylcholine-binding protein, indicating that neonicotinoids could also affect marine gastropods.

Environmental pollution There are no published works regarding the marine environmental contamination of neonicotinoids. Until recently, there has been little public concern of neonicotinoid non-point source pollution of marine environments from land runoff. At least within the USA, this attitude is beginning to change. In the State of Washington 2013, the Willapa-Grays Harbor Oyster Growers Association received a conditional registration from the U.S. Environmental Protection Agency to use imidacloprid to control native burrowing shrimp in Willapa Bay, Washington that may threaten commercial shellfish beds (EPA Reg. no. 88867–1). In Hawaii, there have been public protests and scrutiny over the use of neonicotinoid pesticides in their industrial agricultural practices and their likely negative impacts on coral reefs and sea grass beds (Sergio 2013). For both Hawaii and the U.S. Virgin Islands, there is concern that the use of

neonicotinoids as a method for termite control may be polluting and impacting coastal resources.

Conclusion

At field-realistic levels of pollution, neonicotinoids and fipronil generally have negative effects on physiology and survival for a wide range of non-target invertebrates in terrestrial, aquatic, marine and benthic habitats. Effects are most often found by in vitro testing, using a limited number of test species. This basically means that there is a deficit of information for the grand majority of other invertebrates. In vitro testing to establish safe environmental concentration thresholds is hindered by the fact that most test protocols are based on older methodology, validated for pesticides with very different chemical and toxicological characteristics. New and improved methodologies are needed to specifically address the unique toxicology of these neurotoxic chemicals, including their non-lethal effects and synergistic effects for a variety of terrestrial, aquatic and marine organisms.

The amount of published in vivo field tests is small and experimental setups often suffer from inability to control for variation in (semi)natural circumstances or have insufficient statistical power due to the high financial costs of large robust field experiments. Given the clear body of evidence presented in this paper showing that existing levels of pollution with neonicotinoids and fipronil resulting from presently authorized uses frequently exceed lowest observed adverse effect concentrations and are thus likely to have large-scale and wide ranging negative biological and ecological impacts, the authors strongly suggest that regulatory agencies apply more precautionary principles and tighten regulations on neonicotinoids and fipronil.

Acknowledgments This manuscript benefited from the discussions in the International Task Force on Systemic Pesticides during its plenary meetings in Paris (2010), Bath (2011), Cambridge (2012), Padua (2012), Louvain-la-Neuve (2013) and Legnaro (2013). The authors are organised in alphabetic order, except the first who is also the corresponding author. All authors work for public agencies, except V. Amaral-Rogers who is employed by Buglife, a UK charity devoted to the conservation of invertebrates, and D.A. Noome, whose independent work for the TFSP is financed by the Stichting Triodos Foundation, and N. Simon-Delso working for CARI (association supported by the Belgium government). Contributions of J. Settele and M. Wiemers were part of www.legato-project.net (funded by the BMBF, German Ministry for Education and Research). The work has been funded by the Triodos Foundation's Support Fund for Independent Research on Bee Decline and Systemic Pesticides. This Support Fund has been created from donations by Adessium Foundation (The Netherlands), Act Beyond Trust (Japan), Utrecht University (Netherlands), Stichting Triodos Foundation (The Netherlands), Gesellschaft fuer Schmetterlingsschutz (Germany), M.A. O.C. Gravin van Bylandt Stichting (The Netherlands), Zukunft Stiftung Landwirtschaft (Germany), Study Association Storm (Student Association Environmental Sciences Utrecht University) and citizens. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Conflict of interest The authors declare no conflict of interest.

References

- Abbas N, Shad SA, Razaq M (2012) Fitness cost, cross resistance and realized heritability of resistance to imidacloprid in *Spodoptera litura* (Lepidoptera: Noctuidae). *Pestic Biochem Physiol* 103:181–188
- Abbott VA, Nadeau JL, Higo HA, Winston ML (2008) Lethal and sublethal effects of imidacloprid on *Osmia lignaria* and clothianidin on *Megachile rotundata* (Hymenoptera: Megachilidae). *J Econ Entomol* 101:784–796
- Aebischer NJ (1990) Assessing pesticide effects on non-target invertebrates using long-term monitoring and time-series modelling. *Funct Ecol* 4:369–373
- Ahmad M, Sayyed AH, Saleem MA, Ahmad M (2008) Evidence for field evolved resistance to newer insecticides in *Spodoptera litura* (Lepidoptera: Noctuidae) from Pakistan. *Crop Prot* 27:1367–1372
- Ahmad M, Rafiq M, Arif MI, Sayyed AH (2011) Toxicity of some commonly used insecticides against *Coccinella undecimpunctata* (Coleoptera: Coccinellidae). *Pak J Zool* 43:1161–1165
- Ahmad S, Ansari MS, Ahmad N (2013) Acute toxicity and sublethal effects of the neonicotinoid imidacloprid on the fitness of *Helicoverpa armigera* (Lepidoptera: Noctuidae). *Int J Trop Insect Sci* 33:264–275
- Aizen, Harder (2009) The global stock of domesticated honey bees is growing slower than agricultural demand for pollination. *Curr Biol* 19:915–918
- Alaux C, Brunet JL, Dussaubat C, Mondet F, Tchamitchan S, Cousin M, Brillard J, Baldy A, Belzunces LP, Le Conte Y (2010) Interactions between *Nosema* microspores and a neonicotinoid weaken honeybees (*Apis mellifera*). *Environ Microbiol* 12:774–782
- Albajes R, López C, Pons X (2003) Predatory fauna in cornfields and response to imidacloprid seed treatment. *J Econ Entomol* 96:1805–1813
- Alexander AC, Heard K, Culp JM (2008) Emergent body size of mayfly survivors. *Freshw Biol* 53:171–180
- Aliouane Y, El Hassani AK, Gary V, Armengaud C, Lambin M, Gauthier M (2009) Subchronic exposure of honeybees to sublethal doses of pesticides: effects on behavior. *Environ Toxicol Chem* 28:113–122
- Alves PRL, Cardoso EJBN, Martines AM, Sousa JP, Pasini A (2013) Earthworm ecotoxicological assessments of pesticides used to treat seeds under tropical conditions. *Chemosphere* 90:2674–2682
- Anon (2012) EFSA Statement on the findings in recent studies investigating sub-lethal effects in bees of some neonicotinoids in consideration of the uses currently authorised in Europe. *EFSA Journal* 10: 2752
- Ansari MS, Ali H, Shafqat S (2012) Insecticidal effect on a population of *Spilarctia obliqua* (Lepidoptera: Arctiidae). *Entomol Res* 42:330–338
- Apenet, (2010) <http://www.reterurale.it/flex/cm/pages/ServeBLOB.php/L/IT/IDPagina/3280>
- Araín MS, Hu XX, Li GQ (2014) Assessment of toxicity and potential risk of butene-fipronil using *Drosophila melanogaster*, in comparison to nine conventional insecticides. *Bull Environ Contam Toxicol* 92:190–195
- Arena, M., and Sgolastra, F (2014) A meta-analysis comparing the sensitivity of bees to pesticides. *Ecotoxicol*, 1–11
- Asaro C, Creighton J (2011) Use of systemic fipronil and imidacloprid to control regeneration pests of loblolly pine. *J Econ Entomol* 104: 1272–1279
- Ashauer R, Hintermeister A, Potthoff E, Esche BI (2011) Acute toxicity of organic chemicals to *Gammarus pulex* correlates with sensitivity

- of *Daphnia magna* across most modes of action. *Aquat Toxicol* 103: 38–45
- Aufauvre J, Biron DG, Vidau C, Fontbonne R, Roudel M, Diogon M, Vigues B, Belzunces LP, Delbac F, Blot N (2012) Parasite-insecticide interactions: a case study of *Nosema ceranae* and fipronil synergy on honeybee. *Scientific reports* 2, srep00326-srep00326
- Balderrama N, Nunez J, Giurfa M, Torrealba J, Dealbornoz EG, Almeida LO (1996) A deterrent response in honeybee (*Apis mellifera*) foragers: dependence on disturbance and season. *J Insect Physiol* 42: 463–470
- Barbara G, Grünwald B, Paute S, Gauthier M, Raymond-Delpech V (2008) Study of nicotinic acetylcholine receptors on cultured antennal lobe neurones from adult honeybee brains. *Invert Neurosci* 8:19–29
- Barbieri RF, Lester PJ, Miller AS, Ryan KG (2013) A neurotoxic pesticide changes the outcome of aggressive interactions between native and invasive ants. *Proc R Soc B Biol Sci* 28:20132157
- Bartlett MD, Briones MJ, Neilson R, Schmidt O, Spurgeon D, Creamer RE (2010) A critical review of current methods in earthworm ecology: from individuals to populations. *Eur J Soil Biol* 46:67–73
- Baylay AJ, Spurgeon DJ, Svendsen C, Griffin JL, Swain SC, Sturzenbaum SR, Jones OAH (2012) A metabolomics based test of independent action and concentration addition using the earthworm *Lumbricus rubellus*. *Ecotoxicology* 21:1436–1447
- Beketov MA, Liess M (2008) Potential of 11 pesticides to initiate downstream drift of stream macroinvertebrates. *Arch Environ Contam Toxicol* 55:247–253
- Beketov MA, Kefford BJ, Schäfer RB, Liess M (2013) Pesticides reduce regional biodiversity of stream invertebrates. *Proc Natl Acad Sci U S A* 110:11039–11043. doi:10.1073/pnas.1305618110
- Belzunces LP (2006) Rapport d'étude du programme Etude comparée des impacts de trois classes d'insecticides néonicotinoïdes chez l'abeille. Programme communautaire sur l'apiculture Année 2006
- Belzunces LP, Tchamitchian S, Brunet JL (2012) Neural effects of insecticides in the honey bee. *Apidologie* 43:348–370
- Benzidane Y, Touinsi S, Motte E, Jadas-Hécart A, Communal PY, Leduc L, Thany SH (2010) Effect of thiamethoxam on cockroach locomotor activity is associated with its metabolite clothianidin. *Pest Manag Sci* 66:1351–1359
- Berghahn R, Mohr S, Hübner V, Schmiediche R, Schmiedling I, Svetich-Will E, Schmidt R (2012) Effects of repeated insecticide pulses on macroinvertebrate drift in indoor stream mesocosms. *Aquat Toxicol* 122:56–66
- Biocca M, Conte E, Pulcini P, Marinelli E, Pochi D (2011) Sowing simulation tests of a pneumatic drill equipped with systems aimed at reducing the emission of abrasion dust from maize dressed seed. *J Environ Sci Health B* 46:438–448
- Bloem S, Mizell RF, Bloem KA, Hight SD, Carpenter JE (2005) Laboratory evaluation of insecticides for control of the invasive *Cactoblastis cactorum* (Lepidoptera: Pyralidae). *Fla Entomol* 88: 395–400
- Boggs CL (2003) Environmental variation, life histories, and allocation. In: Boggs CL, Watt WB, Ehrlich PR (eds) *Butterflies: ecology and evolution taking flight*. The University of Chicago Press, Chicago, pp 185–206
- Bonmatin JM, Moineau I, Charvet R, Colin ME, Fleche C, Bengsch ER et al (2005) Behaviour of imidacloprid in fields: toxicity for honey bees. In: Lichtfouse E, Schwarzbauer J, Robert D (eds) *In environmental chemistry: green chemistry and pollutants in ecosystems*. Springer, New York, pp 483–494
- Bonmatin J-M, Giorio C, Girolami V, Goulson D, Kreutzweiser D, Krupke C, Liess M, Long E, Marzaro M, Mitchell E, Noome D, Simon-Delso N, Tapparo A (2014) Environmental fate and exposure; neonicotinoids and fipronil. *Environ Sci Pollut Res*. doi:10.1007/s11356-014-3332-7 (this issue)
- Bordereau-Dubois B, List O, Calas-List D, Marques O, Communal PY, Thany SH, Lapié B (2012) Transmembrane potential polarization, calcium influx and receptor conformational state modulate the sensitivity of the imidacloprid-insensitive neuronal insect nAChR to the neonicotinoid insecticides. *J Pharmacol Exp Ther*. doi:10.1124/jpet.111.188060
- Bortolotti L, Montanari R, Marcelino J, Medrzycki P, Maini S, Porrini C (2003) Effect of sub-lethal imidacloprid doses on the homing rate and foraging activity of honey bees. *Bull Insect* 56:63–67
- Bosch J, Kemp WP (2006) Bee population returns and cherry yields in an orchard pollinated with *Osmia lignaria* (Hymenoptera: Megachilidae). *J Econ Entomol* 99:408–413
- Bostanian NJ, Hardman JM, Ventard E, Racette G (2005) The intrinsic toxicity of several neonicotinoids to *Lygus lineolaris* and *Hyaliodes vitripennis*, a phytophagous and a predaceous mired. *Pest Manag Sci* 61:991–996
- Böttger R, Feibicke M, Schaller J, Dudel G (2013) Effects of low-dosed imidacloprid pulses on the functional role of the caged amphipod *Gammarus roeseli* in stream mesocosms. *Ecotoxicol Environ Saf* 93:93–100
- Braun G, Bicker G (1992) Habituation of an appetitive reflex in the honeybee. *J Neurophysiol* 67:588–598
- Breeze TD, Bailey AP, Balcombe KG et al (2011) Pollination services in the UK: how important are honeybees? *Agric Ecosyst Environ* 142: 137–143
- Brittain C, Bommarco R, Vighi M, Settele J, Potts SG (2010a) Organic farming in isolated landscapes does not benefit flower-visiting insects and pollination. *Biol Conserv* 143:1860–1867. doi:10.1016/j.biocon.2010.04.029
- Brittain CA, Vighi M, Bommarco R, Settele J, Potts SG (2010b) Impacts of a pesticide on pollinator species richness at different spatial scales. *Basic App Ecol* 11:106–115
- Brooks DR, Bater JE, Clark SJ, Monteith DT, Andrews C, Corbett SJ, Beaumont DA, Chapman JW (2012) Large carabid beetle declines in a United Kingdom monitoring network increases evidence for a widespread loss in insect biodiversity. *J Appl Ecol* 49:1009–1019
- Broznic D, Marinic J, Tota M, Juresic GC, Petkovic O, Milin C (2012) Hysteretic behaviour of imidacloprid sorption-desorption in soils of Croatian coastal regions. *Soil Sediment Contam* 21:850–871
- Brunet JL, Badiou A, Belzunces LP (2005) In vivo metabolic fate of [¹⁴C]-acetamiprid in six biological compartments of the honeybee *Apis mellifera* L. *Pest Manag Sci* 61:742–748
- Brunner JF, Beers EH, Dunley JE, Doerr M, Granger K (2005) Role of neonicotinyl insecticides in Washington apple integrated pest management. Part I. Control of lepidopteran pests. *J Insect Sci* 5
- Bryden J, Gill RJ, Mitton RAA, Raine NE, Jansen VAA (2013) Chronic sublethal stress causes bee colony failure. *Ecol Lett* 16:1463–1493
- Byrne FJ, Toscano NC (2007) Lethal toxicity of systemic residues of imidacloprid against *Homalodisca vitripennis* (Homoptera: Cicadellidae) eggs and its parasitoid *Gonatocerus ashmeadi* (Hymenoptera: Mymaridae). *Biol Control* 43:130–135
- Calderone NW (2012) Insect pollinated crops, insect pollinators and us agriculture: trend analysis of aggregate data for the period 1992–2009. *PLoS One* 7:e37235. doi:10.1371/journal.pone.0037235
- Canadian Council of Ministers of the Environment (2007) Canadian water quality guidelines: IMIDACLOPRID. Scientific supporting document. ISBN 978-1-896997-71-1 PDF
- Capowiez Y, Berard A (2006) Assessment of the effects of imidacloprid on the behaviour of two earthworm species (*Aporrectodea nocturna* and *Allolobophora icterica*) using 2D terraria. *Ecotoxicol Environ Saf* 64:198–206
- Capowiez Y, Rault M, Mazzia C, Belzunces L (2003) Earthworm behaviour as a biomarker—a case study using imidacloprid. *Pedobiologia* 47:542–547
- Capowiez Y, Rault M, Costagliola G, Mazzia C (2005) Lethal and sublethal effects of imidacloprid on two earthworm species

- (*Aporrectodea nocturna* and *Allolobophora icterica*). Biol Fertil Soils 41:135–143
- Capowiez Y, Bastardie F, Costagliola G (2006) Sublethal effects of imidacloprid on the burrowing behaviour of two earthworm species: modifications of the 3D burrow systems in artificial cores and consequences on gas diffusion in soil. Soil Biol Biochem 38:285–293
- Capowiez Y, Dittbrener N, Rault M, Triebkorn R, Hedde M, Mazzia C (2010) Earthworm cast production as a new behavioural biomarker for toxicity testing. Environ Pollut 158:388–393
- Casida JE (2011) Neonicotinoid metabolism: compounds, substituents, pathways, enzymes, organisms, and relevance. J Agric Food Chem 59:2923–2931
- Chagnon M, Gingras J, de Oliveira D (1993) Complementary aspects of strawberry pollination by honey and indigenous bees (Hymenoptera). J Econ Entomol 86:416–420
- Charmillot PJ, Pasquier D, Salamin C, Ter-Hovannesian A (2007) Ovicidal and larvicidal effectiveness of insecticides applied by dipping apples on the small fruit tortrix *Grapholita lobarzewskii*. Pest Manag Sci 63:677–681
- Charpentier G, Louat F, Bonmatin JM, Marchand PA, Vannier F, Locker D, Decoville M (2014) Lethal and sublethal effects of imidacloprid, after chronic exposure, on the insect model *Drosophila melanogaster*. Environ Sci Technol 48(7):4096–4102
- Chen RZ, Klein MG (2012) Efficacy of insecticides against the rice stem-borer, *Chilo suppressalis* (Walker) (Lepidoptera: Crambidae), and use of sex pheromones to time accurately the yearly application. Int J Pest Manag 58:353–359
- Chen XD, Culbert E, Herbert V, Stark JD (2010) Mixture effects of the adjuvant R-11 and the insecticide imidacloprid on population growth rate and other parameters of *Ceriodaphnia dubia*. Ecotoxicol Environ Saf 73:132–137
- Cheng XA, Chang C, Dai SM (2010) Responses of striped stem borer, *Chilo suppressalis* (Lepidoptera: Pyralidae), from Taiwan to a range of insecticides. Pest Manag Sci 66:762–766
- Chowdhury S, Mukhopadhyay S, Bhattacharyya A (2012) Degradation dynamics of the insecticide: Clothianidin (Dantop 50 % WDG) in a tea field ecosystem. Bull Environ Contam Toxicol 89:340–343
- Cichon LB, Soleno J, Anguiano OL, Garrido SA, Montagna CM (2013) Evaluation of cytochrome P450 activity in field populations of *Cydia pomonella* (Lepidoptera: Tortricidae) resistant to azinphosmethyl, acetamiprid, and thiacloprid. J Econ Entomol 106:939–944
- Coleman DC, Ingham ER (1988) Carbon, nitrogen, phosphorus and sulphur cycling in terrestrial ecosystems. Biogeochemistry 5:3–6
- Colin ME, Bonmatin JM, Moineau I, Gaimon C, Brun S, Vermandere JP (2004) A method to quantify and analyze the foraging activity of honey bees: relevance to the sublethal effects induced by systemic insecticides. Arch Environ Contam Toxicol 47:387–395
- Couvillon MJ, Barton SN, Cohen JA, Fabricius OK, Kaercher MH, Cooper LS, Silk MJ, Helantera H, Ratnieks FLW (2010) Alarm pheromones do not mediate rapid shifts in honey bee guard acceptance threshold. J Chem Ecol 36:1306–1308
- Cox L, Koskinen WC, Yen PY (1997) Sorption–desorption of imidacloprid and its metabolites in soils. J Agric Food Chem 45:1468–1472
- Cox L, Hermosin MC, Cornejo J (2004) Influence of organic amendments on sorption and dissipation of imidacloprid in soils. Int J Environ Anal Chem 84:95–102
- Cresswell JE (2011) A meta-analysis of experiments testing the effects of neonicotinoid insecticide (imidacloprid) on honey bees. Ecotoxicology 20:149–154
- Cresswell JE, Thompson H (2012) Comment on “A common pesticide decreases foraging success and survival in honey bees”. Science 337:1453
- Cresswell JE, Desneux N, van Engelsdorp D (2012a) Dietary traces of neonicotinoid pesticides as a cause of population declines in honey bees: an evaluation by Hill’s epidemiological criteria. Pest Manag Sci 68:819–827
- Cresswell JE, Page CJ, Uygun MB, Holmbergh M, Li Y, Wheeler JG, Laycock I, Pook CJ, De Ibarra NH, Smimoff N, Tyler CR (2012b) Differential sensitivity of honey bees and bumble bees to a dietary insecticide (imidacloprid). Zoology 115:365–371
- Cutler GC, Scott-Dupree C (2007) Exposure to clothianidin seed-treated canola has no long-term impact on honey bees. J Econ Entomol 100:765–772
- Davis BNK, Lakhani KH, Yates TJ (1991a) The hazards of insecticides to butterflies of field margins. Agric Ecosyst Environ 36:151–161
- Davis BNK, Lakhani KH, Yates TJ, Frost AJ (1991b) Bioassays of insecticide spray drift: the effects of wind speed on the mortality of *Pieris brassicae* larvae (Lepidoptera) caused by diflubenzuron. Agric Ecosyst Environ 36:141–149
- Davis BNK, Lakhani KH, Yates TJ, Frost AJ, Plant RA (1993) Insecticide drift from ground-based, hydraulic spraying of peas and brussels sprouts: bioassays for determining buffer zones. Agric Ecosyst Environ 43:93–108
- Decourtye A, Devillers J (2010) Ecotoxicology of neonicotinoids insecticides in the bees. In: Thany S (ed) Insect nicotinic acetylcholine receptors. Landes Bioscience, St. Austin
- Decourtye A, Lacassie E, Pham-Delegue MH (2003) Learning performances of honeybees (*Apis mellifera* L.) are differentially affected by imidacloprid according to the season. Pest Manag Sci 59:269–278
- Decourtye A, Armengaud C, Renou M, Devillers J, Cluzeau S, Gauthier M, Pham-Delègue MH (2004a) Imidacloprid impairs memory and brain metabolism in the honeybee (*Apis mellifera* L.). Pestic Biochem Physiol 78:83–92
- Decourtye A, Devillers J, Cluzeau S, Charretton M, Pham-Delègue MH (2004b) Effects of imidacloprid and deltamethrin on associative learning in honeybees under semi-field and laboratory conditions. Ecotoxicol Environ Saf 57:410–419
- Decourtye A, Devillers J, Genecque E, Le Menach K, Budzinski H, Cluzeau S, Pham-Delegue MH (2005) Comparative sublethal toxicity of nine pesticides on olfactory learning performances of the honeybee *Apis mellifera*. Arch Environ Contam Toxicol 48:242–250
- Decourtye A, Mader E, Desneux N (2010) Landscape enhancement of floral resources for honey bees in agro-ecosystems. Apidologie 41:264–277
- DEFRA (2007) Assessment of the risk posed to honeybees by systemic pesticides. PS2322, CSL York, UK
- DEFRA (2009) Intermittent exposure in terrestrial invertebrates—a case study with honeybees. PS2341, CSL York, UK
- Dégise P, Grünwald B, Gauthier M (2002) The insecticide imidacloprid is a partial agonist of the nicotinic receptor of honeybee Kenyon cells. Neurosci Lett 321:13–16
- Delbeke E, Vercruysse P, Tirry L, de Clercq P, Degheele D (1997) Toxicity of diflubenzuron, pyriproxyfen, imidacloprid and diafenthiuron to the predatory bug *Orius laevis* (Het.: Anthrenidae). Entomophaga 42:349–358
- Derecka K, Blythe MJ, Malla S, Genereux DP, Guffanti A, Pavan P, Moles A, Snart C, Ryder T, Ortori CA, Barrett DA, Schuster E, Stöger R (2013) Transient exposure to low levels of insecticide affects metabolic networks of honeybee larvae. PLoS One 8:e68191
- Desneux N, Decourtye A, Delpuech JM (2007) The sublethal effects of pesticides on beneficial arthropods. Ann Rev Entomol 52:81–106
- Detzel A, Wink M (1993) Attraction, deterrence or intoxication of bees (*Apis mellifera*) by plant allelochemicals. Chemoecology 4:8–18
- Di Prisco G, Cavaliere V, Annoscia D, Varricchio P, Caprio E, Nazzi F, Gargiulo G, Pennacchio F (2013) Neonicotinoid clothianidin adversely affects insect immunity and promotes replication of a viral pathogen in honey bees. Proc Natl Acad Sci U S A 110:18466–18471. doi:10.1073/pnas.1314923110
- Diez-Rodriguez GI, de Baptista GC, Trevizani LR, Haddad ML, Nava DE (2006) Residues of thiamethoxam, aldicarb and its metabolites

- in coffee leaves and effect on the control of *Leucoptera coffeella* (Guerin-Meneville) (Lepidoptera: Lyonetiidae). *Neotropical Entomol* 35:257–263
- Dilling C, Lambdin P, Grant J, Rhea R (2009) Community response of insects associated with eastern hemlock to imidacloprid and horticultural oil treatments. *Environ Entomol* 38:53–66
- Dittbrenner N, Triebkorn R, Moser I, Capowiez Y (2010) Physiological and behavioural effects of imidacloprid on two ecologically relevant earthworm species (*Lumbricus terrestris* and *Aporrectodea caliginosa*). *Ecotoxicology* 19:1567–1573
- Dittbrenner N, Schmitt H, Capowiez Y, Triebkorn R (2011a) Sensitivity of *Eisenia fetida* in comparison to *Aporrectodea caliginosa* and *Lumbricus terrestris* after imidacloprid exposure. Body mass change and histopathology. *J Soils Sediments* 11:1000–1010
- Dittbrenner N, Moser I, Triebkorn R, Capowiez Y (2011b) Assessment of short and long-term effects of imidacloprid on the burrowing behaviour of two earthworm species (*Aporrectodea caliginosa* and *Lumbricus terrestris*) by using 2D and 3D post-exposure techniques. *Chemosphere* 84:1349–1355
- Dittbrenner N, Capowiez Y, Kohler H, Triebkorn R (2012) Stress protein response (Hsp70) and avoidance behaviour in *Eisenia fetida*, *Aporrectodea caliginosa*, and *Lumbricus terrestris* when exposed to imidacloprid. *J Soils Sediments* 12:198–206
- Dively G, Hooks C (2010) Use patterns of neonicotinoid insecticides on cucurbit crops and their potential exposure to honey bees. Progress report, strategic agricultural initiative grants program, EPA region 3
- Doffou NM, Ochou OG, Kouassi P (2011) Susceptibility of *Pectinophora gossypiella* (Lepidoptera: Gelechiidae) and *Cryptophlebia leucotreta* (Lepidoptera: Tortricidae) to insecticides used on cotton crops in Côte d'Ivoire, West Africa. Implications in insecticide resistance pest management strategies. *Resistant Pest Manag Newsl* 20:10–15
- Dogramaci M, Tingey WM (2008) Comparison of insecticide resistance in a North American field population and a laboratory colony of potato tuberworm (Lepidoptera: Gelechiidae). *J Pest Sci* 81:17–22
- Dondero F, Negri A, Boatti L, Marsano F, Mignone F, Viarengo A (2010) Transcriptomic and proteomic effects of a neonicotinoid insecticide mixture in the marine mussel (*Mytilus galloprovincialis*, Lam.). *Sci Total Environ* 15:3775–3786
- Donnarumma L, Pulcini P, Pochi D, Rosati S, Lusco L, Conte E (2011) Preliminary study on persistence in soil and residues in maize of imidacloprid. *J Environ Sci Health B* 46:469–472
- Dunley JE, Brunner JF, Doerr MD, Beers EH (2006) Resistance and cross-resistance in populations of the leafrollers, *Choristoneura rosaceana* and *Pandemis pyrusana*, in Washington apples. *J Insect Sci* 6
- Dupuis JP, Gauthier M, Raymond-Delpech V (2011) Expression patterns of nicotinic subunits alpha 2, alpha 7, alpha 8, and beta 1 affect the kinetics and pharmacology of ACh-induced currents in adult bee olfactory neuropiles. *J Neurophysiol* 106:1604–1613
- Durham EW, Scharf ME, Siegfried BD (2001) Toxicity and neurophysiological effects of fipronil and its oxidative sulfone metabolite on European corn borer larvae (Lepidoptera: Crambidae). *Pestic Biochem Physiol* 71:97–106
- Durham EW, Siegfried BD, Scharf ME (2002) In vivo and in vitro metabolism of fipronil by larvae of the European corn borer *Ostrinia nubilalis*. *Pest Manag Sci* 58:799–804
- Easton AH, Goulson D (2013) The neonicotinoid insecticide imidacloprid repels pollinating flies and beetles at field-realistic concentrations. *PLoS One* 8:e54819
- Edwards CA, Bohlen PJ (1996) Biology and ecology of earthworms, 3rd edn. Chapman and Hall, London
- EFSA (2006) Draft Assessment Report (DAR). Draft assessment report: initial risk assessment provided by the rapporteur member State Germany for the existing active substance imidacloprid
- Eisenback BM, Salom SM, Kok LT, Lagalante AF (2010) Lethal and sublethal effects of imidacloprid on Hemlock woolly Adelgid (Hemiptera: Adelgidae) and two introduced predator species. *J Econ Entomol* 103:1222–1234
- El Hassani AK, Dacher M, Gauthier M, Armengaud C (2005) Effects of sublethal doses of fipronil on the behavior of the honeybee (*Apis mellifera*). *Pharmacol Biochem Behav* 82:30–39
- El Hassani AK, Dacher M, Gary V, Lambin M, Gauthier M, Armengaud C (2008) Effects of sublethal doses of acetamiprid and thiamethoxam on the behavior of the honeybee (*Apis mellifera*). *Arch Environ Contam Toxicol* 54:653–661
- El Hassani AK, Dupuis JP, Gauthier M, Armengaud C (2009) Glutamatergic and GABAergic effects of fipronil on olfactory learning and memory in the honeybee. *Invert Neurosci* 9:91–100
- Elbert A, Becker B, Hartwig J, Erdelen C (1991) Imidacloprid—a new systemic insecticide. *Plant Prot Nachr Bayer* 44:113–136
- Epstein LH, Robinson JL, Roemmich JN, Marusewski A (2011) Slow rates of habituation predict greater zBMI gains over 12 months in lean children. *Eat Behav* 12:214–218
- European Commission (2010) Commission Directive 2010/21/EU of 12 March 2010 amending Annex I to Council Directive 91/414/EEC as regards the specific provisions relating to clothianidin, thiamethoxam, fipronil and imidacloprid
- Fagin D (2012) The learning curve. *Nature* 490:462–465. doi:10.1038/490462a
- Fang Q, Huang CH, Ye GY, Yao HW, Cheng JA et al (2008) Differential fipronil susceptibility and metabolism in two rice stem borers from China. *J Econ Entomol* 101:1415–1420
- Farooqui T (2013) A potential link among biogenic amines based pesticides, learning and memory, and colony collapse disorder: a unique hypothesis. *Neurochem Int* 62:122–136
- Faucon JP, Aurieres C, Drajnudel P, Mathieu L, Ribiere M, Martel AC, Zeggane S, Chauzat MP, Aubert MFA (2005) Experimental study on the toxicity of imidacloprid given in syrup to honey bee (*Apis mellifera*) colonies. *Pest Manag Sci* 61:111–125
- Feber RE, Firbank LG, Johnson PJ, Macdonald DW (1997) The effects of organic farming on pest and non-pest butterfly abundance. *Agric Ecosyst Environ* 64:133–139
- Feltham H, Park K, Goulson D (2014) Field realistic doses of pesticide imidacloprid reduce bumblebee pollen foraging efficiency. *Ecotoxicology* 37:301–308
- Fischer J, Müller T, Spatz A-K, Greggers U, Grünwald B et al (2014) Neonicotinoids interfere with specific components of navigation in honeybees. *PLoS One* 9:e91364
- Ford KA, Casida JE (2006) Unique and common metabolites of thiamethoxam, clothianidin, and dinotefuran in mice. *Chem Res Toxicol* 19:1549–1556
- Fossen M (2006) Environmental fate of imidacloprid. Environmental monitoring department of pesticide regulation, California, USA
- Franklin MT, Winston ML, Morandin LA (2004) Effects of clothianidin on *Bombus impatiens* (Hymenoptera: Apidae) colony health and foraging behaviour. *J Econ Entomol* 97:369–373
- Frantziou G, Papsiki K, Sidiropoulou B, Lazaridis G, Theophilidis G, Mavragani-Tsipidou P (2008) Evaluation of insecticidal and genotoxic effects of imidacloprid and acetochlor in *Drosophila melanogaster*. *J Appl Entomol* 132:583–590
- Freemark K, Boutin C (1995) Impacts of agricultural herbicide use on terrestrial wildlife in temperate landscapes: a review with special reference to North America. *Agric Ecosyst Environ* 52:67–91
- Funayama K, Ohsumi S (2007) Control effect of neonicotinoid insecticides on apple leaf miner, *Phyllonorycter ringoniella* (Lepidoptera: Gracillariidae). *Annu Rep Soc Plant Protect N Jpn* 58:156–158
- Galvanho JP, Carrera MP, Moreira DDO, Erthal M, Silva CP, Samuels RI (2013) Imidacloprid inhibits behavioral defences of the leaf-cutting ant *Acromyrmex subterraneus subterraneus* (Hymenoptera:

- Formicidae). *J Insect Behav* 26:1–13. doi:10.1007/S10905-012-9328-6
- Gauthier M (2010) State of the art on insect nicotinic acetylcholine receptor function in learning and memory. In: Thany S (ed) *Insect nicotinic acetylcholine receptors*. Springer, Berlin
- Gawleta N, Zimmermann Y, Eltz T (2005) Repellent foraging scent recognition across bee families. *Apidologie* 36:325–330
- Geiger F, Bengtsson J, Berendse F, Weisser WW, Emmerson M, Morales MB, Ceryngier P, Liira J, Tschamtker T, Winqvist C, Eggers S, Bommarco R, Pärt T, Bretagnolle V, Plantegenest M, Clement LW, Dennis C, Palmer C, Oñate JJ, Guerrero I, Hawro V, Aavik T, Thies C, Flohre A, Hånke S, Fischer C, Goedhart PW, Inchausti P (2010) Persistent negative effects of pesticides on biodiversity and biological control potential on European farmland. *Basic Appl Ecol* 11:97–105
- Gill RJ, Ramos-Rodriguez O, Raine NE (2012) Combined pesticide exposure severely affects individual- and colony-level traits in bees. *Nature* 491:105–108
- Girolami V, Mazzon L, Squartini A, Mori N, Marzaro M, Bernardo AD, Tapparo A (2009) Translocation of neonicotinoid insecticides from coated seeds to seedling guttation drops: a novel way of intoxication for bees. *J Econ Entomol* 102:1808–1815
- Girolami V, Marzaro M, Vivan L, Mazzon L, Greatti M, Giorio C, Marton D, Tapparo A (2012) Fatal powdering of bees in flight with particulates of neonicotinoids seed coating and humidity implication. *J Appl Entomol* 136:17–26
- Giurfa M (1993) The repellent scent-mark of the honeybee *Apis mellifera ligustica* and its role as communication cue during foraging. *Insect Soc* 40:59–67
- Gomez-Eyles JL, Svendsen C, Lister L, Martin H, Hodson ME, Spurgeon DJ (2009) Measuring and modelling mixture toxicity of imidacloprid and thiacloprid on *Caenorhabditis elegans* and *Eisenia fetida*. *Ecotoxicol Environ Saf* 72:71–79
- Goulson D (2003) Effects of introduced bees on native ecosystems. *Annu Rev Ecol Syst* 34:1–26
- Goulson D (2010) *Bumblebees: behaviour, ecology and conservation*. Oxford University Press, Oxford
- Goulson D, Chapman JW, Hughes WOH (2001) Discrimination of unrewarding flowers by bees; direct detection of rewards and use of repellent scent marks. *J Insect Behav* 14:669–678
- Gradish AE, Scott-Dupree CD, Shipp L, Harris CR, Ferguson G (2010) Effect of reduced risk pesticides for use in greenhouse vegetable production on *Bombus impatiens* (Hymenoptera: Apidae). *Pest Manag Sci* 66:142–146
- Graystock P, Yates K, Darvill B, Goulson D, Hughes WOH (2013a) Emerging dangers: deadly effects of an emergent parasite in a new pollinator host. *J Invertebr Pathol* 114:114–119
- Graystock P, Yates K, Evison S, Darvill B, Goulson D, Hughes WOH (2013b) The Trojan hives: pollinator pathogens, imported and distributed in bumblebee colonies. *J Appl Ecol* 50:1207–1215
- Greatti M, Sabatini AG, Barbattini R, Rossi S, Stravisi A (2003) Risk of environmental contamination by the active ingredient imidacloprid used for corn seed dressing. Preliminary results. *Bull Insect* 56:69–72
- Greenleaf S, Kremen C (2006) Wild bees enhance honey bees' pollination of hybrid sunflower. *Proc Natl Acad Sci U S A* 103:13890–13895
- Gregor A, Ellis JD (2011) Cell death localization in situ in laboratory reared honey bee (*Apis mellifera* L.) larvae treated with pesticides. *Pestic Biochem Physiol* 99:200–207
- Gross M (2008) Pesticides linked to bee deaths. *Curr Biol* Vol 18:684
- Guez D, Suchail S, Gauthier M, Maleszka R, Belzunces LP (2001) Contrasting effects of imidacloprid on habituation in 7- and 8-day-old honeybees (*Apis mellifera*). *Neurobiol Learn Mem* 76:183–191
- Guez D, Belzunces LP, Maleszka R (2003) Effects of imidacloprid metabolites on habituation in honeybees suggest the existence of two subtypes of nicotinic receptors differentially expressed during adult development. *Pharmacol Biochem Behav* 75:217–222
- Gupta S, Gajbhiye VT (2007) Persistence of acetamiprid in soil. *Bull Environ Contam Toxicol* 78:349–352
- Han P, Niu CY, Lei CL, Cui JJ, Desneux N (2010) Use of an innovative T-tube maze assay and the proboscis extension response assay to assess sublethal effects of GM products and pesticides on learning capacity of the honey bee *Apis mellifera* L. *Ecotoxicology* 19:1612–1619
- Hatjina F, Papaefthymiou C, Charistos L, Dogaroglu T, Bouga M, Emmanouil C, Arnold G (2013) Sublethal doses of imidacloprid decreased size of hypopharyngeal glands and respiratory rhythm of honeybees in vivo. *Apidologie* 44:467–480
- Hayasaka D, Korenaga T, Suzuki K, Saito F, Sánchez-Bayo F, Goka K (2012) Cumulative ecological impacts of two successive annual treatments of imidacloprid and fipronil on aquatic communities of paddy mesocosms. *Ecotoxicol Environ Saf* 80:355–62. doi:10.1016/j.ecoenv.2012.04.004
- Haynes KF (1988) Sublethal effects of neurotoxic insecticides on insect behavior. *Annu Rev Entomol* 33:149–168
- He YP, Gao CF, Cao MZ, Chen WM, Huang LQ et al (2007) Survey of susceptibilities to monosultap, triazophos, fipronil, and abamectin in *Chilo suppressalis* (Lepidoptera: Crambidae). *J Econ Entomol* 100:1854–1861
- He YP, Gao CF, Chen WM, Huang LQ, Zhou WJ et al (2008) Comparison of dose responses and resistance ratios in four populations of the rice stem borer, *Chilo suppressalis* (Lepidoptera: Pyralidae), to 20 insecticides. *Pest Manag Sci* 64:308–315
- He YP, Zhang JF, Gao CF, Su JY, Chen JM et al (2013) Regression analysis of dynamics of insecticide resistance in field populations of *Chilo suppressalis* (Lepidoptera: Crambidae) during 2002–2011 in China. *J Econ Entomol* 106:1832–1837
- Henry M, Beguin M, Requier F, Rollin O, Odoux JF, Aupinel P, Aptel J, Tchamitchian S, Decourtaye A (2012a) A common pesticide decreases foraging success and survival in honey bees. *Science* 336:348–350
- Henry M, Beguin M, Requier F, Rollin O, Odoux J, Aupinel P, Aptel J, Tchamitchian S, Decourtaye A (2012b) Response to comment on “A common pesticide decreases foraging success and survival in honey bees”. *Science* 337:1453
- Hill TA, Foster RE (2000) Effect of insecticides on the diamondback moth (Lepidoptera: Plutellidae) and its parasitoid *Diadegma insulare* (Hymenoptera: Ichneumonidae). *J Econ Entomol* 93:763–768
- Hoi KK, Daborn PJ, Battlay P, Robin C, Batterham P, O'Hair RA, Donald WA (2014) Dissecting the insect metabolic machinery using twin ion mass spectrometry: a single P450 enzyme metabolizes the insecticide imidacloprid in vivo. *Anal Chem* 86(7):3525–3532
- Huang SJ, Xu JF, Han ZJ (2006) Baseline toxicity data of insecticides against the common cutworm *Spodoptera litura* (Fabricius) and a comparison of resistance monitoring methods. *Int J Pest Manag* 52:209–213
- Illarionov AI (1991) Toxic effect of some insecticides on the honeybee. *Agrokhimika* 8:121–125
- Iwasa T, Motoyama N, Ambrose JT, Roe MR (2004) Mechanism for the differential toxicity of neonicotinoid insecticides in the honey bee, *Apis mellifera*. *Crop Prot* 23:371–378
- James DG, Price TS (2002) Fecundity in twospotted spider mite (Acari: Tetranychidae) is increased by direct and systemic exposure to imidacloprid. *J Econ Entomol* 95:729–732
- Jeschke P, Nauen R, Schindler M, Elbert A (2010) Overview of the status and global strategy for neonicotinoids. *J Agric Food Chem* 59:2897–2908
- Jinguji H, Thuyet DQ, Ueda T, Watanabe H (2013) Effect of imidacloprid and fipronil pesticide application on *Sympetrum infuscatum* (Libellulidae: Odonata) larvae and adults. *Paddy Water Environ* 11:277–284

- Jones MM, Robertson JL, Weinzierl RA (2010) Susceptibility of Oriental fruit moth (Lepidoptera: Tortricidae) larvae to selected reduced-risk insecticides. *J Econ Entomol* 103:1815–1820
- Jones MM, Robertson JL, Weinzierl RA (2012) Toxicity of thiamethoxam and mixtures of chlorantraniliprole plus acetamiprid, esfenvalerate, or thiamethoxam to neonates of oriental fruit moth (Lepidoptera: Tortricidae). *J Econ Entomol* 105:1426–1431
- Kalajdzic P, Oehler S, Reczko M, Pavlidi N, Vontas J, Hatzigeorgiou AG, Savakis C (2012) Use of mutagenesis, genetic mapping and next generation transcriptomics to investigate insecticide resistance mechanisms. *PLoS One* 7:e40296. doi:10.1371/journal.pone.0040296
- Kalajdzic P1, Markaki M, Oehler S, Savakis C (2013) Imidacloprid does not induce Cyp genes involved in insecticide resistance of a mutant *Drosophila melanogaster* line. *Food Chem Toxicol* 60:355–359. doi:10.1016/j.fct.2013.07.080
- Kather R, Drijfhout FP, Martin SJ (2011) Task group differences in cuticular lipids in the honey bee *Apis mellifera*. *J Chem Ecol* 37: 205–212
- Khani A, Ahmadi F, Ghadamyari M (2012) Side effects of imidacloprid and abamectin on the Mealybug destroyer, *Cryptolaemus montrouzieri*. *Trakia J Sci* 10:30–35
- Kilpatrick AL, Hagerty AM, Tumipseed SG, Sullivan MJ, Bridges WC (2005) Activity of selected neonicotinoids and dicofol on non-target arthropods in cotton: implications in insect management. *J Econ Entomol* 98:814–820
- Kinkler H, Löser S, Rehnelt K (1987) 10 Jahre Erforschung des Moselapollonfalter (Parnassius apollo vinningensis STICHEL 1899, Lepidoptera, Papilionidae) im modernen Weinbaugebiet der Mosel—ein Beitrag zu seiner Rettung. *Commun Assoc rheinisch-westfälischer Lepidopterologen* 5:74–96
- Klein AM, Vaissie're BE, Cane JH, Steffan-Dewenter I, Cunningham SA, Kremen C, Tscharntke T (2007) Importance of pollinators in changing landscapes for world crops. *Proc R Soc B* 274:303–313
- Kluser S, Neumann P, Chauzat MP, Pettis JS (2011) UNEP emerging issues: global honey bee colony disorder and other threats to insect pollinators. <http://www.unep.org>. Accessed 25 October 2012
- Knight AL (2010) Cross-resistance between azinphos-methyl and acetamiprid in populations of codling moth, *Cydia pomonella* (L.) (Lepidoptera: Tortricidae), from Washington State. *Pest Manag Sci* 66:865–874
- Knight ME, Martin AP, Bishop S, Osborne JL, Hale RJ, Sanderson RA, Goulson D (2005) An interspecific comparison of foraging range and nest density of four bumblebee (*Bombus*) species. *Mol Ecol* 14: 1811–1820
- Knoepf JD, Vose JM, Michael JL, Reynolds BC (2012) Imidacloprid movement in soils and impacts on soil microarthropods in Southern Appalachian eastern hemlock stands. *J Environ Qual* 41:469–478
- Knollengberg WG, Merritt RW, Lawson DL (1985) Consumption of leaf litter by *Lumbricus terrestris* (Oligochaeta) on a Michigan woodland floodplain. *Am Midl Nat J* 113:1–6
- Kramarz P, Stark JD (2003) Population level effects of cadmium and the insecticide, imidacloprid to the Parasitoid, *Aphidius ervi* after exposure through its host, the pea aphid, *Acyrtosiphon pisum*. *Biol Control* 27:310–314
- Kreutzweiser DP, Sibley PK (2013) Aquatic communities: pesticide impacts. In: Jorgensen SE (ed) *Encyclopedia of environmental management*. Taylor and Francis, New York, pp 312–321
- Kreutzweiser DP, Good KP, Chartrand DT, Scarr TA, Thompson DG (2008a) Are leaves that fall from imidacloprid-treated maple trees toxic to control Asian longhorned beetles toxic to non-target decomposer organisms? *J Environ Qual* 37:639–646
- Kreutzweiser DP, Good KP, Chartrand DT, Scarr TA, Holmes SB, Thompson DG (2008b) Effects on litter-dwelling earthworms and microbial decomposition of soil-applied imidacloprid for control of wood-boring insects. *Pest Manag Sci* 64:112–118
- Kreutzweiser DP, Thompson DG, Scarr TA (2009) Imidacloprid in leaves from systemically treated trees may inhibit litter breakdown by non-target invertebrates. *Ecotoxicol Environ Saf* 72:1053–1057
- Krischik VA, Landmark AL, Heimpel GE (2007) Soil-applied imidacloprid is translocated to nectar and kills nectar-feeding *Anagyrus pseudococci* (Girault) (Hymenoptera: Encyrtidae). *Environ Entomol* 36:1238–1245
- Kromp B (1999) Carabid beetles in sustainable agriculture: a review on pest control efficacy, cultivation impacts and enhancement. *Agric Ecosyst Environ* 74:187–228
- Krupke CH, Hunt GJ, Eitzer BD, Andino G, Given K (2012) Multiple routes of pesticide exposure for honey bees living near agricultural fields. *PLoS One* 7:e29268
- Kullik SA, Sears MK et al (2011) Sublethal effects of Cry 1F Bt corn and clothianidin on black cutworm (Lepidoptera: Noctuidae) larval development. *J Econ Entomol* 104:484–93
- Kunkel BA, Held DW, Potter AD (1999) Impact of halofenozide, imidacloprid, and bendiocarb on beneficial invertebrates and predatory activity in turfgrass. *J Econ Entomol* 92:922–930
- Kunkel BA, Held DW, Potter DA (2001) Lethal and sub-lethal effects of bendiocarb, halofenozide, and imidacloprid on *Harpalus pennsylvanicus* (Coleoptera: Carabidae) following different modes of exposure in turfgrass. *J Econ Entomol* 94:60–67
- Kurwadkar ST, Dewinne D, Wheat R, McGahan DG, Mitchell FL (2013) Time dependent sorption behaviour of dinotefuran, imidacloprid and thiamethoxam. *J Environ Sci Health B* 48:237–242
- Lambin M, Armengaud C, Raymond S, Gauthier M (2001) Imidacloprid-induced facilitation of the proboscis extension reflex habituation in the honeybee. *Arch Insect Biochem Physiol* 48:129–134
- Landis DA, Wratten SD, Gurr GM (2000) Habitat management to conserve natural enemies of arthropod pests in agriculture. *Annu Rev Entomol* 45:175–201
- Larson JL, Redmond CT, Potter DA (2012) Comparative impact of an anthranilic diamide and other insecticidal chemistries on beneficial invertebrates and ecosystem services in turfgrass. *Pest Manag Sci* 68:740–748
- Laycock I, Lenthall KM, Barratt AT, Cresswell JE (2012a) Effects of imidacloprid, a neonicotinoid pesticide, on reproduction in worker bumble bees (*Bombus terrestris*). *Ecotoxicology* 21:1937–1945
- Laycock I, Lenthall KM, Barratt AT, Cresswell JE (2012b) Erratum to: Effects of imidacloprid, a neonicotinoid pesticide, on reproduction in worker bumble bees (*Bombus terrestris*). *Ecotoxicology* 21: 1946–1946
- LeBlanc HMK, Culp JM, Baird DJ, Alexander AC, Cessna AJ (2012) Single versus combined lethal effects of three agricultural insecticides on larvae of the freshwater insect *Chironomus dilutes*. *Arch Environ Contam Toxicol* 63:378–390
- Lee KW (1985) *Earthworms: their ecology and relationship with soils and land use*. Academic, Sydney
- Lee JC, Menalled FD, Landis DA (2001) Refuge habitats modify impact of insecticide disturbance on carabid beetle communities. *J Appl Ecol* 38:472–483
- Li AG, Yang YH, Wu SW, Li C, Wu YD (2006) Investigation of resistance mechanisms to fipronil in diamondback moth (Lepidoptera: Plutellidae). *J Econ Entomol* 99:914–919
- Li XT, Huang QC, Yuan JZ, Tang ZH (2007) Fipronil resistance mechanisms in the rice stem borer, *Chilo suppressalis* Walker. *Pestic Biochem Physiol* 89:169–174
- Liang GM, Chen W, Liu TTX (2003) Effects of three neem-based insecticides on diamondback moth (Lepidoptera: Plutellidae). *Crop Prot* 22:333–340
- Liess M, Beketov M (2011) Traits and stress—keys to identify community effects at low toxicant level. *Ecotoxicology* 20:1328–1340
- Liess M, Beketov MA (2012) Rebuttal related to “Traits and stress—keys to identify community effects of low levels of toxicants in test systems”. *Ecotoxicology* 21:300–303

- Liess M, von der Ohe PC (2005) Analyzing effects of pesticides on invertebrate communities in streams. *Environ Toxicol Chem* 24: 954–965
- Liess M, Schäfer RB, Schriever CA (2008) The footprint of pesticide stress in communities—species traits reveal community effects of toxicants. *Sci Total Environ* 406:484–90
- Liess M, Foit K, Becker A, Hassold E, Dolciotti I, Kattwinkel M, Duquesne S (2013) Culmination of low-dose pesticide effects. *Environ Sci Tech* 47:8862–8868
- Lintott DR (1992) NTN 33893 (240 FS formulation): acute toxicity to the mysid, *Mysidopsis bahia*, under flow-through conditions. Toxicon Environmental Sciences, Jupiter, Florida (performing laboratory). Miles Incorporated, Kansas City, Missouri (submitting company). 43 pp. Miles Report No. 103845
- Little EE (1990) Behavioural toxicology; stimulating challenges for a growing discipline. *Environ Toxicol Chem* 9:1–2
- Longley M, Sotherton NW (1997) Factors determining the effects of pesticides upon butterflies inhabiting arable farmland. *Agric Ecosyst Environ* 61:1–12
- Lucas E, Giroux S, Demougeot S, Duchesne RM, Coderre D (2004) Compatibility of a natural enemy, *Coleomegilla maculata lengi* (Col., Coccinellidae) and four insecticides used against the Colorado potato beetle (Col., Chrysomelidae). *J Appl Entomol* 128:233–239
- Lukancic S, Zibrat U, Mezek T, Jerebic A, Simcic T, Brancelj A (2010) Effects of exposing two non-target crustacean species, *Asellus aquaticus* L., and *Gammarus fossarum* Koch., to atrazine and imidacloprid. *Bull Environ Contam Toxicol* 84(1):85–90. doi:10.1007/s00128-009-9854-x
- Luo Y, Zang Y, Zhong Y, Kong Z (1999) Toxicological study of two novel pesticides on earthworm *Eisenia fetida*. *Chemosphere* 39: 2347–2356
- Lydy M, Belden J, Wheelock C, Hammock B, Denton D (2004) Challenges in regulating pesticide mixtures. *Ecol Soc* 9:1
- Magalhaes LC, Walgenbach JF (2011) Life stage toxicity and residual activity of insecticides to codling moth and Oriental fruit moth (Lepidoptera: Tortricidae). *J Econ Entomol* 104:1950–1959
- Maisonnette A, Lenoir JC, Beslay D, Crauser D, Le Conte Y (2010) E-b-Ocimene, a volatile brood pheromone involved in social regulation in the honey bee colony (*Apis mellifera*). *PLoS One* 5
- Mani S, Medrzycki P, Porrini C (2010) The puzzle of honey bee losses: a brief review. *Bull Insect* 63:153–160
- Mann RS, Uppal SK, Sharma S, Mann KK (2009) Soil efficacy of fipronil to early stage pests of sugarcane, and its effect on development on *Chilo infuscatellus* Snellen (Crambidae: Lepidoptera). *Int J Pest Manag* 55:307–315
- Marzaro M, Vivan L, Targa A, Mazzon L, Mori N, Greatti M, Petrucco Toffolo E, Di Bernardo A, Giorio C, Marton D, Tapparo A, Girolami V (2011) Lethal aerial powdering of honey bees with neonicotinoids from fragments of maize seed coat. *Bull Insect* 64:118–125
- Mason R, Tennekes HA, Sanchez-Bayo F, Jepsen PU (2013) Immune suppression by neonicotinoid insecticides at the root of global wildlife declines. *J Environ Immunol Toxicol* 1:3–12
- Maus C, Nauen R (2010) Response to the publication: Tennekes H.A. The significance of the Druckey-Küpfmüller equation for risk assessment—the toxicity of neonicotinoid insecticides is reinforced by exposure time. *Toxicology* 280:176–177. doi:10.1016/j.tox.2010.11.014
- Maxim L, van der Sluijs JP (2007) Uncertainty: cause or effect of stakeholders' debates? Analysis of a case study: the risk for honeybees of the insecticide Gaucho®. *Sci Total Environ* 376:1–17
- Maxim L, van der Sluijs JP (2013) Seed-dressing systemic insecticides and honeybees, chapter 16. In: European Environment Agency (ed.) Late lessons from early warnings: science, precaution, innovation. European Environment Agency (EEA) report 1/2013, Copenhagen. p. 401–438
- Mayer DF, Lunden JD (1997) Effects of imidacloprid insecticide on three bee pollinators. *Hortic Sci* 29:93–97
- McKern JA, Johnson DT, Lewis BA (2007) Biology and control of the raspberry crown borer (Lepidoptera: Sesiidae). *J Econ Entomol* 100: 398–404
- Medrzycki P, Montanari R, Bortolotti L, Sabatini AG, Maini S, Porrini C (2003) Effects of imidacloprid administered in sublethal doses on honey bee behaviour. Laboratory tests. *Bull Insect* 56:59–62
- Medrzycki P, Sgolastra F, Bogo G, Tosi S, Venturi S (2011) Influence of some experimental conditions on the results of laboratory toxicological tests on honeybees. XI International Symposium “Hazards of pesticides to bees” Wageningen, The Netherlands, November 2–4
- Mommaerts V, Reynders S, Boulet J, Besard L, Sterk G, Smagghe G (2010) Risk assessment for side-effects of neonicotinoids against bumblebees with and without impairing foraging behavior. *Ecotoxicology* 19:207–215
- Morandin LA, Winston ML (2003) Effects of novel pesticides on bumble bee (Hymenoptera: Apidae) colony health and foraging ability. *Environ Entomol* 32:555–563
- Moser SE, Obrycki JJ (2009) Non-target effects of neonicotinoid seed treatments: mortality of coccinellid larvae related to zoophytophagy. *Biol Control* 51:487–492
- Mostert MA, Schoeman AS, van der Merwe M (2000) The toxicity of five insecticides to earthworms of the *Pheretima* group, using an artificial soil test. *Pest Manag Sci* 56:1093–1097
- Mostert MA, Schoeman AS, van der Merwe M (2002) The relative toxicities of insecticides to earthworms of the *Pheretima* group (Oligochaeta). *Pest Manag Sci* 58:446–450
- Mota-Sanchez D, Wise JC, Poppen RV, Gut LJ, Hollingworth RM (2008) Resistance of codling moth, *Cydia pomonella* (L.) (Lepidoptera: Tortricidae), larvae in Michigan to insecticides with different modes of action and the impact on field residual activity. *Pest Manag Sci* 64:881–890
- Mullin CA, Frazier M, Frazier JL, Ashcraft S, Simonds R, van Engelsdorp D, Pettis JS (2010) High levels of miticides and agrochemicals in North American apiaries: implications for honey bee health. *PLoS One* 5:e9754
- Nauen R, Ebbinghaus-Kintscher U, Schmuck R (2001) Toxicity and nicotinic acetylcholine receptor interaction of imidacloprid and its metabolites in *Apis mellifera* (Hymenoptera: Apidae). *Pest Manag Sci* 57:577–586
- Nauen R, Ebbinghaus-Kintscher U, Salgado VL, Kaussmann M (2003) Thiamethoxam is a neonicotinoid precursor converted to clothianidin in insects and plants. *Pestic Biochem Physiol* 76:55–69
- Neumann P, Carreck NL (2010) Honey bee colony loss. *J Apic Res* 49:1–6, special issue
- Nguyen BK et al (2009) Does imidacloprid seed-treated maize have an impact on honey bee mortality? *J Econ Entomol* 102:616–623
- Nielsen SA, Brodsgaard CJ, Hansen H (2000) Effects on detoxification enzymes in different life stages of honey bees (*Apis mellifera* L., Hymenoptera: Apidae) treated with a synthetic pyrethroid (flumethrin). *Altern Lab Anim* 28:437–443
- Ninsin KD (2004a) Acetamiprid resistance and cross-resistance in the diamondback moth, *Plutella xylostella*. *Pest Manag Sci* 60:839–841
- Ninsin KD (2004b) Selection for resistance to acetamiprid and various other insecticides in the diamondback moth, *Plutella xylostella* (L.) (Lep., Plutellidae). *J Appl Entomol* 128:445–451
- Ninsin KD, Miyata T (2003) Monitoring acetamiprid resistance in the diamondback moth, *Plutella xylostella* (L.) (Lepidoptera: Yponomeutidae). *Appl Entomol Zool* 38:517–521
- Ninsin KD, Tanaka T (2005) Synergism and stability of acetamiprid resistance in a laboratory colony of *Plutella xylostella*. *Pest Manag Sci* 61:723–727
- Ninsin KD, Mo JC, Miyata T (2000) Decreased susceptibilities of four field populations of the diamondback moth, *Plutella xylostella* (L.)

- (Lepidoptera: Yponomeutidae), to acetamiprid. Appl Entomol Zool 35:591–595
- Nyman AM, Hintermeister A, Schirmer K, Ashauer R (2013) The insecticide imidacloprid causes mortality of the freshwater Amphipod *Gammarus pulex* by interfering with feeding behavior. PLoS One 8: e62472
- Oldroyd PB (2007) What's killing American honey bees? PLoS Biol 5: e168
- Oliveira RA, Roat TC, Carvalho SM, Malaspina O (2013) Side-effects of thiamethoxam on the brain and midgut of the Africanized honeybee *Apis mellifera* (Hymenoptera: Apidae). Environ Toxicol 28 doi: 10.1002/tox.21842
- Osborne JL, Martin AP, Shortall CR, Todd AD, Goulson D, Knight ME, Hale RJ, Sanderson RA (2008) Quantifying and comparing bumblebee nest densities in gardens and countryside habitats. J Appl Ecol 45:784–792
- Osterberg JS, Darnell KM, Blickley TM, Romano JA, Rittschof D (2012) Acute toxicity and sub-lethal effects of common pesticides in post-larval and juvenile blue crabs, *Callinectes sapidus*. J Exp Mar Biol Ecol 424–425:5–14
- Papachristos DP, Milonas PG (2008) Adverse effects of soil applied insecticides on the predatory coccinellid *Hippodamia undecimnotata* (Coleoptera: Coccinellidae). Biol Control 47:77–81
- Paradis D, Bérail G, Bonmatin JM, Belzunces LP (2013) Sensitive analytical methods for 22 relevant insecticides of 3 chemical families in honey by GC-MS/MS and LC-MS/MS. Anal Bioanal Chem 406:621–633
- Peck DC (2009) Comparative impacts of white grub (Coleoptera: Scarabaeidae) control products on the abundance of non-target soil-active arthropods in turfgrass. Pedobiologia 52:287–299
- Peck DC, Olmstead D (2010) Neonicotinoid insecticides disrupt predation on the eggs of turf-infesting scarab beetles. Bull Entomol Res 100:689–700
- Pedibhotla VK, Hall FR, Holmsen J (1999) Deposit characteristics and toxicity of fipronil formulations for tobacco budworm (Lepidoptera: Noctuidae) control on cotton. Crop Prot 18:493–499
- Pestana JLT, Alexander AC, Culp JM, Baird DJ, Cessna AJ, Soares A (2009) Structural and functional responses of benthic invertebrates to imidacloprid in outdoor stream mesocosms. Environ Pollut 157: 2328–2334
- Pettis JS, van Engelsdorp D, Johnson J, Dively G (2012) Pesticide exposure in honey bees results in increased levels of the gut pathogen *Nosema*. Naturwissenschaften 99:153–158
- Pilling E, Campbell P, Coulson M, Ruddle N, Tornier I (2013) A four-year field program investigating long-term effects of repeated exposure of honey bee colonies to flowering crops treated with thiamethoxam. PLoS One 8:e77193
- Pochi D, Biocca M, Fanigliulo R, Pulcini P, Conte E (2012) Potential exposure of bees, *Apis mellifera* L., to particulate matter and pesticides derived from seed dressing during maize sowing. Bull Environ Contam Toxicol 89:354–61
- Prabhaker N, Morse JG, Castle SJ, Naranjo SE, Henneberry TJ, Toscano NC (2007) Toxicity of seven foliar insecticides to four insect parasitoids attacking citrus and cotton pests. J Econ Entomol 100:1053–1061
- Prabhaker N, Castle SJ, Naranjo SE, Toscano NC, Morse JG (2011) Compatibility of two systemic neonicotinoids, imidacloprid and thiamethoxam, with various natural enemies of agricultural pests. J Econ Entomol 104:773–781
- Rahmani S, Bandani AR, Sabahi Q (2013) Effects of thiamethoxam in sublethal concentrations, on life expectancy (ex) and some other biological characteristics of *Hippodamia variegata* (Goeze) (Coleoptera: Coccinellidae). Int Res J App Basic Sci 4:556–560
- Ramasubramanian T (2013) Persistence and dissipation kinetics of clothianidin in the soil of tropical sugarcane ecosystem. Water Air Soil Pollut 224:1468
- Ramirez-Romero R, Chauvaud J, Pham-Delegue MH (2005) Effects of Cry1Ab protoxin, deltamethrin and imidacloprid on the foraging activity and the learning performances of the honeybee *Apis mellifera*, a comparative approach. Apidologie 36: 601–611
- Rasmussen JJ, McKnight US, Loinaz MC, Thomsen NI, Olsson ME, Bjerg PL, Binning PJ, Kronvang B (2013) A catchment scale evaluation of multiple stressor effects in headwater streams. Sci Total Environ 442:420–31
- Reinhard J, Srinivasan MV, Zhang SW (2004) Scent-triggered navigation in honeybees. Nature 427:411–411
- Reyes M, Franck P, Charmillot PJ, Ioriatti C, Olivares J et al (2007) Diversity of insecticide resistance mechanisms and spectrum in European populations of the Codling moth, *Cydia pomonella*. Pest Manag Sci 63:890–902
- Rhainds M, Sadof C (2009) Control of bagworms (Lepidoptera: Psychidae) using contact and soil-applied systemic insecticides. J Econ Entomol 102:1164–1169
- Riaz MA, Chandor-Proust A, Dauphin-Villemant C, Poupardin R, Jones CM, Strode C, Régent-Kloekner M, David JP, Reynaud S (2013) Molecular mechanisms associated with increased tolerance to the neonicotinoid insecticide imidacloprid in the dengue vector *Aedes aegypti*. Aquat Toxicol 126:326–337. doi:10.1016/j.aquatox.2012.09.010
- Richarz N, Neumann D, Wipking W (1989) Untersuchungen zur Ökologie des Apollofalters (Parnassius apollo vinningensis Stichel 1899, Lepidoptera, Papilionidae) im Weinbaugebiet der unteren Mosel. Mitt der Assoc Rheinisch-Westfälischer Lepidopterologen 5:108–259
- Rodrick GB (2008) Effect of temperature, salinity, and pesticides on oyster hemocyte activity. Florida Water Resour J 86:4–14. doi:10.2175/SJWP(2008)1
- Rogers MA, Kirschik VA, Martin LA (2007) Effect of soil application of imidacloprid on survival of adult green lacewing, *Chrysoperla carnea* (Neuroptera: Chrysopidae), used for biological control in greenhouse. Biol Control 42:172–177
- Rondeau G, Sanchez-Bayo F, Tennekes HA, Decourtye A, Ramirez-Romero R, Desneux N (2014) Delayed and time-cumulative toxicity of imidacloprid in bees, ants and termites. Nat Sc Rep 4:5566. doi: 10.1038/srep05566
- Rundlöf M, Bengtsson J, Smith HG (2008) Local and landscape effects of organic farming on butterfly species richness and abundance. J Appl Ecol 45:813–820
- Rust MK, Reiersen DA, Klotz JH (2004) Delayed toxicity as a critical factor in the efficacy of aqueous baits for controlling Argentine ants (Hymenoptera: Formicidae). J Econ Entomol 97:1017–1024
- Sánchez-Bayo F (2006) Comparative acute toxicity of organic pollutants and reference values for crustaceans. I. Branchiopoda, Copepoda and Ostracoda. Environ Pollut 139:385–420
- Sánchez-Bayo F (2009) From simple toxicological models to prediction of toxic effects in time. Ecotoxicology 18:343–354
- Sanchez-Bayo F, Goka K (2014) Pesticide residues and bees—a risk assessment. PLoS One 9:e94482
- Sanchez-Bayo F, Kouchi G (2012) Evaluation of suitable endpoints for assessing the impacts of toxicants at the community level. Ecotoxicology 21:667–680. doi:10.1007/s10646-011-0823-x
- Saour G (2008) Effect of thiacloprid against the potato tuber moth *Phthorimaea operculella* Zeller (Lepidoptera: Gelechiidae). J Pest Sci 81:3–8
- Sardo AM, Soares AMVM (2010) Assessment of the effects of the pesticide imidacloprid on the behaviour of the aquatic Oligochaete *Lumbriculus variegatus*. Arch Environ Contam Toxicol 58:648–656
- Sarkar MA, Roy S, Kole RK, Chowdhury A (2001) Persistence and metabolism of imidacloprid in different soils of West Bengal. Pest Manag Sci 57:598–602

- Sayyed AH, Crickmore N (2007) Selection of a field population of diamondback moth (Lepidoptera: Plutellidae) with acetamiprid maintains, but does not increase, cross-resistance to pyrethroids. *J Econ Entomol* 100:932–938
- Sayyed AH, Wright DJ (2004) Fipronil resistance in the diamondback moth (Lepidoptera: Plutellidae): inheritance and number of genes involved. *J Econ Entomol* 97:2043–2050
- Schäfer RB, vd Ohe P, Rasmussen J, Kefford B, Beketov M, Schulz R, Liess M (2012) Thresholds for the effects of pesticides on invertebrate communities and leaf breakdown in stream ecosystems. *Environ Sci Tech* 46:5134–5142
- Schmidt A (1997) Zur aktuellen situation des mosel-apollofalters *parnassius apollo vinningensis* Stichel, 1899 (Lep., Papilionidae). *Melanargia* 9:38–47
- Schmuck R (2004) Effects of a chronic dietary exposure of the honeybee *Apis mellifera* (Hymenoptera: Apidae) to imidacloprid. *Arch Environ Contam Toxicol* 47:471–478
- Schmuck R, Schoning R, Stork A, Schramel O (2001) Risk posed to honeybees (*Apis mellifera* L. Hymenoptera) by an imidacloprid seed dressing of sunflowers. *Pest Manag Sci* 57:225–238
- Schmuck R, Stadler T, Schmidt HW (2003) Field relevance of a synergistic effect observed in the laboratory between an EBI fungicide and a chloronicotinyl insecticide in the honeybee (*Apis mellifera* L., Hymenoptera). *Pest Manag Sci* 59:279–286
- Schneider CW, Tautz J, Grünwald B, Fuchs S (2012) RFID tracking of sublethal effects of two neonicotinoid insecticides on the foraging behavior of *Apis mellifera*. *PLoS One* 7:e30023
- Scholer J, Krischik V (2014) Chronic exposure of imidacloprid and clothianidin reduce queen survival, foraging, and nectar storing in colonies of *Bombus impatiens*. *PLoS One* 9:e91573
- Scott-Dupree CD, Conroy L, Harris CR (2009) Impact of currently used or potentially useful insecticides for canola agroecosystems on *Bombus impatiens* (Hymenoptera: Apidae), *Megachile rotundata* (Hymenoptera: Megachilidae), and *Osmia lignaria* (Hymenoptera: Megachilidae). *J Econ Entomol* 102:177–182
- Seagraves MP, Lundgren JG (2012) Effects of neonicotinoid seed treatments on soybean aphid and its natural enemies. *J Pest Sci* 85:125–132
- Sergio M (2013) GMO and pesticide experiments in Hawaii: the poisoning of paradise. *Huffington Post*. http://www.huffingtonpost.com/maggie-sergio/gmo-pesticide-experiments_b_3513496.html
- Setamou M, Rodriguez D, Saldana R, Schwarzlose G, Palrang D et al (2010) Efficacy and uptake of soil-applied imidacloprid in the control of Asian citrus psyllid and a citrus leafminer, two foliar-feeding citrus pests. *J Econ Entomol* 103:1711–1719
- Sgolastra F, Renzi T, Draghetti S, Medrzycki P, Lodesani M, Maini S, Pottini C (2012) Effects of neonicotinoid dust from maize seed-dressing on honey bees. *Bull Insect* 65:273–280
- Sheehan C, Kirwan L, Connolly J, Bolger T (2008) The effects of earthworm functional diversity on microbial biomass and the microbial community level physiological profile of soils. *Eur J Soil Biol* 44:65–70
- Shi ZH, Guo SJ, Lin WC, Liu SS (2004) Evaluation of selective toxicity of five pesticides against *Plutella xylostella* (Lep: Plutellidae) and their side-effects against *Cotesia plutellae* (Hym: Braconidae) and *Oomyzus sokolowskii* (Hym: Eulophidae). *Pest Manag Sci* 60:1213–1219
- Siegfried BD, Spencer T, Marcon PCRG (1999) Susceptibility of European corn borer, *Ostrinia nubilalis* (Hubner) (Lepidoptera: Pyralidae) neonate larvae to fipronil. *J Agric Urban Entomol* 16: 273–278
- Simon-Delso N, Amaral-Rogers V, Belzunces LP, Bonmatin J-M, Chagnon M, Downs C, Furlan L, Gibbons DW, Giorio C, Girolami V, Goulson D, Kreutzweiser DP, Krupke C, Liess M, Long E, McField M, Mineau P, Mitchell EAD, Morrissey CA, Noome DA, Pisa L, Settele J, Stark JD, Tapparo A, Van Dyck H, Van Praagh J, Van der Sluijs JP, Whitehorn PR, Wiemers M (2014) Systemic insecticides (neonicotinoids and fipronil): trends, uses, mode of action and metabolites. *Environ Sci Pollut Res* (this issue)
- Sinha SN, Lakhani KH, Davis BNK (1990) Studies on the toxicity of insecticidal drift to the first instar larvae of the large white butterfly *Pieris brassicae* (Lepidoptera: Pieridae). *Ann Appl Biol* 116:27–41
- Smith SF, Kruschick VA (1999) Effects of systemic imidacloprid on *Coleomegilla maculata* (Coleoptera: Coccinellidae). *Environ Entomol* 28:1189–1195
- Smith JF, Catchot AL, Musser FR, Gore J (2013) Effects of aldicarb and neonicotinoid seed treatments on twospotted spider mite on cotton. *J Econ Entomol* 106:807–815
- Smodis Skerl MI, Velikonja Bolta S, Basa Cesnik H, Gregorc A (2009) Residues of pesticides in honeybee (*Apis mellifera carnica*) bee bread and in pollen loads from treated apple orchards. *Bull Environ Contam Toxicol* 83:374–7
- Song MY, Brown JJ (1998) Osmotic effects as a factor modifying insecticide toxicity on *Aedes* and *Artemia*. *Ecotoxicol Environ Saf* 41:195–202
- Song MY, Stark JD, Brown JJ (1997) Comparative toxicity of four insecticides, including imidacloprid and tebufenozide, to four aquatic arthropods. *Environ Toxicol Chem* 16:2494–2500
- Stapel JO, Cortesero AM, Lewis WJ (2000) Disruptive sublethal effects of insecticides on biological control: altered foraging ability and life span of a parasitoid after feeding on extrafloral nectar of cotton treated with systemic insecticides. *Biol Control* 17:243–249
- Stara J, Kocourek F (2007) Insecticidal resistance and cross-resistance in populations of *Cydia pomonella* (Lepidoptera: Tortricidae) in central Europe. *J Econ Entomol* 100:1587–95
- Stark JD, Banks JE (2003) Population-level effects of pesticides and other toxicants on arthropods. *Annu Rev Entomol* 48:505–519
- Stark JD, Jepson PC, Mayer DF (1995) Limitations to use of topical toxicity data for predictions of pesticide side effects in the field. *J Econ Entomol* 88:1081–1088
- Starnes K, Goh KS (2012) Detections of the neonicotinoid insecticide imidacloprid in surface waters of three agricultural regions of California, USA, 2010–2011. *Bull Environ Contam Toxicol*, 1–6
- Stokstad E (2007) The case of the empty hives. *Science* 316:970–972
- Stoughton SJ, Liber K, Culp J, Cessna A (2008) Acute and chronic toxicity of imidacloprid to the aquatic invertebrates *Chironomus tentans* and *Hyalella azteca* under constant- and pulse-exposure conditions. *Arch Environ Contam Toxicol* 54:662–673
- Stygar D, Michalczyk K, Dolezych B, Nakonieczny M, Migula P et al (2013) Digestive enzymes activity in subsequent generations of *Cameraria ohridella* larvae harvested from horse chestnut trees after treatment with imidacloprid. *Pestic Biochem Physiol* 105:5–12
- Suchail S, Guez D, Belzunces LP (2000) Characteristics of imidacloprid toxicity in two *Apis mellifera* subspecies. *Environ Toxicol Chem* 19: 1901–1905
- Suchail S, Guez D, Belzunces LP (2001) Discrepancy between acute and chronic toxicity induced by imidacloprid and its metabolites in *Apis mellifera*. *Environ Toxicol Chem* 20:2482–2486
- Suchail S, De Sousa G, Rahmani R, Belzunces LP (2004a) In vivo distribution and metabolism of [¹⁴C]-imidacloprid in different compartments of *Apis mellifera* L. *Pest Manag Sci* 60:1056–1062
- Suchail S, Debrauwer L, Belzunces LP (2004b) Metabolism of imidacloprid in *Apis mellifera*. *Pest Manag Sci* 60:291–296
- Symington CA (2003) Lethal and sublethal effects of pesticides on the potato tuber moth, *Phthorimaea operculella* (Zeller) (Lepidoptera: Gelechiidae) and its parasitoid *Orgilus lepidus* Muesebeck (Hymenoptera: Braconidae). *Crop Prot* 22:513–519
- Szczepaniec A, Creary SF, Laskowski KL, Nyrop JP, Raupp MJ (2011) Neonicotinoid insecticide imidacloprid causes outbreaks of spider mites on elm trees in urban landscapes. *PLoS One* 6:e20018

- Szczepaniec A, Raupp MJ, Parker RD, Kerns D, Eubanks MD (2013) Neonicotinoid insecticides alter induced defenses and increase susceptibility to spider mites in distantly related crop plants. *PLoS One* 8:e62620
- Tapparo A, Marton D, Giorio C, Zanella A, Solda' L, Marzaro M, Vivan L, Girolami V (2012) Assessment of the environmental exposure of honeybees to the particulate matter containing neonicotinoid insecticides coming from corn coated seeds. *Environ Sci Tech* 46:2592–2599
- Tarmann G (2009) Die Vinschger Trockenrasen - Ein Zustandsbericht auf basis der bioindikatoren Tagfalter und widderchen (Lepidoptera: Rhopalocera, Zygaenidae). *Sci Yearb Tyrolean State Mus* 2:307–350
- Tasei JN, Lerin J, Ripault G (2000) Sub-lethal effects of imidacloprid on bumblebees, *Bombus terrestris* (Hymenoptera: Apidae), during a laboratory feeding test. *Pest Manag Sci* 56:784–788
- Tasei JN, Ripault G, Rivault E (2001) Hazards of imidacloprid seed coating to *Bombus terrestris* (Hymenoptera: Apidae) when applied to sunflower. *J Econ Entomol* 94:623–627
- Taverner PD, Sutton C, Cunningham NM, Dyson C, Lucas N et al (2011) Efficacy of several insecticides alone and with horticultural mineral oils on light brown apple moth (Lepidoptera: Tortricidae) eggs. *J Econ Entomol* 104:220–224
- Taverner PD, Sutton C, Cunningham NM, Myers SW (2012) The potential of mineral oils alone and with reduced rates of insecticides for the control of light brown apple moth, *Epiphyas postvittana* (Walker) (Lepidoptera: Tortricidae), on nursery plants. *Crop Prot* 42:83–87
- Teeters BS, Johnson RM, Ellis MD, Siegfried BD (2012) Using video-tracking to assess sublethal effects of pesticides on honey bees (*Apis mellifera* L.). *Environ Toxicol Chem* 31:1349–1354
- Tennekes HA (2010) The significance of the Druckrey-Küpfmüller equation for risk assessment—the toxicity of neonicotinoid insecticides to arthropods is reinforced by exposure time. *Toxicology* 276:1–4
- Tennekes HA (2011) The significance of the Druckrey-Küpfmüller equation for risk assessment—the toxicity of neonicotinoid insecticides to arthropods is reinforced by exposure time: responding to a letter to the editor by Drs. C. Maus and R. Nauen of Bayer CropScience AG. *Toxicology* 280:173–175
- Tennekes HA, Sánchez-Bayo F (2012) Time-dependent toxicity of neonicotinoids and other toxicants: implications for a new approach to risk assessment. *J Environ Anal Toxicol* S4:001. doi:10.4172/2161-0525.S4-001
- Tennekes HA, Sánchez-Bayo F (2013) The molecular basis of simple relationships between exposure concentration and toxic effects with time. *Toxicology* 309:39–51
- Thany SH, Gauthier M (2005) Nicotine injected into the antennal lobes induces a rapid modulation of sucrose threshold and improves short-term memory in the honeybee *Apis mellifera*. *Brain Res* 1039:216–219
- Thany SH, Lenaers G, Crozatier M, Armengaud C, Gauthier M (2003) Identification and localization of the nicotinic acetylcholine receptor alpha3 mRNA in the brain of the honeybee, *Apis mellifera*. *Insect Mol Biol* 12:255–262
- Thompson HM, Maus C (2007) The relevance of sublethal effects in honey bee testing for pesticide risk assessment. *Pest Manag Sci* 63:1058–1061
- Tomé HVV, Martins GF, Lima MAP, Campos LAO, Guedes RNC (2012) Imidacloprid-induced impairment of mushroom bodies and behavior of the native stingless bee *Melipona quadrifasciata anthidioides*. *PLoS One* 7:e38406
- Tomizawa M, Maltby D, Talley TT, Durkin KA, Medzihradsky KF, Burlingame AL, Taylor P, Casida JE (2008) Atypical nicotinic agonist bound conformations conferring subtype selectivity. *Proc Natl Acad Sci U S A* 105:1728–1732
- Tu C, Wang Y, Duan W, Hertl P, Tradway L, Brandenburg R, Lee D, Snell M, Hu S (2011) Effects of fungicides and insecticides on feeding behavior and community dynamics of earthworms: implications for casting control in turfgrass systems. *Appl Soil Ecol* 47:31–36
- Valdovinos-Núñez J, Quezada-Euán JG, Ancona-Xiu P, Moo-Valle H, Carmona A, Sanchez ER (2009) Comparative toxicity of pesticides to stingless bees (Hymenoptera: Apidae: Meliponini). *J Econ Entomol* 102:1737–1742
- Van den Brink PJ, Roelsma J, van Nes EH, Scheffer M, Brock TCM (2002) PERPEST model: a case-based reasoning approach to predict ecological risks of pesticides. *Environ Toxicol Chem* 21:2500–2506
- Van der Sluijs JP, Simon-Delso S, Goulson D, Maxim L, Bonmatin JM, Belzunces LP (2013) Neonicotinoids, bee disorders and the sustainability of pollinator services. *Curr Opin Environ Sustain* 5:293–305
- Van der Zee R, Pisa L, Andonov S, Brodschneider R, Charriere JD et al (2012) Managed honey bee colony losses in Canada, China, Europe, Israel and Turkey, for the winters of 2008–9 and 2009–10. *J Apic Res* 51:91–114. doi:10.3896/ibra.1.51.1.12
- Van Dijk TC, van Staalduinen MA, van der Sluijs JP (2013) Macro-invertebrate decline in surface water polluted with imidacloprid. *PLoS One* 8:e62374. doi:10.1371/journal.pone.0062374
- Van Engelsdorp D, Meixner MD (2010) A historical review of managed honey bee populations in Europe and the United States and the factors that may affect them. *J Invertebr Pathol* 103:80–95
- Vandame R, Meled M, Colin ME, Belzunces LP (1995) Alteration of the homing-flight in the honey bee *Apis mellifera* L. exposed to sublethal dose of deltamethrin. *Environ Toxicol Chem* 14:855–860
- Vidau C, Diogon M, Aufauvre J, Fontbonne R, Vignes B, Brunet JL, Texier C, Biron DG, Blot N, El Alaoui H, Belzunces LP, Delbac F (2011) Exposure to sublethal doses of fipronil and thiacloprid highly increases mortality of honeybees previously infected by *Nosema ceranae*. *PLoS One* 6:e21550. doi:10.1371/journal.pone.0021550
- Vijver MG, van den Brink PJ (2014) Macro-invertebrate decline in surface water polluted with imidacloprid: a rebuttal and some new analyses. *PLoS One* 9:e89837
- Villanueva-Jimenez JA, Hoy MA (1998) Toxicity of pesticides to the citrus leafminer and its parasitoid *Ageniaspis citricola* evaluated to assess their suitability for an IPM program in citrus nurseries. *Biocontrol* 43:357–388
- Volkov EM, Nurullin LF, Nikolsky E, Vyskocil F (2007) Miniature excitatory synaptic ion currents in the earthworm *Lumbricus terrestris* body wall muscles. *Physiol Res* 56:655–658
- Voudouris CC, Sauphanor B, Franck P, Reyes M, Mamuris Z et al (2011) Insecticide resistance status of the codling moth *Cydia pomonella* (Lepidoptera: Tortricidae) from Greece. *Pestic Biochem Physiol* 100:229–238
- Wang M, Grimm V (2010) Population models in pesticide risk assessment: lessons from assessing population-level effects, recovery, and alternative exposure scenarios from modeling a small mammal. *Environ Toxicol Chem* 29:1292–1300
- Wang AH, Wu JC, Yu YS, Liu JL, Yue JF et al (2005) Selective insecticide-induced stimulation on fecundity and biochemical changes in *Tryporyza incertulas* (Lepidoptera: Pyralidae). *J Econ Entomol* 98:1144–1149
- Wang Y, Cang T, Zhao X, Yu R, Chen L, Wu C, Wang Q (2012a) Comparative acute toxicity of twenty-four insecticides to earthworm, *Eisenia fetida*. *Ecotoxicol Environ Saf* 79:122–128
- Wang Y, Wu S, Chen L, Wu C, Yu R, Wang Q, Zhao X (2012b) Toxicity assessment of 45 pesticides to the epigeic earthworm *Eisenia fetida*. *Chemosphere* 88:484–491
- Ward GS (1990) NTN 33893 technical: acute toxicity to the mysid, *Mysidopsis bahia*, under flow-through test conditions. *Toxicol Environmental Sciences*, Jupiter, Florida (performing laboratory).

- Mobay Corporation, Kansas City, Missouri (submitting laboratory). Mobay Report No. 100355. 46 pp
- Ward GS 1991. NTN 33893 technical: chronic toxicity to the mysid *Mysidopsis bahia* under flow-through conditions. Toxicon Environmental Sciences, Jupiter, FL. 87 pp. Miles Report No. 101347
- Weibull AC, Bengtsson J, Nohlgren E (2000) Diversity of butterflies in the agricultural landscape: the role of farming system and landscape heterogeneity. *Ecography* 23:743–750
- Whitehorn PR, O'Connor S, Wackers FL, Goulson D (2012) Neonicotinoid pesticide reduces bumble bee colony growth and queen production. *Science* 336:351–352
- Wise JC, Jenkins PE, Vander Poppen R, Isaacs R (2010) Activity of broad-spectrum and reduced-risk insecticides on various life stages of cranberry fruitworm (Lepidoptera: Pyralidae) in highbush blueberry. *J Econ Entomol* 103:1720–1728
- Wu JY, Anelli CM, Sheppard WS (2011) Sub-lethal effects of pesticide residues in brood comb on worker honey bee (*Apis mellifera*) development and longevity. *PLoS One* 6:e14720
- Wu JY, Smart MD, Anelli CM, Sheppard WS (2012) Honey bees (*Apis mellifera*) reared in brood combs containing high levels of pesticide residues exhibit increased susceptibility to *Nosema* (Microsporidia) infection. *J Invertebr Pathol* 109:326–329
- Yamada T, Yamada K, Wada N (2012) Influence of dinotefuran and clothianidin on a bee colony. *Japan J Clin Ecol* 21:10–23
- Yang EC, Chuang YC, Chen YL, Chang LH (2008) Abnormal foraging behavior induced by sublethal dosage of imidacloprid in the honey bee (Hymenoptera: Apidae). *J Econ Entomol* 101:1743–1748
- Yang EC, Chang HC, Wu WY, Chen YW (2012) Impaired olfactory associative behavior of honeybee workers due to contamination of imidacloprid in the larval stage. *PLoS One* 7:e49472
- Youn YN, Seo MJ, Shin JG, Jang C, Yu YM (2003) Toxicity of greenhouse pesticides to multicolored Asian lady beetles, *Harmonia axyridis* (Coleoptera: Coccinellidae). *Biol Control* 28:164–170
- Yu YS, Xue S, Wu JC, Wang F, Liu JL et al (2007a) Distribution of imidacloprid residues in different parts of rice plants and its effect on larvae and adult females of *Chilo suppressalis* (Lepidoptera: Pyralidae). *J Econ Entomol* 100:375–380
- Yu YS, Xue S, Wu JC, Wang F, Yang GQ (2007b) Changes in levels of juvenile hormone and molting hormone in larvae and adult females of *Chilo suppressalis* (Lepidoptera: Pyralidae) after imidacloprid applications to rice. *J Econ Entomol* 100:1188–1193
- Yue B, Wilde GE, Arthur F (2003) Evaluation of thiamethoxam and imidacloprid as seed treatments to control European corn borer and Indianmeal moth (Lepidoptera: Pyralidae) larvae. *J Econ Entomol* 96:503–509
- Zang Y, Zhong Y, Luo Y, Kong ZM (2000) Genotoxicity of two novel pesticides for the earthworm, *Eisenia fetida*. *Environ Pollut* 108:271–278
- Zeng CX, Wang JJ (2010) Influence of exposure to imidacloprid on survivorship, reproduction and vitellin content of the carmine spider mite, *Tetranychus cinnabarinus*. *J Insect Sci* 10:20
- Zhou LJ, Huang JG, Xu HH (2011) Monitoring resistance of field populations of diamondback moth *Plutella xylostella* L. (Lepidoptera: Yponomeutidae) to five insecticides in South China: a ten-year case study. *Crop Prot* 30:272–278