

Corrigendum: Global climate targets and future consumption level: an evaluation of the required GHG intensity

This content has been downloaded from IOPscience. Please scroll down to see the full text.

2013 Environ. Res. Lett. 8 029501

(<http://iopscience.iop.org/1748-9326/8/2/029501>)

View [the table of contents for this issue](#), or go to the [journal homepage](#) for more

Download details:

IP Address: 131.211.104.172

This content was downloaded on 31/03/2015 at 13:54

Please note that [terms and conditions apply](#).

Corrigendum: Global climate targets and future consumption level: an evaluation of the required GHG intensity

2013 *Environ. Res. Lett.* **8** 014016

Bastien Girod^{1,2}, **Detlef Peter van Vuuren**^{2,3} and **Edgar G Hertwich**⁴

¹ Swiss Federal Institute of Technology Zurich (ETH Zurich), Department of Management, Technology, and Economics, Chair of Sustainability and Technology, Weinbergstrasse 56/58, 8032 Zurich, Switzerland

² Copernicus Institute of Sustainable Development, Department of Geosciences, Utrecht University, Heidelberglaan 2, 3584 CS Utrecht, The Netherlands

³ PBL—Netherlands Environment Assessment Agency, PO Box 303, 3720 BA Bilthoven, The Netherlands

⁴ Industrial Ecology Program and Department of Energy and Process Engineering, Norwegian University of Science and Technology, NO-7491 Trondheim, Norway

E-mail: bgirod@ethz.ch

Received 20 March 2013

Accepted for publication 22 March 2013

Published 4 April 2013

Online at stacks.iop.org/ERL/8/029501

There is an error in figure 5: for ‘food’ the GHG intensity should be gCO₂/Mcal (instead of gCO₂/kcal). The corrected figure and caption are reproduced below.



Content from this work may be used under the terms of the [Creative Commons Attribution 3.0 licence](http://creativecommons.org/licenses/by/3.0/). Any further distribution of this work must maintain attribution to the author(s) and the title of the work, journal citation and DOI.




Figure 5. Baseline and target GHG intensity for the five main consumption categories. ‘Global’ refers to the global average. Projections for 2050 are shown for the five main world regions. Target GHG intensities in 2050 for the RCP2.6 are shown for different allocation schemes: same cost of reduction, same reduction from the distribution in 2000 and the same reduction in GHG intensity (2050). The different world regions show the target GHG intensity for same reduction and equal per capita allowance. The GHG intensity for the RCP4.5 to RCP8 uses also the same reduction allowance scheme.