

GHGT-11

## CO<sub>2</sub> Storage Capacity of Campos Basin's Oil Fields, Brazil.

Gabriela Camboim ROCKETT<sup>1\*</sup>, João Marcelo Medina KETZER<sup>1</sup>,  
Andrea RAMÍREZ<sup>2</sup>, Machteld van den BROEK<sup>2</sup>

<sup>1</sup> Center of Excellence in Research and Innovation in Petroleum, Mineral Resources and Carbon Storage/Pontifical Catholic University of Rio Grande do Sul (CEPAC/PUCRS). Av. Ipiranga, 6681, Prédio 96J - CEP 90619-900 - Porto Alegre - RS – Brasil

<sup>2</sup> Copernicus Institute of Sustainable Development/Utrecht University. Budapestlaan 6 3584CD Utrecht, The Netherlands.

### Abstract

Large potentials for CO<sub>2</sub> storage were demonstrated in previous studies in Brazil. This study aims to estimate the CO<sub>2</sub> storage capacity in the Campos Basin's oil fields, Southeast Brazil, in order to provide refined values to support CCS planning in the country. The results, based on field/reservoir level data show that there is a large potential for CO<sub>2</sub> storage (ca. 950Mt) in the 17 assessed oil fields in the basin, and 75% of this storage capacity is found in sandstone reservoirs.

© 2013 The Authors. Published by Elsevier Ltd.  
Selection and/or peer-review under responsibility of GHGT

Key Words: CO<sub>2</sub> storage capacity, Campos Basin, Oil fields, Geological storage, Enhanced oil recovery, CCS, CO<sub>2</sub>-EOR

### 1. Introduction

A critical point for the development of carbon dioxide capture and storage (CCS) is the availability of CO<sub>2</sub> storage. A thorough understanding of the available CO<sub>2</sub> storage capacity is required for the implementation of CCS. According to Bachu et al [1], CO<sub>2</sub> storage capacity assessment may be conducted at various scales: country, basin, regional, local and site-specific. Even in different assessment-scales, Bradshaw et al [2] state that the estimation of the capacity of a geological reservoir to store CO<sub>2</sub> is not a straightforward or simple process, and lack of data is still one of the major challenges in CO<sub>2</sub> storage capacity estimations.

There are three different ways to assess storage capacity [1] [3], one for each storage option (coalbeds, aquifers, oil and gas fields), based on specific inherent characteristics of each kind of reservoir. Regarding

---

\* Corresponding author. Tel.: +055-51-30286138.  
E-mail address: [gabriela.rockett@gmail.com](mailto:gabriela.rockett@gmail.com)

CO<sub>2</sub> storage capacity estimation in depleted oil and gas reservoirs, it is straightforward and is based on recoverable reserves, reservoir properties and in situ CO<sub>2</sub> characteristics [1]. The fundamental assumption is that the volume previously occupied by the produced hydrocarbons is available for CO<sub>2</sub> storage. In the case of CO<sub>2</sub>-EOR, the CO<sub>2</sub> storage capacity can be more accurately through numerical simulations [1].

### *1.1. CO<sub>2</sub> storage capacity estimation in Brazil*

In the Brazilian National Plan for Climate Change [4], Carbon Capture and Storage has been recognized as an important option to reduce CO<sub>2</sub> emissions, especially for the oil and gas, and industrial sector. The Brazilian oil and gas industry is engaged in CCS research and development, and CO<sub>2</sub> is already being injected for enhanced oil recovery in the country [5]. Preliminary studies on CO<sub>2</sub> storage potential and source-sink matching in Brazilian sedimentary basins (The CARBMAP Project) have been conducted ([6], [7], [8], [9], [10]), and the findings identify some promising basins for CCS development in Brazil.

A previous study [6] estimated the Brazilian potential storage capacity (in saline aquifers, petroleum fields and coalbeds) and opportunities for CCS in the country. In total, the basin scale estimation indicates that Brazil has a large theoretical storage capacity (ca. 2,000 GtCO<sub>2</sub>). Regarding CO<sub>2</sub> storage in petroleum fields, Campos and Santos Basins appear as good candidates, with theoretical capacities – mostly based on generic basin-scale data (total oil and gas reserves) – of ca. 1,800 and 167 MtCO<sub>2</sub> respectively [6] [7].

In this context, the aim of this study was to estimate CO<sub>2</sub> storage capacity of petroleum fields in the Campos Basin, in order to provide refined capacity values based on field/reservoir level data.

## **2. Methodology**

### *2.1. Study Area: Campos Basin*

The Campos Basin is an offshore basin located in southeastern Brazil, between the geographic parallels 21° and 23° South (Fig.1). The basin covers an area of about 100,000 km<sup>2</sup> [11] [12], and contains 80% of the Brazilian petroleum reserves [11]. The discovered petroleum fields are in water depths ranging from 80 to more than 2,600 meters.

According to PETROBRAS [11], there are 36 petroleum fields in Campos Basin which have already reached peak production and are or will be in the coming years in the mature phase. Because of this and its proximity to large scale stationary CO<sub>2</sub> emission sources, the Campos Basin is considered one of the most promising Brazilian basins for CO<sub>2</sub> storage development [6] [7].




Fig.1: Campos Basin's location

### 2.1.1. Geological setting

The Campos Basin was formed as a result of the breakup of Gondwana paleocontinent, in the Early Cretaceous (~140 Ma), followed by subsequent infilling of the rift basin with as much as 9,000 m of Early Cretaceous–Holocene sediments [12]. According to Moraes et al [13], the sedimentary package that forms the Campos Basin is divided into five megasequences (Fig.2): continental rift (Neocomian); transitional (evaporitic) (Aptian); shallow marine (Albian); marine transgressive (Cenomanian to Upper Eocene); marine regressive (Upper Eocene to Present) megasequences. The Palaeocene–Eocene reservoirs occur mostly within the transgressive megasequence.

Guardado et al [12] described the most important Campos Basin megasequences, as follows: The nonmarine rift megasequence is composed of lacustrine Barremian sedimentary rocks overlying Neocomian basalts. Rift tectonics had influenced the strata deposition paleoenvironments. This sequence contains the calcareous shales of the Lagoa Feia Formation, the most important source rock in the basin. The transitional megasequence was deposited in Aptian time during a non-tectonic period. It represents the beginning of the drift phase and contains a lower sequence (mostly composed of conglomerates and carbonates), and an upper sequence (consisting of evaporites). The marine megasequence is made up of Albian shallow-water carbonates, mudstones, and marls (Macaé Formation) at its base. This basal

sequence grades upward into an Upper Cretaceous–Paleocene bathyal sequence consisting of shales, marls, and sandstone turbidites. A progradational siliciclastic sequence characterizes the remaining Neogene section.

Reservoir rocks with good to excellent permeability and porosity are widespread in the Campos Basin both in time and space [12].




Fig.2: Generalized geological section for the Eastern Brazilian continental margin basins, showing the main megasequences: PR, pre-rift (which does not occur in the Campos Basin); R, rift; T, transitional (which includes the evaporate section); SC, shallow carbonate; MT, marine transgressive; MR, marine regressive (modified from Bruhn [14]).

## 2.2. Geological Reservoir Data

Campos Basin oil fields were selected for this study, because of their well-known structure, proven traps and availability of reservoir data. Specific oil/gas field data were obtained through extensive literature survey. Public available data are very limited, and because of this, the total number of fields analyzed in this study is 17 (from more than 50 oil fields in the basin). The fields included are: Carapeba, Linguado, Marimbá, Marlim, Vermelho, Barracuda, Roncador, Caratinga, Jubarte, Namorado, Badejo, Pampo, Enchova/Bonito, Garoupa, Albacora, Corvina and Malhado. These fields contain about 36% of the Campos Basin's oil reserves (excluding new pre-salt reserves; based on data from ANP [15], Rangel et al [16], Assis et al [17] and Horschutz et al [18]). The assessed reservoirs in these fields are mainly Cretaceous and Tertiary sandstones, and some Cretaceous limestones. For storage capacity estimation purposes, the following data was collected:

- **Reservoir:** For each oil/gas field, data on the type of rock, age and Formation were collected. The reservoirs considered in this study are listed in Table 1.

Table 1. Reservoirs considered in this study, per Oil Field

Oil Field	Reservoir type	Age/Stratigraphic Unit
Carapeba	Turbidite sandstones	Eocene/Carapebus Mb.
	-	Cretaceous/Carapebus Mb.
	Calcirudites	Barremian/Lagoa Feia Fm.
Linguado	Calcarenites and calcirudites	Albian/ Macaé Fm.
	Turbidite sandstones	Maastrichian/Campos Fm.
Marimbá	-	Cretaceous/Campos
Marlim	Turbidite sandstones	Oligocene/Carapebus Mb.
Vermelho		Oligocene
	Medium-grained sandstones	Eocene/Carapebus Mb.
Barracuda	-	Oligocene/Campos
	Turbidite sandstones	Eocene/Campos
Roncador	Sandstones	Maastrichian/Carapebus Fm.
Caratinga	Turbidites	Oligocene
Jubarte	Sandstones	Maastrichian
Namorado	Sandstones (Namorado sandstone)	Cenomanian-Turonian/Macaé Fm.
Badejo	Coquines	Lagoa Feia Fm.
Pampo	Neritic carbonates	Albo-Cenomanian/Macaé Fm.
	Coquines (calcirudites/carcarenites)	Lagoa Feia Fm.
Cherne	Turbidite sandstones	Cretaceous
Enchova/Bonito	Turbidites	Eocene/Carapebus Mb.
Albacora	Turbidite sandstones	Cretaceous-Tertiary
Garoupa	Carbonates	Albian/Macaé Fm.
Albacora Leste	Sandstones	Miocene/Carapebus Fm.
Corvina	Turbidite sandstones	Eocene/Carapebus Mb.
Malhado	Turbidite sandstones	Eocene/Carapebus Mb.

- **Depth:** Reservoir depth data was not always available at the field level, and therefore it is estimated (for each field) by one of the following four indicators: (i) average depth of specific reservoirs in an oil field (ex. Carapeba, Linguado, Vermelho, Namorado, Pampo); (ii) average depth of the reservoir layers in the field (ex. Marlim, Roncador, Caratinga, Jubarte, Badejo, Albacora, Malhado); (iii) average field depth extracted from geological sections (iv) oil-water contact depth extracted from geological sections and articles (ex. Enchova/Bonito, Corvina). All the 17 reservoirs are over 800 meters deep, which is required for CO<sub>2</sub> geological storage in supercritical conditions [19].
- **Original Oil in Place (OOIP):** Recent data on oil reserves are only publicly available at basin level or per federal state and not for specific fields. However, older data on the original oil in place were available in scientific literature dating from 1992 to 2006.

The figures available in the literature and used in this study are from the references in Appendix A.

### 2.3. CO<sub>2</sub> Storage Capacity Estimation

CO<sub>2</sub> storage capacities were estimated based on the methodology proposed by Bachu et al [1], which assumes that the volume previously occupied by the produced hydrocarbons is available for CO<sub>2</sub> storage (Eq. 1).

$$MCO_2 = \rho_{CO_2} r [(Rf \times OOIP/Bf) - V_{iw} + V_{pw}] \quad (\text{Eq.1})$$

Where MCO<sub>2</sub> stands for the theoretical CO<sub>2</sub> storage capacity (in tonnes),  $\rho_{CO_2} r$  is the CO<sub>2</sub> density under reservoir conditions (in kg/m<sup>3</sup>), Rf the recovery factor, OOIP the original oil in place (in m<sup>3</sup>), Bf the formation volume factor, V<sub>iw</sub> the injected water volume and V<sub>pw</sub> the produced water volume.

Note that the analysis in this study is at the field level and some assumptions had to be made when data were only available from specific reservoirs within an oil/gas field.

- **CO<sub>2</sub> Density at Reservoir Condition:** Average CO<sub>2</sub> density of each oil field was determined based on temperature and pressure data. When data were not available, the average temperature was calculated based on its depth and the Campos Basin's average geothermal gradient data (from Jahnert [20]). Furthermore, 4°C was added to each field's average temperature to take into account the sea floor temperature [21]. The average pressure was estimated based on the oil field's depth, assuming hydrostatic pressure of 100 bar per kilometer.
- **Recovery Factor and Original Oil in Place:** The recovery factor is defined as the ratio of the planned oil production and the original oil in place. Since this data is not publicly available, in this study, oil reserve values were used instead. For the oil fields for which also oil reserves data were unavailable (i.e., Badejo, Enchova/Bonito, Albacora, Corvina and Malhado), the original oil in place data were multiplied by a recovery factor of 35.4% (the average recovery factor in Campos Basin with the new technologies application), according to Bosco [22].
- **Formation Volume Factor, Injected Water Volume and Produced Water Volume:** The formation volume factor, which converts the oil volume in normal conditions to the reservoir condition, was assumed to be 1.2 [23] and the sum of injected water volume and produced water volume was assumed to be zero.

### 3. Results and Discussions

The results show that Campos Basin has a storage capacity of approximately 950 MtCO<sub>2</sub>, in the 17 analyzed oil fields. The Roncador field appears as the one with the large storage capacity (~265 MtCO<sub>2</sub>), followed by Marlim, Albacora and Barracuda oil fields with storage capacities of 174, 163, and 118 MtCO<sub>2</sub>, respectively. These 4 oil fields bear more than 75% of the Campos Basin's potential CO<sub>2</sub> storage capacity as it is shown in Fig.3. Some of the analyzed oil fields have limited CO<sub>2</sub> storage capacity (less than 6 MtCO<sub>2</sub>), such as Linguado, Garoupa, Malhado and Badejo.

Figure 4 shows the CO<sub>2</sub> storage capacities contribution to the different assessed reservoirs in the oil fields. Sandstones reservoirs in Roncador field (Carapebus Fm.) contribute with about 28% of Campos Basin's storage capacity, while the siliciclastic reservoirs in Marlim field (Carapebus Mb./Campos Fm.) match about 18%. The Albacora field represents 17% of Campos Basin CO<sub>2</sub> storage capacity, Cretaceous and Tertiary reservoirs are turbidite sandstones from Macaé and Campos Formations, respectively. The Barracuda field's capacity is 50% in Eocene reservoirs (turbidite sandstones) and 50% in Oligocene reservoirs. 100% of Jubarte and Namorado fields storage capacity corresponds to Crataceous sandstones.





Fig.3: CO<sub>2</sub> storage capacity in Campos Basin oil fields

Fig.4: CO<sub>2</sub> storage capacity in Campos Basin oil fields

In Pampo field's storage capacity corresponds to limestone reservoirs (84% in Macaé Formation carbonates and 16% in Lagoa Feia Formation coquinas).

Estimated storage capacities of Enchova/Bonito, Carapeba, Marimbá, Vermelho and Corvina oil fields are in sandstones reservoirs. In Carapeba field, turbidite sandstones from Carapebus Fm. were assessed. Sandstones from Campos Fm. were assessed in Marimbá Field. Linguado's storage capacity is divided into limestone reservoirs and sandstone reservoirs (96% and 4%, respectively). Garoupa's storage capacity corresponds to carbonate reservoirs and Badejo's capacity corresponds to coquina reservoirs. Malhado oil field's storage capacity corresponds to turbidite sandstones reservoirs.

#### 4. Conclusion

In this research, Carbon dioxide storage capacity estimation based on field/reservoir level data was carried out. The findings of this research indicate a CO<sub>2</sub> storage capacity of about 950 Mt in the 17 assessed oil fields in Campos Basin. More than 75% of the storage capacity is in sandstone reservoirs of Roncador, Marlim, Albacora and Barracuda oil fields. These reservoirs can play a significant role for CCS implementation in Brazil due to the large CO<sub>2</sub> storage capacities and because most of them are already mature fields or will be mature in the near future. It is important to note that Campos Basin's storage capacity is potentially larger than the estimated value in this research, since the theoretical storage capacity estimated in this study corresponds to data from only 17 out of 50 oil fields (at least) in Campos Basin, and most data are not updated.

Enhanced Oil Recovery combined with CCS (CO<sub>2</sub>-EOR) is a promising opportunity in Campos Basin, considering that it is a mature basin with a significant amount of data and infrastructure. Further, CO<sub>2</sub> storage capacity, in this case, might be larger due to the oil produced by EOR.

Currently, planned investment in CO<sub>2</sub> storage are focusing in the pre-salt reservoirs (discovered in Santos, Espírito Santo and Campos Basins), considering that in some fields CO<sub>2</sub> content might be >2% vol. and the injection of this amount can be an option to produce residual oil from these fields.

#### Acknowledgements

The authors thank CEPAC/PUCRS, PETROBRAS and the Copernicus Institute/Utrecht University for supporting this research and for research grants. J.M. Ketzer thanks the Brazilian Research Council (CNPq) for research grant.

#### References

- [1] Bachu, S. et al., 2007. CO<sub>2</sub> storage capacity estimation: methodology and gaps. *International Journal of Greenhouse Gas Control*, 1 (4), 430-443.
- [2] Bradshaw, J., Bachu, S., Bonijoly, D., Burruss, R., Holloway, S., Christensen, N.P., Mathiassen, O.M. 2007. CO<sub>2</sub> storage capacity estimation: issues and development of standards. *Int. J. Greenhouse Gas Control* 1, 62–68.
- [3] CSLF (Carbon Sequestration Leadership Forum), 2008: Comparison between Methodologies Recommended for Estimation of CO<sub>2</sub> Storage Capacity in Geological Media – Phase III Report; CSLF-T-2008-04, 21p.
- [4] Brazil, 2008. PNMC – National Plan on Climate Change, 2008. Interministerial Committee on Climate Change. (in Portuguese)
- [5] Lino, U.R.A. Case History of Breaking a Paradigm: Improvement of an Immiscible Gas-Injection Project in Buracica Field by Water Injection at the Gas/Oil Contact. *SPE Latin American and Caribbean Petroleum Engineering Conference*. Rio de



- Janeiro, Brasil, 20-23 de Junho de 2005.
- [6] Ketzer, J.M.M. et al. 2007a. Opportunities for CO<sub>2</sub> capture and geological storage in Brazil: The CARBMAP Project. In: *Annual Conference on Carbon Capture & Sequestration*, 6, Pittsburgh, Pennsylvania. Resource Book. Pittsburgh.
  - [7] Ketzer, J.M.M. et al. The CARBMAP Project: CO<sub>2</sub> Source-Sink Matching and Opportunities for Carbon Sequestration in Brazil. In: *Carbon Sequestration Leadership Forum Annual Meeting*, Paris, 27 mar. 2007b. Poster.
  - [8] Machado, C.X. et al. 2009. Development of a Geographic Information System to match CO<sub>2</sub> stationary sources and geological sinks in Brazil. In: Ziliotto, M.A. (Org.). *Climate Change, Carbon Sequestration and Market in Brazil*. 1 ed. Paraná: Instituto Ecolima/UFPR, pp. 267-279. (in Portuguese)
  - [9] Rockett, G.C. Matching CO<sub>2</sub> sources and potential sinks for geological storage in Campos Basin, Brazil. Porto Alegre. 2010. Master Thesis. Pos-Graduation Program in Materials Engineering and Technology, Pontifical Catholic University of Rio Grande do Sul. 205p. (in Portuguese)
  - [10] Rockett, G.C. et al. 2011. The CARBMAP Project: Matching CO<sub>2</sub> Sources and Geological Sinks in Brazil using Geographic Information System. *Energy Procedia*, v. 4, 2764-2771. DOI: 10.1016/j.egypro.2011.02.179
  - [11] Petrobras, 2010. Petróleo Brasileiro S/A. Campos Basin - The largest oil reserves in Brazil. Available at: <[http://www2.petrobras.com.br/Petrobras/portugues/plataforma/pla\\_bacia\\_campos.htm](http://www2.petrobras.com.br/Petrobras/portugues/plataforma/pla_bacia_campos.htm)>. Accessed on Feb. 2010. (In Portuguese)
  - [12] Guardado, L. R.; Spadini, A. R.; Brandão, J. S. L.; Mello, M. R. Petroleum system of the Campos Basin, Brazil. In: MELLO M. R.; KATZ, B. J. (eds.). *Petroleum systems of South Atlantic margins: AAPG Memoir 73*, 2000, p. 317–324.
  - [13] Moraes, M.A. S.; Maciel, W.B.; Braga, M.S. S.; Viana, A.R. Bottom-current reworked Palaeocene-Eocene deep-water reservoirs of the Campos Basin, Brazil. *Geological Society*, London, Special Publications, 276, p.81 – 94, 2007.
  - [14] Bruhn, C. H. L. Deep-water reservoirs from the eastern Brazilian rift and passive-margin basins. In: *AAPG International Conference and Exhibition*, Rio de Janeiro, 8-11 nov., Course No. 6, Part 2, 1998.
  - [15] ANP, 2011a. Brazilian petroleum, natural gas and biofuels statistic yearbook. National Agency of Petroleum, Natural Gas and Biofuels, Brazil. (in Portuguese).
  - [16] Rangel, H.D., Guimarães, P.T.M.; Spadini, A.R. 2003. Barracuda and Roncador giant oil fields; deep-water Campos Basin, Brazil. *AAPG Memoir*, 78, 123-137.
  - [17] Assis O.C., Becker M.R., Melo J.R.C., Franz E.P., Alves R.R.P., Rodriguez M.R., Maciel W.B., Souza Junior O.G., Johann P.R.S., 1998. Barracuda and Caratinga giant oil fields, deep- water Campos Basin, Brazil. *Proceedings of the Annual Offshore Technology Conference*, Houston, 4-7 May. p.611-617.
  - [18] Horschutz, P.M.C.; Freitas, L.C.S.; Stank, C.V.; Barroso, A.S.; Cruz, W.M., 1992. The Linguado, Carapeba, Vermelho and Marimbá Giant Fields, Campos Basin, Offshore Brazil. In: Halbouty, M.T. (Ed.) *Giant oil and gas fields of the decade, 1978-1988*. Tulsa: American Association of Petroleum Geologists, AAPG Memoir n. 54, p.137-153.
  - [19] IPCC, 2005. IPCC Special Report on Carbon Dioxide Capture and Storage. Prepared by Working Group III of the Intergovernmental Panel on Climate Change, Cambridge University Press, Cambridge, UK and New York, NY, USA.
  - [20] Jahnert, R.J. 1987. Campos Basin's geothermal gradient. *Boletim de Geociências da PETROBRAS*, 1 (2): p.183-189. (in Portuguese)
  - [21] Miller, D.J.; Kowsmann, R.O. 2009. Occurrence of Authigenic Carbonate Chimneys and Crusts in the Campos Basin Continental Slope - SE Brazil. In: *AAPG International Conference and Exhibition*, 15-18 November, Rio de Janeiro, Brazil. Poster presentation.
  - [22] Bosco, F. 2007. More power to your business. *Petro & Química*, Ed. 299. Available at: <[http://www.petroequimica.com.br/edicoes/ed\\_299/299.html](http://www.petroequimica.com.br/edicoes/ed_299/299.html)>. Acesso em: 13 abr. 2010. (In Portuguese)
  - [23] APEC (Asia Pacific Economic Cooperation), 2005. Assessment of Geological Storage Potential of Carbon Dioxide in the APEC Region – Phase I: CO<sub>2</sub> Storage Prospectivity of Selected Sedimentary Basins in the Region of China and South East Asia. APEC Energy Working Group EWG Project 06/2003. Canberra, Australia: Innovative Carbon Technologies Pty Ltd and Geoscience Australia, Jun. 2005.

## Appendix A. Available figures on Campos Basin's reservoirs/oil fields - References used in this study

Figures collected from literature and used in this study (available/published data on depth, original oil in place, oil reserves and reservoir temperature) are from the following references:

- Assis O.C., Becker M.R., Melo J.R.C., Franz E.P., Alves R.R.P., Rodriguez M.R., Maciel W.B., Souza Junior O.G., Johann P.R.S. 1998. Barracuda and Caratinga giant oil fields, deep- water Campos Basin, Brazil. In: 30<sup>th</sup> *Annual Offshore Technology Conference*, OTC 8879, Houston, Texas, 4-7 May, Proceedings..., p.611-617.
- Bacocoli, G.; Morales, R.G.; Campos, O.A.J. 1980. The Namorado oil field: a major oil discovery in the Campos Basin, Brazil. *AAPG Memoir*. 30, Pages: 329-338.
- Barboza, E.G. 2005. Análise estratigráfica do Campo de Namorado (Bacia de Campos) com base na interpretação sísmica tridimensional. 215p. Universidade Federal do Rio Grande do Sul, Programa de Pós-Graduação em Geociências (Doutorate Thesis). Porto Alegre.
- Cândido, A. 1990. Desenvolvimento e estratégia de produção do campo de Albacora. *Boletim de Geociências da PETROBRAS*, Rio de Janeiro, 4 (2): p. 175-181, abr/jun.
- Condessa, L.G., Machado, M.A.P. 1992. Campo de Marlim: analise de colapso em reservatorios arenosos por meio de métodos dinâmicos. (Translated Title: Marlim Field: collapse analysis of sandstone reservoirs using dynamic methods). *Boletim de Geociências da PETROBRAS*, 6; 3-4, Pages: 163-169.
- De Souza, J.M., Scarton, J.C., Cândido, A., Corá, C.A.G. 1990. The Marlim and Albacora Giant Fields, Campos Basin, Offshore Brazil: Abstract. *AAPG Bulletin*, vol. 74, n.9.
- Fontanelli, P.R.; De Ros, L.F., Remus, M.V.D. 2009. Provenance of deep-water reservoir sandstones from the Jubarte oil field, Campos Basin, Eastern Brazilian Margin. *Marine and Petroleum Geology*, Volume 26, Issue 7, August 2009, p.1274-1298, ISSN 0264-8172, 10.1016/j.marpetgeo.2008.09.010.
- Guardado, L.R.; Gamboa, L.A.P.; Lucchesi, C.F. 1989. Petroleum geology of the Campos Basin, Brazil, a model for a producing atlantic type basin. In: EDWARDS, J.D. & SANTOGROSSI, P.A. (Eds.) *Divergent/Passive Margin Basins*. Tulsa: *AAPG Memoir* 48, p.3-79.
- Guimaraes, P.T.; Rangel, M. H.D.; Barroso, A.S.; Mihaguti, M.K.; Stank, C.V.; Castro, D.D.; Santos, M.B.; Alves, R.R.P.; Lemos, W.P. 2001. The Barracuda and Roncador Giant Fields, Deep Water Campos Basin, Brazil. *AAPG Annual Meeting*, Denver, Colorado June 3-6.
- Horschutz, P.M.C.; Freitas, L.C.S.; Stank, C.V.; Barroso, A.S.; Cruz, W.M. 1992. The Linguado, Carapeba, Vermelho and Marimbá Giant Fields, Campos Basin, Offshore Brazil. In: HALBOUTY, M.T. (Ed.) *Giant oil and gas fields of the decade, 1978-1988*. Tulsa: American Association of Petroleum Geologists, *AAPG Memoir* n. 54, p. 137-153.
- Horschutz, P.M.C., Scuta, M. 1992. Fácies-perfis e mapeamento de qualidade do reservatório de coquinas da Formação Lagoa feia do campo de Pampo. *Boletim de Geociências da PETROBRAS*, 6; 1-2, Pages: 45-58.
- Lemos, W.P., Baião de Castro, M.R., Soares, C.M., Rosalba, J.F., Meira, A.A.G. 2006. Albacora Leste Field Development: Reservoir Aspects and Development Strategy. *Offshore Technology Conference*, Houston, Texas, U.S.A., 1-4 May 2006, Proceedings, OTC 18056.
- Mendonça, P. M. M., Milani, E. J., Spadini, A. R. 2004. Oil exploration in Brazil: the 50 years of Petrobras. In: *23rd IAS Meeting of Sedimentology*, Coimbra, 15-17 set. 2004, Abstracts Book, p.27-30.
- Milani, E.J., Araújo, L.M. 2003. Recursos Minerais energéticos: petróleo. In: Bizzi, L.A., Schobbenhaus, C., Vidotti, R.M., Gonçalves, J.H. (orgs.). *Geologia, tectônica e recursos minerais do Brasil: textos, mapas e SIG*. Brasília: CPRM - Serviço Geológico do Brasil, p. 565-568.
- Rangel, H.D., Guimarães, P.T.M.; Spadini, A.R. 2003. Barracuda and Roncador giant oil fields; deep-water Campos Basin, Brazil. *AAPG Memoir*, 78, 123-137.
- Stank, C.V., Rober, F. 1990. Linguado, Carapeba, Vermelho and Marimba giant oil fields, Campos Basin, Offshore Brazil. *AAPG Bulletin*, v.74.
- Varela, C.L., Estever, F.R. 1990. Mapeamento de amplitude na delimitação dos campos de Corvina e Malhado. *Boletim de Geociências da PETROBRAS*, 4; 3, p.263-274.