

# Čech-De Rham theory for leaf spaces of foliations\*

Marius Crainic and Ieke Moerdijk

## Introduction

This paper is concerned with characteristic classes in the cohomology of leaf spaces of foliations. For a manifold  $M$  equipped with a foliation  $\mathcal{F}$  it is well-known that the coarse (naive) leaf space  $M/\mathcal{F}$ , obtained from  $M$  by identifying each leaf to a point, contains very little information. In the literature, various models for a finer leaf space  $M/\mathcal{F}$  are used for defining its cohomology. For example, one considers the cohomology of the classifying space of the foliation [2, 13, 17, 22], the sheaf cohomology of its holonomy groupoid [10, 18, 26], or the cyclic cohomology of its convolution algebra [7, 8]. Each of these methods has considerable drawbacks. E.g. they all involve non-Hausdorff spaces in an essential way. More specifically, the classifying space, which is probably the most common model for the “fine” leaf space, is a space which in general is infinite dimensional and non-Hausdorff, it is not a CW-complex, and it has lost all the smooth structure of the original foliation. In particular, it is not suitable for constructing cohomology theories with compact support. For this reason, the construction of characteristic classes in the cohomology of the classifying space of the foliation proceeds in a very indirect way, and many of the standard geometrical constructions have to be replaced by or supplied with abstract non-trivial arguments. The same applies to the construction of “universal” characteristic classes in the cohomology of the classifying space of the Haefliger groupoid,  $^q$ . It is possible to construct interesting classes of (foliated or transversal) bundles over foliations by explicit geometrical methods [2, 20], but these classes are constructed in the cohomology of the manifold  $M$  rather than that of the leaf space  $M/\mathcal{F}$ .

The purpose of this paper is to present a “Čech-De Rham” model for the cohomology of leaf spaces (Section 2), which circumvents the problems mentioned above. This Čech-De Rham model lends itself to the construction of (known) characteristic classes, now by explicit geometrical constructions which are immediate extensions of the standard constructions for manifolds (Section 3). As a consequence, for any transversal principal bundle over a foliated manifold  $(M, \mathcal{F})$ , we are able to lift the characteristic classes constructed in  $H^*(M)$  by the methods of [21], to the Čech-De Rham cohomology  $H^*(M/\mathcal{F})$ , and establish all the relations, such as the Bott vanishing theorem, at the level of  $H^*(M/\mathcal{F})$  (see Theorem 2 below).

We want to emphasize that the construction of the Čech-De Rham model and of the characteristic classes makes no reference to (holonomy) groupoids or classifying

---

\*Research supported by NWO

spaces. In particular, there are no non-Hausdorffness problems, and these constructions can be understood by anyone having some background in differential geometry, including familiarity with the very basic definitions concerning foliations.

To prove that our Čech-De Rham model gives in fact the same cohomology as the other models (Theorem 1), we use étale groupoids (Section 4). In fact, our model and the associated method for constructing characteristic classes applies to any étale groupoid, not just to holonomy groupoids (see Theorem 3, and 4.6). In particular, when used in the context of the Haefliger groupoid  $\mathcal{H}^q$ , it provides an explicit geometric construction of the universal geometrical characteristic classes (as a map from Gelfand-Fuchs cohomology into the cohomology of  $B, \mathcal{H}^q$  [2]). In this way we rediscover (and explain) the Thurston formula and the Bott formulas for cocycles on diffeomorphism groups [3] (for these explicit formulas, see Section 5). Other groupoids of interest, different from holonomy groupoids, are the monodromy groupoids of foliations. Our methods also show that the characteristic classes of foliated bundles [21] actually live in the cohomology of the monodromy groupoid of the foliation, rather in the cohomology of  $M$  itself.

Our Čech-De Rham cohomology also has a natural version with compact supports, which is related to the one with arbitrary supports by an obvious duality. When passing to the cohomology of holonomy groupoids, this duality becomes the Poincaré duality of [10] (Proposition 3). This new proof of Poincaré duality for leaf spaces appears as a straightforward extension of the standard arguments [4] from manifolds to leaf spaces. Moreover, this duality extends the known one for basic cohomology of Riemannian foliations [30].

There are several other cohomology theories associated to foliations which are easier to describe and are perhaps more familiar, such as basic cohomology (see e.g. [16, 30]) and foliated cohomology (see e.g. [1, 19, 20, 29]). In the last two sections of our paper, we use our Čech-De Rham model to explicitly describe the relations between the cohomology of leaf spaces and the basic and foliated cohomology.

## 1 Transverse structures on foliations

In this section we recall some basic notions concerning the transverse structures of foliations, which formalize the idea of structures over the leaf space. Throughout, we will work in the smooth context.

**1.1 Holonomy** Let  $M$  be a manifold of dimension  $n$ , equipped with a foliation  $\mathcal{F}$  of codimension  $q$ . A *transversal section* of  $\mathcal{F}$  is an embedded  $q$ -dimensional submanifold  $U \subset M$  which is everywhere transverse to the leaves. Recall that if  $\alpha$  is a path between two points  $x$  and  $y$  on the same leaf, and if  $U$  and  $V$  are transversal sections through  $x$  and  $y$ , then  $\alpha$  defines a transport along the leaves from a neighborhood of  $x$  in  $U$  to a neighborhood of  $y$  in  $V$ , hence a germ of a diffeomorphism  $hol(\alpha) : (U, x) \rightarrow (V, y)$ , called the *holonomy* of the path  $\alpha$ . Two homotopic paths always define the same holonomy. The familiar *holonomy groupoid* [6, 17, 32] is the groupoid  $Hol(M, \mathcal{F})$  over  $M$  where arrows  $x \rightarrow y$  are such germs  $hol(\alpha)$ . If the above transport “along  $\alpha$ ” is defined in all of  $U$  and embeds  $U$  into  $V$ , this embedding  $h : U \hookrightarrow V$  is sometimes also denoted by  $hol(\alpha) : U \hookrightarrow V$ . Embeddings of this form will be called *holonomy*

*embeddings*. Note that composition of paths also induces an operation of composition on those holonomy embeddings. (In section 4 below we will present a more general definition of the so-called “embedding category”).

**1.2 Transversal basis** Transversal sections  $U$  through  $x$  as above should be thought of as neighborhoods of the leaf through  $x$  in the leaf space. This motivates the definition of a *transversal basis* for  $(M, \mathcal{F})$  as a family  $\mathcal{U}$  of transversal sections  $U \subset M$  with the property that, if  $V$  is any transversal section through a given point  $y \in M$ , there exists a holonomy embedding  $h : U \hookrightarrow V$  with  $U \in \mathcal{U}$  and  $y \in h(U)$ .

Typically, a transversal section is a  $q$ -disk given by a chart for the foliation. Accordingly, we can construct a transversal basis  $\mathcal{U}$  out of a basis  $\tilde{\mathcal{U}}$  of  $M$  by domains of foliation charts  $\phi_U : \tilde{U} \xrightarrow{\sim} \mathbb{R}^{n-q} \times U$ ,  $\tilde{U} \in \tilde{\mathcal{U}}$ , with  $U = \mathbb{R}^q$ . Note that each inclusion  $\tilde{U} \hookrightarrow \tilde{V}$  between opens of  $\tilde{\mathcal{U}}$  induces a holonomy embedding  $h_{U,V} : U \rightarrow V$  defined by the condition that the plaque in  $\tilde{U}$  through  $x$  is contained in the plaque in  $\tilde{V}$  through  $h_{U,V}(x)$ .

**1.3 Transversal bundles** Let  $G$  be a Lie group and let  $\pi : P \rightarrow M$  be a principal  $G$ -bundle over  $M$ . Recall [21] that  $P$  is said to be *foliated* if  $P$  is equipped with a  $G$ -equivariant foliation  $\tilde{\mathcal{F}}$ , of the same dimension as  $\mathcal{F}$ , whose leaves are transversal to the fibers of  $\pi$  and mapped by  $\pi$  to those of  $\mathcal{F}$ . The vectors tangent to  $\tilde{\mathcal{F}}$  define a flat partial connection on  $P$ . In particular, any path  $\alpha$  in a leaf  $L$  from  $x$  to  $y$  defines a parallel transport  $P_x \rightarrow P_y$  which depends only on the homotopy class of  $\alpha$ . We call  $P$  a *transversal principal bundle* if the transport depends just on the holonomy of  $\alpha$ . A vector bundle  $E$  on  $M$  is said to be foliated (transversal) if the associated principal  $GL_r$ -bundle is foliated (transversal). By the usual relation between Cartan-Ehresmann connections and Koszul connections, we see that a foliated vector bundle is a vector bundle  $E$  over  $M$  endowed with a “flat  $\mathcal{F}$ -connection”, i.e. an operator

$$\nabla : \cdot, (\mathcal{F}) \times \cdot, (E) \rightarrow \cdot, (E)$$

satisfying the usual relations  $\nabla_{fX}(s) = f\nabla_X(s)$ ,  $\nabla_X(fs) = f\nabla_X(s) + X(f)s$ , as well as the flatness relation  $\nabla_{[X,Y]} = [\nabla_X, \nabla_Y]$ , for all  $X, Y \in \cdot, (\mathcal{F})$ ,  $f \in C^\infty(M)$ ,  $s \in \cdot, (s)$ .

Notice that if  $P$  is a transversal (principal or vector) bundle, any holonomy embedding  $h : U \hookrightarrow V$  induces a well-defined map  $h_* : P|_U \rightarrow P|_V$ , which is functorial in  $h$ . We will usually just write  $h : P|_U \rightarrow P|_V$  again for this map.

The basic example of a transversal vector bundle is the normal bundle of the foliation,  $\nu = TM/\mathcal{F}$ . The associated Koszul connection is precisely the Bott connection [2],  $\nabla_X(\overline{Y}) = \overline{[X, Y]}$ . It is a transversal bundle by the very definition of (linear) holonomy.

**1.4 Transversal sheaves** Analogous definitions apply to sheaves. A sheaf  $\mathcal{A}$  on  $M$  is called *foliated* if its restriction to each leaf is locally constant. Thus, (the homotopy class of) a path  $\alpha$  from  $x$  to  $y$  in a leaf  $L$  induces an isomorphism between stalks  $\alpha_* : \mathcal{A}_x \rightarrow \mathcal{A}_y$ . The sheaf is transversal if this isomorphism only depends on the holonomy of  $\alpha$ . A global section  $s \in \cdot, (M, \mathcal{A})$  is called *invariant* if  $s$  is invariant under transport along leaves, i.e.  $\alpha_*s(x) = s(y)$  in the notations above.

If  $\mathcal{A}$  is a transversal sheaf, any holonomy embedding  $h : U \hookrightarrow V$  gives a well-defined restriction  $h^* : \cdot, (V, \mathcal{A}) \rightarrow \cdot, (U, \mathcal{A})$ . The global section  $s$  is invariant if and only if  $h^*(s|_V) = s|_U$  for each such  $h$ .

An example of a transversal sheaf is the sheaf  $\Omega_{bas}^0$  of smooth functions which are locally constant along the leaves. One similarly has the transversal sheaves  $\Omega_{bas}^k$  of germs of basic differential  $k$ -forms. More generally, any foliated vector bundle  $E$  induces a foliated sheaf  $\Gamma(E)$  defined as follows. We denote by  $\Gamma(M; E)$  the space of sections of  $E$  which are  $\nabla$ -constant. Over  $M$ ,  $\Gamma(E)$  is the sheaf whose space of sections over an open  $U$  is  $\Gamma(E|_U)$ . Using the parallel transport with respect to  $\nabla$  we see that this sheaf is locally constant when restricted to leaves, hence it is foliated. Clearly  $\Gamma(E)$  is transversal if  $E$  is. For instance, if  $E = \Lambda^k \nu^*$ , then  $\Gamma(\Lambda^k \nu^*) = \Omega_{bas}^k$ .

Another important example is the (real) transversal orientation sheaf of the foliation, which we denote by  $\mathcal{O}$ . When restricted to a transversal open  $U$ ,  $\Gamma(U; \mathcal{O}) = H_c^q(U)^\vee$ . The foliation is transversally orientable if and only if  $\mathcal{O}$  is constant.

## 2 The transversal Čech-De Rham complex

Let  $(M, \mathcal{F})$  be a foliated manifold and let  $\mathcal{U}$  be a transversal basis. Consider the double complex which in bi-degree  $k, l$  is the vector space

$$C^{k,l} = \check{C}^k(\mathcal{U}, \Omega^l) = \prod_{U_0 \xrightarrow{h_1} \dots \xrightarrow{h_k} U_k} \Omega^l(U_0).$$

Here the product ranges over all  $k$ -tuples of holonomy embeddings between transversal sections from the given basis  $\mathcal{U}$ , and  $\Omega^k(U_0)$  is the space of differential  $k$ -forms on  $U_0$ . For elements  $\omega \in C^{k,l}$ , we denote its components by  $\omega(h_1, \dots, h_k) \in \Omega^l(U_0)$ . The vertical differential  $C^{k,l} \rightarrow C^{k,l+1}$  is  $(-1)^k d$  where  $d$  is the usual De Rham differential. The horizontal differential  $C^{k,l} \rightarrow C^{k+1,l}$  is  $\delta = \sum (-1)^i \delta_i$  where

$$\delta_i(h_1, \dots, h_{k+1}) = \begin{cases} h_1^* \omega(h_2, \dots, h_{k+1}) & \text{if } i = 0 \\ \omega(h_1, \dots, h_{i+1} h_i, \dots, h_{k+1}) & \text{if } 0 < i < k + 1 \\ \omega(h_1, \dots, h_k) & \text{if } i = k + 1 \end{cases} \quad (1)$$

This double complex is actually a bigraded differential algebra, with the usual product

$$(\omega \cdot \eta)(h_1, \dots, h_{k+k'}) = (-1)^{kk'} \omega(h_1, \dots, h_k) h_1^* \dots h_k^* \eta(h_{k+1}, \dots, h_{k+k'})$$

for  $\omega \in C^{k,l}$  and  $\eta \in C^{k',l'}$ . We will also write  $\check{C}(\mathcal{U}, \Omega)$  for the associated total complex, and refer to it as the *Čech-De Rham complex* of the foliation. The associated cohomology is denoted

$$\check{H}_{\mathcal{U}}^*(M/\mathcal{F}),$$

and referred to as the Čech-De Rham cohomology of the leaf space  $M/\mathcal{F}$ , w.r.t. the cover  $\mathcal{U}$ .

Note that, when  $\mathcal{F}$  is the codimension  $n$  foliation by points, then  $\mathcal{U}$  is a basis for the topology of  $M$ , and  $C^{k,l}$  is the usual Čech-De Rham complex [4]. Thus in this case  $\check{H}_{\mathcal{U}}^*(M/\mathcal{F}) = H^*(M)$  is the usual De Rham cohomology of  $M$ .

In general, choosing a transversal basis  $\mathcal{U}$  and a basis  $\check{\mathcal{U}}$  of  $M$  as in 1.2, there is an obvious map of double complexes  $C^{k,l}(\mathcal{U}) \rightarrow C^{k,l}(\check{\mathcal{U}})$  into the Čech-De Rham complex for the manifold  $M$ . Hence a canonical map

$$\pi^* : \check{H}_{\mathcal{U}}^*(M/\mathcal{F}) \rightarrow H^*(M; \mathbb{R}), \quad (2)$$

which should be thought of as the pull-back along the “quotient map”  $\pi : M \longrightarrow M/\mathcal{F}$ .

The standard way [7, 16] to model the leaf space of a foliation  $(M, \mathcal{F})$  is by the classifying space  $BHol(M, \mathcal{F})$  of the holonomy groupoid. Thus, the following theorem can be interpreted as a Čech-De Rham theorem for leaf spaces.

**Theorem 1** *There is a natural isomorphism*

$$\check{H}_{\mathcal{U}}^*(M/\mathcal{F}) \cong H^*(BHol(M, \mathcal{F}); \mathbb{R}),$$

*between the Čech-De Rham cohomology and the cohomology of the classifying space. In particular, the left hand side is independent of the choice of a transversal basis  $\mathcal{U}$ .*

For the proof of this theorem, we choose a complete transversal section  $T$  which contains every  $U \in \mathcal{U}$ , and we consider the “reduced holonomy groupoid”  $Hol_T(M, \mathcal{F})$ , defined as the restriction of  $Hol(M, \mathcal{F})$  to  $T$ . We may assume that  $\mathcal{U}$  is a basis for the topology of  $T$ . By a well known Morita-invariance argument, the classifying spaces  $BHol(M, \mathcal{F})$  and  $BHol_T(M, \mathcal{F})$  are weakly homotopy equivalent. The advantage of passing to a complete transversal is that  $Hol_T(M, \mathcal{F})$  becomes an étale groupoid (see section 4 for the precise definitions). For such groupoids  $\mathcal{G}$  there is a standard cohomology  $H^*(\mathcal{G}; -)$  with coefficients, which was also defined by Haefliger [18] in terms of bar-complexes, and which is known [26] to be isomorphic to the cohomology of the classifying space. In section 4 we will recall all the basic definitions. The theorem will then follow from the following proposition, which is a particular case of the Theorem 3 below.

**Proposition 1** *For any complete transversal  $T$  and any basis  $\mathcal{U}$  of  $T$ , there is a natural isomorphism*

$$\check{H}_{\mathcal{U}}^*(M/\mathcal{F}) \cong H^*(Hol_T(M, \mathcal{F}); \mathbb{R}).$$

We mention here that there are several variations of Theorem 1. For instance, for any transversal sheaf  $\mathcal{A}$  there is a Čech complex  $\check{C}(\mathcal{U}, \mathcal{A})$ . In degree  $k$ ,

$$\check{C}^k(\mathcal{U}; \mathcal{A}) = \prod_{U_0 \xrightarrow{h_1} \dots \xrightarrow{h_k} U_k}, (U_0; \mathcal{A}),$$

with the boundary  $\delta = \sum (-1)^i \delta_i$  given by the formulas (1). A consequence of the more general Theorem 3 says that, if  $\mathcal{A}|_U$  is acyclic for all  $U \in \mathcal{U}$ , then  $\check{C}(\mathcal{U}, \mathcal{A})$  computes the cohomology of the classifying space (of the reduced holonomy groupoid) with coefficients in a sheaf  $\tilde{\mathcal{A}}$  naturally associated to  $\mathcal{A}$ .

Another variation applies to the cohomology with compact supports (see section 4). Note that all these are actually extensions of the usual “Čech-De Rham isomorphisms” [4] from manifolds to leaf space. Accordingly, an immediate consequence will be the Poincaré duality for leaf spaces (see Section 6), which is one of the main results of [10]. With Theorem 1 and its analogue for compact supports available, the new proof of Poincaré duality is this time a rather straightforward extension of the classical proof [4] from manifolds to leaf spaces.

### 3 The transversal Chern-Weil map

To illustrate the usefulness of the transversal Čech-De Rham complex we will adapt the standard geometric construction of characteristic classes of principal bundles to transversal bundles, so as to obtain explicit classes in this complex. We will use the Weil-complex formulation, which we recall first (for an extensive exposition, see [21, 12]).

**3.1 Classical Chern-Weil:** [5] Recall that the Weil algebra of the Lie algebra  $\mathfrak{g}$  (of a Lie group  $G$ ) is the algebra

$$W(\mathfrak{g}) = S(\mathfrak{g}^*) \otimes \Lambda(\mathfrak{g}^*).$$

It is a graded commutative dga (graded as  $W(\mathfrak{g})^n = \bigoplus_{2p+q=n} S^p(\mathfrak{g}^*) \otimes \Lambda^q(\mathfrak{g}^*)$ ), equipped with operations  $i_X$  and  $L_X$  (linear in  $X \in \mathfrak{g}$ ) which satisfy the usual Cartan identities. In the language of [21], this means that  $W(\mathfrak{g})$  is a  $\mathfrak{g}$ -dga. If  $P$  is a principal  $G$ -bundle over a manifold  $M$ , the algebra  $\Omega^*(P)$  of differential forms on  $P$  with its usual operations  $i_X$  and  $L_X$  is another example of a  $\mathfrak{g}$ -dga. A connection  $\nabla$  on  $P$  is uniquely determined by its connection form  $\omega \in \Omega^1(P) \otimes \mathfrak{g}$ . This can be viewed as a map  $\omega : W(\mathfrak{g})^1 = \mathfrak{g}^* \rightarrow \Omega^1(P)$ , which extends uniquely to a map of  $\mathfrak{g}$ -dga's,

$$\tilde{k}(\nabla) : W(\mathfrak{g}) \rightarrow \Omega(P). \quad (3)$$

(On  $\mathfrak{g}^* = S^1(\mathfrak{g}^*) \subset W(\mathfrak{g})^2$ , it restricts to the curvature  $\Omega = d\omega + \frac{1}{2}[\omega, \omega]$ .) The restriction of this map (3) to basic elements (elements annihilated by  $i_X$  and  $G$ -invariant) gives a map of dga's

$$S(\mathfrak{g}^*)^G \rightarrow \Omega^*(M)$$

(zero differential on  $S(\mathfrak{g}^*)^G$ , the usual De Rham differential on  $\Omega(M)$ ), hence a map

$$k(\nabla) : S(\mathfrak{g}^*)^G \rightarrow H^*(M), \quad (4)$$

known as the Chern-Weil map for the principal  $G$ -bundle  $P$ . Because of the  $2p$  in the grading of the Weil algebra,  $k(\nabla)$  maps invariant polynomials of degree  $p$  to degree  $2p$  cohomology classes. Moreover,  $k(\nabla)$  does not depend on  $\nabla$ . This follows from the Chern-Simons construction (see below). A more refined characteristic map is obtained if one uses a maximal compact subgroup  $K$  of  $G$ . Since  $P/K \rightarrow M$  has contractible fibers, the map induced in De Rham cohomology is an isomorphism. Hence, to get down to the base manifold, it suffices to consider the  $K$ -basic elements of (3). Denoting by  $W(\mathfrak{g}, K)$  the subcomplex of  $W(\mathfrak{g})$  of  $K$ -basic elements, one obtains a characteristic map  $H^*(W(\mathfrak{g}, K)) \rightarrow H^*(M)$ .

**3.2 Chern-Simons:** Given  $k$  connections  $\nabla_0, \dots, \nabla_k$  on  $P$ , we consider the convex combination

$$\nabla = t_0 \nabla_0 + \dots + t_k \nabla_k \quad (5)$$

which defines a connection on the principal bundle  $\Delta^k \times P$  over  $\Delta^k \times M$ , where  $\Delta^k = \{(t_0, \dots, t_k) : t_i \geq 0, \sum t_i = 1\}$  is the standard  $k$ -simplex. We define

$$\tilde{k}(\nabla_0, \dots, \nabla_k) = (-1)^k \int_{\Delta^k} \tilde{k}(\nabla) : W(\mathfrak{g}) \rightarrow \Omega^{*-k}(P), \quad (6)$$

where  $\int_{\Delta^k} : \Omega^*(\Delta^k \times P) \rightarrow \Omega^{*-k}(P)$  is the integration along the fibers  $\Delta^k$ . Let us summarize the main properties of this construction:

(i) the map (6) commutes with the action of  $G$ , and with the operators  $i_X$ ,  $L_X$ , and it vanishes on all elements  $\alpha \otimes \beta \in W(\mathfrak{g})$  with  $\alpha$  a polynomial of degree  $> \dim(M)$ .

(ii)

$$[\tilde{k}(\nabla_0, \dots, \nabla_k), d] = \sum_{i=0}^k (-1)^i \tilde{k}(\nabla_0, \dots, \widehat{\nabla}_i, \dots, \nabla_k), \quad (7)$$

(iii) (6) is natural w.r.t. isomorphisms of principal  $G$ -bundles.

*Proof:* (ii) is just a version of Stokes' formula (see also [2]), while (iii) is obvious. We prove the vanishing result of (i). Denote by  $d$  the degree of the polynomial  $\alpha$  and by  $q$  the dimension of  $M$ . We prove that when  $d < k$  or  $2d > q + k$ , our expression

$$\theta = \tilde{k}(\nabla_0, \dots, \nabla_k)(\alpha \otimes \beta)$$

vanishes (note that if  $d > q$ , then at least one of these two equalities holds). First assume that  $d < k$ . We have  $\tilde{k}(\nabla)(\alpha \otimes \beta) = \alpha(\Omega) \wedge \beta(\omega)$ , where  $\nabla$  is the affine combination (5),  $\omega$  is the associated 1-form, and  $\Omega$  is its curvature. Let us say that a homogeneous form  $f dt_{i_1} \dots dt_{i_r} dx_{j_1} \dots dx_{j_s}$  on  $\Delta^k \times P$  has bi-degree  $(r, s)$ . Since  $\omega$  has bi-degree  $(0, 1)$ ,  $\Omega$  is a sum of forms of bi-degree  $(1, 1)$  and  $(0, 2)$ , so  $\int_{\Delta^k} \alpha(\Omega) \wedge \beta(\omega) = 0$  because no bi-degree  $(r, s)$  with  $r = k$  will be involved.

We now turn to the case  $2d > q + k$ . Let  $l$  be the degree of  $\beta$ . Because of the similar property for  $\beta$ , we have  $i_{X_1} \dots i_{X_{l+1}} \theta = 0$  for any vertical vector fields  $X_i$ . On the other hand,  $i_{Y_1} \dots i_{Y_{q+1}} \theta = 0$  for any horizontal vector fields  $Y_i$ . Since  $\deg(\theta) = 2d + l - k > l + q$ , it follows that  $\theta = 0$ .  $\square$

**3.3 Construction of the transversal Chern-Weil map:** Now let  $P$  be a transversal principal  $G$ -bundle on a foliated manifold  $(M, \mathcal{F})$ . Consider the Čech-De Rham complex

$$\check{C}^k(\mathcal{U}, \Omega^l(P)) = \prod_{U_0 \xrightarrow{h_1} \dots \xrightarrow{h_k} U_k} \Omega^l(P|_{U_0}),$$

defined exactly as in section 2 (except that  $\Omega^l(U_0)$  is replaced by  $\Omega^l(P|_{U_0})$ , and hence the horizontal differential  $\delta$  involves the maps  $h_1 : P|_{U_0} \rightarrow P|_{U_1}$  discussed in section 1). Choose a system  $\nabla = \{\nabla_U\}$  of connections, one connection  $\nabla_U$  on  $P|_U$  for each  $U$  in a transversal basis  $\mathcal{U}$ . In general we cannot assume this choice to be respected by holonomy embeddings  $h : U \rightarrow V$ , i.e.  $\nabla_U$  is in general different from the connection on  $P|_U$  induced by  $\nabla_V$  via the isomorphism  $h : P|_U \rightarrow P|_{h(U)}$ . Denote this last connection by  $\nabla_h$ . For a string  $U_0 \xrightarrow{h_1} \dots \xrightarrow{h_k} U_k$  of holonomy embeddings, we consider the map (see 3.1 above)

$$\tilde{k}(\nabla_{U_0}, \nabla_{h_1}, \nabla_{h_2 h_1}, \dots, \nabla_{h_k \dots h_2 h_1}) : W(\mathfrak{g}) \rightarrow \Omega^{*-k}(P|_{U_0}). \quad (8)$$

Doing this for all such strings, we obtain a map into the total complex

$$\tilde{k}(\omega) : W(\mathfrak{g}) \rightarrow \prod_{U_0 \xrightarrow{h_1} \dots \xrightarrow{h_k} U_k} \Omega^{*-k}(P|_{U_0}). \quad (9)$$

This map respects the total degree, and it is obviously compatible with the operations  $i_X$  and the  $G$ -action. So, by restricting to basic elements it yields a map into the transversal Čech-De Rham complex

$$\tilde{k}(\omega) : S(\mathfrak{g}^*)^G \longrightarrow \check{C}^*(\mathcal{U}, \Omega^*) \quad (10)$$

(mapping degree  $p$  polynomials into elements of total degree  $2p$ ).

**Theorem 2** *The Chern-Weil map of a transversal principal  $G$ -bundle  $P$  over  $(M, \mathcal{F})$  has the following properties:*

(i) *The maps (9) and (10) respect the differential, hence they induce a map*

$$k_P := k(\nabla) : S(\mathfrak{g}^*)^G \longrightarrow \check{H}_U^*(M/\mathcal{F}) , \quad (11)$$

(ii) *This map (11) does not depend on the choice of the connections  $\{\nabla_U\}$ , and respects the products.*

(iii) *Composed with the pull-back map  $\pi^* : \check{H}_U^*(M/\mathcal{F}) \longrightarrow H^*(M)$ , see (2), it gives the usual Chern-Weil map (4) of  $P$ .*

(iv) (“Bott vanishing theorem”) *The image of the map (11) is zero in degrees  $> 2q$ , where  $q$  is the codimension of  $\mathcal{F}$ .*

The classical Bott vanishing theorem [2] (for the normal bundle of the foliation) and its extensions to foliated bundles [21] are at the level of  $H^*(M)$ . The point of Theorem 2 is that, using *classical geometrical arguments*, one can prove these vanishing results and construct the resulting characteristic classes at the level of the leaf space, i.e. in the cohomology of the classifying space (cf. Theorem 1).

*Proof of Theorem 2:* (i) and (iv) clearly follow from the main properties of the Chern-Simons construction 3.2. Also (iii) will follow from the independence of the connections. Indeed, it suffices to check that, if  $\mathcal{F}$  is the foliation by points, then the resulting map  $k_\nabla : S(\mathfrak{g}^*)^G \longrightarrow \check{H}_U^*(M)$  composed with Čech-De Rham isomorphism  $\check{H}_U^*(M) \cong H^*(M)$  (induced by the inclusion  $\Omega^*(M) \subset C^*(\mathcal{U}, \Omega^*)$  [4]) gives the usual Chern-Weil map. But this is clear even at the chain level, provided we choose  $\nabla_U = \nabla|_U$  for some globally defined connection  $\nabla$ .

(ii) For two different choices  $\nabla = \{\nabla_U\}$  and  $\nabla' = \{\nabla'_U\}$  of connections, the map  $H : W(\mathfrak{g}) \longrightarrow C^*(\mathcal{U}, \Omega^*)$  defined by

$$H^*(w)(h_1, \dots, h_k) = \sum_{i=0}^k (-1)^i k(\nabla_{h_i \dots h_2 h_1}, \dots, \nabla_{h_k \dots h_2 h_1}, \nabla'_{U_0}, \nabla'_{h_1}, \dots, \nabla'_{h_i \dots h_2 h_1})(w) .$$

provides an explicit chain homotopy. To prove the compatibility with the products, one can either proceed as in [21] using the simplicial Weil complex (see [9] for details), or, since the characteristic map is constructed through the double complex  $\check{C}^p(\mathcal{U}, \Omega^{p+q}(\Delta^q \times P))$  by integration over the simplices, one can use the simplicial De Rham complex and Theorem 2.14 of [13].  $\square$



**3.4 Exotic characteristic classes:** The vanishing result of Theorem 2 shows that the construction of the “exotic” classes also lifts to the Čech-De Rham complex. To describe all the relevant characteristic classes, we consider the complex  $W(\mathfrak{g}, K)$  of  $K$ -basic elements described in 3.1, together with its  $q$ -th truncation  $W_q(\mathfrak{g}, K)$  defined as the quotient by the ideal generated by the elements of polynomial degree  $> q$ . By the vanishing result (more precisely from the proof above), the map (9) induces a chain map  $W_q(\mathfrak{g}, K) \rightarrow \check{C}^*(\mathcal{U}, \Omega^*(P/K))$ . Using the contractibility of  $G/K$  as in 3.1, we obtain the following refinement of the characteristic map of Theorem 2.

**Corollary 1** *The Chern-Weil construction of 3.3 gives a well-defined algebra map*

$$k_P^{ex} := k^{ex}(\nabla) : H^*(W_q(\mathfrak{g}, K)) \rightarrow \check{H}_{\mathcal{U}}^*(M/\mathcal{F}), \quad (12)$$

*again independent of the choice of connections. Moreover, composed with the pull-back map  $\pi^* : \check{H}_{\mathcal{U}}^*(M/\mathcal{F}) \rightarrow H^*(M)$  (see (2)), it gives the exotic characteristic map of the foliated bundle  $P$  [21].*

## 4 The Čech-De Rham complex of an étale groupoid

In this section we prove Theorem 1, as well as some generalizations and variants, in the context of étale groupoids. Our general goal is to describe the (hyper-) homology and cohomology of étale groupoids in terms of the (hyper-) homology and cohomology of small categories. We begin by introducing some standard terminology.

**4.1 Smooth étale groupoids:** A *smooth groupoid* is a groupoid  $\mathcal{G}$  for which the sets  $\mathcal{G}^{(0)}$  and  $\mathcal{G}^{(1)}$  of objects and arrows have the structure of a smooth manifold, all the structure maps are smooth, and the source and the target maps are moreover submersions. The holonomy groupoid  $Hol(M, \mathcal{F})$  of a foliation is an example of a smooth groupoid. Such a smooth groupoid is said to be *étale* if the source and the target maps are local diffeomorphisms. In this case the manifolds  $\mathcal{G}^{(0)}$  and  $\mathcal{G}^{(1)}$  have the same dimension, to which we refer as *the dimension of  $\mathcal{G}$* . An example of an étale groupoid of dimension  $q$  is the universal Haefliger groupoid  $\mathcal{H}_q$  for codimension  $q$  foliations [17]. There is an important notion of *Morita equivalence* between smooth groupoids, see e.g. [7, 17, 24, 28]. For any foliation, the holonomy groupoid  $Hol(M, \mathcal{F})$  is Morita equivalent to an étale groupoid, namely to its restriction to any complete transversal  $T$ , denoted  $Hol_T(M, \mathcal{F})$ . A Morita equivalence between smooth groupoids induces a weak homotopy equivalence between their classifying spaces.

**4.2 Sheaves and cohomology:** For a smooth étale groupoid  $\mathcal{G}$ , a  $\mathcal{G}$ -sheaf is a sheaf  $\mathcal{A}$  over the space  $\mathcal{G}^{(0)}$ , equipped with a continuous  $\mathcal{G}$ -action. For any such sheaf there are natural cohomology groups  $H^n(\mathcal{G}; \mathcal{A})$  whose definition we recall. Denote by  $\mathcal{G}^{(k)}$  the space of composable arrows

$$x_0 \xrightarrow{g_1} \dots \xrightarrow{g_k} x_k \quad (13)$$

of  $\mathcal{G}$ , and by  $\epsilon_k : \mathcal{G}^{(k)} \rightarrow \mathcal{G}^{(0)}$  the map which sends (13) to  $x_0$ . The bar complex of  $\mathcal{A}$  is defined by  $B^k(\mathcal{G}; \mathcal{A}) = \mathcal{G}^{(k)}; \epsilon_k^* \mathcal{A}$ , hence consists on continuous functions  $c$  which associate to a string of arrows (13) an element  $c(g_1, \dots, g_k) \in \mathcal{A}_{x_0}$ . The boundary is

$\delta = \sum (-1)^i \delta_i$  with the same formulas as in (1). If  $\mathcal{A}$  is “good” in the sense that  $\mathcal{A}$  and its pull-backs  $c_k^* \mathcal{A}$  are injective sheaves, then  $H^n(\mathcal{G}; \mathcal{A})$  is computed by the bar complex  $B(\mathcal{G}, \mathcal{A})$ . In general, one chooses a resolution  $\mathcal{S}^*$  of  $\mathcal{A}$  by “good”  $\mathcal{G}$ -sheaves, and  $H^n(\mathcal{G}; \mathcal{A})$  is computed by the double complex  $B^k(\mathcal{G}; \mathcal{S}^l)$ . In general, these cohomology groups coincide with the cohomology groups of the classifying space  $B\mathcal{G}$  [26].

Similarly, using compact supports and direct sums in the definition of the bar complex, one defines the homology groups  $H_*(\mathcal{G}; \mathcal{A})$  [10] (sometimes denoted  $H_c^*(\mathcal{G}; \mathcal{A}) = H_{-*}(\mathcal{G}; \mathcal{A})$ ), which should be thought of as a good model for the compactly supported cohomology of the classifying space.

**4.3 Čech complexes:** Let  $\mathcal{G}$  be an étale groupoid and let  $\mathcal{U}$  be a basis of opens in  $\mathcal{G}^{(0)}$ . A  $\mathcal{G}$ -embedding  $\sigma : U \rightarrow V$  is a smooth family  $\sigma(x)$ ,  $x \in U$ , where each  $\sigma(x) : x \rightarrow y$  is an arrow in  $\mathcal{G}$  from  $x$  to some point  $y \in V$ ; moreover, the map  $x \rightarrow \text{target}(\sigma(x))$  should define an embedding of  $U$  into  $V$ . As in the first section, we can now define the Čech complex  $\check{C}(\mathcal{U}; \mathcal{A})$  for any  $\mathcal{G}$ -sheaf  $\mathcal{A}$ ,

$$\check{C}^k(\mathcal{U}; \mathcal{A}) = \prod_{U_0 \rightarrow \dots \rightarrow U_k} (U_0, \mathcal{A}),$$

where the product is over all strings of  $\mathcal{G}$ -embeddings between opens  $U \in \mathcal{U}$ , and the boundary  $\delta = \sum (-1)^i \delta_i$  is given by the same formulas as in (1).

We say that  $\mathcal{A}$  is  $\mathcal{U}$ -acyclic if  $H^i(U; \mathcal{A}) = 0$  for each  $i > 0$  and each  $U \in \mathcal{U}$ . In this case define  $\check{H}_{\mathcal{U}}^*(\mathcal{G}; \mathcal{A})$  as the cohomology of  $\check{C}(\mathcal{U}; \mathcal{A})$ . In general, we define  $\check{H}_{\mathcal{U}}^*(\mathcal{G}; \mathcal{A})$  as the cohomology of the double complex  $\check{C}^k(\mathcal{U}; \mathcal{S}^l)$ , where  $0 \rightarrow \mathcal{A} \rightarrow \mathcal{S}^0 \rightarrow \dots \rightarrow \mathcal{S}^d \rightarrow 0$  is a bounded resolution by  $\mathcal{U}$ -acyclic sheaves,  $d = \dim(\mathcal{G})$ . By the usual arguments, such resolutions always exist, and the definition does not depend on the choice of the resolution.

**4.4 Examples:** The  $\mathcal{G}$ -sheaf  $\Omega_{\mathcal{G}}^l$  of  $l$ -differential forms with its natural  $\mathcal{G}$ -action is always  $\mathcal{U}$ -acyclic, as is any soft  $\mathcal{G}$ -sheaf. We obtain the Čech-De Rham (double) complex of  $\mathcal{G}$ ,  $\check{C}(\mathcal{U}; \Omega)$ , computing  $\check{H}_{\mathcal{U}}^*(\mathcal{G}; \mathbb{R})$ . If the basis  $\mathcal{U}$  consists of contractible opens (balls), then any locally constant  $\mathcal{G}$ -sheaf  $\mathcal{A}$  is  $\mathcal{U}$ -acyclic, hence  $\check{H}_{\mathcal{U}}^*(\mathcal{G}; \mathcal{A})$  is computed by  $\check{C}(\mathcal{U}; \mathcal{A})$ .

Similarly one defines the Čech complex with compact supports  $\check{C}_c^*(\mathcal{U}; \mathcal{A})$  using

$$\bigoplus_{U_0 \rightarrow \dots \rightarrow U_k} (U_0, \mathcal{A}).$$

In order to get a cochain complex, we associate the degree  $-k$  to the direct sums over strings of  $k$   $\mathcal{G}$ -embeddings. If  $\mathcal{A}$  is  $c$ -soft, then  $\check{H}_{c, \mathcal{U}}^*(\mathcal{G}; \mathcal{A})$  is defined by  $\check{C}_c(\mathcal{U}; \mathcal{A})$ . In general, one uses a resolution  $0 \rightarrow \mathcal{A} \rightarrow \mathcal{S}^0 \rightarrow \dots \rightarrow \mathcal{S}^d \rightarrow 0$  by  $c$ -soft  $\mathcal{G}$ -sheaves, and the double complex  $\check{C}_c^k(\mathcal{U}; \mathcal{S}^l)$ . The resulting cohomology is denoted  $\check{H}_{c, \mathcal{U}}^*(\mathcal{G}; \mathcal{A})$ .

**4.5 The embedding category:** The notion of  $\mathcal{G}$ -embedding originates in [25], where the second author has introduced a small category  $Emb_{\mathcal{U}}(\mathcal{G})$  for each basis  $\mathcal{U}$  of open sets. The objects of  $Emb_{\mathcal{U}}(\mathcal{G})$  are the members  $U$  of  $\mathcal{U}$ , and the arrows are the  $\mathcal{G}$ -embeddings between the opens of  $\mathcal{U}$ . The main result of [25] was that the classifying

space  $B\mathcal{G}$  is weakly homotopy equivalent to the CW-complex  $BEmb_{\mathcal{U}}(\mathcal{G})$ , provided each of the basic opens in  $\mathcal{U}$  is contractible.

Now any  $\mathcal{G}$ -sheaf  $\mathcal{A}$  defines an obvious contravariant functor  $\check{C}(\mathcal{A})$  on  $Emb_{\mathcal{U}}(\mathcal{G})$  sending  $U$  to  $\check{C}(U; \mathcal{A})$ , and  $\check{C}(\mathcal{U}; \mathcal{A})$  is just the usual (bar) complex computing the cohomology  $H^*(Emb_{\mathcal{U}}(\mathcal{G}); \check{C}(\mathcal{A}))$  of the discrete category  $Emb_{\mathcal{U}}(\mathcal{G})$  with coefficients. Hence [25] proves that  $H^*(\mathcal{G}; \mathcal{A}) = \check{H}_{\mathcal{U}}^*(\mathcal{G}; \mathcal{A})$  provided all the opens  $U \in \mathcal{U}$  are contractible and  $\mathcal{A}$  is (locally) constant. We now prove a stronger ‘‘Čech-De Rham isomorphism’’ which applies to more general coefficients, and also to compact supports.

**Theorem 3** *Let  $\mathcal{G}$  be an étale groupoid, and let  $\mathcal{U}$  be a basis for  $\mathcal{G}^{(0)}$  as above. Then for any  $\mathcal{G}$ -sheaf  $\mathcal{A}$ , there are natural isomorphisms*

$$H^n(\mathcal{G}; \mathcal{A}) = \check{H}_{\mathcal{U}}^n(\mathcal{G}; \mathcal{A}), \quad H_c^n(\mathcal{G}; \mathcal{A}) = \check{H}_{c, \mathcal{U}}^n(\mathcal{G}; \mathcal{A}).$$

*Proof:* The proofs of the isomorphisms in the statement are similar, and we only prove the first one (an explicit proof of the second one also occurs in [9]). By comparing resolutions of the  $\mathcal{G}$ -sheaf  $\mathcal{A}$ , it suffices to find a suitable complex  $C(\mathcal{A})$  and explicit quasi-isomorphisms

$$B(\mathcal{G}; \mathcal{A}) \longleftarrow C(\mathcal{A}) \longrightarrow \check{C}(\mathcal{U}; \mathcal{A})$$

natural in  $\mathcal{A}$ , for the case where  $\mathcal{A}$  is ‘‘good’’ in the sense of 4.2. For this we consider the bisimplicial space  $S_{p,q}$ , whose  $p, q$ -simplices are of the form

$$x_0 \xrightarrow{g_1} \dots \xrightarrow{g_q} x_q \xrightarrow{g} U_0 \xrightarrow{\sigma_1} \dots \xrightarrow{\sigma_p} U_p, \quad (14)$$

where  $\sigma_1, \dots, \sigma_p$  are  $\mathcal{G}$ -embeddings, and  $g_1, \dots, g_q, g$  are arrows in the groupoid  $\mathcal{G}$ , the notation  $x_q \xrightarrow{g} U_0$  indicating that the target of  $g$  is in  $U_0$ . The topology on  $S_{p,q}$  is the topology induced from the topology on  $\mathcal{G}$ ,

$$S_{p,q} = \coprod_{U_0 \rightarrow \dots \rightarrow U_p} \mathcal{G}^{(q)}.$$

The  $\mathcal{G}$ -sheaf  $\mathcal{A}$  induces a sheaf  $\mathcal{A}_{p,q}$  on  $S_{p,q}$  by pull-back along the projection which maps (14) to  $x_q$ . Consider the double complex  $C = C(\mathcal{A})$ ,

$$C^{p,q} = \check{C}(S_{p,q}, \mathcal{A}_{p,q}).$$

For a fixed  $p$ , the complex  $C^{p,*}$  is a product of complexes, namely, for each string  $U_0 \rightarrow \dots \rightarrow U_p$ , the bar complex (see 4.2) of the (étale) comma groupoid  $\mathcal{G}/U_0$  with coefficients in the pull-back of the sheaf  $\mathcal{A}$ . Since the groupoid  $\mathcal{G}/U_0$  is Morita equivalent to the space  $U_0$ , this cohomology is  $H^*(U_0; \mathcal{A})$ . Since  $\mathcal{A}$  is assumed to be good,  $H^*(U_0; \mathcal{A})$  vanishes in positive degrees, and we conclude that the canonical map

$$\check{C}^p(\mathcal{U}; \mathcal{A}) \longrightarrow C^{p,*}$$

is a quasi-isomorphism for each fixed  $p$ . Write  $\pi_{p,q} : S_{p,q} \rightarrow \mathcal{G}^{(q)}$  for the projection of (14) to the string  $x_0 \rightarrow \dots \rightarrow x_q$ . Then  $C^{p,q} = \check{C}(\pi_{p,q}, (\pi_{p,q})_*(\mathcal{A}_{p,q}))$ . The stalk of  $(\pi_{p,q})_*(\mathcal{A}_{p,q})$  at  $x_0 \rightarrow \dots \rightarrow x_q$  is

$$\lim_{x_q \in U} \left( \prod_{U \rightarrow U_0 \rightarrow \dots \rightarrow U_p} (U; \mathcal{A}) \right), \quad (15)$$

where the colimit is taken over all basic open neighborhoods  $U$  of  $x_q$ . For a fixed  $U$ , the complex inside the  $\lim$  in (15) computes the cohomology of the (discrete) comma category  $U/Emb_{\mathcal{U}}(\mathcal{G})$  with coefficients in the constant group  $\mathbb{Z}$ ,  $(U, \mathcal{A})$ . Since the comma category is contractible, so is this complex. Taking the colimit, we see that for each  $q$  the map  $\mathcal{A} \rightarrow (\pi_{p,q})_*(\mathcal{A}_{p,q})$  is a quasi-isomorphism of (complexes of) injective sheaves on  $\mathcal{G}^{(q)}$ . Thus the natural map

$$B^q(\mathcal{G}; \mathcal{A}) = \mathbb{Z} \otimes_{\mathbb{Z}} (\mathcal{G}^{(q)}; \epsilon_q^* \mathcal{A}) \longrightarrow \mathbb{Z} \otimes_{\mathbb{Z}} ((\pi_{*,q})_*(\mathcal{A}_{*,q})) = C^{*,q}$$

is a quasi-isomorphism, and the proof is complete.  $\square$

Regarding the relation with the embedding category 4.5 and its cohomology, let us point out the following immediate consequence, which is an improvement of the result of [25].

**Corollary 2** *If  $\mathcal{G}$  is an étale groupoid,  $\tilde{\mathcal{U}}$  is a basis of opens of  $\mathcal{G}^{(0)}$ , and  $\mathcal{A}$  is a  $\mathcal{G}$ -sheaf with the property that  $H^k(U, \mathcal{A}|_U) = 0$  for all  $U \in \mathcal{U}$ ,  $k \geq 1$ , then*

$$H^*(\mathcal{G}; \mathcal{A}) \cong H^*(Emb_{\mathcal{U}}(\mathcal{G}); \mathbb{Z}, (\mathcal{A})) .$$

*Similarly, if  $H_c^k(U, \mathcal{A}|_U) = 0$  for all  $U \in \mathcal{U}$ ,  $k \geq 1$ , then*

$$H_c^*(\mathcal{G}; \mathcal{A}) \cong H_{-*}(Emb_{\mathcal{U}}(\mathcal{G}); \mathbb{Z}, {}_c(\mathcal{A})) .$$

*Also, if each  $U \in \mathcal{U}$  is contractible, and  $\mathcal{A}$  is locally constant as a sheaf on  $\mathcal{G}^{(0)}$ , then*

$$H^*(\mathcal{G}; \mathcal{A}) \cong H^*(Emb_{\mathcal{U}}(\mathcal{G}); \mathbb{Z}, (\mathcal{A})), \quad H_c^*(\mathcal{G}; \mathcal{A}) \cong H_{d-*}(Emb_{\mathcal{U}}(\mathcal{G}); H_c^d(-; \mathcal{A}))$$

*(where  $d$  is the dimension of the base space  $\mathcal{G}^{(0)}$ ).*

**4.6 Chern-Weil for étale groupoids:** Clearly all the constructions of Section 3 apply to any étale groupoid  $\mathcal{G}$ , provided we use the Čech-De Rham complexes mentioned in 4.4. Hence, for any principal  $G$ -bundle  $P$  endowed with a smooth action of  $\mathcal{G}$ , one has an associated Chern-Weil map

$$S(\mathfrak{g}^*)^G \longrightarrow H^*(\mathcal{G}; \mathbb{R})$$

whose image vanishes in degrees  $> 2d$ , where  $d = \dim(\mathcal{G})$ . The refined characteristic map,

$$H^*(W_d(\mathfrak{g}, K)) \longrightarrow H^*(\mathcal{G}; \mathbb{R})$$

defines the exotic characteristic classes. Of particular interest is the (frame bundle of the) tangent space of  $\mathcal{G}^{(0)}$ , which is naturally endowed with an action of  $\mathcal{G}$ , and which induces the exotic characteristic map of  $\mathcal{G}$ ,

$$k_{\mathcal{G}} : H^*(WO_d) \longrightarrow \check{H}_{\mathcal{U}}^*(\mathcal{G}) \cong H^*(B\mathcal{G}; \mathbb{R}) . \quad (16)$$

When  $\mathcal{G} = \text{Hol}_T(M, \mathcal{F})$  this is the map discussed in section 3. But this is not the only interesting example. For instance, if one works with foliated bundles which are not necessarily transversal (as e.g. in [21]), then one has to replace the holonomy groupoid  $\text{Hol}_T(M, \mathcal{F})$  by the monodromy groupoid  $\text{Mon}_T(M, \mathcal{F})$ . The new versions of Theorem 2 and Corollary 1 for foliated bundles then yield characteristic classes in  $H^*(B\text{Mon}_T(M, \mathcal{F}))$ . These classes are refinements of the characteristic classes in  $H^*(M)$ , already constructed in [21].

Another interesting example is when  $\mathcal{G}$  is Haefliger's  $\mathcal{G}^q$ . The importance of this example lies into the fact that  $\mathcal{G}^q$  plays a classifying role for codimension  $q$  foliations, hence its cohomology consists on “universal” classes. We will elaborate this in 5.2 of the next section.

## 5 Explicit formulas

In this section we illustrate our constructions in the case of normal bundles. In particular we deduce Bott's formulas for cocycles associated to group actions [3], as well as Thurston's formula.

**5.1 Explicit formulas for the normal bundle:** We now apply the construction of the exotic characteristic map of Section 3 to the normal bundle  $\nu$ . Corollary 1 applied to the (principal  $GL_q$ -bundle associated to)  $\nu$  provides us with a characteristic map

$$k_{\mathcal{F}} : H^*(WO_q) \longrightarrow \check{H}_{\mathcal{U}}^*(M/\mathcal{F}) ,$$

which, when composed with the pull-back  $\pi^* : \check{H}_{\mathcal{U}}^*(M/\mathcal{F}) \longrightarrow H^*(M)$ , gives the familiar exotic characteristic classes [2] of  $\mathcal{F}$ . Here  $WO_q$  is the standard [2] simplification of the truncated relative Weil complex  $W_q(\mathfrak{gl}_q, O(q))$  that we now recall. The idea is that the relative Weil complex  $W(\mathfrak{gl}_q, O(q))$  (see 3.1) is quasi-isomorphic to a smaller subcomplex, namely the dg algebra  $S[c_1, \dots, c_q] \otimes E(h_1, h_3, \dots, h_{2[\frac{q+1}{2}-1]})$  generated by elements  $c_i$  of degree  $2i$  (namely the polynomials  $c_i(A) = \text{Tr}(A^i)$ ), elements  $h_{2i+1}$  of degree  $4i + 1$  (any elements which transgress  $c_{2i+1}$ ), with the boundary

$$d(c_i) = 0, \quad d(h_{2i+1}) = c_{2i+1}.$$

Truncating by polynomials of degree  $> q$ , the resulting inclusion

$$WO_q := S_q[c_1, \dots, c_q] \otimes E(h_1, h_3, \dots, h_{2[\frac{q+1}{2}-1]}) \longrightarrow W_q(\mathfrak{gl}_q, O(q))$$

induces isomorphism in cohomology. With this simplification, the desired cohomology can be computed explicitly. Apart from the classical Chern elements  $c_i$  (non-trivial only for  $i < q/2$  even) there are new exotic classes. Referring to [14] for the complete description of  $H^*(WO_q)$ , we recall here that the simplest such class is the Godbillon-Vey class  $gv = [h_1 c_1^q] \in H^{2q+1}(WO_q)$ . We denote by  $gv_{\mathcal{F}} \in \check{H}_{\mathcal{U}}^*(M/\mathcal{F})$  the resulting cohomology class  $k_{\mathcal{F}}(gv)$ . Its pull-back to  $H^*(M)$  is the usual Godbillon-Vey class of  $\mathcal{F}$ . More generally, the Bott-Godbillon-Vey classes  $gv^\alpha = [u_1 c_{\alpha_1} \dots c_{\alpha_t}]$  (and their images  $gv_{\mathcal{F}}^\alpha$ ) are defined for any partition  $\alpha = (\alpha_1, \dots, \alpha_t)$  of  $q$  (i.e.  $q = \sum \alpha_i$ ).

For explicit formulas, let us choose a basis  $\mathcal{U}$  so that  $\tilde{\mathcal{U}}$  are also domains of trivialization charts for  $\nu$  (as in 1.2). Let  $J_h : U \rightarrow GL_q$  denote the Jacobian of  $h : U \rightarrow V$  (any holonomy embedding). Then the  $J_h$ 's are the associated transition functions of the transversal bundle  $\nu$ . Locally, we choose the trivial connection  $\nabla_U$  over  $U$ . The corresponding  $\nabla(h)$  are then given by the connection 1-forms:

$$\omega_h := J_h^{-1} dJ_h \in \Omega^1(U; \mathfrak{gl}_q),$$

for  $h : U \rightarrow V$ . We see that the Chern character  $Ch_\nu \in \check{C}^2(\mathcal{U}, \Omega^*)$  is given by:

$$(h_1, \dots, h_p) \mapsto (-1)^p \int_{t_0+t_1+\dots+t_p \leq 1} \exp((t_1\omega_{h_1} + t_2\omega_{h_2 h_1} + \dots + t_p\omega_{h_p \dots h_2 h_1})^2) dt_0 dt_1 \dots dt_p.$$

For instance, the first class  $C_1 = ch_1(\nu) \in \check{C}^*(\mathcal{U}, \Omega^*)$  has the components

$$C_1^{(1,1)}(h) = Tr(J_h^{-1} dJ_h), \quad C_1^{(0,2)} = C_1^{(2,0)} = 0.$$

As we know, this class is cohomologically trivial. This can be seen directly, since  $U_1 \in \check{C}^1(\mathcal{U}, \Omega^*)$ ,

$$U_1^{(0,1)} = 0, \quad U_1^{(1,0)}(h) = \log(|\det(J_h)|)$$

transgresses  $C_1$ . Computing the resulting closed cocycle  $U_1 C_1^q$  we see that

**Corollary 3** *The Godbillon-Vey class  $gv_{\mathcal{F}} \in \check{H}^{2q+1}(M/\mathcal{F})$  is represented in the Čech-De Rham complex by the cocycle  $gv_{\mathcal{F}}$  living in bi-degree  $(q+1, q)$ :*

$$gv_{\mathcal{F}}(h_1, \dots, h_{q+1}) = \log(|\det(J_{h_1})|) h_1^* Tr(\omega_{h_2}) h_1^* h_2^* Tr(\omega_{h_3}) \dots h_1^* \dots h_q^* Tr(\omega_{h_{q+1}}). \quad (17)$$

Similarly, computing  $U_1 C_{\alpha_1} \dots C_{\alpha_t}$  for a partition  $\alpha = (\alpha_1, \dots, \alpha_t)$  of  $q$ , we obtain the following formula, which explains Bott's definition of the cocycles associated to group actions [3].

**Corollary 4** *The Bott-Godbillon-Vey class  $gv_{\mathcal{F}}^\alpha \in \check{H}^{2q+1}(M/\mathcal{F})$  is represented in the Čech-De Rham complex by the closed cocycle  $gv_{\mathcal{F}}^\alpha$  living in bi-degree  $(q+1, q)$ :*

$$gv_{\mathcal{F}}^\alpha(h_1, \dots, h_{q+1}) = \log(|\det(J_{h_1})|) \cdot h_1^* \{ Tr[\omega_{h_2} \cdot h_2^*(\omega_{h_3}) \cdot \dots \cdot (h_{(\alpha_1-1)} \dots h_2)^*(\omega_{h_{\alpha_1}})] \} \cdot \\ (h_{\alpha_1} \dots h_2 h_1)^* \{ Tr[\omega_{h_{(\alpha_1+1)}} \cdot h_{(\alpha_1+1)}^*(h_{(\alpha_1+2)}) \cdot \dots \cdot (h_{(\alpha_1+\alpha_2-1)} \dots h_{(\alpha_1+1)})^*(\omega_{h_{\alpha_2}})] \} \cdot \dots$$

**5.2 Universal formulas.** As pointed out in the previous section, the constructions that we described for foliations apply to any étale groupoid. Due to its classifying properties, the case of the Haefliger groupoid  $,^q$  is of particular interest. We wish to explain how the Čech-De Rham model for  $,^q$  can be used to derive, in an explicit and straightforward way, the known formulas and properties of universal characteristic classes for codimension  $q$  foliations. We emphasize that all these properties are now part of the folklore on characteristic classes for foliations, but they are usually derived by non-trivial abstract arguments at the level of classifying spaces.

First of all we make a slight simplification of the Čech-De Rham complex of  $,^q$ . Choosing the basis  $\mathcal{U}$  of  $\mathbb{R}^q$  by discs, since any such disc is diffeomorphic to  $\mathbb{R}^q$ , we see that the category  $Emb_{\mathcal{U}}(,^q)$  is equivalent to the category which has only one object,

and all the embeddings  $\mathbb{R}^q \longrightarrow \mathbb{R}^q$  as arrows. Accordingly, we define  $\check{C}(\cdot, {}^q; \Omega)$  as in the previous sections, except that we take products only over strings

$$\mathbb{R}^q \xrightarrow{\sigma_1} \dots \xrightarrow{\sigma_k} \mathbb{R}^q$$

of embeddings  $\mathbb{R}^q \longrightarrow \mathbb{R}^q$ . The main theorem of this section implies

**Corollary 5** *The Čech-De Rham complex  $\check{C}(\cdot, {}^q; \Omega)$  computes  $H^*(B, {}^q; \mathbb{R})$ .*

Now we can describe the main (cohomological) universal properties of  $\cdot, {}^q$  in an explicit (and obvious) fashion. First of all, the universal property of  $\cdot, {}^q$  can be seen easily in cohomology: given any codimension  $q$  foliation, choosing a basis  $\tilde{\mathcal{U}}$  for  $M$  and a transversal basis  $\mathcal{U}$  as in 1.2, there is an obvious map  $\check{H}(\cdot, {}^q) \longrightarrow \check{H}_{\tilde{\mathcal{U}}}(M)$ , to be seen as the map induced in cohomology by the classifying map  $M \longrightarrow B, {}^q$  of  $\mathcal{F}$  (well defined up to homotopy). This map is the composition of the pull-back (2) with another obvious map

$$\check{H}^*(\cdot, {}^q) \longrightarrow \check{H}_{\mathcal{U}}^*(M/\mathcal{F}) \quad (18)$$

(compare to [2]). Now, all the characteristic maps for codimension  $q$  foliations are just the composition of the (18)'s with a universal map

$$k_q : H^*(WO_q) \longrightarrow \check{H}^*(\cdot, {}^q) . \quad (19)$$

Again, with the Čech-De Rham complexes at hand this is obvious, and  $k_q$  is not at all abstract: it is just the characteristic map (16) applied to  $\mathcal{G} = \cdot, {}^q$  and can be described in terms of the trivial connection on  $\mathbb{R}^q$  (compare to [2]). In particular, all the formulas of section 3 come from similar universal formulas in  $\check{C}(\cdot, {}^q; \Omega)$ .

At the price of more complicated formulas, we can further simplify the complex  $\check{C}(\cdot, {}^q; \Omega)$ . Indeed, since the cohomology of  $\Omega^*(\mathbb{R}^q)$  is  $\mathbb{R}$  concentrated in degree zero (Poincaré lemma), we see that  $\check{H}(\cdot, {}^q)$  is also computed by the Čech (subcomplex) with constant coefficients

$$\check{C}(\cdot, {}^q) : \quad 0 \longrightarrow \mathbb{R} \longrightarrow \prod_{\mathbb{R}^q \xrightarrow{\sigma_1} \mathbb{R}^q} \mathbb{R} \longrightarrow \prod_{\mathbb{R}^q \xrightarrow{\sigma_1} \mathbb{R}^q \xrightarrow{\sigma_2} \mathbb{R}^q} \mathbb{R} \longrightarrow \dots$$

To pass from  $\check{C}(\cdot, {}^q; \Omega)$  to  $\check{C}(\cdot, {}^q)$  one has to repeatedly apply the Poincaré lemma. After a lengthy but straightforward computation (for the details see Lemma 3.3.8 in [9]) we obtain:

**Lemma 1** *An  $n$ -cocycle in the Čech-De Rham complex:*

$$u = u_0 + u_1 + \dots + u_n, \quad u_k \in \check{C}^{tk}(\cdot, {}^q, \Omega^{n-k})$$

*represents the same class in  $\check{H}^n(\cdot, {}^q)$  as the  $n$ -cocycle  $\tilde{u}$  in the Čech complex  $\check{C}^{ts}(\cdot, {}^q)$ , given by:*

$$\tilde{u}(\sigma_1, \dots, \sigma_n) = \sum_{s=0}^n (-1)^{n(s-1) + \frac{s(s-1)}{2}} \int_{I_{\sigma_1, \dots, \sigma_s}} u_{n-s}(\sigma_{s+1}, \dots, \sigma_n).$$

*Here,  $I_{\sigma_1, \dots, \sigma_s}$  is the  $s$ -cube:*

$$I_{\sigma_1, \dots, \sigma_s}(t_1, \dots, t_s) = \sigma_s(\sigma_{s-1}(\dots \sigma_3(\sigma_2(\sigma_1(0)t_1)t_2)\dots)t_{s-1})t_s.$$

If we apply this to the Godbillon-Vey cocycle (i.e. to the formula (17) in the Čech-De Rham complex  $\check{C}(\cdot, \cdot; \Omega)$ ), we obtain the well-known Thurston's formula:

**Corollary 6** *The universal Godbillon-Vey class  $GV \in H^3(B, \mathbb{R}) \cong \check{H}^3(\cdot, \mathbb{R})$  is represented in  $\check{C}(\cdot, \mathbb{R})$  by the cocycle:*

$$\tilde{g}v_1(\sigma_1, \sigma_2, \sigma_3) = \int_0^{\sigma_1(0)} \log(|\sigma_2'(t)|) \frac{\sigma_3''(\sigma_2(t))}{\sigma_3'(\sigma_2(t))} \sigma_2'(t) dt.$$

## 6 Relations to basic cohomology

In the previous sections we have seen various models for the cohomology of the leaf space, all canonically isomorphic. Let us put

$$H^*(M/\mathcal{F}) = H^*(Hol_T(M, \mathcal{F})), \quad H_c^*(M/\mathcal{F}) = H_c^*(Hol_T(M, \mathcal{F})). \quad (20)$$

The reader may choose one of the many models: Haefliger's model (as indicated by the above notations) i.e. 4.2 applied to the holonomy groupoid reduced to any complete transversal  $T$ , the Čech-De Rham model that we have described in section 2 (cf. Proposition 1), or the classifying-space model (cf. Theorem 1). We emphasize however that the last model only works for the cohomology without restriction on the supports!

Here and in the next section we explain why these cohomology theories are suitable theories for the leaf space. We first compare them to the more familiar *basic cohomology* (see e.g. [16, 30]), which is a different cohomology theory for leaf spaces.

**6.1 Basic cohomology.** Choosing a basis  $\mathcal{U}$  of opens of a complete transversal  $T$  (or any transversal basis for  $\mathcal{F}$ ), one defines  $\Omega_{bas}^k(T/\mathcal{F})$  as the cohomology of  $\check{C}^*(\mathcal{U}, \Omega^k)$  in degree  $* = 0$ . This complex consists on  $k$ -forms on  $T$  which are invariant under holonomy, hence it does not depend on the choice of  $T$  (up to canonical isomorphisms, of course). The resulting cohomology is denoted  $H_{bas}^*(M/\mathcal{F})$ . There is an obvious map (induced by an inclusion of complexes)

$$j : H_{bas}^*(M/\mathcal{F}) \longrightarrow H^*(M/\mathcal{F}). \quad (21)$$

Similarly one defines the basic cohomology with compact supports  $H_{c,bas}^*(M/\mathcal{F})$  [16]. The corresponding complex  $\Omega_{c,bas}^k(T/\mathcal{F})$  is the homology of  $\check{C}_c^*(\mathcal{U}, \Omega^k)$  in degree  $* = 0$ , i.e., as in [16], the quotient of  $\bigoplus_{U \in \mathcal{U}} \Omega_c^k(U)$  by the span of elements of type  $\omega - h^* \omega$  ( $h : U \rightarrow V$  is a holonomy embedding, and  $\omega \in \Omega_c^k(V)$ ). Again, there is an obvious map

$$j_c : H_c^*(M/\mathcal{F}) \longrightarrow H_{c,bas}^*(M/\mathcal{F}). \quad (22)$$

In general, the maps (21) and (22) are not isomorphisms. The basic cohomologies are much smaller than  $H^*(M/\mathcal{F})$ ; for instance  $H_{bas}^*(M/\mathcal{F}) = 0$  in degrees  $* > q$ , and they are finite dimensional if  $\mathcal{F}$  is riemannian and  $M$  is compact. The price to pay is the failure of most of the familiar properties from algebraic topology (e.g., as discussed below, Poincaré duality and characteristic classes). However we point out that (21) and (22) are isomorphisms when the naive leaf space is an orbifold. This was explained



in 4.9 of [10], but the reader should think about the similar statement for actions of finite groups on manifolds, and the fact that the cohomology (over  $\mathbb{R}$ ) of finite groups is trivial. In particular (see also [27]), we have

**Proposition 2** *If  $(M, \mathcal{F})$  is a riemannian foliation with compact leaves, then (21) and (22) are isomorphisms.*

Another fundamental property of our cohomologies (20) is

**Proposition 3** (*Poincaré duality*) *For any codimension  $q$  foliation  $(M, \mathcal{F})$ ,*

$$H^*(M/\mathcal{F}; \mathcal{O}) \cong H_c^{q-*}(M/\mathcal{F})^\vee . \tag{23}$$

This (and the more general Verdier duality) has been proved in [10]. Note however that, with the Čech model in hand, the theorem becomes obvious. This new proof of Poincaré duality can be viewed as a rather straightforward extension of the classical proof for manifolds [4] (and can also be interpreted as the obvious duality between the homology and the cohomology of the discrete category  $Emb_U(\mathcal{G})$ , cf. 4.5). In contrast, the basic cohomologies  $H_{bas}^*(M/\mathcal{F})$  and  $H_{c,bas}^*(M/\mathcal{F})$  satisfy Poincaré duality only in the riemannian case [30]. In this case these dualities are compatible via (21) and (22), and they coincide if the leaves are compact (see Proposition 2).

**6.2 Characteristic classes.** As we have seen, one of the main features of  $H^*(M/\mathcal{F})$  is that it contains the characteristic classes of the bundles over the leaf space (i.e. transversal bundles), and the Bott vanishing theorem and the construction of the exotic classes hold at this level. Regarding the groups  $H_{bas}^*(M/\mathcal{F})$ , again, they are too small to contain these characteristic classes. But, as before, this is not seen in the case of riemannian foliations. The reason is that, if  $\mathcal{F}$  is riemannian, then the transversal metric induces a *transversal connection*, i.e. a connection which is invariant under holonomy. Using this type of connections in the construction the characteristic maps  $k_\nu$  of the normal bundle  $\nu$ , we see that  $k_\nu : S(gl_q^*)^{inv} \rightarrow H^*(M/\mathcal{F})$  vanishes in degrees  $> q$ . This stronger vanishing result (at the level of  $H^*(M)$ ), together with the construction of the refined exotic characteristic map, appears in [22]. Moreover, using the explicit constructions of section 3, we see that  $k_\nu$  (and its exotic versions) factors through the basic cohomology groups. This obviously applies to general transversal bundles. In conclusion,

**Proposition 4** *If  $P$  is a transversal principal  $G$ -bundles over  $(M, \mathcal{F})$  which admits a transversal connection then the characteristic map  $k_P : S(\mathfrak{g}^*)^G \rightarrow H^*(M/\mathcal{F})$  of  $P$  (cf. Theorem 2) vanishes in degrees  $> q$ . Moreover the map  $k_P$  (and its exotic version, cf. Corollary 1) factors through the basic cohomology groups:*

$$\begin{array}{ccc}
 S(\mathfrak{g})^G & \cdots\cdots\cdots & H_{bas}^*(M/\mathcal{F}) \\
 \searrow k_P & & \downarrow j \\
 & & H^*(M/\mathcal{F})
 \end{array}$$

**6.3 Integration along the leaves.** Haefliger’s original motivation [16] for introducing  $H_{c,bas}^*(M/\mathcal{F})$  is the existence of an integration over the leaves map  $\int_{\mathcal{F}} : H_c^*(M) \longrightarrow H_{c,bas}^{*-p}(M/\mathcal{F})$  when the bundle of vectors tangent to the leaves is oriented. We want to point out the existence of a refined integration,

$$\int_{\mathcal{F}} : H_c^*(M) \longrightarrow H_c^{*-p}(M/\mathcal{F}) , \quad (24)$$

which, composed with the canonical map (22), gives precisely Haefliger’s integration. Using the Čech model this map becomes obvious: choosing  $\mathcal{U}, \tilde{\mathcal{U}}$  as in 1.2, the integration over the plaques (with the induced orientation),  $\int : \Omega_c^*(U) \longrightarrow \Omega_c^{*-p}(U)$ , induces a map at the level of the Čech-De Rham complexes associated to  $\mathcal{U}$  and  $\tilde{\mathcal{U}}$ .

An alternative abstract definition of  $\int_{\mathcal{F}}$  follows e.g. from the spectral sequences of [10] by standard methods of algebraic topology (“integration over the fiber” as an edge map). The Hochschild-Serre spectral sequence (i.e. Theorem 4.4 of [10] applied to  $\pi : M \longrightarrow M/\mathcal{F}$ ) takes the form  $H_c^s(M/\mathcal{F}; \mathcal{L}^t) \implies H_c^{s+t}(M)$ , where  $\mathcal{L}^t$  is a transversal sheaf whose stalk above a leaf  $L$  is  $H_c^t(\tilde{L})$ . This second description provides us with qualitative information. E.g., if the holonomy covers of the leaves are  $k$ -connected, we find that  $\int_{\mathcal{F}}$  is isomorphism in degrees  $n - k \leq * \leq n$ . Using Poincaré duality, it follows that the pull-back map  $H^*(M/\mathcal{F}) \longrightarrow H^*(M)$  is isomorphism in degrees  $0 \leq * \leq k$ .

## 7 Relations to foliated cohomology

Another standard cohomology theory in foliation theory is the *foliated cohomology* of foliations (see e.g. [1, 19, 20, 29]). In contrast to the other cohomologies that we have seen so far (transversal cohomologies), the foliated cohomology contains a great deal of longitudinal information. In this section we describe its relation to our Čech-De Rham cohomology.

**7.1 Foliated cohomologies.** The foliated cohomology  $H^*(\mathcal{F})$  is defined in analogy with the De Rham cohomology of  $M$ , which we recover if  $\mathcal{F}$  has only one leaf. The defining complex is  $\Omega^*(M, \mathcal{F}) = \cdot, (\Lambda^* \mathcal{F})$ , with the boundary defined by the usual Koszul-formula

$$\begin{aligned} d(\omega)(X_1, \dots, X_{p+1}) &= \sum_{i < j} (-1)^{i+j-1} \omega([X_i, X_j], X_1, \dots, \hat{X}_i, \dots, \hat{X}_j, \dots, X_{p+1}) \\ &+ \sum_{i=1}^{p+1} (-1)^i L_{X_i}(\omega(X_1, \dots, \hat{X}_i, \dots, X_{p+1})). \end{aligned} \quad (25)$$

Here  $L_X(f) = X(f)$ . For later reference, we note the existence of an obvious (restriction to  $\mathcal{F}$ )

$$r : H^*(M) \longrightarrow H^*(\mathcal{F}) \quad (26)$$

There is also a version with compact supports, as well as versions  $H^*(\mathcal{F}; E)$  with coefficients in any transversal (or foliated) vector bundle  $E$ : one uses  $E$ -valued forms on  $\mathcal{F}$ , and one replaces the  $L_{X_i}$  in the previous formula, by the derivatives  $\nabla_{X_i}$  w.r.t. the Koszul connection of  $E$  (see 1.3).

**7.2 Remarks.** In [29], the cohomology  $H^*(\mathcal{F})$  is called “tangential cohomology”, and is denoted  $H^*_\tau(M)$ . The groups  $H^*(\mathcal{F}; \nu)$  with coefficients in the normal bundle (see 1.3) first appeared in [19] in the study of deformations of foliations, while those with coefficients in the exterior powers  $\Lambda\nu$  show up e.g. in the spectral sequence relating the foliated cohomology with De Rham cohomology [1, 20]. The groups  $H^*(\mathcal{F}; E)$  with general coefficients can also be viewed as an instance of algebroid cohomology [23]. Regarding the characteristic classes, since the Bott connection (see 1.3) is flat, it follows that the characteristic classes of the normal bundle are annihilated by  $r$ . This new vanishing result at the level of foliated cohomology produces new (“secondary”) classes,  $u_{4k-1}(\mathcal{F}) \in H^{4k-1}(\mathcal{F})$ . These appear in [15] and have been described in great detail in [11] in the more general context of algebroids. In particular, these new classes come from the cohomology groups  $H^*(M/\mathcal{F}; \Omega^0_{bas})$  (via the map (27) below). Still related to [11], let us mention that if  $\mathcal{F}$  is the foliation induced by a regular Poisson structure on  $M$ , then one has an induced foliated bundle  $K$  (the kernel of the anchor map), and  $H^2(\mathcal{F}; K)$  contains obstructions to the integrability of the Poisson structure.

As explained in [29] in the case of trivial coefficients, and in [19] in the case of the normal bundle as coefficients, the foliated cohomology can be expressed as the cohomology of certain sheaves on  $M$ . For general coefficients  $E$  we consider the sheaf  $\mathcal{H}_\nabla(E)$  described in 1.4. A version of Poincaré’s lemma with parameters shows that  $H^k(\mathcal{F}; E) = 0$  in degrees  $k > 0$  if  $\mathcal{F}$  is the standard  $p$ -dimensional foliation of  $M = \mathbb{R}^p \times \mathbb{R}^q$ . Since always  $\mathcal{H}_\nabla(M; E) = H^0(\mathcal{F}; E)$ , we deduce that  $U \mapsto \Omega^*(\mathcal{F}|_U; E|_U)$  is a flabby resolution of  $\mathcal{H}_\nabla(E)$ , hence

**Proposition 5** *For any foliated vector bundle  $E$  over  $(M, \mathcal{F})$ ,  $H^*(\mathcal{F}; E)$  is isomorphic to  $H^*(M; \mathcal{H}_\nabla(E))$ , the cohomology of  $M$  with coefficients in the sheaf of  $\nabla$ -constant sections of  $E$ . In particular,  $H^*(\mathcal{F}) \cong H^*(M; \Omega^0_{bas})$ . The same holds for compact supports.*

**7.3 Comparison.** We now note the existence of a canonical map

$$\Phi : H^*(M/\mathcal{F}; \Omega^0_{bas}) \longrightarrow H^*(\mathcal{F}) . \quad (27)$$

Recall that  $\Omega^0_{bas}$ , as a sheaf on  $M$ , is the sheaf of smooth function which are constant on the leaves. This map has various interpretations. First of all, it can be viewed as a version with coefficients of the pull-back map (2) (cf. also Proposition 5). Accordingly, the simplest description is in terms of the Čech-De Rham model. Choosing  $\mathcal{U}$  and  $\check{\mathcal{U}}$  as in 1.2, the left hand side of (27) is computed by the cochain complex  $\check{C}^*(\mathcal{U}; C^\infty(U))$ , which is obviously a subcomplex of the  $t = 0$  column of  $\check{C}^s(\check{\mathcal{U}}; \Omega^t(\check{\mathcal{U}}, \mathcal{F}))$ . Now (27) is the map induced in cohomology. Alternatively, at least when the holonomy groupoid is Hausdorff,  $H^*(M/\mathcal{F}; \Omega^0_{bas})$  coincide with the differentiable cohomology [18] of the holonomy groupoid of  $\mathcal{F}$ , and  $\Phi$  is precisely the associated Van Est map described in [31]. It then follows from one of the main results of [11] (applied to the holonomy groupoid) that  $\Phi$  is an isomorphism in degrees  $\leq k$  provided the leaves (or their holonomy covers) are  $k$ -connected. As in the previous section (see 6.3), the same result follows e.g. from the spectral sequences of [10].

**7.4 Integration along the leaves.** If  $\mathcal{F}$  is oriented, then we have an integration map

$$\int_{\mathcal{F}} : H_c^*(\mathcal{F}) \longrightarrow H_c^{*-p}(M/\mathcal{F}; \Omega_{bas}^0) . \quad (28)$$

This map is dual to the Van Est map (27) and can be viewed as a version of the integration map (24) with coefficients in the normal bundle (accordingly, there are similar maps for any transversal vector bundle  $E$  over  $M$ , cf. also 1.4 and Proposition 5). Again, as in the previous section (see 6.3), this map (28) becomes obvious if one uses the Čech-De Rham model.

We want to point out here that the integration over the fibers that we have described clarifies the construction of the Ruelle-Sullivan current of a measured foliation (cf. e.g. Section 3 of [6], or [29] p 126), and also gives new qualitative information about it. Fix a transversal basis  $\mathcal{U}$  for  $\mathcal{F}$ . A smooth transverse measure  $\mu$  is just a measure on each  $U \in \mathcal{U}$ , which is invariant w.r.t. holonomy embeddings. Hence the integration against  $\mu$  is simply a linear map

$$\int_{\mu} : \Omega_{c,bas}^0(M/\mathcal{F}) = H_c^0(M/\mathcal{F}; \Omega_{bas}^0) \longrightarrow \mathbb{R} .$$

Combining this with the integrations along the leaves (24), (28), we can arrange our maps into a diagram (see also (26))

$$\begin{array}{ccc} H_c^p(M) & \xrightarrow{\int_{\mathcal{F}}} & H_c^0(M/\mathcal{F}) \\ r \downarrow & & \downarrow \\ H_c^p(\mathcal{F}) & \xrightarrow{\int_{\mathcal{F}}} & H_c^0(M/\mathcal{F}; \Omega_{bas}^0) \xrightarrow{\int_{\mu}} \mathbb{R} \end{array}$$

The resulting map  $\int_C : H_c^p(M) \longrightarrow \mathbb{R}$  is precisely the integration of [6] against the Ruelle-Sullivan current  $C = C_{\mu}$  (and this defines  $C$  as a degree  $p$  element in the closed homology of  $M$ ). As pointed out in [29],  $C$  actually comes from the closed homology of  $\mathcal{F}$ . In terms of our diagram this simply means that  $\int_C$  factors through  $H_c^p(\mathcal{F})$ .

**7.5 Spectral sequences.** Almost all of the maps that we have described in the last two sections figure in certain the spectral sequences. First of all, the filtration on  $\Omega^*(M)$  induced by  $\mathcal{F}$  (cf. e.g. [1, 20]) induces a spectral sequence

$$E_1^{s,t} = H^s(\mathcal{F}; \Lambda^t \nu) \implies H^{s+t}(M) .$$

Similarly, the filtration of the Čech-De Rham double complex induces a spectral sequence

$$\bar{E}_1^{s,t} = H^s(M/\mathcal{F}; \Omega_{bas}^t) \implies H^{s+t}(M/\mathcal{F}) .$$

Note that  $E_2^{0,t} = H_{bas}^t(M/\mathcal{F})$ . These two spectral sequences are related by the pull-back map (2), and by the Van Est map (27) with coefficients,  $\Phi : H^s(M/\mathcal{F}; \Omega^t) \longrightarrow H^s(\mathcal{F}; \Lambda^t \nu)$ . With the same arguments as above, these maps are isomorphisms in degrees  $0 \leq s \leq k$ , if the holonomy covers of the leaves are  $k$ -connected. The version with compact supports of this discussion involves (24) and the integrations  $\int_{\mathcal{F}} : H_c^s(\mathcal{F}; \Lambda^t \nu) \longrightarrow H_c^{s-p}(M/\mathcal{F}; \Omega_{bas}^t)$  (cf. 7.4 above).

## References

- [1] JESÚS A. ALVAREZ LÓPEZ, *A decomposition theorem for the spectral sequence of Lie foliations*, Trans. of the A.M.S., Volume **329**, January 1992, 173- 184
- [2] R. BOTT, *Lectures on characteristic classes and foliations*, Springer LNM **279**, 1-94
- [3] R. BOTT, *On the characteristic classes of groups of diffeomorphisms*, L'Enseignement mathématique **XXIII** fasc. 3-4, 209- 220
- [4] R. BOTT and L.W. TU, *Differential forms in algebraic topology*, Graduate Texts in Mathematics 82, Springer (1982/1995)
- [5] H. CARTAN, *Notions d'algèbre différentielle; applications aux groupes de Lie etc.* In "Colloque de Topologie", C.B.R.M. Bruxelles, (1950), 15-27
- [6] A. CONNES, *A survey of foliations and operator algebras*, Proc. Sympos. Pure Math., AMS Providence, **32** (1982), 521-628
- [7] A. CONNES, *Noncommutative Geometry*, Academic Press (1994)
- [8] M. CRAINIC, *Cyclic homology of étale groupoids; The general case*, K-Theory **17** (1999), 319- 362
- [9] M. CRAINIC, PhD Thesis, Utrecht University, April 2 000
- [10] M. CRAINIC and I. MOERDIJK, *A homology theory for étale groupoids*, J. Reine Angew. Math. 521 (2000), 25-46
- [11] M. CRAINIC, *Differentiable and algebroid cohomology, van Est isomorphisms, and characteristic classes*, Preprint 2000
- [12] I.M. DUFLO, M. VERGNE, *Cohomologie équivariante et descente*, Astérisque **215**, (1993)
- [13] J. DUPONT, *Simplicial De Rham cohomology and characteristic classes of flat bundles*, Topology **15** (1976), pp 233-245
- [14] C. GODBILLON, *Cohomologies d'algèbre de Lie de champs de vecteurs formels*, Seminaire Borbaki, LNM **421** (1972), 1-19
- [15] R. GOLDMAN, *The holonomy ring on the leaves of foliated manifolds*, J. Differential Geometry **11** (1976), pp 411-449
- [16] A. HAEFLIGER, *Some remarks on foliations with minimal leaves*, J. Differential Geometry, **15** (1980), 269-284
- [17] A. HAEFLIGER, *Groupoides d'holonomie et espaces classifiants*, Astérisque **116** (1984), 70-97
- [18] A. HAEFLIGER, *Differentiable cohomology*, "Differential Topologi (CIME, Varenna 1976) Ligouri, Naples", 1979, 19-70

- [19] J. HEITSCH, *A cohomology for foliated manifolds*, Bull. Amer. Math. Soc. **79** (1973), 1283-1285
- [20] F. KAMBER and P. TONDEUR, *Foliations and metrics*, Progress in Mathematics **32**, Differential Geometry (1983), 103-152
- [21] F. KAMBER and P. TONDEUR, *Foliated Bundles and Characteristic Classes*, Springer LNM **493**
- [22] C. LAZAROV and J. PASTERNAK, *Secondary characteristic classes for riemannian foliations*, J. Differential Geometry **11** (1976), pp 365- 385
- [23] K. MACKENZIE, *Lie groupoids and Lie algebroids in differential geometry*, Lecture Notes Series **124**, London Mathematical Society, 1987
- [24] I. MOERDIJK, *Classifying topos and foliations*, Ann. Inst. Fourier. Grenoble **41** (1991), 189-209
- [25] I. MOERDIJK, *On the weak homotopy type of étale groupoids*, Integrable Systems and Foliations, Albert, Brouzet, Dufour(eds.), Birkhäuser, 1997
- [26] I. MOERDIJK, *Proof of a conjecture of A. Haefliger*, Topology **37** (1998), 735-741
- [27] P. MOLINO, *Riemannian foliations*, Progress in Mathematics 73, Birkhauser, Boston, 1988
- [28] J. MRCUN, *Functoriality of the Bimodule Associated to a Hilsum-Skandalis Map*, K-theory **18** (1999), 235- 253
- [29] MOORE and SCHOCHET, *Global Analysis on Foliated Spaces*, Springer- Verlag, 1988
- [30] V. SERGIESCU, *Cohomologie basique et dualité des feuilletages riemanniens*, Ann. Inst. Fourier, Grenoble **35** 3, pp 137-158 (1985)
- [31] A. WEINSTEIN and P. XU, *Extensions of symplectic groupoids and quantization*, J. Reine Angew. Math. **417** (1991), 159-189
- [32] H. WINKELNKEMPER, *The graph of a foliation*, Ann. Global Anal. Geom. **1** (1983), 51-75