



Global Biogeochemical Cycles

RESEARCH ARTICLE

10.1002/2015GB005241

Key Points:

- Long-term trends in fossil foraminiferal species are in line with fluctuations in Mg/Casw
- Foraminifera producing a favored biomineral are more likely to recovery after extinction events
- Currently, all low-Mg calcite-producing foraminifera calcify in unfavorable seawater Mg/Ca

Supporting Information:

- Texts S1–S3, Table S1, and Figure S1 Caption
- Figure S1

Correspondence to:

I. van Dijk,
Inge.van.Dijk@nioz.nl

Citation:

van Dijk, I., L. J. de Nooijer, M. B. Hart, and G.-J. Reichart (2016), The long-term impact of magnesium in seawater on foraminiferal mineralogy: Mechanism and consequences, *Global Biogeochem. Cycles*, 30, 438–446, doi:10.1002/2015GB005241.

Received 8 JUL 2015

Accepted 18 FEB 2016

Accepted article online 22 FEB 2016

Published online 8 MAR 2016

The long-term impact of magnesium in seawater on foraminiferal mineralogy: Mechanism and consequences

I. van Dijk¹, L. J. de Nooijer¹, M. B. Hart², and G.-J. Reichart^{1,3}

¹Department of Ocean Systems, NIOZ-Royal Netherlands Institute for Sea Research and Utrecht University, Den Burg, Netherlands, ²School of Geography, Earth and Environmental Sciences, Plymouth University, Plymouth, UK, ³Faculty of Geosciences, Earth Sciences Department, Utrecht University, Utrecht, Netherlands

Abstract Foraminifera are unicellular protists, primarily known for their calcium carbonate shells that provide an extensive fossil record. This record, ranging from Cambrian to present shows both major shifts and gradual changes in the relative occurrence of taxa producing different polymorphs of carbonate. Here we present evidence for coupling between shifts in calcite- versus aragonite-producing species and periods with, respectively, low and high seawater Mg/Ca throughout the Phanerozoic. During periods when seawater Mg/Ca is <2 mol/mol, low-Mg calcite-producing species dominate the foraminiferal community. Vice versa, high-Mg calcite- and aragonite-producing species are more abundant during periods with relatively high seawater Mg/Ca. This alteration in dominance of the phase precipitated is due to selective recovery of groups producing the favorable polymorph after shifts from calcite to aragonite seas. In addition, relatively high extinction rates of species producing the mineral phase not favored by the seawater Mg/Ca of that time may be responsible for this alteration. These results imply that the current high seawater Mg/Ca will, in the long term, favor prevalence of high-Mg and aragonite-producing foraminifera over calcite-producing taxa, possibly shifting the balance toward a community in which calcite production is less dominant.

1. Introduction

In 1983, Sandberg proposed that the mineral phase of inorganic carbonates that precipitated from seawater reflects Phanerozoic seawater chemistry [Sandberg, 1983]. Shifts between so-called aragonite and calcite seas are primarily caused by long-term changes in the ratio of $[Mg^{2+}]$ over $[Ca^{2+}]$ in seawater [Dickson, 2002; Hardie, 1996; Lowenstein et al., 2001] (i.e., Mg/Ca_{sw}). This ratio varies over geological timescales due to changes in the balance between weathering, oceanic crust production, dolomite formation, and rate of carbonate burial [Arvidson et al., 2006, 2011; Berner, 2004; Farkaš et al., 2007; Hardie, 1996; Holland, 2005; Wilkinson and Algeo, 1989]. Dominance of organisms producing calcite (e.g., rugose and tabulate corals) or aragonite (e.g., corals and dasycladacean algae) as well as abiotically precipitated calcite and aragonite follows trends in Mg/Ca_{sw} [Stanley, 2006; Stanley and Hardie, 1998]. The impact of Mg/Ca_{sw} on biogenic mineralogy can be explained by differences in affinity of Mg ions for calcite versus aragonite crystal surfaces [De Choudens-Sánchez and González, 2009; Fallini et al., 1994; Fernández-Díaz et al., 1996; Nancollas and Sawada, 1982; Reddy and Wang, 1980]. Magnesium ions adsorb relatively well to calcite surfaces compared to those of aragonite [Mucci and Morse, 1985], resulting in a relative decrease in calcite growth rate at high seawater $[Mg^{2+}]$ [Berner, 1975]. The boundary between nucleation fields for inorganic aragonite and calcite lies at a Mg/Ca_{sw} of 2 mol/mol [Sun et al., 2015] and depends on temperature and pressure [Berner, 1975]. Recently, it is shown that seawater sulfate concentrations have an (additional) effect on the seawater Mg/Ca threshold [Bots et al., 2011]. Periods with $Mg/Ca_{sw} < 2$ (e.g., Devonian and Silurian) are associated with calcite-producing organisms and are therefore termed “calcite seas,” whereas “aragonite seas” are defined by periods with $Mg/Ca_{sw} > 2$ and are characterized by dominance of aragonite and/or high-magnesium calcite producers [Dickson, 2004; Stanley, 2008] (e.g., Permian and Triassic). The biological mechanisms affected by changes in Mg/Ca_{sw} may include decreasing growth rates and increased energy costs for calcification by organisms producing an unfavored $CaCO_3$ crystal phase [Ries, 2010]. Both of these biological impediments result in a dominance of calcite-producing organisms during periods with low Mg/Ca_{sw} , while favoring those precipitating aragonite when Mg/Ca_{sw} is high.

Although the relationship between Phanerozoic changes in seawater Mg/Ca and the evolutionary history of many marine calcifiers has been analyzed [Kiessling et al., 2008; Porter, 2010; Stanley, 2008], the response of

foraminifera to changes in $\text{Mg}/\text{Ca}_{\text{sw}}$ is only sparsely investigated [Martin, 1995]. Most low-Mg calcite-producing or hyaline taxa are known to actively control the amount of magnesium incorporated [Bentov and Erez, 2006; De Nooijer et al., 2014]. Since the Mg/Ca of the calcifying fluid is, in this case, dissimilar from that of ambient seawater, foraminifera might potentially be unaffected by changes in $\text{Mg}/\text{Ca}_{\text{sw}}$. High concentrations of seawater Mg^{2+} , however, may require allocation of more energy to maintain biomineralization and thereby link long-term changes in $\text{Mg}/\text{Ca}_{\text{sw}}$ to the success of aragonite- versus calcite-producing foraminifera. Geological longevity, areal and depth distribution, composition of the tests (aragonite or calcite), and importance in the global carbon cycle, however, make foraminifera particularly suitable to study the interplay between seawater chemistry and biogenic mineralogy. Here we present an analysis of foraminiferal abundances through the Phanerozoic and relate their mineralogy to changes in $\text{Mg}/\text{Ca}_{\text{sw}}$ and mass extinction events. Past shifts in dominance of calcite- versus aragonite-producing foraminifera serve as an analogue to evaluate future foraminiferal success under ongoing climate and environmental stress.

2. Methods

2.1. Foraminiferal Occurrences Over Geological Timescales

In order to investigate major trends in foraminiferal distribution, we analyzed foraminiferal occurrences in the Paleobiology database of Fossilworks (<http://fossilworks.org>). In general, we used the database in default mode, with some small exceptions, and this input data are listed in the supporting information. In short, we extracted 6270 collections with 37,586 species occurrences covering the last 500 Ma [Kiessling et al., 2015], when queried the database for *Foraminifera*, *Foraminiferida*, and *Foraminifera* (genus level or above). All foraminiferal species occurrences were grouped into ~5 Myr time bins ("Fossil Record 2 bin"). Occurrences were extracted from the database on a species level, excluding contributions with open nomenclature (aff., cf., "...," and "?") [Bengston, 1988] which take into account different levels of uncertainty during the determination on species level. To estimate global foraminiferal abundances, the full geographical range was used. By enabling the database's "scale option" we also extracted composition and life habitat information of every species. The assignments of these mineralogical and ecological attributes were based on a number of sources reporting mineralogy and habitat preferences for the different foraminiferal (sub)orders [Loeblich and Tappan, 1984, 1988]. The composition of the tests was aragonite or calcite with low (LMC; <8%wt MgCO_3) and high magnesium content (HMC; >8%wt MgCO_3). So-called intermediate-Mg calcite (4–8%wt MgCO_3) was already labeled LMC by the authors of the database. This distinction coincides with known differences in biomineralization strategies (e.g., hyaline versus porcelaneous foraminifera, respectively [De Nooijer et al., 2009; Hemleben et al., 1986]). Life habitat information was either planktonic or benthic (infaunal, semi-infaunal, and epifaunal). For 10.5% of all species present in the database, the CaCO_3 mineralogy is unknown or not recorded and was therefore excluded from further statistical analysis. Also, species which do not (actively) calcify, like agglutinated and naked foraminifera, or precipitate carbonate but opaline silica [Gupta, 2002] (order Silicolocinida), for example, were not used for this analysis. Occurrences of species from collections older than 320 Ma were excluded from analysis due to scarcity of collections and consequently very low foraminiferal species occurrences (<25) per time bin. Expression of Fossilworks-derived abundances as proportional occurrences circumvents potential biases introduced when using absolute abundances [Valentine et al., 2013]. The resulting four groups (HMC, aragonite, LMC benthic, and LMC planktonic) generally follow the "classic" orders [Loeblich and Tappan, 1988]. However, there are some exceptions. For instance, the "aragonite" group comprises the orders Involutinida and Robertinida and also includes the aragonitic taxon Favusellacea, which is actually part of the calcite-producing order Globigerinida, according to Loeblich and Tappan [1988]. Another example is the aragonite precipitating *Hoegludina elegans* from the low-Mg calcite-producing order Rotaliina, which is also placed in the aragonite group in our study. Because of such exceptions, we decided not to generalize Fossilworks to solely looking at the species occurrence of different orders. However, our assigned groups still overlap strongly with the orders. Diversity graphs were created for these four compiled groups by using the diversity graph function of FW. As a result, the compiled data provide an assessment of rational occurrences at species level and thereby indicate the dominance or rareness of the four groups relative to each other. In some time bins (e.g., around ~70 and 100 Ma), planktonic foraminifera make up > 90% of the total occurrences. This could be due to a possible bias of Fossilworks toward planktonic foraminifera. However, when performing the analysis without the planktonic LMC group, we still observe the same general pattern for all the other groups (see supporting information).

2.2. Aragonite Versus Calcite Sea Events

Since different $\text{Mg}/\text{Ca}_{\text{sw}}$ reconstructions yield slightly different timings of calcite-aragonite sea events [Evans and Müller, 2012], we choose to define periods in which reconstructions [Farkaš et al., 2007; Stanley and Hardie, 1998] agree as the “true” calcite and aragonite sea events. To test whether occurrences of LMC versus aragonite/HMC producers are significantly different in these calcite and aragonite seas or before/after mass extinction events [Raup and Sepkoski, 1982], two-way Student’s t tests assuming equal variances were performed. We compared the average $p(\text{aragonite} + \text{HMC})$ between adjacent calcite and aragonite seas, as well as between all calcite and aragonite sea intervals combined. Finally, the three bins before and after three mass extinction events with student’s t tests to determine whether these average occurrences were different. Our analyses include the end-Permian (252 Ma), end-Triassic (200 Ma), end-Cretaceous (65 Ma), and Paleocene Eocene Thermal Maximum (PETM; ~55 Ma) mass extinctions. Due to the resolution of our database, the end Cretaceous and PETM were regarded as one event, despite known differences in their causes and consequences. We in this case compared $p(\text{aragonite} + \text{HMC})$ of the three time bins before the end Cretaceous (65 Ma) with the three time bins after the PETM (55 Ma). All of the aforementioned events are associated with minor or major perturbations in marine inorganic chemistry [Hart et al., 2014; Hönisch et al., 2012] and are characterized by a decrease in biodiversity, including that of foraminifera.

3. Results

3.1. Seawater Mg/Ca Reconstruction

Long-term patterns in $\text{Mg}/\text{Ca}_{\text{sw}}$ can be reconstructed from model studies taking into account mid-ocean ridge spreading rates, weathering, and dolomite formation [Hardie, 1996; Wilkinson and Algeo, 1989]. Over the last few decades, these reconstructions have been improved by, e.g., new insights in biochemical cycles. One of the most recent reconstructions [Farkaš et al., 2007] (Figure 1) combines a model of coupled calcium, carbon, and magnesium global cycles [Hansen and Wallmann, 2003; Heuser et al., 2005; Wallmann, 2001, 2004] with reconstructed seafloor spreading rates based on $^{87}\text{Sr}/^{86}\text{Sr}$ [Veizer et al., 1999; Wallmann, 2004]. The resulting reconstruction is in close agreement with a number of independent proxies for $\text{Mg}/\text{Ca}_{\text{sw}}$, including Mg/Ca of fluid inclusions [Horita et al., 2002; Lowenstein et al., 2001] and that of rudists [Steuber and Rauch, 2005] and echinoderms [Dickson, 2002]. This relatively recent $\text{Mg}/\text{Ca}_{\text{sw}}$ reconstruction is largely in agreement with those from earlier studies [Hardie, 1996], whereas deviations between models is likely caused by recent updates in the assumptions for the earlier models [Holland et al., 1996; Holland and Zimmermann, 2000; Horita et al., 2002]. The model by Farkaš et al. [2007] also includes more recent data on global major elements cycling and refined estimates of seafloor spreading rates, resulting in slightly different $[\text{Ca}^{2+}]$ and $[\text{Mg}^{2+}]$ trends. The transitions between these calcite and aragonite seas deviate somewhat from those of the often-used reconstruction [Stanley and Hardie, 1998], which is therefore included in our discussion. This divided the past 500 Ma into two periods with a relatively low $\text{Mg}/\text{Ca}_{\text{sw}}$ (end Cambrian to Devonian and end Triassic to mid-Cretaceous, CI and CII, respectively) and two with a relatively high $\text{Mg}/\text{Ca}_{\text{sw}}$ (Carboniferous to end Triassic and mid-Cretaceous to recent, AII and AIII, respectively).

3.2. The Response of Foraminifera to Changes in $\text{Mg}/\text{Ca}_{\text{sw}}$

Although it has been suggested that foraminifera are “sophisticated biomineralizers” and are, therefore, able to cope with relatively variable Mg concentrations in seawater [Stanley, 2008], there is a significant ($p < 0.01$) correlation between average seawater Mg/Ca of combined aragonite versus calcite sea events (CII versus AII + AIII) and the ratio between calcite- versus aragonite-producing foraminifera (Figure 2). Aragonite sea species (taxa-producing aragonite or HMC) are more abundant during periods with high seawater Mg/Ca values. In periods with low $\text{Mg}/\text{Ca}_{\text{sw}}$ values, there is an increase in abundance of LMC-producing taxa. However, after the shift from CII to AIII, there appears to be a lag in the response of the foraminiferal community. Aragonite sea species only start to increase after the end Cretaceous and PETM, not directly when $\text{Mg}/\text{Ca}_{\text{sw}}$ increases. Nevertheless, the effect of mass extinction events on the dominating foraminiferal mineralogy is less clear (Table 1).

Only the end-Triassic extinction, coinciding with a transition from aragonite to calcite sea interval (AII/CII), correlates to a significant ($p < 0.05$) shift in abundance from aragonite to calcite sea species. Furthermore, first appearance of aragonite precipitating foraminifera, i.e., the orders Involutinida and Robertinida, occurs




Figure 1. (top) Reconstruction of calcite (CI and CII) and aragonite sea (AII and AIII) intervals based on modeled seawater Mg/Ca reconstructions (solid line [Stanley and Hardie, 1998] and dashed line [Farkaš et al., 2007]), supported by Mg/Ca from fluid inclusions (bars [Lowenstein et al., 2001] and pentagons [Horita et al., 2002]) and that of echinoderm ossicles and rudists (arrows [Steuber and Rauch, 2005] and crosses [Dickson, 2002]). The horizontal line divides calcite sea ($\text{Mg}/\text{Ca} < 2$) and aragonite sea ($\text{Mg}/\text{Ca} > 2$) intervals [Berner, 2004; Stanley, 2008]. Accordingly, aragonite sea intervals are indicated by blue panels and calcite sea events with red panels. The grey panels indicate periods where the two models deviate in the timing of the division between calcite and aragonite seas. (bottom) Phanerozoic reconstruction of SO_4^{2-} (dotted line) and Ca^{2+} (solid line) concentrations over geological time [after Demicco et al., 2005]. Cm = Cambrian, O = Ordovician, S = Silurian, D = Devonian, Ca = Carboniferous, P = Permian; Tr = Triassic, J = Jurassic, C = Cretaceous, Pg = Palaeogene, and N = Neogene.




Figure 2. Proportional foraminiferal species occurrences through the Phanerozoic Era. The $p(\text{occurrences})$ is the number of times that a species of one of the four groups is present per time bin, relative to the other three groups. Low-Mg calcite-producing (LMC) benthic and planktonic foraminifera (light and dark red areas, respectively) and of high-Mg calcite (HMC) and aragonite foraminifera (light and dark blue areas, respectively) are divided by a dashed black line. The three mass extinction events (P-Tr, Tr-J, PETM, and C-Pg) are indicated by vertical dotted lines. Light shaded data before 320 Ma are not statistically supported due to a limited amount of foraminiferal occurrences in this part of the fossil record. Abbreviations as in Figure 1.

Table 1. Changes in Aragonite Versus Calcite Sea Species Over Mass Extinction Events and Aragonite/Calcite Sea Intervals^a

Transition or Event	Δ mean	t Value	d.f.	p
C/A ^b	0.394	9.85	20	<0.00
CII/AIII	0.306	4.91	10	<0.00
AII/CII	0.483	7.39	10	<0.00
End Permian (252 Ma)	0.310	3.39	2	0.08
End Triassic (200 Ma)	0.444	7.63	2	0.02
End Cretaceous (65 Ma) and PETM (55)	0.411	3.07	2	0.09

^aBold values are significant. Aragonite (AII and AIII) and calcite (CII) sea intervals. Δ mean is the difference between the means of the compared groups.

^bAll Calcite sea periods versus all aragonite sea periods = CII/(AII + AIII).

during aragonite sea II (approximately 250 Ma ago), while planktonic calcite precipitating foraminifera evolve during calcite sea II.

4. Discussion and Conclusion

4.1. Foraminiferal Trends Are Controlled by Seawater Mg/Ca

Long-term trends in fossil foraminiferal occurrences are in line with fluctuations in Mg/Ca_{sw} and community changes of other marine calcifiers

[Stanley, 2008; Stanley and Hardie, 1998; Porter, 2010]. However, $[\text{SO}_4^{2-}]$ has an additional effect on preferential CaCO_3 polymorph formation due to its effect on the Mg/Ca threshold separating the nucleation fields for aragonite and calcite. Especially when $[\text{SO}_4^{2-}]$ decreases below <10 mM [Bots et al., 2011], the Mg/Ca threshold for preferential aragonite increases, resulting in an overall increase of calcite precipitated. Hence, high $[\text{SO}_4^{2-}]$ promotes the precipitation of aragonite, similar to high seawater Mg/Ca. Since Phanerozoic $[\text{SO}_4^{2-}]$ reconstructions [Demichio et al., 2005] show a covariation of $[\text{SO}_4^{2-}]$ with seawater Mg/Ca, it is difficult to disentangle the contribution of these two controls. During biomineralization, however, the $[\text{SO}_4^{2-}]$ of the calcifying fluid of a foraminifera is much lower than that of the surrounding seawater, as indicated by the low partition coefficient for sulfur [Berry, 1998; I. van Dijk et al., manuscript in preparation, 2016]. This implies that despite geological fluctuations in $[\text{SO}_4^{2-}]$, its effect on the boundary between nucleation fields for calcite versus aragonite precipitation is similar through time.

Nevertheless, periods with relatively low Mg/Ca_{sw} are dominated by calcite-producing foraminifera, whereas aragonite producers are found predominantly during periods with high Mg/Ca_{sw}. For other organisms, shifts in dominance of marine calcite and aragonite producers are explained by increased extinction and speciation rates after transitions of Mg/Ca_{sw}, or by adaptation to preferred mineralogy by switching from calcite to aragonite precipitation or vice versa [Ries, 2004, 2010; Stanley et al., 2002]. For foraminifera, extinction and radiation seem more likely since there are significant differences in community composition before and after mass extinction events. Moreover, no foraminiferal species has yet been found to produce both calcium carbonate phases, nor have culturing experiments with varying Mg/Ca_{sw} [Evans et al., 2015; Mewes et al., 2014; Segev and Erez, 2006] suggested that foraminifera can change or adapt to another CaCO_3 phase, kinetically favored, e.g., coralline algae [Stanley et al., 2002] and echinoids, crabs, shrimps, and calcareous serpulid worms [Ries, 2004]. Since foraminifera do not switch from calcite to aragonite production (or vice versa) as a response to changes in Mg/Ca_{sw}, they will eventually be calcifying at unfavorable Mg/Ca_{sw} conditions as a consequence of changing ocean chemistry. For other organisms, this is shown to negatively impact calcification through decreased growth rates and/or increased energy spent on calcification [Ries, 2010]. Hyaline and porcelainous foraminifera have different calcification strategies to produce their shells [De Nooijer et al., 2009, 2014; Hemleben et al., 1986]. The costs involved in calcification are poorly constrained but will depend on biomineralization strategy [Zeebe and Sanyal, 2002]. In case Mg^{2+} is selectively removed from the calcifying fluid [Bentov and Erez, 2006], increased seawater $[\text{Mg}^{2+}]$ will likely increase energy requirements for calcification due to increased pumping rates. When calcification is driven by active, transmembrane transport of Ca^{2+} [Nehrke et al., 2013; De Nooijer et al., 2014], the involved energy may also increase by Mg ions blocking selective Ca transporters. The latter may be overcome by, e.g., increasing the Ca transporter density, which will also increase the energy requirements when calcifying under high $[\text{Mg}^{2+}]$.

4.2. Preferential Dissolution

Geological trends in fossil foraminiferal shells are potentially biased by preferential dissolution as a result of changes in seawater saturation state with respect to calcium carbonate. Saturation state (Ω) is the product of seawater carbonate and calcium concentrations ($[\text{Ca}^{2+}] \times [\text{CO}_3^{2-}]$), divided by the solubility product K_{sp} . This solubility product differs between aragonite ($-\log K_{sp} = 6.19$) and low-Mg calcite ($-\log K_{sp} = 6.37$) and therefore results in relatively poor preservation of aragonite shells compared to calcite ones. Such a preservation bias could have affected fossil occurrences of the different foraminiferal groups, thereby compromising our

results, discussion, and conclusions. Although preferential dissolution may have affected foraminiferal taphonomy, it is unlikely to impact overall shifts in dominance of calcite versus aragonite producers since reconstructed seawater saturation states [Riding and Liang, 2005] are anticorrelated to changes in Mg/Ca_{sw} . Low Ω states co-occur with high Mg/Ca_{sw} on global and geological scales and vice versa. Even though periods with a low Ω may be reflected by increased aragonite dissolution and therefore decreased preservation, the opposite is found (Figure 2). Hence, the effect of elevated Mg/Ca_{sw} on aragonite production apparently outweighs any potential bias due to preferential dissolution. Depending on the magnitude of increased aragonite dissolution during periods with a low Ω , the observed correlation between Mg/Ca_{sw} and calcite versus aragonite foraminiferal species abundances might be stronger than reported here. However, we cannot account for postburial diagenetic change, which may also cause the preferential dissolution of aragonite and/or replacement by calcite.

4.3. The Effect of Mass Extinction Events

The importance of mass extinction events on the balance between calcite and aragonite producers, in general, has been analyzed by Kiessling *et al.* [2008], who proposed that major environmental perturbations rather than changes in ocean chemistry determine the dominance of calcite versus aragonite producers. This conclusion is based on a study also relying on fossil abundances and included a group “foraminifera + algae,” representing a minor portion of the total species occurrences. In Kiessling *et al.*’s analysis, trends in the combination of all marine-calcifying organisms are mainly driven by mass extinction events, while Mg/Ca_{sw} seems to have a minor impact. This is seemingly in contrast with our results, which suggest the relation between long-term seawater chemical changes and evolution of calcifying organisms is not always straightforward. An explanation for these contrasting results may point to the fact that different controls shape the evolution of marine calcifiers on different levels and/or that each group (e.g., corals, bivalves, and foraminifera) responds differently to changes in Mg/Ca_{sw} . It is therefore crucial to analyze the evolution of (and within) marine-calcifying groups in relation to seawater chemistry separately. It is likely that mass extinction events impact the mineralogy of marine calcifiers, in general, while within these orders, changes in seawater chemistry influence evolutionary success and extinction rates.

When analyzing foraminiferal community shifts as a function of mass extinction events, only the end-Triassic event shows a significant shift from a high-Mg and aragonite-dominated community to one with more low-Mg calcite-producing species. Due to differences in the causes, duration, and magnitude of all events analyzed, in combination with the resolution of our data, the impact of mass extinctions on overall foraminiferal community cannot be unambiguously assessed based on this data set. Nevertheless, for other major groups of calcifying organisms, mass extinction events play an important role in aragonite versus calcite dominance [Kiessling *et al.*, 2008] mainly due to selective recovery of the organisms which precipitated the favored polymorph, which may also be the case for foraminifera.

4.4. First Occurrence, Extinction, and Recovery of Foraminiferal Taxa

The effect of ocean chemistry on foraminiferal evolution is likely caused by (1) timing of the first occurrences of newly formed taxa and (2) selective recoveries and extinctions of taxa before and after Mg/Ca_{sw} transitions and/or major carbon perturbations. This would explain, for instance, the observed lag of dominance by HMC and aragonite producers after the CII/AIII transition.

The importance of Mg/Ca_{sw} on foraminiferal mineralogy is reflected by the first occurrences of newly evolved taxa, generally exhibiting the mineralogy which is favored by the seawater Mg/Ca of that time. For example, all taxa from the orders Involutinida and Robertinida produce aragonite shells and both orders originated in Aragonite sea II (at 303 and 250 Ma, respectively). After establishing their mineralogy, these taxa did not change their mineralogical preference within their calcification mechanism and continued producing aragonite, even during periods of lower Mg/Ca_{sw} . This was already suggested by Martin [1995] that Mg/Ca_{sw} plays a key role in determining the mineralogy of clades at their moment of origination [Porter, 2007, 2010].

There might be an increased recovery or diversification of taxa producing the polymorph which is supported by seawater Mg/Ca . For instance, high Mg/Ca_{sw} could have played a part in the postextinction recovery by aragonitic trochospiral Robertinida (genera *Oberhauserella* and *Reinholdella*) following the end-Triassic extinction [Cl  mence and Hart, 2013; Fuchs, 1967, 1970], which occurred during relatively high Mg/Ca_{sw} . In the Lower Jurassic, calcitic taxa (*Lenticulina*, *Lingulina*, *Paralingulina*, *Nodosaria*, *Pseudonodosaria*, etc.)

gradually began to dominate the assemblage in the developing calcite II ocean [Copestake and Johnson, 2014], with the aragonitic taxa (especially *Reinholdella* and *Epistomina*) reduced to a low-diversity lineage throughout the mid–Late Jurassic and Early Cretaceous [Hart, 1984; Hart et al., 2009; Oxford et al., 2004].

After the end-Cretaceous extinction event, occurrences of high-Mg foraminiferal taxa increased (Figure 2) and aragonitic pteropods and heteropods became a significant part of the oceanic plankton [Wall-Palmer et al., 2012] while seawater Mg/Ca rose to its current value.

An example of the interplay between the mineralogy of newly originated taxa and selective recovery by subsequent changes in oceanic state can be found in the evolution of planktonic foraminifera. The early planktonic foraminifera, *Conoglobigerina*, were aragonite producers and evolved from aragonitic benthic ancestors in the high Mg/Ca_{sw} ocean of the Early Jurassic [Hart et al., 2003, 2012; Wernli, 1988, 1995]. This *Conoglobigerina*–*Globuligerina*–*Favusella* lineage appeared to be unsuccessful and did not diversify in the developing calcite II ocean and then became extinct in the mid-Cretaceous, after which the new calcite-producing taxa became highly diverse and successful. These new calcite-producing planktonic foraminifera emerged in the Early Cretaceous, and their evolution can be traced throughout the Cretaceous, despite a number of significant changes in the foraminiferal community at oceanic anoxic events [BouDagher-Fadel et al., 1997; Hart, 1999; Hart et al., 2002; Premoli Silva and Sliter, 1999].

4.5. Consequences for (Future) Foraminifera

Currently, high seawater Mg/Ca favors occurrence of organisms producing aragonite and high-Mg calcite [e.g., Stanley, 2006; Stanley and Hardie, 1998]. Following fossil alterations in dominant foraminiferal mineral phases, all extant low-Mg calcite-producing foraminifera, including all planktonic species, are currently living in an ocean that is not favorable to the mineral phase of their shells. Ongoing anthropogenic CO₂ emissions and subsequent ocean acidification are currently resulting in a major marine carbon perturbation by decreasing seawater pH and Ω , which will likewise hamper marine calcification [Van de Waal et al., 2013; Ilyina and Zeebe, 2012] as it did in the geological past [e.g., Hönisch et al., 2012]. Since ongoing oceanic CO₂ uptake will cause seawater to become undersaturated with respect to aragonite sooner than it will be for calcite, organisms producing aragonite and high-Mg calcite (scleractinian corals, echinoderms, and some foraminiferal taxa) are generally thought to be most vulnerable to ocean acidification [Orr et al., 2005]. However, foraminiferal species producing this polymorph of carbonate (i.e., aragonite or HMC) are supported by the current high seawater Mg/Ca. Even though they may be relatively vulnerable to ongoing ocean acidification, subsequent recovery and diversification may favor their long-term success. Our results highlight the potential role of seawater Mg/Ca as a possible controller on the success of calcite versus aragonite producers in a changing ocean. We suggest that on longer timescales, high-Mg calcite- and aragonite-producing foraminifera dominance will not be negatively impacted by ongoing ocean acidification due to rapid recovery and diversification after the event, which will be perhaps even at the expense of low-Mg calcite-producing foraminifera.

Acknowledgments

This research is funded by the NIOZ–Royal Netherlands Institute for Sea Research and the Darwin Centre for Biogeosciences project “Double Trouble: Consequences of Ocean Acidification—Past, Present and Future—Evolutionary changes in calcification mechanisms.” The data set (“Taxonomic occurrences of to 600 Foraminifera, Foraminiferida, and Foraminifera recorded in Fossilworks, the Evolution of Terrestrial Ecosystems database, and the Paleobiology Database”) used in this study is freely available at <http://fossilworks.org>. The input parameters necessary to reproduce the results are also available in the supporting information. This is Paleobiology Database publication.

References

- Arvidson, R. S., F. T. Mackenzie, and M. Guidry (2006), MAGic: A Phanerozoic model for the geochemical cycling of major rock-forming components, *Am. J. Sci.*, 306(3), 135–190.
- Arvidson, R. S., M. W. Guidry, and F. T. Mackenzie (2011), Dolomite controls on Phanerozoic seawater chemistry, *Aquat. Geochem.*, 17, 735–747.
- Bengston, P. (1988), Open nomenclature, *Palaeontology*, 31, 223–227.
- Bentov, S., and J. Erez (2006), The impact of biomineralization processes on the Mg content of foraminiferal shells: A biological perspective, *Geochem. Geophys. Geosyst.*, 7, Q01P08, doi:10.1029/2005GC001015.
- Berner, R. A. (1975), The role of magnesium in the crystal growth of calcite and aragonite from seawater, *Geochim. Cosmochim. Acta*, 39, 489–504.
- Berner, R. A. (2004), A model for calcium, magnesium and sulfate in seawater over Phanerozoic time, *Am. J. Sci.*, 304, 438–453.
- Berry, J. N. (1998), Sulfate in foraminiferal calcium carbonate: Investigating a potential proxy for sea water carbonate ion concentration, MS thesis, 88 pp., Mass. Inst. of Technol., Cambridge, Mass.
- Bots, P., L. G. Benning, R. E. M. Rickaby, and S. Shaw (2011), The role of SO₄ in the switch from calcite to aragonite seas, *Geology*, 39(4), 331–334.
- BouDagher-Fadel, M. K., F. T. Banner, and J. E. Whittaker (1997), *The Early Evolutionary History of Planktonic Foraminifera*, 269 pp., Chapman and Hall, London.
- Clémence, E.-M., and M. B. Hart (2013), Proliferation of Oberhauserellidae during the recovery following the late Triassic extinction: Paleocological implications, *J. Paleontol.*, 87(6), 1004–1015.
- Copestake, P., and B. Johnson (2014), *Lower Jurassic Foraminifera From the Llanbedr (Mochras Farm) Borehole, North Wales, UK*, Monograph of the Palaeontographical Society, vol. 167, 403 pp., Palaeontogr. Soc., London.
- De Choudens-Sánchez, V., and L. A. González (2009), Calcite and aragonite precipitation under controlled instantaneous supersaturation: Elucidating the role of CaCO₃ saturation state and Mg/Ca ratio on calcium carbonate polymorphism, *J. Sediment. Res.*, 79, 363–376.

- De Nooijer, L. J., T. Toyofuku, and H. Kitazato (2009), Foraminifera promote calcification by elevating their intracellular pH, *Proc. Natl. Acad. Sci. U.S.A.*, **106**, 15,374–15,378.
- De Nooijer, L. J., H. J. Spero, J. Erez, J. Bijma, and G. J. Reichart (2014), Biomineralization in perforate foraminifera, *Earth Sci. Rev.*, **135**, 48–58.
- Demicco, R. V., T. K. Lowenstein, L. A. Hardie, and R. J. Spencer (2005), Model of seawater composition for the Phanerozoic, *Geology*, **33**(11), 877–880.
- Dickson, J. A. D. (2002), Fossil echinoderms as monitor of the Mg/Ca ratio of Phanerozoic oceans, *Science*, **298**, 1222–1224.
- Dickson, J. A. D. (2004), Echinoderm skeletal preservation: Calcite-aragonite seas and the Mg/Ca ratio of Phanerozoic oceans, *J. Sediment. Res.*, **74**, 355–365.
- Evans, D., and W. Müller (2012), Deep time foraminifera Mg/Ca paleothermometry: Nonlinear correction for secular change in seawater Mg/Ca, *Paleoceanography*, **27**, PA4205, doi:10.1029/2012PA002315.
- Evans, D., J. Erez, S. Oron, and W. Müller (2015), Mg/Ca-temperature and seawater-test chemistry relationships in the shallow-dwelling large benthic foraminifera *Operculina ammonoides*, *Geochim. Cosmochim. Acta*, **148**, 325–342.
- Fallini, G., M. Gazzano, and A. Ripamonti (1994), Crystallization of calcium carbonate in presence of magnesium and polyelectrolytes, *J. Cryst. Growth*, **137**, 577–584.
- Farkaš, J., F. Böhm, K. Wallmann, J. Blenkinsop, A. Eisenhauer, R. van Geldern, A. Munnecke, S. Voigt, and J. Veizer (2007), Calcium isotope record of Phanerozoic oceans: Implications for chemical evolution of seawater and its causative mechanisms, *Geochim. Cosmochim. Acta*, **71**, 5117–5134.
- Fernández-Díaz, L., A. Putnis, M. Prieto, and C. V. Putnis (1996), The role of magnesium in the crystallization of calcite and aragonite in a porous medium, *J. Sediment. Res.*, **66**, 482–491.
- Fuchs, W. (1967), *Ober Ursprung und Phylogenie der Trias-"Globigerinen" und die Bedeutung Dieses Formenkreises für das Echte Plankton*, pp. 135–177, Verh. der Geol. Bundesanst., Wien.
- Fuchs, W. (1970), *Eine Alpine, Tiefliassische Foraminiferenfauna von Hernstein in Niederösterreich*, pp. 66–145, Verh. der Geol. Bundesanst., Wien.
- Gupta, S. B. K. (2002), *Modern Foraminifera*, edited by S. B. K. Gupta, pp. 23–33, Kluwer Acad., Dordrecht, Netherlands.
- Hansen, K. W., and K. Wallmann (2003), Cretaceous and Cenozoic evolution of seawater composition, atmospheric O₂ and CO₂: A model perspective, *Am. J. Sci.*, **303**, 94–148.
- Hardie, L. A. (1996), Secular variation in seawater chemistry: An explanation for the coupled secular variation in the mineralogies of marine limestones and potash evaporites over the past 600 Myr, *Geology*, **24**, 279–283.
- Hart, M. B. (1984), *Benthos '83: 2nd International Symposium on Benthic Foraminifera*, Pau, April 1983, edited by H. J. Oertli, pp. 289–298, Elf Aquitaine, Esso REP and Total CFP, Pau and Bordeaux.
- Hart, M. B. (1999), The evolution and biodiversity of Cretaceous planktonic Foraminifera, *Géobios*, **32**, 247–255.
- Hart, M. B., M. J. Oxford, and W. Hudson (2002), *Palaeobiogeography and Biodiversity Change: The Ordovician and Mesozoic-Cenozoic Radiations*, edited by J. A. Crame and A. W. Owen, *Geol. Soc. London, Spec. Publ.*, **194**, 115–125.
- Hart, M. B., M. D. Hylton, M. J. Oxford, G. D. Price, W. Hudson, and C. W. Smart (2003), The search for the origin of the planktonic foraminifera, *J. Geol. Soc. London*, **160**, 341–343.
- Hart, M. B., G. D. Price, and C. W. Smart (2009), Foraminifera and sequence stratigraphy of the lower part of the Speeton Clay Formation (Lower Cretaceous), N. E. England, *Ann. Naturhist. Mus. Wien*, **110A**, 423–442.
- Hart, M. B., W. Hudson, C. W. Smart, and J. Tyska (2012), A reassessment of "Globigerina bathoniana" Pazdrowa, 1969 and the palaeoceanographic significance of Jurassic planktic foraminifera from southern Poland, *J. Micropaleontol.*, **31**, 97–109.
- Hart, M. B., A. D. Leighton, C. W. Smart, L. R. Pettit, I. N. Medina-Sánchez, P. J. Harries, A. L. Cárdenas, J. M. Hall-Spencer, and R. M. Prol-Ledesma (2014), Ocean acidification in modern seas and its recognition in the geological record: The Cretaceous/Paleogene boundary in Texas and Alabama, *Gulf Coast Assoc. Geol. Soc. Trans.*, **64**, 193–213.
- Hemleben, C., O. R. Anderson, W. Berthold, and M. Spindler (1986), Calcification and chamber formation in foraminifera—A brief overview, in *The Systematics Association, Special*, vol. 30, edited by B. S. Leadbeater and R. Riding, pp. 237–249, Clarendon Press, Oxford.
- Heuser, A., A. Eisenhauer, F. Böhm, K. Wallmann, N. Gussone, P. N. Pearson, T. F. Nägler, and W.-C. Dullo (2005), Calcium isotope ($\delta^{44}\text{Ca}$) variations of Neogene planktonic foraminifera, *Paleoceanography*, **20**, PA2013, doi:10.1029/2004PA001048.
- Holland, H. D. (2005), Sea level, sediments and the composition of seawater, *Am. J. Sci.*, **305**, 220–239.
- Holland, H. D., and H. Zimmermann (2000), The dolomite problem revisited, *Int. Geol. Rev.*, **42**, 481–490.
- Holland, H. D., J. Horita, and W. E. Seyfried Jr. (1996), On the secular variations in the composition of Phanerozoic marine potash evaporites, *Geology*, **24**, 993–996.
- Hönisch, B., et al. (2012), The geological record of ocean acidification, *Science*, **335**, 1058–1063.
- Horita, J., H. Zimmermann, and H. D. Holland (2002), Chemical evolution of seawater during the 288 Phanerozoic: Implications from the record of marine evaporites, *Geochim. Cosmochim. Acta*, **66**, 3733–3756.
- Ilyina, T., and R. E. Zeebe (2012), Detection and projection of carbonate dissolution in the water column and deep-sea sediments due to ocean acidification, *Geophys. Res. Lett.*, **39**, L06606, doi:10.1029/2012GL051272.
- Kiessling, W., M. Aberhan, and L. Villier (2008), Phanerozoic trends in skeletal mineralogy driven by mass extinctions, *Nat. Geosci.*, **1**(8), 527–530.
- Kiessling, W., M. E. Clapham, A. J. W. Hendy, F. T. Fursich, L. Villier, J. Alroy, and A. I. Miller (2015), Taxonomic occurrences of to 600 Foraminifera, Foraminifera, and Foraminifera recorded in, Fossilworks: The evolution of terrestrial ecosystems database, and the paleobiology database, Fossilworks. [Available at <http://fossilworks.org>.]
- Loeblich, A. R., and H. Tappan (1984), Suprageneric classification of the Foraminifera (Protozoa), *Micropaleontology*, **30**, 1–70.
- Loeblich, A. R., and H. Tappan (1988), *Foraminiferal Genera and Their Classification*, Van Nostrand Reinhold, New York.
- Lowenstein, T. K., M. N. Timofeeff, S. T. Brennan, L. A. Hardie, and R. V. Demicco (2001), Oscillations in Phanerozoic seawater chemistry: Evidence from fluid inclusions, *Science*, **294**, 1086–1088.
- Martin, R. E. (1995), Cyclic and secular variation in microfossil biomineralization: Clues to the biogeochemical evolution of Phanerozoic oceans, *Global Planet. Change*, **11**, 1–23.
- Mewes, A., G. Langer, L. J. De Nooijer, J. Bijma, and G. J. Reichart (2014), Effect of different seawater Mg²⁺ concentrations on calcification in two benthic foraminifera, *Mar. Micropaleontol.*, **113**, 56–64.
- Mucci, A., and J. W. Morse (1985), Auger spectroscopy determination of the surface-most adsorbed layer composition on aragonite, calcite, dolomite and magnesite in synthetic seawater, *Am. J. Sci.*, **285**, 306–317.
- Nancollas, G. H., and K. Sawada (1982), Formation of scales of calcium carbonate polymorphs: The influence of magnesium ion and inhibitors, *J. Pet. Technol.*, **34**, 645–652.
- Nehrke, G., N. Keul, G. Langer, L. J. de Nooijer, J. Bijma, and A. Meibom (2013), A new model for biomineralization and trace-element signatures of Foraminifera tests, *Biogeosciences*, **10**, 6759–6767, doi:10.5194/bg-10-6759-2013.

- Orr, J. C., et al. (2005), Anthropogenic ocean acidification over the twenty-first century and its impact on calcifying organisms, *Nature*, **437**, 681–686.
- Oxford, M. J., M. B. Hart, and M. P. Watkinson (2004), Foraminiferal characterization of mid-Upper Jurassic sequences in the Wessex Basin (United Kingdom), *Riv. Ital. Paleontol. Stratigr.*, **110**, 209–218.
- Porter, S. M. (2007), Seawater chemistry and early carbonate biomineralization, *Science*, **316**, 1302.
- Porter, S. M. (2010), Calcite and aragonite seas and the de novo acquisition of carbonate skeletons, *Geobiology*, **8**, 256–277.
- Premoli Silva, I., and W. V. Sliter (1999), *Evolution of the Cretaceous Ocean-Climate System, Spec. Pap.*, vol. 232, edited by E. Barrera and C. Johnson, pp. 301–328, Geol. Soc. of Am., Boulder, Colo.
- Raup, D. M., and J. J. Sepkoski Jr. (1982), Mass extinctions in the marine fossil record, *Science*, **215**, 1501–1503.
- Reddy, M., and K. K. Wang (1980), Crystallization of calcium carbonate in the presence of metal ions. I. Inhibition by magnesium ion at pH8.8 and 25°C, *J. Cryst. Growth*, **50**, 470–480.
- Riding, R., and L. Liang (2005), Seawater chemistry control of marine limestone accumulation over the past 550 million years, *Rev. Esp. Micropaleontol.*, **37**, 1–11.
- Ries, J. B. (2004), The effect of ambient Mg/Ca on Mg fractionation in calcareous marine invertebrates: A record of Phanerozoic Mg/Ca in seawater, *Geology*, **32**(11), 981–984.
- Ries, J. B. (2010), Geological and experimental evidence for secular variation in seawater Mg/Ca (calcite-aragonite seas) and its effects on marine biological calcification, *Biogeosciences*, **7**, 2795–2849.
- Sandberg, P. A. (1983), An oscillating trend in Phanerozoic non-skeletal carbonate mineralogy, *Nature*, **305**, 19–22.
- Segev, E., and J. Erez (2006), Effect of Mg/Ca ratio in seawater on shell composition in shallow benthic foraminifera, *Geochem. Geophys. Geosyst.*, **7**, Q02P09, doi:10.1029/2005GC000969.
- Stanley, S. M. (2006), Influence of seawater chemistry on biomineralization throughout phanerozoic time: Paleontological and experimental evidence, *Palaeogeogr. Palaeoclimatol. Palaeoecol.*, **232**, 214–236.
- Stanley, S. M. (2008), Effects of global seawater chemistry on biomineralization: Past, present, and future, *Chem. Rev.*, **108**, 4483–4498.
- Stanley, S. M., and L. A. Hardie (1998), Secular oscillations in the carbonate mineralogy of reef-building and sediment-producing organisms driven by tectonically forced shifts in seawater chemistry, *Palaeogeogr. Palaeoclimatol. Palaeoecol.*, **144**, 3–19.
- Stanley, S. M., J. B. Ries, and L. A. Hardie (2002), Low-magnesium calcite produced by coralline algae in seawater of Late Cretaceous composition, *Proc. Natl. Acad. Sci. U.S.A.*, **99**, 15,323–15,326.
- Steuber, M., and M. Rauch (2005), Evolution of the Mg/Ca ratio of Cretaceous seawater: Implications from the composition of biological low-Mg calcite, *Mar. Geol.*, **217**, 199–213.
- Sun, W., S. Jayaraman, W. Chein, K. A. Persson, and G. Ceder (2015), Nucleation of metastable aragonite CaCO₃ in seawater, *Proc. Natl. Acad. Sci. U.S.A.*, **112**(11), 3199–3204.
- Valentine, J. W., D. Jablonski, A. Z. Krug, and S. K. Berke (2013), The sampling and estimation of marine paleodiversity patterns: Implications of a Pliocene model, *Paleobiology*, **39**, 1–20.
- Van de Waal, D. B., U. John, P. Ziveri, G.-J. Reichert, M. Hoins, A. Sluijs, and B. Rost (2013), Ocean acidification reduces growth and calcification in a marine dinoflagellate, *PLoS One*, **8**, e65987.
- Veizer, J., et al. (1999), ⁸⁷Sr/⁸⁶Sr, δ¹³C and δ¹⁸O evolution of Phanerozoic seawater, *Chem. Geol.*, **161**, 59–88.
- Wallmann, K. (2001), Controls on the Cretaceous and Cenozoic evolution of seawater composition, atmospheric CO₂ and climate, *Geochim. Cosmochim. Acta*, **65**, 3005–3025.
- Wallmann, K. (2004), Impact of atmospheric CO₂ and galactic cosmic radiation on Phanerozoic climate change and the marine δ¹⁸O record, *Geochem. Geophys. Geosyst.*, **5**, Q06004, doi:10.1029/2003GC000683.
- Wall-Palmer, D., M. B. Hart, C. W. Smart, R. S. J. Sparks, A. Le Friant, G. Boudon, C. Deplus, and J. C. Komorowski (2012), Pteropods from the Caribbean Sea: Variations in calcification as an indicator of past ocean carbonate saturation, *Biogeosciences*, **9**, 309–315.
- Wernli, R. (1988), Les protoglobigérines (foraminifères) du Toarcien et de l'Aalenien du Domuz Dag (Taurus Occidental, Turquie), *Eclogae Geol. Helv.*, **81**, 661–668.
- Wernli, R. (1995), Les foraminifères globigériniformes (Oberhauserellidae) du Toarcien inférieur de Teysachaux (Préalpes médianes, Fribourg, Suisse), *Rev. Paléobiol.*, **14**, 257–269.
- Wilkinson, B. H., and T. J. Algeo (1989), Sedimentary carbonate record of calcium-magnesium cycling, *Am. J. Sci.*, **289**, 1158–1194.
- Zeebe, R. E., and A. Sanyal (2002), Comparison of two potential strategies of planktonic foraminifera for house building: Mg²⁺ or H⁺ removal?, *Geochim. Cosmochim. Acta*, **66**, 159–169.