

**Title:**

**Light signaling, root development and plasticity**

Kasper van Gelderen<sup>1</sup>, Chiakai Kang<sup>1</sup> & Ronald Pierik<sup>\*</sup>

<sup>1</sup>These authors contributed equally

<sup>\*</sup>Author for correspondence

Author affiliation:

Plant Ecophysiology, Utrecht University. Padualaan 8, 3584CH Utrecht, the Netherlands.

**Short title:**

**Light signaling, root development and plasticity**

Corresponding author: Prof. Dr. Ronald Pierik, [r.pierik@uu.nl](mailto:r.pierik@uu.nl), Plant Ecophysiology, Utrecht University. Padualaan 8, 3584CH Utrecht, the Netherlands

Author contributions: K.v.G., C.K. and R.P. wrote the manuscript and composed the figures.

**Funding Information**

K.v.G. is funded by the Netherlands Organisation for Scientific Research, open competition grant 823.01.013 to R.P., C.K. is funded by the scholarship of government sponsorship for overseas study, Taiwan, admission number 0991167-2-UK-004.

**Single sentence summary**

Light signaling can affect root development and plasticity, either direct, or through shoot-root communication via sugars, hormones, light, or other mobile factors.

**Competing interests**

The authors declare no competing interests

33 **Corresponding author e-mail:**

34 [r.pierik@uu.nl](mailto:r.pierik@uu.nl)

35

Light is the energy source for plants as it drives photosynthesis to produce sugars. Given the obvious fact that light mostly occurs aboveground and not in the soil, most interactions of plants with light have been studied in shoot parts of the plant. Research over more than a century has yielded tremendous insights into how light not only drives photosynthesis, but also acts as an environmental cue that informs plants about their environment. Light quality and duration for example, drive major developmental changes such as photomorphogenesis, photoperiodic induction of flowering, phototropism, and shade avoidance (see for example the following recent reviews: (Wu, 2014; Fankhauser and Christie, 2015; Xu et al., 2015; Ballaré and Pierik, 2017). The picture that has emerged is that plants have very detailed light signaling mechanisms, with photoreceptors dedicated to different wavelengths in the light spectrum and interactions between these photoreceptors themselves and their downstream signal transduction pathways. Studies have accumulated over the past 15 years, and intensified in recent years showing pronounced effects of light on root physiology and development. Although some effects of light availability on root growth will be the simple consequence of differential sugar availability to the roots due to photosynthesis in the shoot, there is substantial evidence for more sophisticated signaling impacts of different aspects of the light environment. In this Update, we will briefly review the core light signaling mechanisms, their impact on root development and plasticity and the functional implications of these above-belowground interactions.

## **LIGHT SIGNALING**

Different wavelengths of light are associated with various functions in plant development and plants have a range of photoreceptors to detect these wavelengths. UVR8 is sensitive to UV-B light, cryptochromes (CRY) and phototropins (PHOT) detect UV-A and blue light, whereas phytochromes (PHY) sense red (R) and far-red (FR) light. Photoreceptors occur all over the plant body and although they are most abundant in the shoot, they are also expressed in the roots (Fig. 1). Photoreceptors share downstream signaling hubs, notably the COP1/SPA complex and PIFs, which will be discussed first, followed by the properties of each photoreceptor group.

### Shared signaling hubs


#### *COP1-SPA-HY5*

### BOX 1. Root Development

Plant roots take up water and nutrients that are essential for plant growth. The Arabidopsis root grows from a set of stem cells at the root tip governed by the organizing cells of the quiescent center (QC), four cells that can be traced back to the triangular embryo development stage (Scheres et al., 1994). Around the QC, stem cells divide to form all the layers of the Arabidopsis root, which are from outer to inner: the root cap, epidermis, cortex, endodermis, pericycle, and vasculature, including the xylem and phloem (Petricka et al., 2012). In the meristem zone above the QC, stem cells divide and are not yet differentiated, while at roughly 0.2 mm above the QC the cells differentiate and elongate. It is at the boundary of these two zones where the lateral roots are primed in the xylem pole pericycle layer through an auxin signal (De Smet et al., 2007). Lateral roots develop and emerge at a later time by anticlinal and periclinal divisions of the primed xylem pole pericycle cells (De Rybel et al., 2010), after which the growing lateral root primordium actively modifies the overlying endodermis, cortex, and epidermis tissue to emerge (Péret et al., 2013; Vermeer et al., 2014).

The plant hormone auxin is necessary for all aspects of root development. In the root tip, there is a high concentration of auxin that is achieved by the active transport of auxin towards the tip and QC (Blilou et al., 2005) and local auxin biosynthesis in the root stem cells (Stepanova et al., 2008). The auxin in the root tip drives the expression of PLETHORA transcription factors, which regulate differentiation via a degradation-based PLETHORA gradient (Galinha et al., 2007; Mähönen et al., 2014). In lateral root development, auxin is similarly important and facilitates the first divisions of the primordium, followed by the development through cell division and the emergence through the overlying tissue (Lavenus et al., 2013). Cytokinin acts antagonistically to auxin, and together these two plant hormones determine root patterning. Briefly, auxin induces cell division, while cytokinin promotes differentiation (Dello Iorio et al., 2007, 2008).

69 CONSTITUTIVE PHOTOMORPHOGENESIS1 (COP1) functions as a ubiquitin E3 ligase that  
70 targets proteins required for photomorphogenesis for degradation, such as the bZIP transcription  
71 factor ELONGATED HYPOCOTYL 5 (HY5) (Osterlund et al., 2000; Saijo et al., 2003) and HY5-  
72 HOMOLOGUE (HYH) (Holm et al., 2002). Especially HY5 is a key regulator in  
73 photomorphogenesis, and it serves as the center of a transcriptional network hub of hormone  
74 and light signaling (Gangappa and Botto, 2016). Besides HY5, COP1 also targets factors like the  
75 MYB transcription factor LONG AFTER FAR-RED LIGHT 1 (LAF1) (Seo et al., 2003), the bHLH  
76 protein LONG HYPOCOTYL IN FAR-RED 1 (HFR1) (Duek et al., 2004) and even phytochrome  
77 A (phyA) and phyB photoreceptors (Seo et al., 2004; Jang et al., 2010). COP1 activity relies on  
78 physical interaction with SUPPRESSOR OF PHYTOCHROME A proteins (SPAs) in the COP1-  
79 SPA complex. The cryptochrome photoreceptors CRY1 and CRY2, upon blue light activation,  
80 can bind to SPAs and this results in reduced COP1-SPA interaction, leading to stabilization of  
81 COP1 targets (Lian et al., 2011; Liu et al., 2011; Zuo et al., 2011). Comparable interaction



**Figure 1. Relative expression levels of photoreceptors in *Arabidopsis* tissues.**

The graph displays the relative expression of photoreceptors across seedling tissues. Both gene expression and protein (when data are available) abundance are shown. The box that groups classes of receptors indicates that intensities within can be compared. In the phytochrome box 'L' stands for protein levels in light and 'D' for protein levels in the dark. Source data: For all, BAR eFP browser (<http://bar.utoronto.ca/efp/cgi-bin/efpWeb.cgi>), phytochromes (Somers and Quail, 1995; Goosey et al., 1997; Toth et al., 2001; Sharrock and Clack, 2002; Salisbury et al., 2007), phototropins (Sakamoto and Briggs, 2002; Moni et al., 2015), cryptochromes (Toth et al., 2001), and UVR8 (Rizzini et al., 2011).

mechanisms exist between phytochromes and SPAs (Zheng et al., 2013; Sheerin et al., 2015), and UVR8 and SPAs (Favory et al., 2009; Huang et al., 2013) Another mechanism that controls COP1 activity is its light-based nuclear exclusion that affects HY5 degradation (von Arnim and Deng, 1994; Pacin et al., 2014). COP1 and HY5 are well-known regulators of the transition from skoto- to photomorphogenesis and for a detailed overview we refer to (Huang et al., 2014).

### *PHYTOCHROME INTERACTING FACTORS (PIFs)*

PIFs are a group of basic helix-loop-helix (bHLH) transcription factors that mediate various physiological responses, such as seed germination, seedling photomorphogenesis, shade avoidance and shoot architectural responses to elevated temperature (Koini et al., 2009; Chen and Chory, 2011; Hornitschek et al., 2012; Jeong and Choi, 2013; Xu et al., 2015). PIFs are regulated by light, temperature and the circadian clock and these cues can lead to PIF degradation by concerted action of the phytochromes and the E3-ligase Light-Response Bric-a-Brack/Tramtrack/Broad (BTB) (LRB) (Leivar and Monte, 2014; Ni et al., 2014; Jung et al., 2016; Legris et al., 2016). The phytochrome and cryptochrome photoreceptors can physically interact with specific sets of PIF proteins and thereby regulate PIF phosphorylation. The phosphorylation

status of PIFs subsequently determines their activity and in most instances their stability (Paik et al., 2017). ChIP-seq studies on PIFs have identified a very broad range of target genes, including those associated with cell wall modifications, auxin biology and several other transcription factors. This broad range of targets combined with the multitude of signals that impact on PIF stability makes PIF proteins a signaling hub to integrate environmental conditions (Leivar and Monte, 2014; Paik et al., 2017).

## Photoreceptors

### *UV-B RESISTANCE LOCUS8 (UVR8)*

The effect of UV-B radiation on plant growth is dual. High fluence UV-B exposure causes photodamage, but low fluence UV-B contributes to photomorphogenesis and it increases the resistance to herbivorous insects and pathogens (Ballaré et al., 2012; Galvão and Fankhauser, 2015). UV-B (280-315 nm) is perceived by UVR8, which uses a structure based on tryptophans and a complex salt bridge network. Upon sensing UV-B light, the salt bridges are disrupted, inducing dimer dissociation into UVR8 monomers and initiating signal transduction (Christie et al., 2012). The UV-B induced responses include hypocotyl growth inhibition, altered leaf morphogenesis, stomatal closure and compound synthesis associated with the prevention and repair of UV damage (Binkert et al., 2014; Galvão and Fankhauser, 2015). UVR8 regulates UV-B responses by negatively affecting COP1 function through the UVR8-SPA interaction (Osterlund et al., 2000; Favory et al., 2009; Huang et al., 2013; Hayes et al., 2014). Formation of UVR8-SPA leads to inhibition of COP1 function and this results in stabilization of amongst others HY5 and HYH, thereby inducing genes downstream associated with UV-B signaling (Osterlund et al., 2000; Rizzini et al., 2011; Hayes et al., 2014).

### *Cryptochromes*

Blue light (320-500 nm) is used for photosynthesis and also serves as a signal for shade, photoperiodism and directional light. Blue light sensing is performed by three groups of light receptors: cryptochromes, phototropins, and other LOV-domain containing receptors such as ZEITLUPE. Cryptochromes (CRYs) are blue light receptors found in a broad range of organisms, including bacteria, fungi, animals, and plants. Three CRYs are encoded in the Arabidopsis genome, *CRY1*, *CRY2*, and *CRY3*. *CRY1* and *CRY2* act redundantly in promoting flower induction, sensing blue light as input to circadian clock, and stomatal opening in *Arabidopsis* (Li and Yang, 2007; Galvão and Fankhauser, 2015; Mo et al., 2015). Both *CRY1* and *CRY2* regulate primary root elongation, but *CRY1* promotes primary root elongation in blue light,

whereas CRY2 has the opposite effect (Canamero et al., 2006). Mainly CRY2 is expressed in *Arabidopsis* roots (Fig. 1) and CRY1 and CRY2 inhibit root growth via modulation of free auxin levels and polar auxin transport (Mo et al., 2015). The biological function of CRY3 is unclear yet (Song et al., 2006; Klar et al., 2007; Mo et al., 2015). CRYs are nuclear flavoproteins, composed of two domains, an N-terminal photolyase-related (PHR) region and a C-terminal domain of varying size. The PHR region binds the chromophore FAD and serves as the light sensing part. Upon absorbing blue/UV-A light, CRY is phosphorylated and can interact with other protein partners, such as COP1, SPAs and PIFs (Lian et al., 2011; Liu et al., 2011; Zuo et al., 2011; Pedmale et al., 2016; de Wit et al., 2016).

### *Phototropins*

Phototropins are blue light receptors that mediate phototropism, chloroplast movement and stomatal opening in *Arabidopsis* (Briggs and Christie, 2002). There are two phototropins identified in *Arabidopsis*, phot1 and phot2, with similar function and structure. Besides their function in the shoot, *phot1* is also expressed in the roots, regulating root bending (Wan et al., 2012; Zhang et al., 2013). In contrast to other photoreceptors, which are present in the nucleus or cytoplasm, phototropins are located in the plasma membrane (Sakamoto and Briggs, 2002). When phot1 is activated by blue light, it phosphorylates PHYTOCHROME KINASE SUBSTRATE 4 (PKS4), and interacts with NON-PHOTOTROPIC HYPOCOTYL 3 (NPH3) to form a phot1-PKS-NPH3 protein complex (Pedmale and Liscum, 2007; Demarsy et al., 2012). This modulates auxin transport and thus underlies plant phototropism (Fankhauser and Christie, 2015). Phototropins have N-terminal flavin mononucleotide (FMN) chromophore-binding light oxygen voltage (LOV1 and LOV2) domains for light absorption, and a C-terminal AGC-type Ser/Thr protein kinase signaling domain (Briggs and Christie, 2002).

The ZEITLUPE (ZTL), FLAVIN-BINDING, KELCH REPEAT, F-BOX (FKF1), and LOV KELCH PROTEIN2 (LKP2), are blue light photoreceptors involved in circadian clock and photoperiodic flowering regulation, with a structure similar to Phototropins, but with only one LOV domain followed by an F-box and six Kelch repeats (Galvão and Fankhauser, 2015).

### *Phytochromes*

Phytochromes are photoreceptors sensing R and FR light that occur in plants as well as in fungi and prokaryotes (Burgie and Vierstra, 2014). Plants use R/FR light signaling through phytochromes to regulate germination, de-etiolation, stomatal development, flowering transition, senescence, and shade avoidance (Franklin and Quail, 2010). Five phytochromes are identified

in *Arabidopsis*, phyA-phyE. Phytochromes use the N-terminal covalently linked phytochromobilin to sense light and the C-terminus to transmit the light signal. Phytochromes undergo reversible conformation changes: The inactive Pr form absorbs R light (max=660nm) that leads to its photoconversion into the active Pfr form that can then absorb FR light (max=730nm) to be inactivated. Upon activation, Pfr translocates from the cytosol to the nucleus, where it interacts with PIFs, modulating their activity (Chen and Chory, 2011; Leivar and Quail, 2011; Xu et al., 2015). PhyA, PhyB, PhyD and PhyE are expressed in the root, while PhyC expression in the root is hardly detectable (Fig. 1). *Phyb* mutants have been shown to produce fewer lateral roots (Salisbury et al., 2007) and *Phyb* and *Phya* single and double mutants have reduced root elongation compared to wild-type (Correll and Kiss, 2005; Silva-Navas et al., 2015). Recently, PhyB has been identified to act as a temperature sensor (Jung et al., 2016; Legris et al., 2016), which is interesting since heat stress is known to lead to increased main root growth (Hanzawa et al., 2013). If this occurs through PhyB signaling is yet to be resolved, but since high temperatures destabilize PhyB and *Phyb* mutants have reduced root lengths, this may not be likely.

## IMPACT OF DIRECT LIGHT SIGNALING ON ROOT DEVELOPMENT AND PLASTICITY

### Root development in dark versus light

Root development (Box 1) starts at embryogenesis (Scheres et al., 1994) and all root layers are continuously formed from stem cells at the tip (Petricka et al., 2012) that are maintained by a constant auxin flow (Blilou et al., 2005; Galinha et al., 2007; Stepanova et al., 2008; Mähönen et al., 2014), while cytokinin promotes differentiation of these root layers (Dello Iorio et al., 2007; Dello Iorio et al., 2008). Lateral roots form after embryogenesis and are primed in the differentiation zone of the meristem (De Smet et al., 2007; De Rybel et al., 2010). Subsequently they emerge through the outer layers of the main root (Lavenus et al., 2013) upon modifications of these layers (Péret et al., 2013; Vermeer et al., 2014). In total darkness, the *Arabidopsis* seedling elongates its hypocotyl in an attempt to penetrate the soil, however its root stays very small (Fig. 2). Roots of dark-grown seedlings are much shorter and have a much thinner diameter than those of light-grown seedlings (Laxmi et al., 2008; Dyachok et al., 2011). When etiolated seedlings are exposed to light they inhibit hypocotyl elongation, develop their cotyledons and start to photosynthesize (Wu, 2014). Cotyledon-derived sugars are essential for the start of root growth and when young seedlings are decapitated root growth is slowed




## BOX 2. Shade Avoidance

When shade-intolerant plants are surrounded by neighbors, they typically engage in direct competition by inducing the “shade avoidance syndrome” (see fig.). Shade avoidance responses include elongation of stems and petioles, upward leaf movement (hyponasty), enhanced apical dominance (reduced branching), and early flowering (Gommers et al., 2013; Ballaré and Pierik, 2017). These responses help plants to escape from shade caused by neighboring leaves and enable access to direct sunlight. Shade avoidance responses are adaptive, but may come at the costs of the economic yield of crops or at the cost of pathogen and herbivore resistance (Ballaré et al., 2012; Ballaré, 2014), and therefore are not always desirable in agricultural systems. Shade avoidance is induced by the sensing of light signals: First, due to the reflection of far-red (FR) light of neighboring leaves, which lowers the R:FR ratio, and second, by the depletion of blue (B) light when light limitation through canopy shading occurs (Casal, 2013; Ballaré and Pierik, 2017). The low R:FR ratio is perceived by phytochromes, which are subsequently inactivated and leading to PIF stabilization, as described in the main text. Mainly PIF4, PIF5, and PIF7 are involved in the shade avoidance response, and their stabilization and transcriptional responses lead to auxin accumulation, induction of cell wall-remodeling genes, and subsequent cell elongation, ultimately resulting in hypocotyl and petiole elongation and hyponastic leaf movement (Lorrain et al., 2008; Keuskamp et al., 2010; Kohnen et al., 2016; Michaud et al., 2017; Pantazopoulou et al., 2017). Low B is perceived by the cryptochromes and their signaling pathway is partly converged with phytochromes during shade avoidance

by acting on PIFs (Pedmale et al., 2016) and on the degradation of the negative regulator of shade avoidance HFR1 (Keuskamp et al., 2011; de Wit et al., 2016). The combination of low B and low R:FR represents the actual canopy shade, and the shoot elongation response to this signal combination is stronger than the response to low R:FR alone (de Wit et al., 2016), indicating that plants adjust shade avoidance responses to signal intensity.

200 dramatically, also when grown in the light (Kircher and Schopfer, 2012). Besides sucrose, the  
201 basipetal flow of auxin is necessary to facilitate root growth in seedlings (Bhalerao et al., 2002)  
202 and in the dark this basipetal auxin transport is very low due to the depletion of PIN-FORMED  
203 (PIN) auxin efflux carriers from the plasma membrane (Laxmi et al., 2008; Sassi et al., 2012).



**Figure 2. Root growth is affected by light quantity and quality.**

(A) 8d old seedlings grown on  $\frac{1}{2}$  MS medium in darkness and white light (WL) ( $140\mu\text{M m}^{-2} \text{s}^{-1}$  photosynthetically active radiation, PAR). (B) 8d old seedling grown in WL ( $140\mu\text{M m}^{-2} \text{s}^{-1}$ ) or WL+FR (R:FR 0.1).

The arrows point to the root-shoot junctions.

204 Light induces root growth by providing sugars and auxin to the young root and specifically red  
 205 and blue light exhibit a positive effect on root elongation when compared to darkness (Sweere,  
 206 2001; Canamero et al., 2006; Costigan et al., 2011)(Fig. 2). However, the addition of sucrose to  
 207 the agar medium can sometimes reverse this effect (Correll and Kiss, 2005). Supplementation of  
 208 white light with FR light reduces root growth compared to normal white light (Salisbury et al.,

2007)(Fig. 2) and UVB has a strong inhibiting effect on root growth, either when supplied to the whole seedling or only the root (Tong et al., 2008; Leasure et al., 2009; Silva-Navas et al., 2015)

Many experiments addressing root development have been performed in the presence of light on the whole seedling since root development is typically studied in vertical agar plate set-ups with transparent medium. In field conditions the top part of the root system will grow in minimal light and the lower part will even develop in darkness, while the shoot can be exposed to various different light conditions (Smith, 1982). Several solutions to these undesirable lab conditions have been postulated, including dark agar plugs (Sassi et al., 2012), black-colored vertical plates (Xu et al., 2013) and the D-root system consisting of plate inserts plus cover slips (Silva-Navas et al., 2015). In the D-root system, seedlings grow on medium in vertical square plates, but an insert at the root-shoot junction and an external cover slip prevents light exposure of the roots, whereas the shoot is exposed to the ambient light conditions. Compared to plants with roots grown in fully exposed light conditions, plants with roots in darkness have increased main root length and lateral root number, while root hair length decreased. When dark-grown roots were exposed to light for a duration longer than 8 hours, their growth rate and root meristem size declined (Silva-Navas et al., 2016). Interestingly, shielding roots from light decreased the sensitivity of the main root to the plant hormones abscisic acid (ABA), brassinolide (BL) and 6-benzyladenine (BAP), while the sensitivity to indole-3-acetic acid (IAA) was increased (Silva-Navas et al., 2015). Interestingly, dark-exposed roots were less sensitive to salt- and low nitrogen conditions, compared to light-exposed roots (Silva-Navas et al., 2015). These data show that direct perception of light by the root is physiologically relevant in *Arabidopsis* seedlings.

#### Light affects root developmental plasticity

The root system can change the direction of growth in response to stimuli such as gravity (Morita, 2010) or light (Kutschera and Briggs, 2012). The movement of the root away from light sources or 'root negative phototropism' is dependent upon blue-light perception by phototropins (Wan et al., 2012). Contrary to the negative response to blue light, a positive growth response of the root to red light has also been observed (Kiss et al., 2001), but this response is weak and can only be observed in the absence of gravity sensing (Ruppel et al., 2001; Kiss, 2003). Downstream of light perception, root phototropism impinges on elements involved in root gravitropism (Kiss, 2003; Kutschera and Briggs, 2012).

#### *Root-negative phototropism*

Root negative phototropism is induced by blue and white light and the photoreceptors involved are mainly phot1 and phot2, with minor roles for phyA and cry1/cry2 (Boccalandro et al., 2007; Wan et al., 2012; Silva-Navas et al., 2016). NON-PHOTOTROPIC HYPOCOTYL3 (NPH3), a target of phot1 and phot2 involved in hypocotyl phototropism, is also an important player in root negative phototropism (Wan et al., 2012). Downstream of directional light perception, polar auxin transport directs auxin away from the illuminated side. This auxin transport gradient requires rootward plasma membrane localization of the auxin efflux carrier PIN1 in the stele, lateral relocalization of PIN3 in the columella and rootward relocalization of PIN2 in the epidermis (Wan et al., 2012; Zhang et al., 2013; Zhang et al., 2014). The relocalization of PIN2 and PIN3 is dependent upon the recycling and targeted degradation (vacuolar targeting) of these PIN carriers from one side of the membrane to the other (Wan et al., 2012; Zhang et al., 2013). The initial perception of gravitropism and phototropism is different, but the downstream signaling events of negative phototropism are very similar to those found in root gravitropism, involving comparable PIN protein and auxin transport dynamics (Friml et al., 2002; Abas et al., 2006; Baster et al., 2012). Most root phototropism experiments have been performed under conditions where the whole seedling was directionally illuminated. When roots, which had been grown in the dark, were stimulated by a one-sided white-light stimulus, while the shoot still remained in the dark, the negative phototropism persisted and an important role for flavonols was uncovered in the regulation of this response (Silva-Navas et al., 2016). White light, together with cytokinins, stimulates the accumulation of flavonols on the lighted side of the root, which induces cell elongation and stimulates PIN1 plasma membrane abundance (Buer and Muday, 2004; Silva-Navas et al., 2016).

#### *Interaction of light and gravitropism*

As mentioned above, root-negative phototropism and gravitropism both rely on polar auxin transport. Light influences the direction of polar auxin transport by controlling the plasma membrane abundance of PIN proteins (Laxmi et al., 2008; Sassi et al., 2012; Wan et al., 2012; Zhang et al., 2013; Zhang et al., 2014). An interesting example of how light interacts with the gravitropic output is the U-turn that an inverted maize seedling root makes when growing in a glass tube exposed to light (Burbach et al., 2012; Suzuki et al., 2016). When these seedlings are inverted in the dark, they do not show this strong gravitropic response (U-turn), indicating that light can increase gravitropism. In accordance with this, *Arabidopsis* root slanting (on agar plates) and the root gravitropic response are reduced in the D-root system where roots are kept in darkness (Silva-Navas et al., 2015). Light stimulates the gravitropic response through the

increase of flavonol biosynthesis, which increases root auxin levels (Buer and Muday, 2004; Silva-Navas et al., 2016). Interestingly, FR light enrichment of *Arabidopsis* seedlings grown fully in the light leads to reduced activity of the auxin reporter pDR5::GUS (Salisbury et al., 2007), showing how light quality can markedly change root auxin homeostasis. Another way that light signaling influences auxin transport in the root is by controlling the removal of PIN proteins from the plasma membrane via the process of vacuolar degradation. Since PINs are transmembrane proteins, their degradation occurs through vacuolar targeting and subsequent degradation of Multi-Vesicular-Bodies containing PINs (Korbei and Luschig, 2013). This vacuolar degradation of PINs is an important process in changing PIN polarity and is essential for the regulation of gravitropism (Baster et al., 2012). PIN2-GFP vacuolar targeting is controlled by the COP1/CSN complex in a light dependent manner (Laxmi et al., 2008; Sassi et al., 2012). When a seedling is incubated in darkness for several hours PIN2-GFP is targeted towards the vacuole (Kleine-Vehn et al., 2008), however this does not occur when only the seedling shoot is given a light treatment, or in the *cop1-6* mutant (Sassi et al., 2012), showing how (COP1-mediated) light signaling in the shoot can affect root development.

## **ABOVEGROUND LIGHT REGULATES ROOT DEVELOPMENT: MECHANISMS AND CONSEQUENCES.**

As mentioned above, it is critical to research the effects of light signaling on root developmental plasticity under conditions where roots are not directly exposed to the light environment as they are typically shielded from by the soil under natural conditions. Indeed, root systems develop differently between dark and light conditions of the roots themselves. Since plants constantly coordinate growth and development of root and shoot in response to their highly dynamic environment, they do need to translate important information about their light environment to the root system. This is of particular importance under dense planting conditions, such as in most agricultural fields, where both above and below ground competition for resources occurs (Pierik et al., 2013; Gundel et al., 2014). Belowground, plants compete for water and nutrients whereas aboveground they struggle for light. Belowground competition is size-symmetric, which means that resource acquisition is proportional to the size of the root system of a given individual in a dense stand. Competition for light is size-asymmetric: A plant that is only slightly taller than its neighbors can put its leaves above these neighbors, thereby severely limiting their total access to light, whilst itself not being affected at all by its neighbors (Weiner, 1985; Weiner and Thomas, 1986). Aboveground, plants maximize their competitive performance against neighbors by

activating the so-called shade avoidance response (Box 2). Briefly, FR light reflected by neighbor plants inactivates the phytochrome photoreceptors and this relieves their repression of PIF 4, 5 and 7. PIFs then accumulate, induce auxin biosynthesis and transport, and auxin subsequently induces elongation of hypocotyls, stems and petioles as well as upward leaf movement (Lorrain et al., 2008; Keuskamp et al., 2010; Keuskamp et al., 2011; Casal, 2013; Gommers et al., 2013; Kohnen et al., 2016; de Wit et al., 2016; Ballaré and Pierik, 2017; Michaud et al., 2017; Pantazopoulou et al., 2017). In this section, we briefly review the currently established mechanisms of light signal information transfer from the shoot to the root, followed by a discussion on its functional importance under dense planting conditions and plant competition.

## Mechanisms of light signal information transfer from shoot to root


### *Stem-piped light*

Stem-piped light refers to the light transmitted through the internal tissues of the plants from the shoot to the root. This manner of light transmission through the interiors of the plant has been described for woody and herbaceous species. Light piping is wavelength-specific and long wavelengths such as FR and near infra-red light are transmitted relatively well while shorter wavelengths such as blue and green light are less effectively transmitted (Sun et al., 2003; Sun et al., 2005). Stem-piped light can activate root-expressed phyB, which on its turn regulates HY5 in the Arabidopsis root (Lee et al., 2016). HY5 is involved in root growth in response to light, modulating e.g. root gravitropism and nitrogen uptake (Cluis et al., 2004; Lee et al., 2007; Huang et al., 2015; Chen et al., 2016). Therefore, stem-piped light might communicate information about the aboveground light environment to the root (Fig. 3). However, the light transmission goes down to 1% when conduction distances increase to 3 cm in herbaceous plants (Sun et al., 2005), which suggests that this mechanism might play a relatively modest role in mature planting systems.

### *Mobile (signaling) chemicals*

Although various components are transported from the shoot to the root, only few of them have been directly linked to light signaling in the shoot. These include sugars derived from photosynthesis and transported through the phloem, plant hormones, and HY5. These will be briefly discussed below.

### *Sugars*



**Figure 3. The aboveground regulation of root development in dense vegetation involves photosynthesis, light piping, hormones and mobile factors.**

The primary effect of light is to enable photosynthesis which leads to the production of sugars (sucrose), that enable the root to grow. Photomorphogenic development is associated with the production of auxins in the shoot, which are transported rootward and enable root development. HY5 is stabilized in the shoot during photomorphogenesis and is transported rootwards where it regulates nitrate uptake and root development.

Light is used as a cue to detect neighboring plant competition via sensing of the R:FR ratio. Plant tissues reflect FR, which lowers the nearby R:FR ratio, leading to shade avoidance responses mediated by, amongst others, auxin and GA. These hormones can be transported rootward where they affect root development. There are also indications that shade avoidance responses, in a negative feedback mechanism, decrease the amount of rootward auxin transport. FR light itself can also be transmitted directly through woody, vascular, tissues from the shoot to the root, where it can affect root-localized phytochromes.

Once a seed is germinated, it first invests energy in hypocotyl growth. Upon penetrating the soil and perceiving light, photomorphogenesis is initiated and the seedling acquires the ability to conduct photosynthesis, followed by root growth (Kircher and Schopfer, 2012). It was shown in *Arabidopsis* seedlings that blocking photosynthesis inhibits root growth, just like in darkness, and that adding sucrose to the growth medium can rescue root growth. Interestingly, sucrose addition could not induce additional root growth in photosynthesis-positive seedlings (Kircher and Schopfer, 2012). These data confirm that sucrose is needed for root growth and development, and indicate that sucrose can serve as a long-distance signal from the shoot to inform the root system about light availability aboveground (Fig. 3).

### Auxin

Young shoot tissues are the main source of the plant hormone auxin that is then transported to the root system where it regulates root development, including lateral root formation (Reed et al., 1998; Bhalerao et al., 2002)(Fig. 3). Interestingly, light signaling affects auxin biosynthesis and transport, implicating this hormone as a potential integrator of light signaling and root development. Phytochrome inactivation in low R:FR light triggers auxin biosynthesis in the shoot



(Hornitschek et al., 2012), which is then transported laterally through the hypocotyl during low R:FR conditions by PIN3 as well as PIN4 and PIN7 (Keuskamp et al., 2010; Kohnen et al., 2016) and rootward by PIN1 and PIN2 in response to light (Sassi et al., 2012). In darkness, the expression of PIN1 is largely reduced, thus reducing auxin delivery to the root system (Sassi et al., 2012).

### Gibberellic Acid (GA)

The plant hormone GA mediates various growth and developmental processes, such as seed germination, cell elongation and reproductive development (Hedden and Sponsel, 2015). Low R:FR induces GA biosynthesis, at least partly through elevated expression of *GA20OX* genes (Hisamatsu et al., 2005). GAs are diterpenoid tetracyclic carboxylic acids, but only a few are biological active, such as GA<sub>1</sub> and GA<sub>4</sub> (Hedden and Thomas, 2012). Interestingly, biologically inactive GA<sub>12</sub> has been identified as a long-distance growth signal that is transported from the shoot to the root. Here GA<sub>12</sub> is converted into active form by GA20ox and GA3ox enzymes and subsequently promotes root growth in *Arabidopsis* (Regnault et al., 2015)(Fig. 3). It is yet unknown if low R:FR induced production of bioactive GAs in the shoot affects downward GA<sub>12</sub> transport.

### HY5

The transcription factor HY5 is a key integrator of photomorphogenesis and is involved in light, hormone and stress signaling (Cluis et al., 2004; Lee et al., 2007; Gangappa and Botto, 2016). HY5 was recently shown to be transported upon light activation from the shoot to the root (Chen et al., 2016). In the root system, HY5 activates its own expression and that of HY5 HOMOLOGUE (HYH), creating a positive feedback loop after shoot-to-root transport of HY5 (Zhang et al., 2017). HY5 and HYH modulate lateral root development and lateral root gravitropism (Oyama et al., 1997; Sibout et al., 2006). Interestingly, R:FR ratio's influence the expression and stability of HY5 and this transcription factor is, therefore, together with auxin and gibberellin a candidate regulator of root developmental adjustments to shoot-sensed R:FR light ratio's, which indicate the presence of neighboring competitors.

### Functional implications in dense vegetation

R:FR light conditions above-ground are a reliable cue for the presence of competing neighbors and effectively trigger shade avoidance (Box 2). Interestingly, if the R:FR ratio signals neighbor proximity, it would also signal the likely presence of neighbors belowground competing for water



and nutrients (Gundel et al., 2014). Competition experiments in *Arabidopsis* have shown that root biomass decreases when plants are competing in dense stands and that there is a separate transcriptomic regulation in the root versus the shoot (Masclaux et al., 2012). In controlled growth on medium, seedlings experiencing a low R:FR ratio have a lower lateral root number and main root length (Salisbury et al., 2007). However, as argued previously, in soil conditions the roots are not directly exposed to low R:FR light, and the shoot might thus relay information about nearby competitors to the root system either by direct piping, or through secondary mobile messengers.

An important question that remains is what are the functional implications of light signal transmission from the shoot to the root? One clear functional example is that HY5 can stimulate nitrate uptake via NRT2;1, and will do so upon light-activation in the shoot (Chen et al., 2016)(Fig. 3). This will help keep the carbon and nitrogen acquisition in tune. In accordance with these findings, direct shading in different phases of the growth period of field-grown maize was shown to suppress root mass, length and absorptive area, suggesting reduced nutrient and/or water uptake rates (Gao et al., 2017). In addition to true shade, a reduced R:FR light ratio also affects root growth in different species (Kasperbauer and Hunt, 1992; Kasperbauer and Hunt, 1994; Salisbury et al., 2007). The effects of reduced R:FR ratios on lateral root formation (Salisbury et al., 2007) may go at the expense of nutrient uptake rates, and might reflect a resource prioritization strategy at the whole plant level, ensuring that resources are invested to consolidate light capture in a growing vegetation with increasing competition for light. It is presently unknown if the root architecture responses to low R:FR affect nutrient acquisition at all, but the resulting reduced root length would suggest so. Nevertheless, much more detailed research into nutrient and water uptake rates of plants under high and low R:FR conditions is needed to assess the precise functional implications of shoot-to-root communication of the aboveground light climate.

Alternatively, plants may also sense neighbors belowground through for example nutrient depletion zones and chemical exudates. It is unknown if these cues are transferred to the shoot to prepare for upcoming competition or to adjust shoot responses to light cues from neighbors. Finally, abiotic stresses, e.g. drought, occur primarily in the belowground environment and may interact with aboveground light responses. Although little is known about interactions between abiotic stresses and shade responses of the shoot, there is ample information about the interaction of biotic stress and neighbor detection through light cues. The emerging picture is that shade avoidance responses and phytochrome signaling dominate plant defenses against pathogenic microorganisms and against insects (De Wit et al., 2013; Ballaré, 2014), thus

prioritizing light interception. It will be interesting to study if next to biotic stresses, responses against abiotic stresses are suppressed to accommodate shade avoidance responses at high plant density.

### **Concluding remarks**

Plants have a variety of photoreceptors that control many different aspects of plant life, including root development. Recently, several novel mechanisms have been discovered that allow plants to relay information from the shoot and its environment to the root system. These include direct light transmission, hormones and mobile proteins. Although research into these different mechanisms will continue to identify novel mechanisms, it is also crucial to establish the functional implications of this information transfer. It is therefore, pertinent that ecologists, agronomists and plant scientists join forces to unravel both the mechanisms and functional implications of shoot-root communication induced by light cues from the environment.

## Figure Legends

### Figure 1. Relative amounts of photoreceptors in *Arabidopsis* tissues.

The graph displays the relative expression of photoreceptors across seedling tissues. Both gene expression and protein (when data are available) abundance are shown. The box that groups classes of receptors indicates that intensities within can be compared. In the phytochrome box 'L' stands for protein levels in light and 'D' for protein levels in the dark. Source data: For all, BAR eFP browser (<http://bar.utoronto.ca/efp/cgi-bin/efpWeb.cgi>), phytochromes (Somers and Quail, 1995; Goosey et al., 1997; Toth et al., 2001; Sharrock and Clack, 2002; Salisbury et al., 2007), phototropins (Sakamoto and Briggs, 2002; Moni et al., 2015), cryptochromes (Toth et al., 2001), and UVR8 (Rizzini et al., 2011).

### Figure 2. Root growth is affected by light quantity and quality.

(A) 8d old seedlings grown on ½ MS medium in darkness and white light (WL) ( $140\mu\text{M m}^{-2} \text{s}^{-1}$  photosynthetically active radiation, PAR). (B) 8d old seedling grown in WL ( $140\mu\text{M m}^{-2} \text{s}^{-1}$ ) or WL+FR (R:FR 0.1). The arrows point to the root-shoot junctions.

### Figure 3. The aboveground regulation of root development in dense vegetation involves photosynthesis, light piping, hormones and mobile factors.

The primary effect of light is to enable photosynthesis which leads to the production of sugars (sucrose), that enable the root to grow. Photomorphogenic development is associated with the production of auxins in the shoot, which are transported rootward and enable root development. HY5 is stabilized in the shoot during photomorphogenesis and is transported rootwards where it regulates nitrate uptake and root development.

Light is used as a cue to detect neighboring plant competition via sensing of the R:FR ratio. Plant tissues reflect FR, which lowers the nearby R:FR ratio, leading to shade avoidance responses mediated by, amongst others, auxin and GA. These hormones can be transported rootward where they affect root development. There are also indications that shade avoidance responses, in a negative feedback mechanism, decrease the amount of rootward auxin transport. FR light itself can also be transmitted directly through woody, vascular, tissues from the shoot to the root, where it can affect root-localized phytochromes.

480 **Box 2 Figure.** Shade avoidance induced by low R:FR. Carrot (*Daucus carota*) plants grown in white light  
481 (WL; left) and WL+FR (right). Plants were grown in WL or WL+FR for 8 weeks. PAR was  $\sim 120 \mu\text{mol}$   
482  $\text{m}^{-2} \text{s}^{-1}$ , and the FR ratio was 1.8 (WL) and 0.2 (WL+FR). Scale bar = 1 cm.

483





**WL**

**WL+FR**

**BOX2, Figure A. Shade avoidance induced by low R:FR.** Carrot (*Daucus carota*) plants grown in white light (WL; left) and WL+FR (right). Plants were grown in WL or WL+FR for 8 weeks. PAR was  $\sim 120 \mu\text{mol m}^{-2} \text{s}^{-1}$ , and the FR ratio was 1.8 (WL and 0.2 (WL+FR). Scale bar = 1 cm.

## ADVANCES

- New modes of information transfer between the shoot and root stems have been identified, including light piping and the light-sensitive transcription factor HY5.
- Light signaling interacts with root-environment interactions, including nutrient acquisition and gravitropism.
- A vertical agar plate system for Arabidopsis growth, D-root, was developed to keep roots in darkness, while the shoot experiences light. This system has allowed probing direct and indirect light effects on root system architecture.

## OUTSTANDING QUESTIONS

- Do plants use light cues associated with density in vegetation to adjust root development, architecture, and nutrient acquisition physiology to neighbors?
- Do root responses to soil (stress) conditions affect aboveground (architectural) responses to light signals?
- Which mobile signals control root system architecture upon photoreceptor activation by different wavebands in the shoot, and through which tissue(s) do these factors travel towards the root system?
- Is there strong genetic variation for light (quality and quantity) effects on root development, or are these broadly occurring responses throughout the higher plant kingdom?



## Parsed Citations

**Abas L, Benjamins R, Malenica N, Paciorek T, Wiśniewska J, Wirniewska J, Moulinier-Anzola JC, Sieberer T, Friml J, Luschnig C (2006) Intracellular trafficking and proteolysis of the Arabidopsis auxin-efflux facilitator PIN2 are involved in root gravitropism. Nat Cell Biol 8: 249–56**

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

**von Arnim a G, Deng XW (1994) Light inactivation of Arabidopsis photomorphogenic repressor COP1 involves a cell-specific regulation of its nucleocytoplasmic partitioning. Cell 79: 1035–45**

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

**Ballaré CL (2014) Light Regulation of Plant Defense. Annu Rev Plant Biol 65: 335–363**

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

**Ballaré CL, Mazza C a, Austin AT, Pierik R (2012) Canopy light and plant health. Plant Physiol 160: 145–55**

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

**Ballaré CL, Pierik R (2017) The shade-avoidance syndrome: multiple signals and ecological consequences. Plant Cell Environ. doi: 10.1111/pce.12914**

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

**Baster P, Robert S, Kleine-Vehn J, Vanneste S, Kania U, Grunewald W, De Rybel B, Beeckman T, Friml J (2012) SCF(TIR1/AFB)-auxin signalling regulates PIN vacuolar trafficking and auxin fluxes during root gravitropism. EMBO J 32: 260–274**

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

**Bhalerao RP, Eklof J, Ljung K, Marchant A, Bennett M, Sandberg G (2002) Shoot-derived auxin is essential for early lateral root emergence in Arabidopsis seedlings. Plant J 29: 325–332**

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

**Binkert M, Kozma-Bognár L, Terecskei K, De Veylder L, Nagy F, Ulm R, Kozma-Bognar L, Terecskei K, De Veylder L, Nagy F, et al (2014) UV-B-responsive association of the Arabidopsis bZIP transcription factor ELONGATED HYPOCOTYL5 with target genes, including its own promoter. Plant Cell 26: 4200–13**

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

**Blilou I, Xu J, Wildwater M, Willemsen V, Paponov I, Heidstra R, Aida M, Palme K, Scheres B, Friml J, et al (2005) The PIN auxin efflux facilitator network controls growth and patterning in Arabidopsis roots. Nature 433: 39–44**

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

**Boccalandro HE, De Simone SN, Bergmann-Honsberger A, Schepens I, Fankhauser C, Casal JJ (2007) PHYTOCHROME KINASE SUBSTRATE1 Regulates Root Phototropism and Gravitropism. Plant Physiol 146: 108–115**

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

**Briggs WR, Christie JM (2002) Phototropins 1 and 2: Versatile plant blue-light receptors. Trends Plant Sci 7: 204–210**

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

**Buer CS, Muday GK (2004) The transparent testa4 Mutation Prevents Flavonoid Synthesis and Alters Auxin Transport and the Response of Arabidopsis Roots to Gravity and Light. Plant Cell 16: 1191–1205**

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

**Burbach C, Markus K, Zhang Y, Schlicht M, Baluška F (2012) Photophobic behavior of maize roots. Plant Signal Behav 7: 876–880**

Pubmed: [Author and Title](#)  
CrossRef: [Author and Title](#)  
Google Scholar: [Author Only Title Only Author and Title](#)

**Burgie ES, Vierstra RD (2014) Phytochromes: An Atomic Perspective on Photoactivation and Signaling. Plant Cell 1–17**

Pubmed: [Author and Title](#)  
CrossRef: [Author and Title](#)  
Google Scholar: [Author Only Title Only Author and Title](#)

**Canamero RC, Bakrim N, Bouly JP, Garay A, Dudkin EE, Habricot Y, Ahmad M (2006) Cryptochrome photoreceptors cry1 and cry2 antagonistically regulate primary root elongation in Arabidopsis thaliana. Planta 224: 995–1003**

Pubmed: [Author and Title](#)  
CrossRef: [Author and Title](#)  
Google Scholar: [Author Only Title Only Author and Title](#)

**Casal JJ (2013) Photoreceptor signaling networks in plant responses to shade. Annu Rev Plant Biol 64: 403–27**

Pubmed: [Author and Title](#)  
CrossRef: [Author and Title](#)  
Google Scholar: [Author Only Title Only Author and Title](#)

**Chen M, Chory J (2011) Phytochrome signaling mechanisms and the control of plant development. Trends Cell Biol 21: 664–671**

Pubmed: [Author and Title](#)  
CrossRef: [Author and Title](#)  
Google Scholar: [Author Only Title Only Author and Title](#)

**Chen X, Yao Q, Gao X, Jiang C, Harberd NP, Fu X (2016) Shoot-to-Root Mobile Transcription Factor HY5 Coordinates Plant Carbon and Nitrogen Acquisition. Curr Biol 26: 640–646**

Pubmed: [Author and Title](#)  
CrossRef: [Author and Title](#)  
Google Scholar: [Author Only Title Only Author and Title](#)

**Christie JM, Arvai AS, Baxter KJ, Heilmann M, Pratt AJ, O'Hara A, Kelly SM, Hothorn M, Smith BO, Hitomi K, et al (2012) Plant UVR8 photoreceptor senses UV-B by tryptophan-mediated disruption of cross-dimer salt bridges. Science 335: 1492–6**

Pubmed: [Author and Title](#)  
CrossRef: [Author and Title](#)  
Google Scholar: [Author Only Title Only Author and Title](#)

**Cluis CP, Ha C, Mouchel CF, Hardtke CS, Ha C, Mouchel CF, Hardtke CS (2004) The Arabidopsis transcription factor HY5 integrates light and hormone signaling pathways. Plant J 38: 332–347**

Pubmed: [Author and Title](#)  
CrossRef: [Author and Title](#)  
Google Scholar: [Author Only Title Only Author and Title](#)

**Correll MJ, Kiss JZ (2005) The roles of phytochromes in elongation and gravitropism of roots. Plant Cell Physiol 46: 317–323**

Pubmed: [Author and Title](#)  
CrossRef: [Author and Title](#)  
Google Scholar: [Author Only Title Only Author and Title](#)

**Costigan SE, Warnasooriya SN, Humphries BA, Montgomery BL (2011) Root-Localized Phytochrome Chromophore Synthesis Is Required for Photoregulation of Root Elongation and Impacts Root Sensitivity to Jasmonic Acid in Arabidopsis. Plant Physiol 157: 1138–1150**

Pubmed: [Author and Title](#)  
CrossRef: [Author and Title](#)  
Google Scholar: [Author Only Title Only Author and Title](#)

**Demarsy E, Schepens I, Okajima K, Hersch M, Bergmann S, Christie J, Shimazaki K-I, Tokutomi S, Fankhauser C (2012) Phytochrome Kinase Substrate 4 is phosphorylated by the phototropin 1 photoreceptor. EMBO J 31186: 3457–3467**

Pubmed: [Author and Title](#)  
CrossRef: [Author and Title](#)  
Google Scholar: [Author Only Title Only Author and Title](#)

**Duek PD, Elmer M V., Van Oosten VR, Fankhauser C (2004) The degradation of HFR1, a putative bHLH class transcription factor involved in light signaling, is regulated by phosphorylation and requires COP1. Curr Biol 14: 2296–2301**

Pubmed: [Author and Title](#)  
CrossRef: [Author and Title](#)  
Google Scholar: [Author Only Title Only Author and Title](#)

**Dyachok J, Zhu L, Liao F, He J, Huq E, Blancaflor EB (2011) SCAR Mediates Light-Induced Root Elongation in Arabidopsis through Photoreceptors and Proteasomes. Plant Cell 23: 3610–3626**

Pubmed: [Author and Title](#)  
CrossRef: [Author and Title](#)  
Google Scholar: [Author Only Title Only Author and Title](#)

**Fankhauser C, Christie JM (2015) Plant phototropic growth. Curr Biol 25: R384–R389**

Pubmed: [Author and Title](#)  
CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

**Favory J-J, Stec A, Gruber H, Rizzini L, Oravecz A, Funk M, Albert A, Cloix C, Jenkins GI, Oakeley EJ, et al (2009) Interaction of COP1 and UVR8 regulates UV-B-induced photomorphogenesis and stress acclimation in Arabidopsis. EMBO J 28: 591–601**

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

**Franklin KA, Quail PH (2010) Phytochrome functions in Arabidopsis development. J Exp Bot 61: 11–24**

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

**Friml J, Wiśniewska J, Benková E, Mendgen K, Palme K (2002) Lateral relocation of auxin efflux regulator PIN3 mediates tropism in Arabidopsis. Nature 415: 806–9**

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

**Galinha C, Hoffhuis H, Luijten M, Willemsen V, Blilou I, Heidstra R, Scheres B (2007) PLETHORA proteins as dose-dependent master regulators of Arabidopsis root development. Nature 449: 1053–1057**

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

**Galvão VC, Fankhauser C (2015) Sensing the light environment in plants: Photoreceptors and early signaling steps. Curr Opin Neurobiol 34: 46–53**

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

**Gangappa SN, Botto JF (2016) The multifaceted roles of HY5 in plant growth and development. Mol Plant 9: 1353–1365**

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

**Gao J, Shi J, Dong S, Liu P, Zhao B, Zhang J (2017) Grain yield and root characteristics of summer maize (Zea mays L.) under shade stress conditions. J Agron Crop Sci 1–12**

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

**Gommers CMM, Visser EJW, St Onge KR, Voeseek L a CJ, Pierik R (2013) Shade tolerance: when growing tall is not an option. Trends Plant Sci 18: 65–71**

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

**Goosey L, Palecanda L, Sharrock R a (1997) Differential patterns of expression of the Arabidopsis PHYB, PHYD, and PHYE phytochrome genes. Plant Physiol 115: 959–969**

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

**Gundel PE, Pierik R, Mommer L, Ballare CL (2014) Competing neighbors: Light perception and root function. Oecologia 176: 1–10**

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

**Hanzawa T, Shibasaki K, Numata T, Kawamura Y, Gaude T, Rahman A (2013) Cellular Auxin Homeostasis under High Temperature Is Regulated through a SORTING NEXIN1 – Dependent Endosomal Trafficking Pathway. Plant Cell 25: 3424–3433**

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

**Hayes S, Velanis CN, Jenkins GI, Franklin K a. (2014) UV-B detected by the UVR8 photoreceptor antagonizes auxin signaling and plant shade avoidance. Proc Natl Acad Sci U S A 111: 11894–9**

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

**Hedden P, Sponsel V (2015) A Century of Gibberellin Research. J Plant Growth Regul 34: 740–760**

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

**Hedden P, Thomas SG (2012) Gibberellin biosynthesis and its regulation. Biochem J 444: 11–25**

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

**Hisamatsu T, King RW, Helliwell C a, Koshioka M (2005) The involvement of gibberellin 20-oxidase genes in phytochrome-regulated petiole elongation of Arabidopsis. Plant Physiol 138: 1106–1116**

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

**Holm M, Ma L-G, Qu L-J, Deng X-W (2002) Two interacting bZIP proteins are direct targets of COP1-mediated control of light-dependent gene expression in Arabidopsis. Genes Dev 16: 1247–59**

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

**Hornitschek P, Kohnen M V, Lorrain S, Rougemont J, Ljung K, López-Vidriero I, Franco-Zorrilla JM, Solano R, Trevisan M, Pradervand S, et al (2012) Phytochrome interacting factors 4 and 5 control seedling growth in changing light conditions by directly controlling auxin signaling. Plant J 71: 699–711**

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

**Huang L, Zhang HH, Zhang HH, Deng XW, Wei N (2015) HY5 regulates nitrite reductase 1 (NIR1) and ammonium transporter1;2 (AMT1;2) in Arabidopsis seedlings. Plant Sci 238: 330–339**

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

**Huang X, Ouyang X, Deng XW (2014) Beyond repression of photomorphogenesis: role switching of COP/DET/FUS in light signaling. Curr Opin Plant Biol 21C: 96–103**

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

**Huang X, Ouyang X, Yang P, Lau OS, Chen L, Wei N, Deng XW (2013) Conversion from CUL4-based COP1-SPA E3 apparatus to UVR8-COP1-SPA complexes underlies a distinct biochemical function of COP1 under UV-B. Proc Natl Acad Sci U S A 110: 16669–74**

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

**Dello Ioio R, Linhares FS, Scacchi E, Casamitjana-Martinez E, Heidstra R, Costantino P, Sabatini S (2007) Cytokinins Determine Arabidopsis Root-Meristem Size by Controlling Cell Differentiation. Curr Biol 17: 678–682**

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

**Dello Ioio RD, Nakamura K, Moubayidin L, Perilli S, Taniguchi M, Morita MT, Aoyama T, Costantino P, Sabatini S (2008) A Genetic Framework for the Control of Cell Division and Differentiation in the Root Meristem. Science (80- ) 322: 1380–1384**

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

**Jang I-C, Henriques R, Seo HS, Nagatani A, Chua N-H (2010) Arabidopsis PHYTOCHROME INTERACTING FACTOR proteins promote phytochrome B polyubiquitination by COP1 E3 ligase in the nucleus. Plant Cell 22: 2370–83**

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

**Jeong J, Choi G (2013) Phytochrome-interacting factors have both shared and distinct biological roles. Mol Cells 35: 371–380**

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

**Jung J-H, Domijan M, Klose C, Biswas S, Ezer D, Gao M, Khattak AK, Box MS, Charoensawan V, Cortijo S, et al (2016) Phytochromes function as thermosensors in Arabidopsis. Science 354: 886–889**

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

**Kasperbauer MJ, Hunt PG (1992) Root size and shoot/root ratio as influenced by light environment of the shoot. J Plant Nutr 15: 685–697**

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

**Kasperbauer MJ, Hunt PG (1994) Shoot/root assimilate allocation and nodulation of *Vigna unguiculata* seedlings as influenced by shoot light environment. Plant Soil 161: 97–101**

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

**Keuskamp DH, Keller MM, Ballaré CL, Pierik R (2012) Blue light regulated shade avoidance. Plant Signal Behav 7: 514–7**

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

**Keuskamp DH, Pollmann S, Voeselek LACJ, Peeters AJM, Pierik R (2010) Auxin transport through PIN-FORMED 3 (PIN3) controls shade avoidance and fitness during competition. Proc Natl Acad Sci 3: 22740–4**

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

**Kircher S, Schopfer P (2012) Photosynthetic sucrose acts as cotyledon-derived long-distance signal to control root growth during early seedling development in *Arabidopsis*. Proc Natl Acad Sci U S A 109: 11217–21**

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

**Kiss JZ (2003) Phytochromes A and B Mediate Red-Light-Induced Positive Phototropism in Roots. Plant Physiol 131: 1411–1417**

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

**Kiss JZ, Ruppel NJ, Hangarter RP (2001) Phototropism in *Arabidopsis* roots is mediated by two sensory systems. Adv Sp Res 27: 877–885**

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

**Klar T, Pokorny R, Moldt J, Batschauer A, Essen LO (2007) Cryptochrome 3 from *Arabidopsis thaliana*: Structural and Functional Analysis of its Complex with a Folate Light Antenna. J Mol Biol 366: 954–964**

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

**Kleine-Vehn J, Leitner J, Zwiewka M, Sauer M, Abas L, Luschig C, Friml J (2008) Differential degradation of PIN2 auxin efflux carrier by retromer-dependent vacuolar targeting. Proc Natl Acad Sci U S A 105: 17812–7**

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

**Kohnen M, Schmid-Siegert E, Trevisan M, Alenbach Petrolati L, Sénéchal F, Müller-Moulé P, Maloof JN, Xenarios I, Fankhauser C (2016) Neighbor Detection Induces Organ-specific Transcriptomes, Revealing Patterns Underlying Hypocotyl-specific Growth. Plant Cell 28: 1–53**

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

**Koini MA, Avey L, Allen T, Tilley CA, Harberd NP, Whitelam GC, Franklin KA, Le L (2009) High temperature-mediated adaptations in plant architecture require the bHLH transcription factor PIF4. Curr Biol 19: 408–413**

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

**Korbei B, Luschig C (2013) Plasma membrane protein ubiquitylation and degradation as determinants of positional growth in plants. J Integr Plant Biol 55: 809–23**

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

**Kutschera U, Briggs WR (2012) Root phototropism: From dogma to the mechanism of blue light perception. Planta 235: 443–452**

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

**Lavenus J, Goh T, Roberts I, Guyomarc'h S, Lucas M, De Smet I, Fukaki H, Beeckman T, Bennett M, Laplace L (2013) Lateral root development in *Arabidopsis*: fifty shades of auxin. Trends Plant Sci 18: 450–8**

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)



Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

**Laxmi A, Pan J, Morsy M, Chen R (2008) Light plays an essential role in intracellular distribution of auxin efflux carrier PIN2 in *Arabidopsis thaliana*. PLoS One 3: e1510**

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

**Leasure CD, Tong H, Yuen G, Hou X, Sun X, He Z-H (2009) ROOT UV-B SENSITIVE2 Acts with ROOT UV-B SENSITIVE1 in a Root Ultraviolet B-Sensing Pathway. Plant Physiol 150: 1902–1915**

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

**Lee H-J, Ha J-H, Kim S-G, Choi H-K, Kim ZH, Han Y-J, Kim J-I, Oh Y, Fragoso V, Shin K, et al (2016) Stem-piped light activates phytochrome B to trigger light responses in *Arabidopsis thaliana* roots. Sci Signal 9: ra106–ra106**

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

**Lee J, He K, Stolc V, Lee H, Figueroa P, Gao Y, Tongprasit W, Zhao H, Lee I, Deng XW (2007) Analysis of transcription factor HY5 genomic binding sites revealed its hierarchical role in light regulation of development. Plant Cell 19: 731–49**

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

**Legris M, Klose C, Burgie ES, Rojas CCR, Neme M, Hiltbrunner A, Wigge PA, Schäfer E, Vierstra RD, Casal JJ (2016) Phytochrome B integrates light and temperature signals in *Arabidopsis*. Science 354: 897–900**

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

**Leivar P, Monte E (2014) PIFs: Systems Integrators in Plant Development. Plant Cell 26: 56–78**

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

**Leivar P, Quail PH (2011) PIFs: Pivotal components in a cellular signaling hub. Trends Plant Sci 16: 19–28**

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

**Li Q-H, Yang H-Q (2007) Cryptochrome Signaling in Plants†. Photochem Photobiol 83: 94–101**

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

**Lian HL, He SB, Zhang YC, Zhu DM, Zhang JY, Jia KP, Sun SX, Li L, Yang HQ (2011) Blue-light-dependent interaction of cryptochrome 1 with SPA1 defines a dynamic signaling mechanism. Genes Dev 25: 1023–1028**

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

**Liu B, Zuo Z, Liu H, Liu X, Lin C (2011) *Arabidopsis* cryptochrome 1 interacts with SPA1 to suppress COP1 activity in response to blue light. Genes Dev 25: 1029–34**

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

**Lorrain S, Allen T, Duek PD, Whitelam GC, Fankhauser C (2008) Phytochrome-mediated inhibition of shade avoidance involves degradation of growth-promoting bHLH transcription factors. Plant J 53: 312–323**

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

**Mähönen AP, Tusscher K ten, Siligato R, Smetana O, Díaz-Triviño S, Salojärvi J, Wachsman G, Prasad K, Heidstra R, Scheres B (2014) PLETHORA gradient formation mechanism separates auxin responses. Nature 515: 125–9**

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

**Masclaux FG, Bruessow F, Schweizer F, Gouhier-Darimont C, Keller L, Reymond P (2012) Transcriptome analysis of intraspecific competition in *Arabidopsis thaliana* reveals organ-specific signatures related to nutrient acquisition and general stress response pathways. BMC Plant Biol 12: 227**

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

**Michaud O, Fiorucci A-S, Xenarios I, Fankhauser C (2017) Local auxin production underlies a spatially restricted neighbor-detection response in Arabidopsis. Proc Natl Acad Sci 114: 7444–7449**

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

**Mo M, Yokawa K, Wan Y, Baluška F (2015) How and why do root apices sense light under the soil surface? Front Plant Sci 6: 1–8**

**Moni A, Lee AY, Briggs WR, Han IS (2015) The blue light receptor Phototropin 1 suppresses lateral root growth by controlling cell elongation. Plant Biol 17: 34–40**

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

**Morita MT (2010) Directional gravity sensing in gravitropism. Annu Rev Plant Biol 61: 705–20**

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

**Ni W, Xu S-L, Tepperman JM, Stanley DJ, Maltby DA, Gross JD, Burlingame AL, Wang Z-Y, Quail PH (2014) A mutually assured destruction mechanism attenuates light signaling in Arabidopsis. Science 344: 1160–4**

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

**Osterlund MT, Hardtke CS, Wei N, Deng XW (2000) Targeted destabilization of HY5 during light-regulated development of Arabidopsis. Nature 405: 462–6**

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

**Oyama T, Shimura Y, Okada K (1997) The Arabidopsis HY5 gene encodes a bZIP protein that regulates stimulus- induced development of root and hypocotyl. Genes Dev 11: 2983–2995**

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

**Pacin M, Legris M, Casal JJ (2014) Rapid Decline in Nuclear COSTITUTIVE PHOTOMORPHOGENESIS1 Abundance Anticipates the Stabilization of Its Target ELONGATED HYPOCOTYL5 in the Light. Plant Physiol 164: 1134–1138**

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

**Paik I, Kathare PK, Kim J-I, Huq E (2017) Expanding Roles of PIFs in Signal Integration from Multiple Processes. Mol Plant. doi: 10.1016/j.molp.2017.07.002**

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

**Pantazopoulou CK, Bongers FJ, Kopers JJ, Reinen E, Das D, Evers JB, Anten NPR, Pierik R (2017) Neighbor detection at the leaf tip adaptively regulates upward leaf movement through spatial auxin dynamics. Proc Natl Acad Sci 114: 201702275**

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

**Pedmale U V, Liscum E (2007) Regulation of phototropic signaling in Arabidopsis via phosphorylation state changes in the phototropin 1-interacting protein NPH3. J Biol Chem 282: 19992–20001**

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

**Pedmale U V., Huang SC, Zander M, Cole BJ, Hetzel J, Ljung K, Reis PAB, Sridevi P, Nito K, Nery JR, et al (2016) Cryptochromes Interact Directly with PIFs to Control Plant Growth in Limiting Blue Light. Cell 164: 233–245**

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

**Péret B, Middleton AM, French AP, Larrieu A, Bishopp A, Njo M, Wells DM, Porco S, Mellor N, Band LR, et al (2013) Sequential induction of auxin efflux and influx carriers regulates lateral root emergence. Mol Syst Biol 9: 699**

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

**Petricka JJ, Winter CM, Benfey PN (2012) Control of Arabidopsis Root Development. Annu Rev Plant Biol 63: 563–590**

Copyright © 2017 American Society of Plant Biologists. All rights reserved.

Pubmed: [Author and Title](#)  
CrossRef: [Author and Title](#)  
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

**Pierik R, Mommer L, Voesenek LA (2013) Molecular mechanisms of plant competition: Neighbour detection and response strategies. *Funct Ecol* 27: 841–853**

Pubmed: [Author and Title](#)  
CrossRef: [Author and Title](#)  
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

**Reed RC, Brady SR, Muday GK (1998) Inhibition of auxin movement from the shoot into the root inhibits lateral root development in *Arabidopsis*. *Plant Physiol* 118: 1369–1378**

Pubmed: [Author and Title](#)  
CrossRef: [Author and Title](#)  
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

**Regnault T, Davière J-M, Wild M, Sakvarelidze-Achard L, Heintz D, Carrera Bergua E, Lopez Diaz I, Gong F, Hedden P, Achard P (2015) The gibberellin precursor GA12 acts as a long-distance growth signal in *Arabidopsis*. *Nat Plants* 1: 15073**

Pubmed: [Author and Title](#)  
CrossRef: [Author and Title](#)  
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

**Rizzini L, Favory JJ, Cloix C, Faggionato D, O'Hara A, Kaiserli E, Baumeister R, Schafer E, Nagy F, Jenkins GI, et al (2011) Perception of UV-B by the *Arabidopsis* UVR8 protein. *Science* (80- ) 332: 103–106**

Pubmed: [Author and Title](#)  
CrossRef: [Author and Title](#)  
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

**Ruppel NJ, Hangarter RP, Kiss JZ (2001) Red-light-induced positive phototropism in *Arabidopsis* roots. *Planta* 212: 424–430**

Pubmed: [Author and Title](#)  
CrossRef: [Author and Title](#)  
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

**De Rybel B, Vassileva V, Parizot B, Demeulenaere M, Grunewald W, Audenaert D, Van Campenhout J, Overvoorde P, Jansen L, Vanneste S, et al (2010) A novel aux/IAA28 signaling cascade activates GATA23-dependent specification of lateral root founder cell identity. *Curr Biol* 20: 1697–706**

Pubmed: [Author and Title](#)  
CrossRef: [Author and Title](#)  
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

**Saijo Y, Sullivan J a, Wang H, Yang J, Shen Y, Rubio V, Ma L, Hoecker U, Deng XW (2003) The COP1-SPA1 interaction defines a critical step in phytochrome A-mediated regulation of HY5 activity. *Genes Dev* 17: 2642–7**

Pubmed: [Author and Title](#)  
CrossRef: [Author and Title](#)  
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

**Sakamoto K, Briggs WR (2002) Cellular and subcellular localization of phototropin 1. *Plant Cell* 14: 1723–1735**

Pubmed: [Author and Title](#)  
CrossRef: [Author and Title](#)  
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

**Salisbury FJ, Hall A, Grierson CS, Halliday KJ, Buildings K, Road M, Building B, Street C (2007) Phytochrome coordinates *Arabidopsis* shoot and root development. *Plant J* 50: 429–38**

Pubmed: [Author and Title](#)  
CrossRef: [Author and Title](#)  
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

**Sassi M, Lu Y, Zhang Y, Wang J, Dhonukshe P, Blilou I, Dai M, Li J, Gong X, Jaillais Y, et al (2012) COP1 mediates the coordination of root and shoot growth by light through modulation of PIN1- and PIN2-dependent auxin transport in *Arabidopsis*. *Development* 139: 3402–12**

Pubmed: [Author and Title](#)  
CrossRef: [Author and Title](#)  
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

**Scheres B, Wolkenfelt H, Willemsen V, Terlouw M, Lawson E, Dean C, Weisbeek P (1994) Embryonic origin of the *Arabidopsis* primary root and root meristem initials. *Development* 2487: 2475–2487**

Pubmed: [Author and Title](#)  
CrossRef: [Author and Title](#)  
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

**Seo HS, Watanabe E, Tokutomi S, Nagatani A, Chua NH (2004) Photoreceptor ubiquitination by COP1 E3 ligase desensitizes phytochrome A signaling. *Genes Dev* 18: 617–622**

Pubmed: [Author and Title](#)  
CrossRef: [Author and Title](#)  
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)



Seo HS, Yang J-Y, Ishikawa M, Bolle C, Ballesteros ML, Chua N-H (2003) LAF1 ubiquitination by COP1 controls photomorphogenesis and is stimulated by SPA1. *Nature* 423: 995–999

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Sharrock RA, Clack T (2002) Patterns of expression and normalized levels of the five Arabidopsis phytochromes. *Plant Physiol* 130: 442–56

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Sheerin DJ, Menon C, zur Oven-Krockhaus S, Enderle B, Zhu L, Johnen P, Schleifenbaum F, Stierhof Y-D, Huq E, Hiltbrunner A (2015) Light-activated phytochrome A and B interact with members of the SPA family to promote photomorphogenesis in Arabidopsis by reorganizing the COP1/SPA complex. *Plant Cell* 27: 189–201

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Sibout R, Sukumar P, Hettiarachchi C, Holm M, Muday GK, Hardtke CS (2006) Opposite root growth phenotypes of hy5 versus hy5 hyh mutants correlate with increased constitutive auxin signaling. *PLoS Genet* 2: e202

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Silva-Navas J, Moreno-Risueno MA, Manzano C, Pallero-Baena M, Navarro-Neila S, Tellez-Robledo B, Garcia-Mina JM, Baigorri R, Gallego FJ, Del Pozo JC (2015) D-Root: A system for cultivating plants with the roots in darkness or under different light conditions. *Plant J* 84: 244–255

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Silva-Navas J, Moreno-Risueno MA, Manzano C, Tellez-Robledo B, Navarro-Neila S, Carrasco V, Pollmann S, Gallego FJ, del Pozo JC (2016) Flavonols Mediate Root Phototropism and Growth through Regulation of Proliferation-to-Differentiation Transition. *Plant Cell* 28: 1372–1387

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

De Smet I, Tetsumura T, De Rybel B, Frey NF d., Laplace L, Casimiro I, Swarup R, Naudts M, Vanneste S, Audenaert D, et al (2007) Auxin-dependent regulation of lateral root positioning in the basal meristem of Arabidopsis. *Development* 134: 681–690

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Smith H (1982) Light Quality, Photoperception, and Plant Strategy. *Annu Rev Plant Physiol* 481–518

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Somers DE, Quail PH (1995) Temporal and spatial expression patterns of PHYA and PHYB genes in Arabidopsis. *Plant J* 7: 413–427

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Song S-H, Dick B, Penzkofer A, Pokorny R, Batschauer A, Essen L-O (2006) Absorption and fluorescence spectroscopic characterization of cryptochrome 3 from Arabidopsis thaliana. *J Photochem Photobiol B* 85: 1–16

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Stepanova AN, Robertson-Hoyt J, Yun J, Benavente LM, Xie D-Y, Dolezal K, Schlereth A, Jürgens G, Alonso JM (2008) TAA1-mediated auxin biosynthesis is essential for hormone crosstalk and plant development. *Cell* 133: 177–91

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Sun Q, Yoda K, Suzuki H (2005) Internal axial light conduction in the stems and roots of herbaceous plants. *J Exp Bot* 56: 191–203

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Sun Q, Yoda K, Suzuki M, Suzuki H (2003) Vascular tissue in the stem and roots of woody plants can conduct light. *J Exp Bot* 54: 1627–1635

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)  
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

**Suzuki H, Yokawa K, Nakano S, Yoshida Y, Fabrisin I, Okamoto T, Baluska F, Koshiba T (2016) Root cap-dependent gravitropic U-turn of maize root requires light-induced auxin biosynthesis via the YUC pathway in the root apex. J Exp Bot 67: 4581–4591**

Pubmed: [Author and Title](#)  
CrossRef: [Author and Title](#)  
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

**Sweere U (2001) Interaction of the Response Regulator ARR4 with Phytochrome B in Modulating Red Light Signaling. Science (80- ) 294: 1108–1111**

Pubmed: [Author and Title](#)  
CrossRef: [Author and Title](#)  
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

**Tong H, Leasure CD, Hou X, Yuen G, Briggs W, He Z-H (2008) Role of root UV-B sensing in Arabidopsis early seedling development. Proc Natl Acad Sci U S A 105: 21039–44**

Pubmed: [Author and Title](#)  
CrossRef: [Author and Title](#)  
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

**Toth R, Kevei E, Hall A, Millar AJ, Nagy F, Kozma-Bognar L (2001) Circadian Clock-Regulated Expression of Phytochrome and Cryptochrome Genes in Arabidopsis. Plant Physiol 127: 1607–1616**

Pubmed: [Author and Title](#)  
CrossRef: [Author and Title](#)  
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

**Vermeer JEM, von Wangenheim D, Barberon M, Lee Y, Stelzer EHK, Maizel A, Geldner N (2014) A spatial accommodation by neighboring cells is required for organ initiation in Arabidopsis. Science 343: 178–83**

Pubmed: [Author and Title](#)  
CrossRef: [Author and Title](#)  
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

**Wan Y, Jasik J, Wang L, Hao H, Volkmann D, Menzel D, Mancuso S, Baluska F, Lin J (2012) The Signal Transducer NPH3 Integrates the Phototropin1 Photosensor with PIN2-Based Polar Auxin Transport in Arabidopsis Root Phototropism. Plant Cell 24: 551–565**

Pubmed: [Author and Title](#)  
CrossRef: [Author and Title](#)  
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

**Weiner J (1985) Size hierarchies in experimental populations of annual plants. Ecology 66: 743–752**

Pubmed: [Author and Title](#)  
CrossRef: [Author and Title](#)  
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

**Weiner J, Thomas SC (1986) Size variability and competition in plant monocultures. Oikos 47: 211–222**

Pubmed: [Author and Title](#)  
CrossRef: [Author and Title](#)  
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

**de Wit M, Keuskamp DH, Bongers FJ, Hornitschek P, Gommers CMM, Reinen E, Martinez-Ceron C, Fankhauser C, Pierik R (2016) Integration of Phytochrome and Cryptochrome Signals Determines Plant Growth during Competition for Light. Curr Biol 26: 3320–3326**

Pubmed: [Author and Title](#)  
CrossRef: [Author and Title](#)  
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

**De Wit M, Spoel SH, Sanchez-Perez GF, Gommers CMM, Pieterse CMJ, Voeselek L a CJ, Pierik R (2013) Perception of low red: Far-red ratio compromises both salicylic acid- and jasmonic acid-dependent pathogen defences in Arabidopsis. Plant J 75: 90–103**

Pubmed: [Author and Title](#)  
CrossRef: [Author and Title](#)  
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

**Wu S-H (2014) Gene expression regulation in photomorphogenesis from the perspective of the central dogma. Annu Rev Plant Biol 65: 311–33**

Pubmed: [Author and Title](#)  
CrossRef: [Author and Title](#)  
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

**Xu W, Ding G, Yokawa K, Baluška F, Li Q-F, Liu Y, Shi W, Liang J, Zhang J (2013) An improved agar-plate method for studying root growth and response of Arabidopsis thaliana. Sci Rep 3: 1273**

Pubmed: [Author and Title](#)  
CrossRef: [Author and Title](#)  
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

**Xu X, Paik I, Zhu L, Huq E (2015) Illuminating Progress in Phytochrome-Mediated Light Signaling Pathways. Trends Plant Sci 20: 641–650**

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

**Zhang K-X, Xu H-H, Gong W, Jin Y, Shi Y-Y, Yuan T-T, Li J, Lu Y-T (2014) Proper PIN1 Distribution Is Needed for Root Negative Phototropism in Arabidopsis. PLoS One 9: e85720**

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

**Zhang K-X, Xu H-H, Yuan T-T, Zhang L, Lu Y-T (2013) Blue-light-induced PIN3 polarization for root negative phototropic response in Arabidopsis. Plant J 76: 308–321**

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

**Zhang Y, Li C, Zhang J, Wang J, Yang J, Lv Y, Yang N, Liu J, Wang X, Palfalvi G, et al (2017) Dissection of HY5/HYH expression in Arabidopsis reveals a root-autonomous HY5-mediated photomorphogenic pathway. PLoS One 12: e0180449**

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

**Zheng X, Wu S, Zhai H, Zhou P, Song M, Su L, Xi Y, Li Z, Cai Y, Meng F, et al (2013) Arabidopsis phytochrome B promotes SPA1 nuclear accumulation to repress photomorphogenesis under far-red light. Plant Cell 25: 115–33**

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

**Zuo Z, Liu H, Liu B, Liu X, Lin C (2011) Blue light-dependent interaction of CRY2 with SPA1 regulates COP1 activity and floral initiation in Arabidopsis. Curr Biol 21: 841–7**

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)