

Environmental and mucosal microbiota and their role in childhood asthma

L. T. Birzele^{1,*}, M. Depner^{1,*}, M. J. Ege^{1,2}, M. Engel^{3,4}, S. Kublik⁴, C. Bernau⁵, G. J. Loss^{1,6}, J. Genuneit⁷, E. Horak⁸, M. Schloter⁴, C. Braun-Fahländer^{9,10}, H. Danielewicz¹¹, D. Heederik¹², E. von Mutius^{1,2} & A. Legatzki¹

¹Dr. von Hauner Children's Hospital, LMU Munich, Munich, Germany; ²Member of the German Center for Lung Research (DZL), Munich, Germany; ³Research Unit Scientific Computing, Helmholtz Center Munich, Neuherberg, Germany; ⁴Research Unit for Environmental Genomics, Helmholtz Center Munich, Neuherberg, Germany; ⁵Leibniz Supercomputing Center of the Bavarian Academy of Sciences and Humanities, Garching, Germany; ⁶Department of Pediatrics, School of Medicine, University of California, San Diego, CA, USA; ⁷Institute of Epidemiology and Medical Biometry, Ulm University, Ulm, Germany; ⁸Division of Cardiology and Pulmonology, Department of Pediatrics and Adolescents, Innsbruck Medical University, Innsbruck, Austria; ⁹Swiss Tropical and Public Health Institute, Basel, Switzerland; ¹⁰Department of Epidemiology and Public Health, University of Basel, Basel, Switzerland; ¹¹Department of Pediatrics, Allergology and Cardiology, Wrocław Medical University, Wrocław, Poland; ¹²Division of Environmental Epidemiology, Institute for Risk Assessment Sciences, University of Utrecht, Utrecht, The Netherlands

To cite this article: Birzele LT, Depner M, Ege MJ, Engel M, Kublik S, Bernau C, Loss GJ, Genuneit J, Horak E, Schloter M, Braun-Fahländer C, Danielewicz H, Heederik D, von Mutius E, Legatzki A. Environmental and mucosal microbiota and their role in childhood asthma. *Allergy* 2017; **72**: 109–119.

Keywords

asthma; epidemiology; microbiome; pediatrics.

Correspondence

Martin Depner, PhD, Dr von Hauner Children's Hospital, LMU Munich, Lindwurmstrasse 4, 80337 Munich, Germany.

Tel.: +49 89 4400 52776

Fax: +49 89 4400 54452

E-mail: martin.depner@med.uni-muenchen.de

*Both authors contributed equally to this study.

Accepted for publication 6 August 2016

DOI:10.1111/all.13002

Edited by: Hans-Uwe Simon

Abstract

Background: High microbial diversity in the environment has been associated with lower asthma risk, particularly in children exposed to farming. It remains unclear whether this effect operates through an altered microbiome of the mucosal surfaces of the airways.

Methods: DNA from mattress dust and nasal samples of 86 school age children was analyzed by 454 pyrosequencing of the 16S rRNA gene fragments. Based on operational taxonomic units (OTUs), bacterial diversity and composition were related to farm exposure and asthma status.

Results: Farm exposure was positively associated with bacterial diversity in mattress dust samples as determined by richness ($P = 8.1 \times 10^{-6}$) and Shannon index ($P = 1.3 \times 10^{-5}$). Despite considerable agreement of richness between mattress and nasal samples, the association of richness with farming in nasal samples was restricted to a high gradient of farm exposure, that is, exposure to cows and straw vs no exposure at all. In mattress dust, the genera *Clostridium*, *Facklamia*, an unclassified genus within the family of *Ruminococcaceae*, and six OTUs were positively associated with farming. Asthma was inversely associated with richness [aOR = 0.48 (0.22–1.02)] and Shannon index [aOR = 0.41 (0.21–0.83)] in mattress dust and to a lower extent in nasal samples [richness aOR 0.63 = (0.38–1.06), Shannon index aOR = 0.66 (0.39–1.12)].

Conclusion: The stronger inverse association of asthma with bacterial diversity in mattress dust as compared to nasal samples suggests microbial involvement beyond mere colonization of the upper airways. Whether inhalation of metabolites of environmental bacteria contributes to this phenomenon should be the focus of future research.

Abbreviations

16S rRNA, 16S ribosomal RNA; aOR, adjusted odds ratio; CI, confidence interval; GABRIELA, multidisciplinary study to identify the Genetic and Environmental Causes of Asthma in the European Community—Advanced Studies; GMR, geometric mean ratio; NMDS, nonmetric multidimensional scaling; OTU, operational taxonomic unit; SSCP, single-strand conformation polymorphism.

Living on a working farm is characterized by intensive contact of the farmers and their families with animals and plants. The farm environment offers various specific habitats and niches for microbes; for example, animals, manure, feeding, and bedding material are responsible for a higher microbial diversity at farms compared with nonfarm environments (1). Also the indoor environment of farm dwellings is

characterized by a higher microbial diversity as shown in mattress dust samples (2, 3). It remains, however, unclear whether the environmental microbiome affects the nasal microbiome. A key function of the nose is filtering inhaled air (4). Therefore, the nose may harbor microbes originating from the surrounding environment in addition to its local microflora.

A high diversity of environmental microorganisms has been found to be inversely associated with asthma in rural populations (5). Growing up on traditional farms consistently showed a reduced risk of childhood asthma (reviewed in Ref. 6, 7). Different farm exposures such as contact with cows and straw, which obviously harbor a set of different microbes, have been identified as protective factors for asthma (8). Furthermore, several studies have demonstrated an association between specific environmental microbes or their cell wall components and a reduced risk of allergic diseases including childhood asthma (2, 9, 10). In addition, experimental mouse studies suggest protective effects of several bacterial strains isolated from the farm environment (11–13).

The aim of this study was a comprehensive analysis of the bacterial exposure related to farming using a cultivation-independent approach based on directly extracted DNA and high-throughput sequencing of amplified 16S rRNA gene fragments.

Diversity and composition of bacterial microbiota of mattress dust and nasal swab samples of the same children were characterized and related to farming and asthma status.

Methods

Study design and subjects

The cross-sectional multidisciplinary study to identify the Genetic and Environmental Causes of Asthma in the European Community – Advanced Studies (GABRIELA) included children 6–12 years of age from five rural areas in Europe (8) and was approved by the ethics committees of the participating universities and the regional data protection authorities.

From the Austrian arm of the GABRIELA study, a stratified random sample of 102 children was drawn for microbial analyses (Fig. S1). Children living on a farm run by the family were defined as farm children. Nonfarm children were defined as children not living on a farm and not exposed to farm environments or cow's milk directly bought from a farm (8, 14). As a more specific farm-related exposure, contact to cow and straw during the last 12 months *vs* no contact to either was defined.

Asthma was defined as either parent-reported wheeze during the last 12 months, a positive answer to the question 'Did your child ever use an asthma spray?' or a doctor's diagnosis of asthma at least once or of wheezy bronchitis more than once (8, 14). Medication was based as intake of any medication in the 4 weeks prior to nasal sampling.

Sample collection and processing

Mattress dust was collected by the parents of the participating children using a standardized dust collection protocol (see Supporting information) (15). Nasal samples were

collected by trained field workers using sterile dry cotton-headed swabs (MASTASWAB MD 559; MAST Diagnostica GmbH, Reinfeld, Germany). The samples were collected between May and July 2007 (Fig. S2). The two different specimens from each child were obtained on average within 2 days. After arrival at the study center, the samples were stored at -80°C until further processing.

Mattress dust DNA was extracted from 30 mg of dust using the FastDNA SPIN Kit for Soil (MP Biomedicals GmbH, Eschwege, Germany) including a bead beating step according to the protocol of the manufacturer.

Nasal swab DNA was extracted with the QIAmp DNA Mini Kit (Qiagen GmbH, Hilden, Germany) using an adapted protocol (16). At the beginning of this procedure, a 20-s bead beating step with 50 mg of acid-washed UV-treated glass beads (Sigma-Aldrich GmbH, Taufkirchen, Germany) was included. With every extraction run, a negative control was processed. Extracted DNA was stored at -20°C until further processing.

The V3–V5 region of the bacterial 16S rRNA gene was amplified with the primers 357F (5'-CCTACGGGAGGCAG CAG-3') and 926R (5'-CCGTCAATTCMTTTRAGT-3') (17), tagged with sequencing adapters and barcodes (18) using the Fast Start High Fidelity PCR System (Roche Diagnostics Deutschland GmbH, Mannheim, Germany) (see Supporting information). After purification of the amplicons with Agencourt AMPure XP beads (Beckman Coulter GmbH, Krefeld, Germany), concentrations were determined using Quant-iT PicoGreen (Thermo Fisher Scientific GmbH, Dreieich, Germany). Subsequently, the amplicons were pooled for sequencing at equimolar concentrations.

Unidirectional sequencing of the 16S rRNA gene fragments starting from the reverse primer 926R was performed using 454-GS FLX Titanium (Roche Diagnostics Deutschland GmbH) protocol. Raw data were processed by the 454 GS Run Processor v2.8 for shotgun reads. Denoising and removal of chimera was achieved by AmpliconNoise (19), resulting in about 400-bp-long fragments. Following, files from all mattress dust and nasal samples were concatenated and sequences clustered together in operational taxonomic units (OTUs) using UCLUST (20) applying the *de novo* picking method in QIIME version 1.8 (21). Operational taxonomic units were defined as clusters of the respective 16S rRNA sequences with at least 97% sequence similarity. For taxonomy assignment, OTUs were aligned against the SILVA database version 111 NR. Operational taxonomic units classified as *Chloroplast* were excluded from further analysis as well as singletons.

Negative extraction controls were performed for both extraction protocols and evaluated (see Table S1). A negligible read count close to zero validated the experiment. Following filtering, children with <1000 sequences in either the mattress dust or nasal samples were excluded, resulting in 86 children (Fig. S1). Unclassified sequences were excluded from downstream analyses on phylum and genus levels.

Data analysis

Alpha diversity, that is the diversity of the community within one sample, was calculated as number of OTUs (richness). In

addition, the Shannon index was calculated, which combines information on number and proportional abundance of taxa.

Differences in α -diversity between mattress dust and nasal swab were calculated by geometric mean ratios, and agreement between richness in mattress dust and the nose on an individual level was assessed by a Bland–Altman plot.

Descriptive analyses were based on relative abundance of the main taxa, which were defined as being represented in at least 1% of all reads of all children. The remaining taxa (<1% relative abundance) were combined in one variable termed 'rare group'. In addition, all samples were rarefied to 1252 sequences, the minimum sequence number of all mattress dust and nasal swab samples. This process was performed in 1000 replicates per sample and subsequently averaged for α -diversity analysis. α -diversity indices such as species richness and Shannon index were calculated with the R package *vegan* (22). Survey-weighted Wilcoxon tests were used to compare α -diversity between exposure groups. Moreover, presence of common genera and OTUs prevalent in at least 10% of the children was defined as the occurrence of the respective taxon in the rarefied sample.

To assess β -diversity, that is, the diversity of the community between samples, we used the UniFrac measure (23) representing dissimilarity on a phylogenetic level and the Bray–Curtis dissimilarity index on a nonphylogenetic level. Survey-weighted regression models were used to relate differences in relative abundances or presence of single taxa to farm exposure and asthma status. Association analyses with asthma are adjusted for farming if not further specified. Correction for multiple comparisons was made by the Bonferroni method or the false discovery rate (FDR) as indicated. A mediation analysis was performed by multiplying the effect estimates of the single paths and tested for significance on the basis of a bootstrapped standard error.

All statistical analyses were performed with R 3.02 (24) and the *phyloseq* package (25) if not further specified. Survey-weighted analyses were applied to account for the stratified sampling design of the GABRIELA study (8, 14) by the R package 'survey' (<https://cran.r-project.org/web/packages/survey/index.html>).

Results

Study population

The analyzed study population of 86 children did not differ from the original Austrian part of the GABRIELA study population with respect to farm exposure, asthma, and general demographic variables except for parental education (Table S2).

Bacterial community composition in mattress dust and nasal samples

Of all 1 730 479 reads, 212 030 reads from mattress dust and 29 179 reads from nasal samples aligned to *Chloroplast* sequences with one predominant OTU (>90% abundance in both sample types) being classified within the plant family

Pinaceae. These were considered contaminants and excluded leaving 618 635 reads from mattress dust and 870 635 reads from nasal samples for further analysis.

The mean number of reads was 7193 for mattress dust and 10 120 for nasal samples (Fig. S3). Rarefaction analysis showed an adequate saturation except for richness in mattress dust samples (Fig. S4).

The two sampling sources separated well by phylogenetic and nonphylogenetic distance measures (Fig. S5). On the phylum level, 0.48% of the mattress dust reads and 0.05% of nasal reads were unclassified; on the genus level, the proportions were slightly higher (2.72% and 0.74%, respectively).

Bacterial mattress dust microbiota were dominated by the phyla *Firmicutes* (53.2%), *Proteobacteria* (16.8%), *Actinobacteria* (16.0%), and *Bacteroidetes* (10.9%), whereas in nasal samples *Proteobacteria* (44.8%), *Firmicutes* (37.1%), *Actinobacteria* (12.3%), and *Bacteroidetes* (4.1%) prevailed (Fig. 1A). On the genus level, 23 main genera (genera with $\geq 1\%$ relative abundance on average) were observed in mattress dust, led by *Streptococcus* with 15.5% (Fig. 1B). For the nasal samples, 12 main genera were detected (Fig. 1B) with *Moraxella* prevailing at 34.2%. Common to both sample sources were the genera *Streptococcus*, *Staphylococcus*, *Corynebacterium*, *Propionibacterium*, *Prevotella*, *Peptoniphilus*, *Neisseria*, *Actinobacillus*, *Porphyromonas*, and *Moraxella*. Most of the other main genera detected in the mattress dust were also present in the nose, however at frequencies below 1% on average. These genera defined as 'rare' accounted for 32.9% in mattress dust and 14.7% in the nose. The corresponding variable correlated with richness (mattress dust, $r = 0.77$; nose, $r = 0.89$). Mattress dust showed a higher α -diversity compared with nasal samples (richness: GMR = 3.42; Shannon index GMR = 2.56). On an individual level, richness values of mattress dust and nasal samples agreed reasonably (Fig. 2).

Farm exposure and bacterial communities

As already suggested by Fig. 2, α -diversity was significantly higher in mattress dust of farm children compared with non-farm children (richness $P = 8.1 \times 10^{-6}$; Shannon index $P = 1.3 \times 10^{-5}$, Fig. 3A). Similarly, exposure to cow and straw was associated with higher bacterial α -diversity in mattress dust (Fig. 3B). In nasal samples, α -diversity was not associated with farming but with exposure to cow and straw (Fig. 3A,B).

In mattress dust, farming was significantly associated with relative abundance of the 'rare' genera, a not further classified genus of the family *Ruminococcaceae* (Fig. 4A) and six OTUs with sequence similarities to taxa previously related to cow feces (Table 1). In addition, farming was associated with the presence of the genera *Clostridium* and *Facklamia* and one OTU (OTU2597) belonging to the family of *Ruminococcaceae* (Table 1). For exposure to cow and straw, similar associations emerged (Fig. 4B, Table S3).

For the nasal samples, no significant association of farm exposures with relative abundance was observed except for a weak association of *Streptococcus* with cow/straw exposure (Fig. 4). When not correcting for multiple testing, farming

was associated with the presence of the genus *Prevotella* [aOR = 4.84 (1.39–16.89), $P = 0.016$], an unclassified genus of the *Ruminococcaceae* family [aOR = 4.05 (1.22–13.43), $P = 0.025$], and the *Ruminococcaceae* OTU 2597 [aOR = 7.57 (1.94–29.55), $P = 4.6 \times 10^{-3}$]. Also with respect to its relative abundance, this OTU was related to farming [$\beta = 0.37$ (0.14–0.59), $P = 1.9 \times 10^{-3}$], thereby mirroring the associations observed in mattress dust.

Asthma and bacterial communities

Asthma was inversely associated with richness and Shannon index in mattress dust [aOR = 0.48 (0.22–1.02), $P = 0.060$; aOR = 0.41 (0.21–0.83), $P = 0.015$, respectively] and nasal samples [aOR = 0.63 (0.38–1.06), $P = 0.087$; aOR = 0.66 (0.39–1.12), $P = 0.129$, respectively]; the effects of nasal and mattress diversity were independent of each other (Fig. 5) and not confounded by medication or atopy status (Table S4). Exclusion of atopic controls increased the effect size only in nasal swabs (Table S4).

Richness in mattress dust ($P = 0.028$), but not in nasal swabs ($P = 0.430$), mediated the protective farm effect on asthma as shown by the indirect effect of the path from farm on asthma via richness.

Among specific genera, inverse associations of asthma with the relative abundance of bacteria belonging to the genus *Prevotella* in the nasal swabs [aOR = 0.44 (0.21–0.93), $P = 0.0345$] and with the rare group (<1% relative abundance) in mattress dust [aOR = 0.43 (0.22–0.87), $P = 0.0216$] emerged, although the latter might be a proxy for bacterial diversity.

A presence vs absence analysis on genus and OTU level revealed several candidates for asthma protection both in mattress dust and nasal samples (Table S5). All significant hits for the specific taxa require confirmation in larger population as they were above the FDR.

Discussion

This analysis revealed differences in the microbial composition and diversity between mattress dust and nasal samples.




Figure 1 Taxonomic composition of the mattress dust and nasal microbiota on (A) phyla level and (B) genera level. *Genus is unclassified, and only family is reported.




Figure 2 Bland-Altman plot for bacterial richness of mattress dust and nasal microbiota. Richness is shown as mean of 1000 rarefying steps to 1252 reads per subject.




Figure 3 Bacterial richness and Shannon index in the rarefied sample of mattress dust and nasal microbiota stratified for (A) farming or (B) regular exposure to cow/straw. Shown are the P-values of survey-weighted Wilcoxon tests.

Despite agreement of richness between mattress and nasal samples on an individual level, growing up on a farm was related to bacterial diversity and specific taxa indicative of animal exposure primarily in mattress dust samples. In contrast, asthma was associated with bacterial diversity in both mattress dust and nasal samples. In mattress dust, bacterial richness operated as an intermediate factor in the association of farm exposure with asthma.

Indoor dust samples in general are a mix of microbial species originating from the house occupants and from the

outside environment (26–28). The main genera from our analysis, those are, *Staphylococcus*, *Propionibacterium*, *Corynebacterium*, and *Streptococcus*, are typical for human skin (29, 30). Most likely, the house occupants shed these microbes throughout the indoor environment including the mattresses, where they eventually were detected. There is also strong evidence for outdoor environmental bacteria in the mattress dust samples such as *Sphingomonas*, which is known from samples of soil, water, and sediments (31). Further, *Acinetobacter* and *Lactococcus* were previously detected in




Figure 4 Effects of (A) farming and (B) regular exposure to cow/straw on relative abundance of the main bacterial genera in mattress dust and nasal samples. In case of unclassified genus, family is reported, and this is indicated by (F).

raw milk (32, 33) and isolated from cow sheds (11). We also detected a *Chloroplast* OTU related to the plant family *Pinaceae*, a group of trees and scrubs native to northern temperate regions. As *Chloroplast* DNA of *Pinaceae* is inherited via pollen (34) and our sample collection occurred during the pollen season of *Pinaceae* (35), we take this OTU as evidence for transfer of particles from the outside environment to the mattress dust.

Although this *Pinaceae Chloroplast* OTU was occasionally detected in nasal samples as well, the main nasal taxa were local bacteria such as *Moraxella* and *Streptococcus*, which are well known from clinical experience and previous studies (36–38), particularly in children (38).

This predominance of local bacteria may explain the less pronounced signature of environmental microbes in nasal samples. Nevertheless, mattress dust and nasal samples

agreed reasonably with respect to bacterial richness as illustrated by Fig. 2 although mucosal samples are characterized by a large interindividual variability (39). Nasal samples also reflect other exposures beyond the home environment; however, the frequent clearance of the nose may result in a shorter time of residence of airborne bacteria, thereby constraining the effect of the environmental exposure quantitatively. This may explain the generally less pronounced associations of nasal microbiota with environmental determinants in the present analysis.

In contrast, mattress dust is a long-term reservoir for settled material reflecting continuous exposure also to an outer environment. Indeed, transfer of bacteria from animal sheds to the children’s mattresses was shown before (3). Bacterial dispersal from the outdoor to the indoor environment by air-flow or direct transport via the child and other family

Table 1 Genera and operational taxonomic units (OTUs) from mattress dust significantly associated with farming. Listed are only associations significant ($P < 0.05$) after Bonferroni correction

Taxa (classification)		Effect	P-value	Sample site of BLAST hit*
Relative abundance	Genus	β (95% CI)		
	rare group	0.97 (0.57–1.36)	6.3×10^{-6}	n/a
	g_unclassified (f_Ruminococcaceae)	0.98 (0.42–1.55)	1.0×10^{-3}	n/a
	OTU			
	OTU12986 (g_unclassified/f_Ruminococcaceae)	1.08 (0.67–1.48)	1.2×10^{-6}	Bovine reproductive tract/fecal sample from <i>Bos taurus</i> ; gayal (<i>Bos frontalis</i>) breeding farm/cattle herd feces/fecal sample in taurus (neonatal calf)
	OTU9879 (g_Planococcus/f_Planococcaceae)	1.01 (0.62–1.40)	2.7×10^{-6}	Bioaerosol emitted from wastewater treatment plant/ <i>Planomicrobium stacke-brandtii</i> —strain collection/Chandra Tal Lake water/soil/soil from Qinghai lake
	OTU13554 (g_Pseudomonas/f_Pseudomonadaceae)	0.94 (0.57–1.32)	3.7×10^{-6}	Biogas plant/leaf cutter ant refuse dump/cow manure/Japan/soil
	OTU2597 (g_unclassified/f_Ruminococcaceae)	0.98 (0.58–1.38)	6.6×10^{-6}	Bovine reproductive tract/manure/cow manure/descending colon ingesta from <i>Bos taurus</i> /cow teat skin
	OTU7854 (g_Bacteroides/f_Bacteroidaceae)	0.84 (0.48–1.19)	1.6×10^{-5}	descending colon ingesta from <i>Bos taurus</i> /sheep feces Ireland/Okapi feces/fecal sample <i>Procapra przewalskii</i> /fecal sample <i>Bos taurus</i> /
	OTU1017 (g_Alistipes/f_Rikenellaceae)	0.60 (0.34–0.86)	2.1×10^{-5}	descending colon mucosa of <i>Bos taurus</i> /Argali sheep feces/fecal sample from beef cattle/fecal cattle sample/Okapi feces
Presence/absence	Genus	OR (95% CI)		
	g_Clostridium (f_Clostridiaceae)	27.71 (6.03–127.45)	5.3×10^{-5}	n/a
	g_Facklamia (f_Aerococcaeaceae)	13.43 (3.77–47.84)	1.4×10^{-4}	n/a
	OTU			
OTU2597 (g_unclassified/f_Ruminococcaceae)	50.81 (10.9–236.6)	3.2×10^{-6}	Bovine reproductive tract/manure/cow manure/descending colon ingesta from <i>Bos taurus</i> /cow teat skin	

g_, genus; f_, family; n/a, not applicable.

*A BLAST search of the OTU sequences against NCBI database was performed (2016-03-09). Listed are up to five different sites where similar sequences with $\geq 97\%$ were found.

members (by hair or cloths) or pets (by fur or paws) most likely accounts for the increased bacterial diversity in homes of farm children and the detection of farm- and animal-specific taxa in these samples.

When performing a BLAST search for the mattress dust OTUs that were associated with farming or cow/straw exposure, we found nearly all hits to be known from previous farm- or cow-related samples (Tables 1 and S3).

In this regard, the association of farming with the genera *Clostridium* and *Facklamia* is notable since representatives of the genus *Clostridium* are commonly found in soil, sewage,

and intestines of animals and humans, whereas *Facklamia* is known particularly from milk and cow samples (40, 41). *Trichococcus* was associated with current cow/straw exposure and was previously isolated from raw milk (42), dairy waste (43), and sewage fluent (44).

More interestingly, an association of farming with bacteria from a not further classified genus of the family *Ruminococcaceae* emerged and was even paralleled in nasal samples by a more prevalent occurrence of OTU 2597 in farm children. This OTU belongs also to the family *Ruminococcaceae*, which is typically found in the mammalian gut, also of cows and




Figure 5 Effects of richness and Shannon index of mattress dust and nasal microbiota, respectively, on asthma. Diversity measurements are shown unadjusted, adjusted for farming only, and adjusted for farming + mutually adjusted. Mutually adjusted means

humans. In another study on airborne microbes in a rural area of the USA, the family of *Ruminococcaceae* was identified as an indicator for cow feces (45).

The identification of these animal- and feces-specific taxa supports the notion that the identified microorganisms are specific for the farm exposure and may originate from animal sheds. Alternatively, the detection of fecal microorganisms, which are also known from human studies, may point toward another important body surface where host-microbial interaction is quite intensive, that is, the intestinal mucosa. As the gut microbiome is highly diverse, the detected signal by bacterial richness or rare taxa might also be a proxy for intestinal microorganisms.

Anyway, the pronounced gradient in bacterial diversity between farm and nonfarm children we detected in this study is paralleled by findings in a comparable study including school children from rural areas of five European countries (5). In that study, microbial diversity was assessed in mattress dust by the gel-based single-strand conformation polymorphism (SSCP) method. Now we rediscovered this gradient in bacterial diversity between farm and nonfarm children by a more elaborate method offering a much higher resolution of bacterial diversity down to an OTU level. In that earlier study (5), the higher bacterial diversity of farm children also partially explained their lower asthma prevalence. Hence, we tested whether this effect was also observed in the GABRIELA population and, more importantly, whether it was related to nasal microbiota, as well.

Indeed, in the present analysis α -diversity was inversely associated with asthma occurrence also after adjustment for the farm exposure. Of note, the effect was observed not only in mattress dust but also independently in nasal samples, albeit with a lower effect size (Fig. 5). An important difference of these effects between both sampling sites is illustrated by the mediation analysis: Only richness in mattress dust

additional adjustment for the respective diversity measurement in nasal swabs or mattress dust. Richness is shown on the z-transformed scale of the log₁₀ (of richness) and Shannon on the z-transformed scale of the original Shannon index.

operated as an intermediate factor in the association of farming with asthma, whereas richness in nasal swabs was hardly related to farming but still contributed to a lower asthma risk.

One may speculate that diversity of the nasal microbiome is influenced by other factors, which independently of the farm exposure lowers the risk of asthma. For nasal microbiome, obviously airborne bacteria might be relevant, but also skin bacteria might play a role since we found typical skin in the nasal samples abundantly (Fig. 1). Ultimately, a host factor itself may impact on the microbial diversity of the nasal cavity. This might be governed by the host immune system, which again might be shaped by genetic as well as environmental factors such as (viral) infections possibly transmitted by siblings in the classical sense of the hygiene hypothesis.

As bacteria trigger a rather limited number of different receptors of the innate immune system, microbial diversity has been interpreted as a proxy for a cocktail of specific environmental bacteria (5). Therefore, we assessed also individual bacterial genera and OTUs for associations with asthma. The inverse association of asthma with the genus *Prevotella* in nasal samples is interesting as this genus belongs to the phylum Bacteroidetes. This phylum produces immunomodulatory metabolites such as short-chain fatty acids (46), which have been implied in asthma protection in mouse models (47). In addition, *Prevotella* may alleviate detrimental effects of other airway bacteria such as *Haemophilus* (48).

However, the detection of individual bacterial taxa with asthma-protective potential (Table S5) is an exploratory approach, as these protective effects did not withstand correction for multiple testing. This is most likely limited by the sample size of this pilot study and warrants replication in larger studies in the future. The moderate sample size and the stratification for farm exposure also precluded a more

detailed analysis of further exposures beyond farming. Moreover, this study was performed in healthy children attending school. Very few of them (5.8%) were taking any drug at time of nasal sampling, thereby rendering specific medication effects difficult.

In addition, the present analyses were based on extracted DNA. As microbial DNA is not necessarily derived from living organisms but can persist freely or associated with cell wall particles of dead organisms for years (49), the detected taxa do not unambiguously represent living or active bacteria.

On the other side, the in-depth analysis of samples from two rather different sampling sites, but related to the same individuals, is a major strength of this study. Samples were taken only 2 days apart and processed following identical protocols except for DNA extraction. This situation provided the unique opportunity to study the microbiome of the upper airways in the context of the pertinent environmental microbiota.

Taken together, we found environmental, or more precisely farm-related, bacteria in mattress dust and to a lower extent in nasal microbiota. The effect of microbial exposure on asthma was stronger in mattress dust as compared to nasal samples. This may seem unexpected as the nasal cavity as part of the upper airway mucosa is likely to reflect interactions of environmental microorganisms with the host mucosa more adequately. This paradox thus challenges the concept of colonization of the (upper) airways by asthma-protective bacteria. Other hypotheses such as asthma protection by colonization of the intestine by beneficial microorganisms or inhalation of microbial metabolites with immunomodulatory properties may come to the focus of future research.

Acknowledgments

This work was supported by the European Commission (Grant LSHB-CT-2006-018996) and the European Research Council (Grant 250268). We thank the Leibniz Supercomputing Centre (LRZ) of the Bavarian Academy of Sciences and Humanities (BAdW) for the provisioning and support of Cloud computing infrastructure essential to this publication. Furthermore, we thank the children, who participated in the study, and their families and the field workers participated in GABRIELA.

Author contributions

HD, MJE, CBF, JG, DH, EH, and EvM involved in conception and design of the GABRIELA study and sample acquisition. LTB, MJE, ME, MS, AL, and SK involved in design of the study and planning of experiments. LTB, ME, and SK involved in processing of samples. CB, LTB, MD, MJE, AL, and GL involved in analysis and interpretation of the data.

References

- Andersson AM, Weiss N, Rainey F, Salkinoja-Salonen MS. Dust-borne bacteria in animal sheds, schools and children's day care centres. *J Appl Microbiol* 1999;**86**:622–634.
- Braun-Fahrlander C, Riedler J, Herz U, Eder W, Waser M, Grize L et al. Environmental exposure to endotoxin and its relation to asthma in school-age children. *N Engl J Med* 2002;**347**:869–877.
- Normand AC, Sudre B, Vacheyrou M, Depner M, Wouters IM, Noss I et al. Airborne cultivable microflora and microbial transfer in farm buildings and rural dwellings. *Occup Environ Med* 2011;**68**:849–855.

MD, MJE, and AL wrote the manuscript. All authors involved in editing and final approval of the paper.

Conflicts of interest

Dr. Depner reports grants from European Research Council, during the conduct of the study; grants from German Research Foundation, outside the submitted work; Dr. Ege reports a patent EP000002361632A1 pending; Dr. Genuneit reports grants from European Commission during the conduct of the study; Dr. Heederik reports grants from European Commission, during the conduct of the study; Dr. von Mutius reports grants from European Research Council, during the conduct of the study; personal fees from American Academy of Allergy, Asthma & Immunology, personal fees from Ökosoziales Forum Oberösterreich, personal fees from Mundipharma, personal fees from HAL Allergie GmbH, personal fees from DOC Congress SRL, personal fees from American Thoracic Society, personal fees from University of Tampere, personal fees from GBS RE HEFCE, personal fees from Novartis Pharma, personal fees from OM Pharma SA, personal fees from AbbVie Deutschland GmbH & Co. KG, personal fees from medUpdate GmbH, personal fees from System Analytic Ltd., outside the submitted work. The other authors have no conflicts of interest to disclose.

Supporting Information

Additional Supporting Information may be found in the online version of this article:

Figure S1. Study design.

Figure S2. Distribution of sample collection in spring/summer 2007.

Figure S3. Number of reads per sample for mattress dust and nasal samples.

Figure S4. Multiple rarefaction curves for (A) richness and (B) Shannon-index for the nasal and mattress dust samples.

Figure S5. NMDS of (A) unweighted UniFrac and (B) Bray-Curtis analysis of mattress dust and nasal samples, further stratified for farming and non-farming.

Table S1. Read counts for negative controls.

Table S2. Description of the Austrian part of the GABRIELA recruitment population and the analyzed study population.

Table S3. Genera and OTUs from mattress dust significantly associated with current regular exposition to cow/straw.

Table S4. Sensitivity analyses for medication and atopy.

Table S5. Association tests of presence/absence and asthma in the rarefied datasets for genera or OTUs.

Appendix S1. Methods.

4. Lippmann M, Yeates DB, Albert RE. Deposition, retention, and clearance of inhaled particles. *Br J Ind Med* 1980;**37**:337–362.
5. Ege MJ, Mayer M, Normand AC, Genuneit J, Cookson WO, Braun-Fahrlander C et al. Exposure to environmental microorganisms and childhood asthma. *N Engl J Med* 2011;**364**:701–709.
6. Genuneit J. Exposure to farming environments in childhood and asthma and wheeze in rural populations: a systematic review with meta-analysis. *Pediatr Allergy Immunol* 2012;**23**:509–518.
7. von Mutius E, Vercelli D. Farm living: effects on childhood asthma and allergy. *Nat Rev Immunol* 2010;**10**:861–868.
8. Illi S, Döpner M, Genuneit J, Horak E, Loss G, Strunz-Lehner C et al. Protection from childhood asthma and allergy in Alpine farm environments—the GABRIEL Advanced Studies. *J Allergy Clin Immunol* 2012;**129**:1470–1477.
9. Ege MJ, Mayer M, Schwaiger K, Mattes J, Pershagen G, van Hage M et al. Environmental bacteria and childhood asthma. *Allergy* 2012;**67**:1565–1571.
10. Valkonen M, Wouters IM, Taubel M, Rintala H, Lenters V, Vasara R et al. Bacterial exposures and associations with atopy and asthma in children. *PLoS One* 2015;**10**: e0131594.
11. Debarry J, Garn H, Hanuszkiewicz A, Dickgreber N, Blumer N, von Mutius E et al. *Acinetobacter hwoffii* and *Lactococcus lactis* strains isolated from farm cowsheds possess strong allergy-protective properties. *J Allergy Clin Immunol* 2007;**119**:1514–1521.
12. Hagner S, Harb H, Zhao M, Stein K, Holst O, Ege MJ et al. Farm-derived Gram-positive bacterium *Staphylococcus sciuri* W620 prevents asthma phenotype in HDM- and OVA-exposed mice. *Allergy* 2013;**68**:322–329.
13. Vogel K, Blumer N, Korhals M, Mittelstadt J, Garn H, Ege M et al. Animal shed *Bacillus licheniformis* spores possess allergy-protective as well as inflammatory properties. *J Allergy Clin Immunol* 2008;**122**:307–312.
14. Genuneit J, Buchele G, Waser M, Kovacs K, Debinska A, Boznanski A et al. The GABRIEL advanced surveys: study design, participation and evaluation of bias. *Paediatr Perinat Epidemiol* 2011;**25**:436–447.
15. Schram-Bijkerk D, Doeke G, Douwes J, Boeve M, Riedler J, Ublagger E et al. Bacterial and fungal agents in house dust and wheeze in children: the PARSIFAL study. *Clin Exp Allergy* 2005;**35**:1272–1278.
16. Cardenas PA, Cooper PJ, Cox MJ, Chico M, Arias C, Moffatt MF et al. Upper airways microbiota in antibiotic-naive wheezing and healthy infants from the tropics of rural Ecuador. *PLoS One* 2012;**7**:e46803.
17. Sim K, Cox MJ, Wopereis H, Martin R, Knol J, Li MS et al. Improved detection of bifidobacteria with optimised 16S rRNA-gene based pyrosequencing. *PLoS One* 2012;**7**:e32543.
18. Fierer N, Hamady M, Lauber CL, Knight R. The influence of sex, handedness, and washing on the diversity of hand surface bacteria. *Proc Natl Acad Sci USA* 2008;**105**:17994–17999.
19. Quince C, Lanzen A, Davenport RJ, Turnbaugh PJ. Removing noise from pyrosequenced amplicons. *BMC Bioinformatics* 2011;**12**:38.
20. Edgar RC. Search and clustering orders of magnitude faster than BLAST. *Bioinformatics* 2010;**26**:2460–2461.
21. Caporaso JG, Kuczynski J, Stombaugh J, Bittinger K, Bushman FD, Costello EK et al. QIIME allows analysis of high-throughput community sequencing data. *Nat Methods* 2010;**7**:335–336.
22. Oksanen J, Blanchet FG, Kindt R, Legendre P, Minchin PR, O'Hara RB et al. vegan: Community Ecology Package. R package version 2.3-5. 2016.
23. Lozupone C, Lladser ME, Knights D, Stombaugh J, Knight R. UniFrac: an effective distance metric for microbial community comparison. *ISME J* 2011;**5**:169–172.
24. R-Development-Core-Team. *R: a language and environment for statistical computing*. Vienna, Austria: R Foundation for Statistical Computing; 2008.
25. McMurdie PJ, Holmes S. phyloseq: an R package for reproducible interactive analysis and graphics of microbiome census data. *PLoS One* 2013;**8**:e61217.
26. Barberan A, Dunn RR, Reich BJ, Pacifici K, Laber EB, Menninger HL et al. The ecology of microcosmic life in household dust. *Proc R Soc Lond B Biol Sci* 2015;**282**:20151139.
27. Dunn RR, Fierer N, Henley JB, Leff JW, Menninger HL. Home life: factors structuring the bacterial diversity found within and between homes. *PLoS One* 2013;**8**:e64133.
28. Taubel M, Rintala H, Pitkaranta M, Paulin L, Laitinen S, Pekkanen J et al. The occupant as a source of house dust bacteria. *J Allergy Clin Immunol* 2009;**124**:834–840.
29. Belkaid Y, Segre JA. Dialogue between skin microbiota and immunity. *Science* 2014;**346**:954–959.
30. Grice EA, Segre JA. The skin microbiome. *Nat Rev Microbiol* 2011;**9**:244–253.
31. White DC, Sutton SD, Ringelberg DB. The genus *Sphingomonas*: physiology and ecology. *Curr Opin Biotechnol* 1996;**7**:301–306.
32. Hagi T, Sasaki K, Aso H, Nomura M. Adhesive properties of predominant bacteria in raw cow's milk to bovine mammary gland epithelial cells. *Folia Microbiol* 2013;**58**:515–522.
33. Quigley L, McCarthy R, O'Sullivan O, Beresford TP, Fitzgerald GF, Ross RP et al. The microbial content of raw and pasteurized cow milk as determined by molecular approaches. *J Dairy Sci* 2013;**96**:4928–4937.
34. Neale DB, Marshall KA, Sederoff RR. Chloroplast and mitochondrial DNA are paternally inherited in *Sequoia sempervirens* D. Don Endl. *Proc Natl Acad Sci USA* 1989;**86**:9347–9349.
35. Berger UE. Kiefer (*Pinus*). Available from: www.polleninfo.org/AT/de/allergie-infos/allgemein/steckbriefe/kiefer.html (cited 11 March 2016).
36. Bassis CM, Tang AL, Young VB, Pynnonen MA. The nasal cavity microbiota of healthy adults. *Microbiome* 2014;**2**:27.
37. Frank DN, Feazel LM, Bessesen MT, Price CS, Janoff EN, Pace NR. The human nasal microbiota and *Staphylococcus aureus* carriage. *PLoS One* 2010;**5**:e10598.
38. Oh J, Conlan S, Polley EC, Segre JA, Kong HH. Shifts in human skin and nares microbiota of healthy children and adults. *Genome Med* 2012;**4**:77.
39. Consortium THMP. Structure, function and diversity of the healthy human microbiome. *Nature* 2012;**486**:207–214.
40. Rasolofoa EA, St-Gelais D, LaPointe G, Roy D. Molecular analysis of bacterial population structure and dynamics during cold storage of untreated and treated milk. *Int J Food Microbiol* 2010;**138**:108–118.
41. Takamatsu D, Ide H, Osaki M, Sekizaki T. Identification of *Facklamia soureikii* from a lactating cow. *J Vet Med Sci* 2006;**68**:1225–1227.
42. Hantsis-Zacharov E, Halpern M. Culturable psychrotrophic bacterial communities in raw milk and their proteolytic and lipolytic traits. *Appl Environ Microbiol* 2007;**73**:7162–7168.
43. McGarvey JA, Miller WG, Zhang R, Ma Y, Mitloehner F. Bacterial population dynamics in dairy waste during aerobic and anaerobic treatment and subsequent storage. *Appl Environ Microbiol* 2007;**73**:193–202.
44. VandeWalle JL, Goetz GW, Huse SM, Morrison HG, Sogin ML, Hoffmann RG et al. *Acinetobacter*, *Aeromonas* and *Trichococcus* populations dominate the microbial community within urban sewer infrastructure. *Environ Microbiol* 2012;**14**:2538–2552.
45. Bowers RM, Clements N, Emerson JB, Wiedinmyer C, Hannigan MP, Fierer N. Seasonal variability in bacterial and fungal diversity of the near-surface atmosphere. *Environ Sci Technol* 2013;**47**:12097–12106.

46. Reichardt N, Duncan SH, Young P, Belenguer A, McWilliam Leitch C, Scott KP et al. Phylogenetic distribution of three pathways for propionate production within the human gut microbiota. *ISME J* 2014;**8**:1323–1335.
47. Trompette A, Gollwitzer ES, Yadava K, Sichelstiel AK, Sprenger N, Ngom-Bru C et al. Gut microbiota metabolism of dietary fiber influences allergic airway disease and hematopoiesis. *Nat Med* 2014;**20**:159–166.
48. Larsen JM, Steen-Jensen DB, Laursen JM, Sondergaard JN, Musavian HS, Butt TM et al. Divergent pro-inflammatory profile of human dendritic cells in response to commensal and pathogenic bacteria associated with the airway microbiota. *PLoS One* 2012;**7**:e31976.
49. Nielsen KM, Johnsen PJ, Bensasson D, Dafonchio D. Release and persistence of extracellular DNA in the environment. *Environ Biosafety Res* 2007;**6**:37–53.