

PLATELETS AND THROMBOPOIESIS

Rare variants in *GP1BB* are responsible for autosomal dominant macrothrombocytopenia

Suthesh Sivapalaratnam,^{1-3,*} Sarah K. Westbury,^{4,*} Jonathan C. Stephens,^{1,2,5,*} Daniel Greene,^{1,2,6,*} Kate Downes,^{1,2,5} Anne M. Kelly,⁷ Claire Lentaingne,^{8,9} William J. Astle,^{1,2,6} Eric G. Huizinga,¹⁰ Paquita Nurden,¹¹ Sofia Papadia,^{1,2} Kathelijne Peerlinck,¹² Christopher J. Penkett,^{1,2} David J. Perry,¹³ Catherine Roughley,¹⁴ Ilenia Simeoni,^{1,2,5} Kathleen Stirrups,^{1,2} Daniel P. Hart,³ R. Campbell Tait,¹⁵ Andrew D. Mumford,^{4,16} NIHR BioResource,² Michael A. Laffan,^{8,9} Kathleen Freson,^{12,†} Willem H. Ouwehand,^{1,2,5,17,†} Shinji Kunishima,^{18,†} and Ernest Turro^{1,2,5,6,†}

¹Department of Haematology, University of Cambridge, Cambridge, United Kingdom; ²National Institute for Health Research BioResource–Rare Diseases, Cambridge University Hospitals, Cambridge, United Kingdom; ³Department of Haematology, Barts Health National Health Service Trust, London, United Kingdom; ⁴School of Clinical Sciences, University of Bristol, Bristol, United Kingdom; ⁵National Health Service Blood and Transplant, Cambridge Biomedical Campus, Cambridge, United Kingdom; ⁶Medical Research Council Biostatistics Unit, Cambridge Institute of Public Health, Cambridge Biomedical Campus, Cambridge, United Kingdom; ⁷Department of Haematology, Great Ormond Street Hospital for Children National Health Service Trust, London, United Kingdom; ⁸Centre for Haematology, Hammersmith Campus, Imperial College Academic Health Sciences Centre, Imperial College London, London, United Kingdom; ⁹Imperial College Healthcare National Health Service Trust, London, United Kingdom; ¹⁰Crystal and Structural Chemistry, Department of Chemistry, Faculty of Science, Bijvoet Center for Biomolecular Research, Utrecht University, Utrecht, The Netherlands; ¹¹Institut Hospitalo-Universitaire LIRYC, Hôpital Xavier Arnoz, Pessac, France; ¹²Department of Cardiovascular Sciences, Center for Molecular and Vascular Biology, University of Leuven, Leuven, Belgium; ¹³Department of Haematology, Addenbrooke's Hospital, Cambridge University Hospitals National Health Service Foundation Trust, Cambridge Biomedical Campus, Cambridge, United Kingdom; ¹⁴Kent Haemophilia Thrombosis Centre at East Kent Hospitals University NHS Foundation Trust, Canterbury, United Kingdom; ¹⁵Department of Haematology, Royal Infirmary, Glasgow, United Kingdom; ¹⁶School of Cellular and Molecular Medicine, University of Bristol, Bristol, United Kingdom; ¹⁷Wellcome Trust Sanger Institute, Wellcome Trust Genome Campus, Hinxton, Cambridge, United Kingdom; and ¹⁸Department of Advanced Diagnosis, Clinical Research Center, National Hospital Organization Nagoya Medical Center, Nagoya, Japan

Key Points

- Variants in *GP1BB* can cause autosomal dominant macrothrombocytopenia.

The von Willebrand receptor complex, which is composed of the glycoproteins Ib α , Ib β , GPV, and GPIX, plays an essential role in the earliest steps in hemostasis. During the last 4 decades, it has become apparent that loss of function of any 1 of 3 of the genes encoding these glycoproteins (namely, *GP1BA*, *GP1BB*, and *GP9*) leads to autosomal recessive macrothrombocytopenia complicated by bleeding. A small number of variants in *GP1BA* have been reported to cause a milder and dominant form of macrothrombocytopenia, but

only 2 tentative reports exist of such a variant in *GP1BB*. By analyzing data from a collection of more than 1000 genome-sequenced patients with a rare bleeding and/or platelet disorder, we have identified a significant association between rare monoallelic variants in *GP1BB* and macrothrombocytopenia. To strengthen our findings, we sought further cases in 2 additional collections in the United Kingdom and Japan. Across 18 families exhibiting phenotypes consistent with autosomal dominant inheritance of macrothrombocytopenia, we report on 27 affected cases carrying 1 of 9 rare variants in *GP1BB*. (*Blood*. 2017;129(4):520-524)

Introduction

The earliest step in hemostasis is the tethering of platelets to the damaged endothelium through their cell surface receptor for von Willebrand factor (VWF). The glycoproteins (GPs) Ib α and Ib β , in association with GPV and GPIX, form the transmembrane receptor complex on platelets for VWF.^{1,2} The leucine-rich repeat (LRR) domain of GPIb α contains the binding site for VWF, whereas the β chain contributes to the surface expression of the complex, which, by phosphorylating its intracellular domain, also participates in downstream signaling. GPIb β is synthesized from a 1.5-kb mRNA transcribed from *GP1BB* and is highly expressed in megakaryocytes, but absent from other blood cell progenitors.³ The mRNA encodes a

206–amino-acid-long transmembrane protein with a 22-kD molecular mass and an extracellular LRR domain.⁴⁻⁶

Bernard and Soulier were the first to describe a patient with a rare autosomal recessive syndrome (Bernard-Soulier syndrome; BSS) characterized by giant platelets with concomitant thrombocytopenia complicated by severe bleeding.⁷ Platelets from patients with this syndrome do not agglutinate in response to ristocetin and show a subtly reduced aggregation response to thrombin. Over the course of 4 decades, 45, 39, and 28 variants have been identified in *GP1BA*, *GP1BB*, and *GP9*, respectively, that, if present in compound heterozygous or homozygous form, cause the VWF receptor to be

Submitted 10 August 2016; accepted 3 November 2016. Prepublished online as *Blood* First Edition paper, 14 November 2016; DOI 10.1182/blood-2016-08-732248.

*S.S., S.K.W., J.C.S., and D.G. contributed equally to this study.

†K.F., W.H.O., S.K., and E.T. contributed equally to this study.

The online version of this article contains a data supplement.

There is an Inside *Blood* Commentary on this article in this issue.

The publication costs of this article were defrayed in part by page charge payment. Therefore, and solely to indicate this fact, this article is hereby marked "advertisement" in accordance with 18 USC section 1734.

© 2017 by The American Society of Hematology

absent or reduced in function.⁸ Only 4 variants in *GP1BA* with dominant effects have been reported, one being the “Bolzano variant,”⁹ and a further 3 observed in 3 isolated pedigrees.⁸ Only 2 variants in *GP1BB* have been reported as being possibly responsible for dominant macrothrombocytopenia. One encoding Y113C was identified in 2 Japanese families and led to initial speculation that variants in the subunits of the GPIb/IX complex could span a spectrum of platelet and bleeding phenotypes.^{10,11} Another, encoding R42C, was observed in only 1 Japanese patient, and data from family or functional studies supporting an association with a dominant phenotype were lacking.¹² There are no reports of variants in *GP9* having a dominant effect, and BSS cases resulting from variants in *GP5* do not exist because GPV is not required for receptor expression.¹³

We found a significant statistical association between rare monoallelic nonsynonymous variants in *GP1BB* and macrothrombocytopenia. Family history and cosegregation data from 18 pedigrees, in which 27 affected cases carry 1 of 9 rare variants in *GP1BB*, are consistent with autosomal dominant inheritance.

Study design

Study population

The study population comprises a discovery collection and 2 validation collections. The discovery collection consists of cases with an assumed inherited bleeding or platelet disorder (BPD) of unknown molecular etiology or their relatives enrolled in the National Institute for Health Research (NIHR) BioResource after providing informed written consent.¹⁴ The BioResource cohort was enrolled during a 4-year period and is composed of 1542 patients with a BPD or their close relatives, with a further 5422 patients with other rare inherited disorders or their close relatives (<https://bioresource.nihr.ac.uk/rare-diseases/clinicians/>). The validation collections contain data from 75 patients with thrombocytopenia assessed with the ThromboGenomics diagnostic platform,¹⁵ and 301 Japanese probands¹⁶ referred during an 8-year period for a suspected diagnosis of inherited macrothrombocytopenia and their relatives. Institutional review board approval was obtained for this international multicenter study.

Data analysis

Coding of clinical and laboratory phenotypes with Human Phenotype Ontology (HPO) terms and collection of numerical and family history data were performed as described previously.¹⁴ Variants from high-throughput sequencing were called and annotated using Isaac (Illumina, Inc; whole-genome sequencing) or as described previously¹⁵ (capture-based sequencing). We used phenotype similarity regression¹⁷ (SimReg) to identify statistical associations between presence of a variant affecting protein sequence in a gene and similarity to a latent HPO-coded phenotype in the discovery collection, subsequently corroborated using Fisher’s exact test. The variants obtained by high-throughput DNA sequencing of the probands were genotyped by Sanger sequencing in relatives who agreed to participate in this study. The variants identified in all members of the Japanese collection were obtained by Sanger sequencing. Where possible, complete blood counts were obtained using automated hematology analyzers, VWF receptor expression levels on platelets were determined by cytometry, and platelet function was tested by light transmission aggregometry. Macrothrombocytopenia was deemed present if the platelet count was below $150 \times 10^9/L$, the mean platelet volume was above 12 fL, or there was clear platelet anisocytosis, with a subset of platelets being abnormally large.

Results and discussion

We identified a strong statistical association between the presence of rare nonsynonymous monoallelic DNA variants (population allele frequency less than 1/10 000) in *GP1BB* and macrothrombocytopenia in the

discovery collection (SimReg posterior probability = 0.93 with inferred characteristic phenotype preferentially included the HPO term “Increased mean platelet volume”; Fisher’s $P = 2.10 \times 10^{-6}$). All 8 probands with macrothrombocytopenia from the discovery collection had a family history suggestive of autosomal dominant inheritance. This mode of transmission was corroborated by results from cosegregation studies (Figure 1; $P = 1.95 \times 10^{-3}$). Systematic review of rare variants in these 8 cases within the 15 established genes implicated in macrothrombocytopenia,¹⁸ including *GP1BA* and *GP9*, did not reveal any alternative variants that could plausibly explain the platelet phenotype. We searched for further cases in the validation collection and identified 4 further probands in the ThromboGenomics data set and 6 further probands in the Japanese collection who carried a variant specific to people of Japanese ancestry (Figure 1).

Altogether, we identified 9 unique variants in *GP1BB*: 1 resulting in a disruption of the canonical methionine start codon; 1 encoding a premature stop at residue 46; 5 missense variants encoding L16P, G43W, T68M, Y113C, and L132Q; a 3-amino acid deletion P79_L81del; and a frameshift in the codon for residue A150 leading to an alternative reading frame with a premature stop after 43 further residues (Figure 2A). Assessment of the potential effect of the missense and in-frame variants on GPIb β structure, supported by prior work by McEwan et al,¹⁹ is available in supplemental Table 2 and supplemental Figure 1 on the *Blood* Web site. Only the variant encoding G43W is present in the ExAC database²⁰ of more than 60 000 exome-sequenced individuals, where it is heterozygous in just 1 individual. It is noteworthy that the variant underlying the premature stop at residue 46, the variant leading to a frameshift at residue 150, and the missense variants encoding L16P, G43W, and Y113C have been previously implicated in BSS.^{8,10,21,22} We reason that the effects of these 5 variants in heterozygosity, except for Y113C,^{10,11} went unnoticed because the counts of the parents of these BSS probands were not measured or were not deemed notably low in isolation from other monoallelic cases.

The count and mean volume of platelets of the 18 probands were $107.9 \times 10^9/L$ (range, $47\text{--}172 \times 10^9/L$) and 12.74 fL (range, 10.7–14.3 fL), respectively (Figure 2B; supplemental Table 1). In several cases, May-Grünwald-Giemsa-stained blood smears revealed platelet anisocytosis, with a subset of platelets being abnormally large; however, blood films were not available for all cases (Figure 2C; supplemental Table 1). Electron micrographs of the platelets of B-3 show significantly increased platelet surface area that is intermediate between healthy controls and BSS cases, but includes a small number of giant platelets (Figure 2D). Thus, the mean platelet volumes obtained by automated complete blood count analysis may obscure an increased spread of platelet volumes. Bleeding diathesis, including menorrhagia, epistaxis, spontaneous bleeding, and postpartum bleeding, was reported in 9 of 23 females but in none of the 8 males for whom platelet count and bleeding phenotype information was available, which suggests that the overall propensity to bleeding is, at most, only marginally increased in these patients relative to the general population. There was a reduction of at least 30% of GPIb α on platelets, as measured by flow cytometry in cases from 8 of the 9 families for which measurements were available (supplemental Table 1).

Each VWF receptor complex has 4 GPIb β molecules, which are covalently bonded with 2 copies of GPIb α through cysteines at residues 147 and 526/527, respectively,²³ and noncovalently paired with 2 molecules each of GPV and GPIX.²⁴ This stoichiometry may explain why a single allele encoding a mutated GPIb β molecule may exert a dominant negative effect on the function of the VWF complex. It could also exacerbate haploinsufficiency in the cases with disruption of the methionine start codon and truncation at residue 46. It is important to note that although the overall association between




Figure 1. Graphic representation of the 18 pedigrees with autosomal dominant inheritance of macrothrombocytopenia in the discovery and validation collections. No genome-wide excess relatedness among individuals subjected to high-throughput sequencing from different pedigrees sharing the same variant (pedigrees B and I, C and J, D and E, and H and L) could be detected by genetic analysis,²⁵ although the shared variants in *GP1BB* may have been co-inherited from a distant common ancestor. Filled symbols, macrothrombocytopenia; gray symbols, unknown; blank symbols, normal platelet count and volume and absence of macrothrombocytes. Squares, males; circles, females; +/M, heterozygous; +/+ wild-type. A second variant encoding L175P was identified by Sanger sequencing in K-3 absent from K-6, which might influence PLT and MPV by affecting the cytoskeleton. The probability of the genotyping results under the null hypothesis of random segregation, conditional on the genotypes of the index cases, is $0.5^9 = 1.95 \times 10^{-3}$.

monoallelic variants in *GP1BB* and macrothrombocytopenia is robust, it is possible that some of the 9 variants reported here are individually not causal. Therefore, further work and the continued sharing of genotype

and phenotype data from many patients with macrothrombocytopenia will be required to establish the molecular consequences of these variants definitively.




Figure 2. (A) Localization and consequence of the 9 distinct rare variants identified in *GP1BB* (shown above the diagram) in the context of all 39 variants published previously, excluding the 22q11.2 deletion. Novel variants are shown in red. Known variants were obtained from a recent review,⁸ 2 publications on BSS published since the review,^{26,27} and an article reporting a single case with dominant macrothrombocytopenia possibly because of a heterozygous variant encoding R42C.¹² The variant encoding T68M was observed in an unaffected relative of a patient with non-BPD rare disease for whom a complete blood count was unavailable. Only the variant encoding G43W is present (in 1 individual) in the ExAC database. The domains shown in colors, from left to right, are signal peptide (1–25), LRR N-terminal (27–55), LRR (60–83), LRR C-terminal (89–143), transmembrane domain (148–172), and cytoplasmic domain (173–206). (B) Gender-stratified histograms of platelet count measurements obtained using a Sysmex hematology analyzer from 48 345 blood donors from the INTERVAL randomized controlled trial²⁸ after adjustment for technical artifacts. The red arrows superimposed on the histograms indicate the gender and values for patients with 1 of the variants in *GP1BB* reported herein. For individuals I-3 and L-3, with a range of platelet counts, the midpoints were used. The green arrow indicates the gender and value for individual E-1, who is homozygous for the wild-type allele. Any superimposed arrows have been nudged slightly along the vertical axis so they can be distinguished. (C) Images of blood films from cases E-5, F-5, and H-7, with different variants in *GP1BB* (encoding W46*, T68M, and A150Rfs*43). Original magnification, $\times 100$; May-Grünwald-Giemsa stain. (D) Images show representative electron micrographs of platelets from patient B-3, a patient with BSS (homozygous for a variant encoding Y131C) and an unaffected control, whereas the dot plot shows area measurements of 30 randomly selected platelets from a patient with BSS, 30 from case B-3, and 88 from 3 controls (30+30+28 platelets) with similar distributions. There was a significant difference among the means of the 3 groups ($P < .0001$, one-way analysis of variance). There was a significant difference between the mean area of platelets from the BSS case and patient B-3 ($P = .0021$), between the BSS case and controls ($P < .0001$), and between B-3 and controls ($P = .004$; Tukey's method for multiple comparisons).

Acknowledgments

We thank research nurse Amy Frary at the University of Cambridge and Alice Glaser at Imperial College London for their contributions to the clinical phenotyping of NIHR BioResource participants. We thank Ryoji Kobayashi, Masaaki Noguchi, Koji Miyazaki, Junichi

Watanabe, Yoshiyuki Ogawa and Tomohiro Kajiguchi for providing patient samples.

M.A.L. acknowledges support from the NIHR Imperial College Biomedical Research Centre. The NIHR BioResource–Rare Diseases is funded by the National Institute for Health Research of England (award number RG65966). Research in the W.H.O. laboratory is supported by the British Heart Foundation, European Commission,

Medical Research Council, NIHR, and the Wellcome Trust, and also receives support from National Health Service Blood and Transplant. C.L. and S.K.W. are supported by Medical Research Council Clinical Training Fellowships (MR/K023489/1).

Authorship

Contribution: S.S. and E.T. wrote the paper with assistance from J.C.S., M.A.L., K.F. and W.H.O.; J.C.S. maintained pedigrees, managed clinical data, and performed genotyping; S.K.W., A.M.K., C.L., P.N., K.P., D.J.P., C.R., D.P.H., R.C.T., A.D.M., NIHR BioResource, M.A.L., K.F., and S.K. provided samples and clinical data; E.G.H. conducted structure analysis; C.J.P. and K.S. managed the

NIHR BioResource sequencing pipeline; S.P. coordinated the NIHR BioResource–Rare Diseases BPD project; K.D. and I.S. managed ThromboGenomics; D.P.H. supervised S.S.; S.K. managed the Japanese collection; W.J.A. analyzed the INTERVAL dataset; K.F. analyzed microscopy data; and D.G. analyzed data under the supervision of E.T.

Conflict-of-interest disclosure: The authors declare no competing financial interests.

A complete list of the members of NIHR BioResource appears in the online appendix.

ORCID profiles: S.K.W., 0000-0002-0950-8148; E.T., 0000-0002-1820-6563.

Correspondence: Ernest Turro, National Health Service Blood and Transplant, Long Rd, Cambridge CB2 0PT, United Kingdom; e-mail: et341@cam.ac.uk.

References

- Ozaki Y, Asazuma N, Suzuki-Inoue K, Berndt MC. Platelet GPIb-IX-V-dependent signaling. *J Thromb Haemost*. 2005;3(8):1745-1751.
- López JA, Andrews RK, Afshar-Kharghan V, Berndt MC. Bernard-Soulier syndrome. *Blood*. 1998;91(12):4397-4418.
- Watkins NA, Gusnanto A, de Bono B, et al; Bloodomics Consortium. A HaemAtlas: characterizing gene expression in differentiated human blood cells. *Blood*. 2009;113(19):e1-e9.
- Kelly MD, Essex DW, Shapiro SS, et al. Complementary DNA cloning of the alternatively expressed endothelial cell glycoprotein Ib beta (GPIb beta) and localization of the GPIb beta gene to chromosome 22. *J Clin Invest*. 1994;93(6):2417-2424.
- Lopez JA, Chung DW, Fujikawa K, Hagen FS, Davie EW, Roth GJ. The alpha and beta chains of human platelet glycoprotein Ib are both transmembrane proteins containing a leucine-rich amino acid sequence. *Proc Natl Acad Sci USA*. 1988;85(7):2135-2139.
- Zieger B, Hashimoto Y, Ware J. Alternative expression of platelet glycoprotein Ib(beta) mRNA from an adjacent 5' gene with an imperfect polyadenylation signal sequence. *J Clin Invest*. 1997;99(3):520-525.
- Bernard J, Soulier JP. Sur une nouvelle variété de dystrophie thrombocytaire hémorragique congénitale. *Sem Hop*. 1948;24(Spec. No.):3217-3223.
- Savoia A, Kunishima S, De Rocco D, et al. Spectrum of the mutations in Bernard-Soulier syndrome. *Hum Mutat*. 2014;35(9):1033-1045.
- De Marco L, Mazzucato M, Fabris F, et al. Variant Bernard-Soulier syndrome type bolzano. A congenital bleeding disorder due to a structural and functional abnormality of the platelet glycoprotein Ib-IX complex. *J Clin Invest*. 1990;86(1):25-31.
- Kunishima S, Lopez JA, Kobayashi S, et al. Missense mutations of the glycoprotein (GP) Ib beta gene impairing the GPIb alpha/beta disulfide linkage in a family with giant platelet disorder. *Blood*. 1997;89(7):2404-2412.
- Kurokawa Y, Ishida F, Kamijo T, et al. A missense mutation (Tyr88 to Cys) in the platelet membrane glycoprotein Ibbeta gene affects GPIb/IX complex expression—Bernard-Soulier syndrome in the homozygous form and giant platelets in the heterozygous form. *Thromb Haemost*. 2001;86(5):1249-1256.
- Kunishima S, Naoe T, Kamiya T, Saito H. Novel heterozygous missense mutation in the platelet glycoprotein Ib beta gene associated with isolated giant platelet disorder. *Am J Hematol*. 2001;68(4):249-255.
- López JA, Leung B, Reynolds CC, Li CQ, Fox JE. Efficient plasma membrane expression of a functional platelet glycoprotein Ib-IX complex requires the presence of its three subunits. *J Biol Chem*. 1992;267(18):12851-12859.
- Westbury SK, Turro E, Greene D, et al; BRIDGE-BPD Consortium. Human phenotype ontology annotation and cluster analysis to unravel genetic defects in 707 cases with unexplained bleeding and platelet disorders. *Genome Med*. 2015;7(1):36.
- Simeoni I, Stephens JC, Hu F, et al. A high-throughput sequencing test for diagnosing inherited bleeding, thrombotic, and platelet disorders. *Blood*. 2016;127(23):2791-2803.
- Kunishima S, Okuno Y, Yoshida K, et al. *ACTN1* mutations cause congenital macrothrombocytopenia. *Am J Hum Genet*. 2013;92(3):431-438.
- Greene D, Richardson S, Turro E; NIHR BioResource. Phenotype similarity regression for identifying the genetic determinants of rare diseases. *Am J Hum Genet*. 2016;98(3):490-499.
- Lentaingne C, Freson K, Laffan MA, Turro E, Ouwehand WH; BRIDGE-BPD Consortium and the ThromboGenomics Consortium. Inherited platelet disorders: toward DNA-based diagnosis. *Blood*. 2016;127(23):2814-2823.
- McEwan PA, Yang W, Carr KH, et al. Quaternary organization of GPIb-IX complex and insights into Bernard-Soulier syndrome revealed by the structures of GPIbβ and a GPIbβ/GPIX chimera. *Blood*. 2011;118(19):5292-5301.
- Lek M, Karczewski KJ, Minikel EV, et al; Exome Aggregation Consortium. Analysis of protein-coding genetic variation in 60,706 humans. *Nature*. 2016;536(7616):285-291.
- Moran N, Morateck PA, Deering A, et al. Surface expression of glycoprotein Ib alpha is dependent on glycoprotein Ib beta: evidence from a novel mutation causing Bernard-Soulier syndrome. *Blood*. 2000;96(2):532-539.
- Kunishima S, Sako M, Yamazaki T, Hamaguchi M, Saito H. Molecular genetic analysis of a variant Bernard-Soulier syndrome due to compound heterozygosity for two novel glycoprotein Ibbeta mutations. *Eur J Haematol*. 2006;77(6):501-512.
- Luo S-Z, Mo X, Afshar-Kharghan V, Srinivasan S, López JA, Li R. Glycoprotein Ibalph forms disulfide bonds with 2 glycoprotein Ibbeta subunits in the resting platelet. *Blood*. 2007;109(2):603-609.
- Dong JF, Sae-Tung G, López JA. Role of glycoprotein V in the formation of the platelet high-affinity thrombin-binding site. *Blood*. 1997;89(12):4355-4363.
- Astle WJ, Balding DJ. Population structure and cryptic relatedness in genetic association studies. *Stat Sci*. 2009;24(4):451-471.
- Bragadottir G, Birgisdottir ER, Gudmundsdottir BR, et al. Clinical phenotype in heterozygote and biallelic Bernard-Soulier syndrome—a case control study. *Am J Hematol*. 2015;90(2):149-155.
- Qiao J, Davis AK, Morel-Kopp M-C, Ward CM, Gardiner EE, Andrews RK. Low levels of CD9 coincidental with a novel nonsense mutation in glycoprotein Ibβ in a patient with Bernard-Soulier syndrome. *Ann Hematol*. 2015;94(12):2069-2071.
- Moore C, Sambrook J, Walker M, et al. The INTERVAL trial to determine whether intervals between blood donations can be safely and acceptably decreased to optimise blood supply: study protocol for a randomised controlled trial. *Trials*. 2014;15:363.



blood[®]

2017 129: 520-524

doi:10.1182/blood-2016-08-732248 originally published
online November 14, 2016

Rare variants in *GP1BB* are responsible for autosomal dominant macrothrombocytopenia

Suthesh Sivapalaratnam, Sarah K. Westbury, Jonathan C. Stephens, Daniel Greene, Kate Downes, Anne M. Kelly, Claire Lentaigne, William J. Astle, Eric G. Huizinga, Paquita Nurden, Sofia Papadia, Kathelijne Peerlinck, Christopher J. Penkett, David J. Perry, Catherine Roughley, Ilenia Simeoni, Kathleen Stirrups, Daniel P. Hart, R. Campbell Tait, Andrew D. Mumford, NIHR BioResource, Michael A. Laffan, Kathleen Freson, Willem H. Ouwehand, Shinji Kunishima and Ernest Turro

Updated information and services can be found at:

<http://www.bloodjournal.org/content/129/4/520.full.html>

Articles on similar topics can be found in the following Blood collections

[Brief Reports](#) (1967 articles)

[Free Research Articles](#) (4889 articles)

[Platelets and Thrombopoiesis](#) (774 articles)

[Thrombocytopenia](#) (245 articles)

[Thrombosis and Hemostasis](#) (1134 articles)

Information about reproducing this article in parts or in its entirety may be found online at:

http://www.bloodjournal.org/site/misc/rights.xhtml#repub_requests

Information about ordering reprints may be found online at:

<http://www.bloodjournal.org/site/misc/rights.xhtml#reprints>

Information about subscriptions and ASH membership may be found online at:

<http://www.bloodjournal.org/site/subscriptions/index.xhtml>