

Quantifying Transmission of *Campylobacter jejuni* in Commercial Broiler Flocks^{∇†}

Twan van Gerwe,^{1*} Jeanette K. Mifflin,² Jillian M. Templeton,² Annemarie Bouma,¹
Jaap A. Wagenaar,³ Wilma F. Jacobs-Reitsma,⁴
Arjan Stegeman,¹ and Don Klinkenberg¹

Department of Farm Animal Health, Faculty of Veterinary Medicine, Utrecht University, P.O. Box 80151, 3508 TD Utrecht, The Netherlands¹; Department of Primary Industries and Fisheries (Queensland), Animal Research Institute, Yeerongpilly, Australia²; Department of Infectious Disease and Immunology, Faculty of Veterinary Medicine, Utrecht University, P.O. Box 80165, 3508 TD Utrecht, The Netherlands³; and Rikilt, Institute of Food Safety, Wageningen UR, P.O. Box 230, 6700 AE Wageningen, The Netherlands⁴

Received 18 August 2008/Accepted 23 November 2008

Since meat from poultry colonized with *Campylobacter* spp. is a major cause of bacterial gastroenteritis, human exposure should be reduced by, among other things, prevention of colonization of broiler flocks. To obtain more insight into possible sources of introduction of *Campylobacter* into broiler flocks, it is essential to estimate the moment that the first bird in a flock is colonized. If the rate of transmission within a flock were known, such an estimate could be determined from the change in the prevalence of colonized birds in a flock over time. The aim of this study was to determine the rate of transmission of *Campylobacter* using field data gathered for 5 years for Australian broiler flocks. We used unique sampling data for 42 *Campylobacter jejuni*-colonized flocks and estimated the transmission rate, which is defined as the number of secondary infections caused by one colonized bird per day. The estimate was 2.37 ± 0.295 infections per infectious bird per day, which implies that in our study population colonized flocks consisting of 20,000 broilers would have an increase in within-flock prevalence to 95% within 4.4 to 7.2 days after colonization of the first broiler. Using Bayesian analysis, the moment of colonization of the first bird in a flock was estimated to be from 21 days of age onward in all flocks in the study. This study provides an important quantitative estimate of the rate of transmission of *Campylobacter* in broiler flocks, which could be helpful in future studies on the epidemiology of *Campylobacter* in the field.

Campylobacter spp. are a common cause of diarrhea in humans, and many cases of campylobacteriosis are associated with the handling and consumption of contaminated poultry meat (5). Studies of the epidemiology of *Campylobacter* have resulted in the implementation of biosecurity and hygienic measures on poultry farms and slaughterhouses with the ultimate goal of reducing human exposure (6, 16). These measures likely contributed to a reduction in the number of *Campylobacter*-positive broiler flocks. Nevertheless, contaminated meat is still on the market (10), and a further reduction in the prevalence of *Campylobacter*-positive flocks is considered necessary by public health authorities in many countries (4).

Clearly, more knowledge concerning the mechanism of introduction of *Campylobacter* into a flock is essential for improving the current control programs. This, in turn, requires an estimate of the moment that a flock becomes colonized. However, it does not seem feasible to detect the first bird that is colonized in a commercial broiler flock because of flock size and the necessary sampling frequency.

An alternative approach is to determine the transmission

rate (β) of *Campylobacter* within a flock. β , which is defined as the number of secondary infections caused by one colonized bird per day, determines the rate of increase in the number of colonized birds over time. It can be used to determine the moment of introduction from field data for increasing *Campylobacter* prevalence over time. The estimates for β that have been obtained in experimental studies (15, 17) are 1.04 to 1.13 per day. However, experimental conditions differ substantially from the field situation, which implies that the β in commercial flocks should also be estimated.

A series of field studies in Australia were carried out between 1999 and 2004 in which broiler flocks were sampled daily to weekly. The aim of these studies was to develop an understanding of the epidemiology of *Campylobacter* in Australian broiler flocks. We analyzed data from the unique data set obtained in these studies to estimate the rate of *Campylobacter* transmission in commercial broiler flocks. Additionally, we estimated the moment that the first bird in a flock was colonized with *Campylobacter* (for reasons of convenience, we refer to this event as the moment of introduction of *Campylobacter* into a flock) and assessed how accurately moments of introduction can be estimated.

MATERIALS AND METHODS

Data set. Three longitudinal studies were carried out in southeast Queensland, Australia, between 1999 and 2004. A subset of the full data set was selected for this study; for flocks to be included in this study, they had to have been sampled at least twice, and at least one *Campylobacter*-positive dropping had to be de-

* Corresponding author. Mailing address: Department of Farm Animal Health, Faculty of Veterinary Medicine, Utrecht University, Yalelaan 7, 3584 CL Utrecht, The Netherlands. Phone: (31) 30 2531248. Fax: (31) 30 2521887. E-mail: tvgerwe@gmail.com.

† Supplemental material for this article may be found at <http://aem.asm.org/>.

∇ Published ahead of print on 1 December 2008.

tected. Flocks with migration barriers separating groups of birds within a shed and flocks whose sampling age was not stated were excluded. A total of 42 flocks met all the inclusion criteria (see the supplemental material for details).

At each sampling age randomly selected individual fecal or cecal droppings were collected immediately after the droppings were produced. Birds that produced these droppings were not marked, as the likelihood of sampling a single bird more than once a day was considered limited for the large populations used. Care was taken to ensure that each sample was collected without any contaminating material. The samples were collected with a sterile swab, which was immediately placed in a sterile container. Samples were kept on ice during transport to the laboratory and were streaked on Karmali *Campylobacter* agar base (Oxoid CM935; Oxoid, Melbourne, Australia) containing *Campylobacter* selective supplement (Oxoid SR167E) immediately after arrival in the laboratory. Agar plates were incubated at 42°C for 48 h in an incubator with an atmosphere consisting of 85% N₂, 10% CO₂, and 5% O₂. Colony morphology and cell motility as determined by phase-contrast microscopy were used to confirm identification to the genus level. Single colonies from a number of positive samples from each positive flock were subcultured on sheep blood agar and incubated as described above before they were identified to the species level by oxidase, catalase production, and hippurate hydrolysis tests.

The final data set consisted of data for sampling events for each flock, including the age of the flock (t), sample size (n_t), flock size at the start of the rearing period (N), and number of positive samples (x_t) (see the supplemental material for details).

Modeling of *Campylobacter* transmission in broiler flocks. It is generally assumed that following the onset of *Campylobacter* colonization of the gut, broilers shed these bacteria for the rest of their lives (1). Therefore, we assumed that birds were either susceptible (noncolonized) or infectious (colonized) and that an increase in prevalence could be described by a susceptible-infectious type of mathematical model (12, 17). In this model, susceptible chicks can be colonized upon contact with an infectious chick, which occurs at rate $\beta i(t)$, where $i(t)$ is the proportion of infectious chicks in the shed. β is considered the mean number of chickens that can be colonized by one infectious chicken per day in a susceptible population.

In the large populations of broilers usually present in commercial flocks, the change in $i(t)$ can be approximated by the deterministic differential equation:

$$\frac{di(t)}{dt} = \beta[1 - i(t)]i(t) \quad (1)$$

whose the solution is the logistic curve described by

$$i(t) = \frac{e^{\beta(t - \tau)}}{1 + e^{\beta(t - \tau)}} \quad (2)$$

in which τ is the time at which 50% of the birds are infected. From τ , β , and N , the time when transmission starts (t_0) is calculated as follows:

$$t_0 = \tau - \log(N - 1)/\beta \quad (3)$$

Estimation of β in the study population. Because the increase in $i(t)$ follows a logistic curve, β is estimated by a logistic regression analysis of x_t . However, this is possible only if for at least two samples only some of the swabs are positive, because only then is the steepness of the increase in prevalence (determined by β) observed. Eight flocks met this criterion. The susceptible-infectious model was fitted for each of these eight flocks by logistic regression of the binomially distributed x_t (and n_t) with t as covariate and $\exp(\tau)$ as the intercept. The model fits resulted in eight estimates for β and τ , from which t_0 could be estimated by using equation 3. Confidence intervals for β and t_0 were derived by profile likelihood analysis (3), which was carried out using Mathematica (version 6.0; Wolfram Research, Inc. [http://www.wolfram.com]).

Because flocks with low β have been more likely to meet the inclusion criteria for the logistic regression analysis than flocks with high β , bias would be introduced if the eight estimates were considered representative of all commercial broiler flocks. Therefore, in the next step, we estimated the mean (μ_β) and standard deviation (σ_β) of a normal distribution of β among all flocks by maximizing the likelihood function

$$l(\mu_\beta, \sigma_\beta) = \prod_{\text{flocks}} \int_0^\infty p(\beta) \prod_t i(t)^{x_t} [1 - i(t)]^{n_t - x_t} d\beta \quad (4)$$

where $p(\beta)$ is the density of the normal distribution of β , $i(t)$ is the logistic curve for equation 2, and t , x_t , and n_t are the data for all flocks. This maximum likelihood estimation method does not result in any flock-specific estimates for β




FIG. 1. (Top) Eight estimates of *Campylobacter* β (with 95% confidence intervals), obtained by logistic regression. (Bottom) Estimated density of the normal distribution of β in the Australian study population, obtained by maximum likelihood estimation ($\mu_\beta = 2.37 \text{ day}^{-1}$; $\sigma_\beta = 0.295 \text{ day}^{-1}$).

or t_0 ; estimation results are limited to μ_β and σ_β , with 95% confidence intervals derived by profile likelihood (3). Although β cannot be negative, the assumption of a normal distribution is valid, because the distribution is well above zero (see Results). The analysis was carried out using Mathematica.

Estimation of the t_0 for the study population. Because no flock-specific estimates of the moment of introduction (t_0) could be obtained with the analysis described above, we used Markov chain Monte Carlo integration to obtain Bayesian posterior distributions of β and τ , and therefore t_0 , for each flock separately (7). The prior distribution of τ was uninformative (flat), whereas the prior distribution of β was the normal distribution resulting from maximum likelihood estimation (with $\mu_\beta = 2.37 \text{ day}^{-1}$ and $\sigma_\beta = 0.295 \text{ day}^{-1}$). The likelihood function was

$$l(\beta, \tau) = \prod_t i(t)^{x_t} [1 - i(t)]^{n_t - x_t} \quad (5)$$

We used the maximum likelihood results for all 42 flocks as input for estimation of t_0 in the same flocks, but it would have been more correct to divide the data into two mutually exclusive subsets and use each subset only once. However, the approach which we used resulted in more precise estimates of μ_β and σ_β and estimates of t_0 for all flocks instead of only a subset. The possible errors resulting from our approach were minor, as determined by a separate sensitivity analysis of estimation of t_0 to correctness of the prior distribution of β (see below).

The posterior distribution was sampled 10,000 times by single-component Metropolis-Hastings sampling, after 100 samples were used for burn-in (8). The means of the (normal) proposal distributions for the $(i + 1)$ th samples of both β and τ were the i th samples. The standard deviation of the proposed distribution of β was 0.295 day^{-1} ; the standard deviation of the proposal for τ was determined from the data set, and it was one-quarter of the time interval between the last 0% prevalence sample and first 100% prevalence sample. This resulted in means and 95% credible intervals for t_0 for 40 of the 42 flocks (for 2 flocks, the flock size was not known). The sampling algorithm was programmed using Mathematica.

Accuracy of t_0 estimation. Although Bayesian estimation of t_0 does provide 95% credible intervals, these intervals are based on a logistic curve which does not take account of the stochastic nature of transmission in the early phase of a *Campylobacter* outbreak. Therefore, we assessed the accuracy of the method by estimating t_0 using simulated outbreaks in which the estimates could be compared to the real value.

We simulated 10,000 outbreaks in flocks consisting of 20,000 chicks with values for β sampled from the estimated distribution ($\mu_\beta = 2.37 \text{ day}^{-1}$; $\sigma_\beta = 0.295 \text{ day}^{-1}$), with a t_0 of 0. Simulations were carried out as described previously (17) with three sample sizes (10, 20, and 100 birds) and three sampling frequencies (once every day, every third day, or every seventh day). This resulted in 10,000 simulated data sets for all combinations of sample sizes and sample frequencies, which were subsequently analyzed by the Bayesian method described above to obtain posterior means for 1,000 samples of the posterior distribution.

In addition to these 10,000 simulations, where the population distribution of β (used in the simulations) was identical to the prior distribution in the Bayesian




FIG. 2. Epidemic curves for three different values of the estimated distribution of β (lower limit, point estimate, and upper limit) and an experimentally derived β estimate (17) for flocks consisting of 20,000 broilers.

analysis, we assessed the accuracy by simulation with other distributions, based on the confidence intervals for μ_β (2.19 to 2.58 day⁻¹) and σ_β (0.144 to 0.488 day⁻¹). Four new data sets were simulated, including two data sets with different μ_β (2.19 and 2.58 day⁻¹, with σ_β kept at 0.295 day⁻¹) and two data sets with different σ_β (0.144 and 0.488 day⁻¹, with μ_β kept at 2.37 day⁻¹). This resulted in four sets of 1,000 simulated data sets for 10 samples every 1, 3, or 7 days. These data were used to estimate t_0 as described above.

RESULTS

Data set. The flock age at the time of first detection of *Campylobacter*-positive droppings varied from 24 to 54 days. For 14 of 42 flocks the prevalence was more than 95% at the time of first detection. For an additional 10 flocks in which the prevalence was up to 10% at the time of first detection, sampling was continued until the flocks were found to be fully colonized (>95% prevalence), which took 4.4 days on average (range, 3 to 6 days) (see the supplemental material for details).

For 38 of 42 flocks, all *Campylobacter* isolates were identified as *Campylobacter jejuni*. Both *C. jejuni* and *Campylobacter coli* were found in four flocks. Because insufficient data concerning *C. coli* outbreaks were available, our analyses were based solely on *C. jejuni* outbreaks.




FIG. 3. Histogram showing the t_0 for 40 Australian *C. jejuni* outbreaks.

TABLE 1. Accuracy of Bayesian estimation (with prior distribution $\mu_\beta = 2.37$ day⁻¹ and $\sigma_\beta = 0.295$ day⁻¹) of the posterior mean time of *Campylobacter* introduction into a flock of 20,000 broilers, with different sample sizes and sampling intervals

μ_β (day ⁻¹)	σ_β (day ⁻¹)	No. of data sets	Sample size	90% Interval of estimation error		
				Each day	Each 3rd day	Each 7th day
2.37	0.295	10,000	10	-0.9 to 1.7	-1.0 to 1.8	-2.1 to 2.7
		10,000	20	-0.8 to 1.6	-0.9 to 1.7	-1.8 to 2.4
		10,000	100	-0.6 to 1.4	-0.8 to 1.6	-1.3 to 2.1
2.19	0.295	1,000	10	-0.7 to 2.2	-0.8 to 2.4	-1.7 to 3.1
2.58	0.295	1,000	10	-1.1 to 1.3	-1.4 to 1.3	-2.5 to 2.3
2.37	0.144	1,000	10	-0.8 to 1.4	-0.8 to 1.4	-1.9 to 2.5
2.37	0.488	1,000	10	-1.0 to 2.3	-1.3 to 2.7	-2.4 to 3.3

Estimation of β and t_0 for the study population. For eight flocks with sufficient data, flock-specific β and t_0 were estimated by logistic regression. The point estimates for β ranged from 1.3 to 3.1 day⁻¹ (Fig. 1), and t_0 ranged from 21 to 35 days (see the supplemental material for details).

The estimates for the distribution of β , which were based on 42 outbreaks, were as follows: $\mu_\beta = 2.37$ day⁻¹ (95% confidence interval, 2.19 to 2.58 day⁻¹) and $\sigma_\beta = 0.295$ (95% confidence interval, 0.144 to 0.488 day⁻¹) (Fig. 1). Figure 2 shows how fast flocks of 20,000 broilers were colonized for the mean β , a low β , and a high β within the distribution of β .

For 40 *Campylobacter* outbreaks in which the population size was known we estimated t_0 by Bayesian analysis. The results (Fig. 3) show that colonization did not take place before chickens reached an age of 3 weeks. On average, t_0 was 4.8 days (range, 2.2 to 9.3 days) earlier than the first detection of *Campylobacter*-positive samples (see the supplemental material for details).

The estimates obtained by logistic regression and Bayesian analysis did not differ more than 0.6 day except for one flock (3.4-day difference).

Accuracy of t_0 estimation. The accuracy of the Bayesian method was assessed by analyzing simulated outbreaks; the results are shown in Table 1. If the prior distribution used for the analysis reflected the underlying β in the population cor-

rectly, the t_0 was estimated within 3 days even with weekly sampling of only 10 birds.

The Bayesian estimation method was not very sensitive to the correctness of the prior distribution (within the margins of the 95% confidence intervals), and the sensitivity was greatest if μ_β was lower or σ_β was higher, resulting in a margin of 3.3 days with weekly sampling of 10 birds.

DISCUSSION

The first aim of this study was to determine the β of *Campylobacter* in commercial broiler flocks. β , based on the field data, was estimated to be $2.37 \pm 0.295 \text{ day}^{-1}$, which means that one colonized bird could, on average, infect 2.37 birds per day. This implies that in a flock of 20,000 broilers the within-flock prevalence of *Campylobacter* would increase from a prevalence of one colonized bird to a prevalence of 95% within 1 week.

Previous studies showed that flocks determined to be *Campylobacter* negative at one moment could appear to be fully colonized within 1 week, suggesting that *Campylobacter* was introduced in the intervening week (2, 9, 11). This seems consistent with our findings, but the sample size used in these studies did not allow detection of positive flocks at the start of an epidemic. This implies that bacteria could have been introduced at some previous time.

The second aim of this study was to estimate the moment of colonization of the first bird in a flock (t_0) by using the estimated β . The estimates of t_0 were all greater than a flock age of 21 days. The estimate of t_0 was accurate within 3 days in 90% of the cases, even if only 10 birds were sampled weekly. The sensitivity to incorrectness of the prior distribution was low. The apparent lack of *Campylobacter* transmission in the first 3 weeks of life, which seemed to have occurred in our study, has been described previously (11). It is difficult or probably impossible to determine whether in the first 3 weeks no introduction occurred or whether introduction of bacteria did occur but did not result in colonization of the birds. The risk of introduction of *Campylobacter* can be high throughout the rearing period, but it could also be hypothesized that this risk may increase over time. Another possibility is that young chicks are less susceptible, as indicated by Ringoir et al. (13) and Sahin et al. (14), who demonstrated that age and maternally derived immunity had an effect on susceptibility. More insight into the underlying mechanism of this phenomenon may provide clues for prevention of *Campylobacter* colonization.

Conclusions. A unique data set describing the change in prevalence of *Campylobacter*-colonized birds in commercial broiler flocks was used to quantify *C. jejuni* transmission under field conditions. The estimated β implies that in a flock of 20,000 broilers the within-flock prevalence of *Campylobacter* increases to 95% within 1 week after colonization of the first bird. Since such rapid spread has not been described previously on the basis of experimental transmission experiments, this study provides important new quantitative information on the epidemiology of *Campylobacter* in broilers.

This study also showed how the β can be used to estimate when the first bird is colonized. We showed that this method

for estimation was accurate, and therefore this method is a promising method for further studies of mechanisms of *Campylobacter* introduction, because it allows focusing on the chronology of events.

Additionally, t_0 estimation can result in an accurate description of the period in which no transmission of *Campylobacter* occurs, like the first 3 weeks for our study population. Consequently, interventions aimed at prevention of introduction of and subsequent colonization by *Campylobacter* might better be targeted at the second half of the rearing period, which in our study population could be considered a high-risk period.

ACKNOWLEDGMENT

This work was funded in part by the Australian funding body RIRDC.

REFERENCES

- Achen, M., T. Y. Morishita, and E. C. Ley. 1998. Shedding and colonization of *Campylobacter jejuni* in broilers from day-of-hatch to slaughter age. *Avian Dis.* **42**:732–737.
- Allen, V. M., H. Weaver, A. M. Ridley, J. A. Harris, M. Sharma, J. Emery, N. Sparks, M. Lewis, and S. Edge. 2008. Sources and spread of thermophilic *Campylobacter* spp. during partial depopulation of broiler chicken flocks. *J. Food Prot.* **71**:264–270.
- Dohoo, I., W. Martin, and H. Stryhn. 2003. *Veterinary epidemiologic research*. AVC Inc., Charlottetown, Canada.
- EFSA 27 January 2005, posting date. Opinion of the Scientific Panel on Biological Hazards (BIOHAZ) related to *Campylobacter* in animals and foodstuffs. http://www.efsa.europa.eu/EFSA/efsa_locale-1178620753812_1178620776955.htm.
- Friedman, C. R., R. M. Hoekstra, M. Samuel, R. Marcus, J. Bender, B. Shiferaw, S. Reddy, S. D. Ahuja, D. L. Helfrick, F. Hardnett, M. Carter, B. Anderson, and R. V. Tauxe. 2004. Risk factors for sporadic *Campylobacter* infection in the United States: a case-control study in FoodNet sites. *Clin. Infect. Dis.* **38**:S285–S296.
- Gellynck, X., W. Messens, D. Halet, K. Grijspeerdt, E. Hartnett, and J. Viaene. 2008. Economics of reducing *Campylobacter* at different levels within the Belgian poultry meat chain. *J. Food Prot.* **71**:479–485.
- Gelman, A., J. B. Carlin, H. S. Stern, and D. B. Rubin. 2003. *Bayesian data analysis*, 2nd ed. CRC Press, Boca Raton, FL.
- Gilks, W. R., S. Richardson, and D. J. Spiegelhalter. 1996. *Markov chain Monte Carlo in practice*. CRC Press, Boca Raton, FL.
- Hald, B., E. Rattenborg, and M. Madsen. 2001. Role of batch depletion of broiler houses on the occurrence of *Campylobacter* spp. in chicken flocks. *Lett. Appl. Microbiol.* **32**:253–256.
- Hansson, I., L. P. Forshell, P. Gustafsson, S. Boqvist, J. Lindblad, E. O. Engvall, Y. Andersson, and I. Vagsholm. 2007. Summary of the Swedish *Campylobacter* program in broilers, 2001 through 2005. *J. Food Prot.* **70**:2008–2014.
- Jacobs-Reitsma, W. F., A. W. van de Giessen, N. M. Bolder, and R. W. Mulder. 1995. Epidemiology of *Campylobacter* spp. at two Dutch broiler farms. *Epidemiol. Infect.* **114**:413–421.
- Keeling, M. J., and P. Rohani. 2008. *Modeling infectious diseases in humans and animals*. Princeton University Press, Princeton, NJ.
- Ringoir, D. D., D. Szylo, and V. Korolik. 2007. Comparison of 2-day-old and 14-day-old chicken colonization models for *Campylobacter jejuni*. *FEMS Immunol. Med. Microbiol.* **49**:155–158.
- Sahin, O., N. Luo, S. Huang, and Q. Zhang. 2003. Effect of *Campylobacter*-specific maternal antibodies on *Campylobacter jejuni* colonization in young chickens. *Appl. Environ. Microbiol.* **69**:5372–5379.
- Stern, N. J., N. A. Cox, and M. T. Musgrove. 2001. Incidence and levels of *Campylobacter* in broilers after exposure to an inoculated seeder bird. *J. Appl. Poult. Res.* **10**:315–318.
- van den Giessen, A. W., J. J. H. C. Tilburg, W. S. Ritmeester, and J. van der Plas. 1998. Reduction of *Campylobacter* infections in broiler flocks by application of hygiene measures. *Epidemiol. Infect.* **121**:57–66.
- Van Gerwe, T. J. W. M., A. Bouma, W. F. Jacobs-Reitsma, J. van den Broek, D. Klinkenberg, J. A. Stegeman, and J. A. P. Heesterbeek. 2005. Quantifying transmission of *Campylobacter* spp. among broilers. *Appl. Environ. Microbiol.* **71**:5765–5770.