

Quantitative and qualitative analysis of antimicrobial usage patterns in 180 selected farrow-to-finish pig farms from nine European countries based on single batch and purchase data

Steven Sarrazin¹, Philip Joosten^{1*}, Liese Van Gompel ², Roosmarijn E. C. Luiken², Dik J. Mevius^{3,4}, Jaap A. Wagenaar^{3,4}, Dick J. J. Heederik² and Jeroen Dewulf ¹ on behalf of the EFFORT consortium†

¹Veterinary Epidemiology Unit, Department of Reproduction, Obstetrics and Herd Health, Faculty of Veterinary Medicine, Ghent University, Salisburylaan 133, 9820 Merelbeke, Belgium; ²Institute for Risk Assessment Sciences, Faculty of Veterinary Medicine, Utrecht University, Yalelaan 2, 3584 CM Utrecht, The Netherlands; ³Department of Infectious Diseases and Immunology, Faculty of Veterinary Medicine, Utrecht University, Yalelaan 1, 3584 CL Utrecht, The Netherlands; ⁴Wageningen Bioveterinary Research, Houribweg 39, 8221 RA Lelystad, The Netherlands

*Corresponding author. Tel: +32 92647548; Fax: +32 92647534; E-mail: Philip.Joosten@UGent.be
†Members are listed in the Acknowledgements section.

Received 22 May 2018; returned 9 August 2018; revised 10 October 2018; accepted 8 November 2018

Objectives: Farm-level quantification of antimicrobial usage (AMU) in pig farms.

Methods: In a cross-sectional study, AMU data on group treatments administered to a single batch of fattening pigs from birth to slaughter (group treatment data) and antimicrobials purchased during 1 year (purchase data) were collected at 180 pig farms in nine European countries. AMU was quantified using treatment incidence (TI) based on defined (DDDvet) and used (UDDvet) daily doses and defined (DCDvet) and used (UCDvet) course doses.

Results: The majority of antimicrobial group treatments were administered to weaners (69.5% of total TI_{DDDvet}) followed by sucklers (22.5% of total TI_{DDDvet}). AMU varied considerably between farms with a median TI_{DDDvet} of 9.2 and 7.1 for a standardized rearing period of 200 days based on group treatment and purchase data, respectively. In general, UDDvet and UCDvet were higher than DDDvet and DCDvet, respectively, suggesting that either the defined doses were set too low or that group treatments were often dosed too high and/or administered for too long. Extended-spectrum penicillins (31.2%) and polymyxins (24.7%) were the active substances most often used in group treatments, with the majority administered through feed or water (82%). Higher AMU at a young age was associated with higher use in older pigs.

Conclusions: Collecting farm-level AMU data of good quality is challenging and results differ based on how data are collected (group treatment data versus purchase data) and reported (defined versus used daily and course doses).

Introduction

During their general assemblies of 2015, WHO, the World Organisation for Animal Health (OIE) and the Food and Agriculture Organization (FAO) re-emphasized the importance of the public health crisis posed by antimicrobial resistance (AMR).^{1–3} Since there are strong indications of animal–human transmission of AMR^{4,5} and that antimicrobial usage (AMU) is the strongest driver for selection of AMR,^{6,7} it is important to reduce veterinary AMU.⁸

To target AMU in food-producing animals, WHO and the OIE recommend that the quantities of antimicrobials used in food-producing animals are monitored.^{9,10} In Europe, sales data of antimicrobials are monitored in a standardized manner by the

European Surveillance of Veterinary Antimicrobial Consumption (ESVAC) project and reports show that considerable differences exist between countries.¹¹ In ESVAC reports, no distinction is made between livestock species, whereas it is known that there are differences in AMU between species. Pig production accounts for a large proportion of AMU in animal production.^{12–14} Furthermore, ESVAC sales figures at the country level do not take into account the large differences in dosage between different antimicrobial compounds nor show how antimicrobials are used between different age categories of livestock species.⁸

Standardized AMU quantification will allow for benchmarking between farms in different geographical areas, regions, etc., and

will also allow the study of its association with animal health, production characteristics and AMR. However, before the aforementioned associations with AMU can be explored, more detailed data such as the assessment of exposure (e.g. exposure duration) are required at the farm level.¹⁵

Therefore, this study aimed to quantify AMU for pigs at the farm level in nine European countries in a standardized manner. This study was conducted within the European research project EFFORT (Ecology from Farm to Fork Of microbial drug Resistance and Transmission), which investigates the epidemiology and ecology of AMR in food-producing animals, the environment and humans to quantify AMR exposure pathways for humans.

Materials and methods

Ethics

Written informed consent from the participating farmers was obtained.

Selection of farms

A cross-sectional study was conducted in Belgium, Bulgaria, Denmark, France, Germany, Italy, Poland, Spain and the Netherlands. To obtain a sample of farms that were comparable within and between countries, 20 conventional farrow-to-finish pig farms, with no other livestock for commercial intentions, were selected in each country according to the following inclusion criteria: a minimum of 150 sows and 600 finishing places and using a batch management system. The farms were required not to have contact with each other through trade and each farm had only one owner. Farms were selected based on these criteria in agreement with local farming organizations, and also partially based on convenience (e.g. distances to farms). As a result, the sample of farms in each country cannot be considered representative for the pig sector in that country. The country-specific selection procedure (Appendix A) together with characteristics of the sampled farms (Table S1) and an illustration of the swine industry in each participating country (Figure S1) are available as [Supplementary data](#) at JAC Online and elsewhere.¹⁶

Collection of AMU records

The farms were visited between June 2014 and December 2015. From each farm, data were collected on antimicrobial group treatments from birth to slaughter during the rearing of one batch of fattening pigs (further referred to as group treatment data). A group treatment was defined as each treatment applied simultaneously to all animals present in, at least, the smallest housing unit. Individual treatments were not included as these are prone to recall bias as they are often not well recorded.¹⁷ Furthermore, data on antimicrobials purchased during the year preceding the visit were acquired (further referred to as purchase data). Data collection was based on a questionnaire developed within the EFFORT consortium (Appendix B). The collected variables applied in the AMU quantification are described in Appendix C.

Data processing

Data were entered into EpiData 3.1 (EpiData Association, Denmark) and checked for quality. Further data quality checks were performed using ActivePerl 5.24.1 (ActiveState Software Inc.) and SAS 9.4 (SAS Institute Inc., Cary, NC, USA). Ten percent of all questionnaires were entered twice and compared in SAS to check for data entry quality. All inconsistencies were corrected and datasets from countries with >5% of inconsistencies were thoroughly re-evaluated. Data required for AMU quantification were imported into Microsoft Excel and a second, more in-depth, data quality control was performed. Finally, an online database was created which

consisted of all study data and metadata. All farms were anonymized to ensure that results cannot be traced back to individual farms. Countries were anonymized as required by one participating country.

AMU quantification

Based on Defined Daily Doses Animal (DDDvet), Used Daily Doses Animal (UDDvet), Defined Course Doses Animal (DCDvet), Used Course Doses Animal (UCDvet) and long-acting (LA) factor (a value to represent the duration of activity of an LA product), the treatment incidence (TI) indicator was used to quantify AMU, in accordance with Timmerman et al.¹⁸:

$$TI = \frac{\text{total amount of antimicrobials administered or purchased}}{\text{DDDvet or UDDvet or DCDvet or UCDvet} \times \text{number of days at risk} \times \text{kg of animal at risk} \times \text{LA factor} \times 100 \text{ pigs at risk}}$$

TI_{DDDvet} and TI_{UDDvet} express the percentage of pigs that received a dose of antimicrobials each day or, equivalently, during which percentage of time a pig was treated with antimicrobials in a certain production phase or its entire life.¹⁸ TI_{DCDvet} and TI_{UCDvet} express the percentage of pigs for which a treatment was initiated.

UDDvet values were derived from the data on the truly administered products and UCDvet values were obtained by multiplying UDDvet with the treatment duration. DDDvet and DCDvet values were used as provided by ESVAC.¹⁹ Whenever, for a given combination of active substance and administration route, no DDDvet and DCDvet values were defined by ESVAC, the values as described by Postma et al.²⁰ were used or the data were obtained from the summary of product characteristics of that specific product. LA factors were used as described by Postma et al.²⁰ To obtain the treatment duration for LA formulations the number of days the treatment was applied was multiplied by the LA factor. The detailed calculation of TI is described in Appendix C, Tables S2 and S3.

TI calculations were performed for each administered or purchased product per age category. In accordance with Sjölund et al.⁸, the TI of sucklers, weaned piglets and finishers were combined and recalculated into a standardized lifespan of 200 days to correct for possible differences in ages at slaughter between farms and countries and to express AMU from birth to slaughter (TI₂₀₀):

$$TI_{200} = \frac{TI_{\text{suckler}} \times \text{suckling period} + TI_{\text{weaner}} \times \text{nursery period} + TI_{\text{finisher}} \times \text{finishing period}}{\text{life span in a farm}} \times \frac{200 \text{ days life span}}{\text{life span in a farm}}$$

To obtain the total TI with respect to a categorical variable (e.g. country), all TIs for the different antimicrobials were summed within the respective category of this variable.

Statistical analysis

Since TIs deviated from normality, median results are reported together with 10th and 90th percentiles. Relative deviations between TI_{DDDvet} and TI_{UDDvet} [(TI_{DDDvet}-TI_{UDDvet})/TI_{DDDvet}] and between TI_{DCDvet} and TI_{UCDvet} [(TI_{DCDvet}-TI_{UCDvet})/TI_{DCDvet}] were calculated to explore to what extent UDD_{vet} and UCD_{vet} matched with DDDvet and DCDvet, respectively. Positive deviations indicated that a higher dose was used than defined by ESVAC. Associations between TI_{DDDvet} and TI_{UDDvet} and between TI_{DCDvet} and TI_{UCDvet} were estimated with Spearman correlation coefficients. Relationships between TIs for different age categories were compared by fitting Loess curves after TI_{DDDvet} variables were log₁₀-transformed to deal with the problem of a skewed distribution. Before transforming the data, a value of 1 was added to all observations




Figure 1. This histogram shows the number of antimicrobial group treatments per week applied to a batch of fattening pigs from birth to slaughter based on 750 treatments (30 instances of treatments missing). The stacked bars show the distribution of treatments between countries. The full black and grey lines denote the frequency distribution of the weaning age and the age of the weaners at transfer to the grow-finishing barns, respectively.

to overcome the problem of zero use in a particular age category.⁸ The relationships between TIs of fattening pigs (sucklers, weaners and finishers) were based on group treatment data, while the relationship between TIs of sows and sucklers was based on purchase data. Cross-tabulation was used and graphs were prepared to visualize the results with R version 3.3.1 (<https://cran.r-project.org>).

Results

AMU quantification using TI_{DDvet}

Antimicrobials were administered as group treatments on 780 occasions by 159 farms, meaning that 21 farms (11.7%) did not report antimicrobial group treatments from birth to slaughter for the batch of pigs from which the data were collected (Table S4, Appendix D). The majority of the treatments were applied in the beginning of the rearing period, with peaks at week 1 (16.9%), week 4 (10.9%) and week 9 (7.3%) (Figure 1). Based on group treatment data, TI_{200} varied considerably between the samples of farms within a country (Figure 2, Table 1), with a median TI_{200} of 9.2 (10th–90th percentile: 0.0–46.6) and a maximum of 113.6. Regarding age categories, the highest AMU was observed in weaners (69.5% of total TI_{DDvet}), followed by sucklers (22.5% of total TI_{DDvet}) and finishers (8.0% of total TI_{DDvet}). This observation was also confirmed at the country level, except for country C, where AMU in sucklers was higher compared with weaners (Table S5, Appendix D).

Antimicrobial products were purchased by 168 farms, meaning that 12 (6.7%) farms did not report any purchased products during the year preceding the visit (Table S4, Appendix D). For 30% of the farms purchasing antimicrobial products, the period for which they reported to have purchased antimicrobials was less than 6 months. Furthermore, for 52 purchased products (3.1%), mainly from country F ($n = 45$), the age category for which the antimicrobial was purchased was missing and these missing values were imputed based on other observations for the same product and on the knowledge about the product.

Based on purchase data, the median TI_{200} was 7.1 (10th–90th percentile: 0.2–35.6), with a maximum TI_{200} of 96.4, and the median TI for sows was 0.2 (10th–90th percentile: 0.0–3.5) (Table 2). The highest TIs were observed in weaners for all countries, except for countries C and F, where AMU was the highest in sucklers.

For corresponding age categories [sucklers, weaners, finishers and from birth to slaughter (TI_{200})] only moderate correlations between group treatment and purchase data were observed for TI_{DDvet} and TI_{DCDvet} , ranging from 0.39 to 0.50 (all $P < 0.001$). Removing the 43 farms for which the period they reported to have purchased antimicrobials was less than 6 months did not improve the associations (data not shown). A very large difference between group treatment and purchase data for country H was observed: antimicrobials administered through premedicated feed were not included in the purchase data because the feed company could




Figure 2. Country-level comparison of TI per 100 pig-days at risk between group treatment data (white boxplots) and purchase data (grey boxplots) for different age categories (sucklers, weaners and finishers) and combined for a standardized lifespan of 200 days (TI200) based on DDDvet. To enhance the comparison between the boxplots some outlying observations are not shown. The full black line within each boxplot represents the median. The whiskers (dashed lines) extend to the most extreme data point within the range of $1.5 \times \text{IQR}$ from the box. Outliers, data points outside the range of the whiskers, are shown as open circles. Group treatment data were obtained from a single batch of fattening pigs from birth to slaughter, while purchase data were data on antimicrobials purchased during 1 year. Whenever the period during which antimicrobials were purchased deviated from 1 year, the amount of antimicrobials purchased was corrected to 1 year. It should be taken into account that for the following farms the period for which the purchase of antimicrobials was reported was <6 months: country B ($n = 2$), country H ($n = 4$), country G ($n = 11$), country C ($n = 17$) and country I ($n = 20$). For country H, antimicrobials administered through premedicated feed were not included in the purchase data because the feed company could not exactly distinguish what proportion of feed was delivered to each specific farm.

Table 1. Overview of the quantification of antimicrobial group treatments administered to a single batch of fattening pigs from birth to slaughter expressed in TI per 100 pigs at risk, based on DDDvet, UDDvet, DCDvet and UCDvet

Country	TI200 _{DDDvet}	TI200 _{UDDvet}	Relative deviation		Relative deviation	
			TI200 _{DDDvet} VS TI200 _{UDDvet} ^a	TI200 _{DCDvet}	TI200 _{UCDvet}	TI200 _{DCDvet} VS TI200 _{UCDvet} ^a
A	9.2 (3.1–31.9)	7.5 (3.4–20.8)	12 (–31 to 52)	1.7 (0.3–7.9)	1.0 (0.5–3.4)	40 (–61 to 69)
B	6.5 (0.8–24.9)	5.3 (0.8–17.6)	19 (–30 to 53)	1.2 (0.4–5.3)	1.1 (0.7–5.2)	–1 (–70 to 41)
C	9.1 (1.1–23.8)	7.1 (1.2–13.1)	17 (–11 to 44)	2.5 (0.2–6.8)	2.4 (0.4–5.8)	6 (–94 to 54)
D	17.5 (9.3–50.1)	13.7 (5.7–34.2)	25 (4–36)	3.6 (1.4–8.1)	3.3 (1.3–4.9)	14 (–18 to 50)
E	5.5 (0.0–11.7)	6.3 (0.0–13.3)	–7 (–81 to 39)	0.6 (0.0–1.6)	0.7 (0.0–1.6)	12 (–71 to 49)
F	0.0 (0.0–4.9)	0.0 (0.0–5.8)	10 (–19 to 39)	0.0 (0.0–0.9)	0.0 (0.0–0.9)	25 (–34 to 44)
G	12.9 (5.4–19.6)	9.8 (6.9–16.9)	20 (–35 to 41)	2.5 (1.3–6.7)	2.0 (1.3–3.5)	28 (–51 to 63)
H	56.7 (13.5–94.1)	55.6 (9.4–83.3)	8 (–30 to 25)	9.2 (4.1–13.1)	5.6 (2.9–7.7)	39 (20–55)
I	0.5 (0.01–3.2)	0.4 (0.01–2.2)	2 (–57 to 68)	0.2 (0.003–0.6)	0.1 (0.002–0.3)	36 (–2 to 75)
Overall	9.2 (0.0–46.6)	7.3 (0.0–44.2)	14 (–37 to 45) 0.96 ^b	1.6 (0.0–9.2)	1.3 (0.0–5.3)	25 (–50 to 60) 0.94 ^b

TI200 summarizes the TIs for the age categories (sucklers, weaners and finishers) for a standardized rearing period of 200 days. TI200_{DDDvet} and TI200_{UDDvet} express the percentage of pigs that received a dose of antimicrobials each day or, equivalently, during which percentage of time a pig was treated with antimicrobials in its entire life. TI200_{DCDvet} and TI200_{UCDvet} express the percentage of pigs for which a treatment was initiated. The relative deviation between TI200_{DDDvet} and TI200_{UDDvet} $[(\text{TI200}_{\text{DDDvet}} - \text{TI200}_{\text{UDDvet}}) / \text{TI200}_{\text{DDDvet}}]$ and between TI200_{DCDvet} and TI200_{UCDvet} $[(\text{TI200}_{\text{DCDvet}} - \text{TI200}_{\text{UCDvet}}) / \text{TI200}_{\text{DCDvet}}]$ is also shown. A positive deviation indicates that a higher dose was used than defined by ESVAC. Median values are shown, with 10th and 90th percentiles in brackets, per country and overall, as well as the overall Spearman correlation for the relative deviations, with *P* value in brackets.

^aValues shown are the median (10th–90th percentile).

^bSpearman correlation coefficients ($P < 0.001$).

Table 2. Overview of the quantification of antimicrobials purchased during 1 year for fattening pigs from birth to slaughter and sows expressed in TI per 100 pigs at risk, based on DDDvet values per age category (sucklers, weaners, finishers and sows) and for fattening pigs combined from birth to slaughter for a standardized rearing period of 200 days (TI200)

Country	TI _{DDvet} sucklers	TI _{DDvet} weaners	TI _{DDvet} finishers	TI _{200DDvet}	TI _{DDvet} SOWS
A	9.1 (0.4–45.5)	23.5 (1.9–67.8)	1.5 (0.0–14.9)	8.1 (2.7–31.4)	0.4 (0.02–3.7)
B	2.4 (0.0–16.8)	13.2 (0.0–45.3)	0.9 (0.01–7.2)	6.3 (0.4–15.0)	0.3 (0.0–3.6)
C	0.3 (0.0–19.4)	0.0 (0.0–24.7)	0.0 (0.0–2.4)	0.6 (0.0–16.1)	0.0 (0.0–0.9)
D	32.5 (8.9–118.0)	135.0 (48.3–254.4)	11.7 (1.8–24.9)	30.1 (14.2–58.5)	0.8 (0.4–6.2)
E	6.3 (0.8–23.6)	14.1 (2.9–94.4)	0.8 (0.0–3.9)	8.4 (2.8–32.1)	0.7 (0.2–4.4)
F	1.6 (0.0–33.7)	1.0 (0.0–4.8)	0.8 (0.0–3.1)	1.8 (0.3–5.9)	0.3 (0.05–4.4)
G	10.4 (0.2–102.9)	60.1 (1.4–150.0)	5.3 (0.1–11.9)	17.3 (3.3–42.7)	0.2 (0.0–4.6)
H ^a	1.8 (0.0–10.2)	15.4 (30.0–37.9)	4.4 (0.9–10.3)	7.6 (4.0–12.0)	0.0 (0.0–0.6)
I	0.0 (0.0–2.1)	0.6 (0.0–40.3)	0.3 (0.0–26.8)	0.6 (0.1–35.1)	0.0 (0.0–0.0)
Overall	4.5 (0.0–36.7)	13.5 (0.0–132.4)	1.2 (0.0–13.5)	7.1 (0.2–35.6)	0.2 (0.0–3.5)

Median values are shown, with 10th and 90th percentiles in brackets, per country. TI_{DDvet} expresses the percentage of pigs that received a dose of antimicrobials each day or, equivalently, during which percentage of time a pig was treated with antimicrobials in a certain production phase or its entire life.

It should be taken into account that for the following farms the period for which the purchase of antimicrobials was reported was <6 months: country B ($n = 2$), country H ($n = 4$), country G ($n = 11$), country C ($n = 17$) and country I ($n = 20$).

^aFor country H, antimicrobials administered through premedicated feed were not included in the purchase data, because the feed company could not exactly distinguish what proportion of feed was delivered to each specific farm.

not exactly distinguish what proportion of feed was delivered to each specific farm (Figure 2).

Relationship between indicators for AMU quantification

Overall a higher daily dose of antimicrobials was administered as group treatment to a batch of fattening pigs from birth to slaughter than doses defined by ESVAC: the median deviation for TI₂₀₀ between TI_{DDvet} and TI_{UDvet} was 14% (10th–90th percentile: –37% to 45%) (Table 1). Except for country E, this median deviation was positive at the country level for all countries (Figure 3). Furthermore, the used course dose was in general higher than the defined course dose, with a median deviation between TI_{DCvet} and TI_{UCvet} of 25% (10th–90th percentile: –50% to 60%). Since the latter deviation was in general larger than the deviation between TI_{DDvet} and TI_{UDvet}, this indicates that besides the use of higher doses the duration of group treatments was also longer than recommended. This was also observed at the country level, since the median deviations between TI_{DCvet} and TI_{UCvet} were larger than between TI_{DDvet} and TI_{UDvet} for all countries, except for B, C and D. Regarding age categories, the largest deviations in used daily and course doses were observed for sucklers and weaners (Tables S5 and S6, Appendix D). Despite large deviations between different indicators for AMU quantification, results were highly correlated at the farm level, with Spearman correlation coefficients of 0.91 or higher (all $P < 0.001$). TI_{DCvet} results for purchase data are presented in Table S7 (Appendix D).

Positive associations between age categories were noticed, indicating that higher AMU at a young age was associated with higher use in older pigs (Figure 4). This was also reflected in Spearman correlation coefficients of 0.48 (sows–sucklers), 0.46

(sucklers–weaners), 0.47 (sucklers–finishers) and 0.37 (weaners–finishers) for the relationship between TIs.

Qualitative results on group treatment data

Based on TI_{DDvet}, the majority of the antimicrobials were administered through feed or water (82%), 18% were administered parenterally and only 0.007% consisted of topical treatments. Considerable differences between countries were observed regarding the administration route: country C administered most of the antimicrobials parenterally (59%), followed by countries G (36%), D (22%), B (18%), I (19%), A (12%), E (11%), F (9%) and H (7%). In terms of TI_{DDvet}, 61% of the treatments of sucklers were parenteral, while the majority of the treatments of weaners and finishers were oral (95% and 90%, respectively).

The average treatment duration for parenterally administered antimicrobials was 2.6 days (minimum 1; median 3; maximum 14) for non-LA formulations and 5.2 days (1.8; 5; 28) for LA formulations, while for oral treatments this was 10.6 days on average (1; 7; 80) and 2.2 days (1; 1; 5) for topical treatments. Moreover, 10% of the oral treatments were applied for a period of at least 21 consecutive days.

In terms of TI_{DDvet}, extended-spectrum penicillins (31.2%) and polymyxins (24.7%) were the most frequently used antimicrobial classes, followed by tetracyclines (15.3%) and macrolides (9.7%) (Table 3). Extended-spectrum penicillins, polymyxins and tetracyclines were mainly used for weaners, while cephalosporins, fluoroquinolones and β -lactamase-sensitive penicillins were more frequently administered to sucklers.

The most frequent indications for treatment were general (37.5%), intestinal (24.4%) and respiratory disorders (20.1%). Less frequently reported indications for treatment were locomotor

Table 3. Proportion (as a percentage) of amount of antimicrobials used as group treatment (based on TI with DDDvet values for a batch of fattening pigs observed from birth to slaughter) by antimicrobial class: total use, within-country and by age category^a

Class	ATCvet code	Total	A	B	C	D	E	F	G	H	I	Suckler	Weaner	Finisher
Aminoglycosides	QJ01G	1.3	—	2.3 (2)	5.5 (3)	1.2 (3)	—	—	2.0 (5)	0.7 (11)	12.7 (2)	50.5	44.7	4.8
Amphenicols	QJ01BA	0.4	—	—	0.01 (1)	1.1 (13)	—	—	0.1 (2)	0.1 (5)	12.3 (3)	0.1	46.1	53.8
Cephalosporins	QJ51DD	4.4	3.5 (2)	1.0 (3)	—	10.2 (12)	—	—	7.4 (2)	1.9 (5)	0.9 (3)	93.5	5.9	0.6
	QJ51DE													
Fluoroquinolones ^b	QJ01MA	1.3	—	3.4 (5)	—	2.3 (10)	—	—	1.9 (5)	0.5 (16)	11.0 (2)	66.6	13.6	19.8
	QJ01MB													
Linco & spec	QJ01FF52	0.2	0.9 (2)	—	1.2 (4)	0.002 (1)	2.0 (1)	—	0.005 (1)	—	—	13.1	82.4	4.5
Lincosamides	QJ01FF02	2.1	—	1.1 (1)	0.5 (5)	6.7 (5)	—	—	—	0.9 (6)	6.2 (2)	—	67.2	32.8
Macrolides	QJ01FA	9.7	2.5 (4)	16.1 (9)	26.5 (12)	6.1 (15)	20.7 (5)	—	21.7 (8)	6.5 (20)	0.6 (2)	30.3	55.5	14.2
Penicillins (β-lactamase-sensitive)	QJ01CE	2.2	0.9 (2)	0.5 (3)	8.9 (12)	0.02 (3)	—	—	12.6 (11)	0.6 (11)	—	84.6	11.5	3.9
Penicillins (extended-spectrum)	QJ01CA	31.2	57.7 (17)	46.9 (15)	20.6 (16)	27.3 (20)	22.9 (10)	50.8 (3)	25.9 (17)	27.0 (16)	4.5 (4)	29.3	66.6	4.0
Pleuromutilins	QJ01XQ	4.5	—	—	7.3 (6)	2.6 (6)	—	—	2.1 (8)	8.9 (13)	0.9 (1)	2.8	87.5	9.6
Polymyxins	QJ01XB	24.7	11.1 (6)	15.0 (4)	0.8 (2)	32.6 (17)	40.3 (11)	26.7 (2)	7.1 (6)	31.5 (18)	50.0 (3)	7.9	89.2	2.9
Sulphonamides	QJ01EQ	0.3	—	—	—	1.5 (4)	—	—	—	—	—	91.1	8.9	—
Tetracyclines	QJ01AA	15.3	12.2 (8)	13.7 (4)	24.5 (14)	7.2 (13)	10.8 (5)	3.1 (2)	18.3 (14)	20.8 (14)	0.9 (2)	2.0	83.3	14.7
Trim & sulfa	QJ01EW	2.4	11.2 (7)	—	4.3 (4)	1.3 (4)	3.4 (2)	19.5 (3)	0.9 (1)	0.7 (5)	—	10.2	66.0	23.9

Linco & spec, lincomycin and spectinomycin; Trim & sulfa, trimethoprim and sulphonomides.

^aThe number of farms within a country using a specific antimicrobial class is shown in brackets.

^bOther quinolones (e.g. flumequine) consisted of 0.03% of the treatments and therefore they were included in the class of fluoroquinolones.




Figure 3. Country-level overview of the relative deviation between TIs calculated with DDDvet, UDDvet, DCDvet or UCDvet based on group treatment data from birth to slaughter (TI200) for a batch of fattening pigs. $TI_{200_{DDvet}}$ and $TI_{200_{UDvet}}$ express the percentage of pigs that received a dose of antimicrobials each day, or equivalently, during which percentage of time a pig was treated with antimicrobials in its entire life. The left plot compares $TI_{200_{DDvet}}$ and $TI_{200_{UDvet}}$ [$(TI_{200_{DDvet}} - TI_{200_{UDvet}}) / TI_{200_{DDvet}}$] and the right plot compares $TI_{200_{DCvet}}$ and $TI_{200_{UCvet}}$ [$(TI_{200_{DCvet}} - TI_{200_{UCvet}}) / TI_{200_{DCvet}}$]. A positive deviation indicates that a higher dose was used than defined by ESVAC. To enhance the comparison between the boxplots some outlying observations are not shown. The full black horizontal line at 0% represents no deviation between TIs. The full black line within each box plot represents the median. The whiskers of each box plot (dashed lines) extend to the most extreme data point within the range of $1.5 \times IQR$ from the box. Outliers, data points outside the range of the whiskers, are shown as open circles.

(4.4%), nervous (8.1%), dermatological (0.8%) and other disorders (4.8%) (Table S8, Appendix D).

Discussion

With a standardized methodology at the farm level this study gained more insight into the AMU patterns of 180 farrow-to-finish farms in nine countries. Given the fact that the 20 sampled farms in each country were not necessarily representative for that country, the results do not necessarily give a good description of country-level AMU, but rather a detailed description of farm-level AMU taking into account variation throughout nine European countries. This study was able to confirm on a larger scale the main findings of Sjölund *et al.*⁸, i.e. large differences in AMU between farms and age groups; mainly oral administration; positive associations between age categories; and considerable use of highest priority critically important antimicrobials for human medicine (HPCIA). Furthermore, by collecting detailed data on antimicrobial group treatments to a batch of fattening pigs, additional results were obtained, such as the moment and duration of treatment and used daily and course doses.

Only moderate correlations in TIs between the group treatment and purchase data were observed. The group treatment data covered only a single batch and might therefore under- or over-estimate AMU due to seasonal influences in the occurrence of

clinical diseases, or due to incidental disease episodes.⁸ They may also be influenced by recall bias with respect to the truly administered products. The latter is why individual treatments were not included, as these are even more prone to recall bias as they are often not well recorded.¹⁷ It is described that the majority of antimicrobial treatments in pig production are group treatments.^{17,21} Purchase data were presumed to provide a more robust overview of the actual use, since they covered multiple batches and included individual treatments. Yet, for each type of data, challenges in data collection were encountered that might influence data quality. For group treatment data, farmers had to make a good estimation of the live weight at treatment. Furthermore, for 20% of the treatments, assumptions had to be made on daily feed and water intake, since doses were expressed per amount of feed or water. For purchase data, the exact moment of AMU is not known and standard live weights were therefore used. When an antimicrobial was purchased for multiple age categories, assumptions had to be made with respect to what proportion of the antimicrobial was used in each age group. This could result in an under- or over-estimation of AMU in a specific age group. Moreover, 30% of the farms were only able to report purchase data for a period up to 6 months and in country H the feed company could not exactly distinguish what proportion of feed was delivered to each specific farm. From this study it became clear that collecting good quality farm-level AMU data is




Figure 4. After \log_{10} transformation of TIs based on DDDvet values, Loess curves (with 95% confidence band) were fitted between different age categories (sucklers, weaners, finishers and sows). TI_{DDvet} expresses the percentage of pigs that received a dose of antimicrobials each day or, equivalently, during which percentage of time a pig was treated with antimicrobials in a certain production phase or its entire life. The relationships between TIs of fattening pigs (sucklers, weaners and finishers) were based on group treatment data, while the relationship between TIs of sows and sucklers was based on purchase data. Each farm is represented with a point, the shape of which denotes the country where the sampled farm was from. The figure shows positive associations between TIs in different age categories.

challenging and shows the need for user-friendly data collection systems.^{15,22}

When comparing used daily and course doses with defined doses, very strong correlations at the farm level were observed. Nonetheless, considerable deviations between the used and defined doses were present. Firstly, UDDvet values were generally higher than DDDvet values. Although DDDvet values were not specifically developed to describe under- or overdosing, the deviations suggest that either the administered doses were too high or the DDDvet values are set too low. The latter are based on detailed dosing data from only nine European Union Member States¹⁹ and might therefore not cover all variability in suggested doses of antimicrobial products. In a study where defined doses were assigned in four countries, considerable deviations from the consensus defined dose were observed for individual products with the same active substance and administration route.²⁰ Secondly, UCDvet values were generally higher than DCDvet values with larger deviations compared with the daily doses. Together with the observation that 10% of the oral group treatments were applied for a period of at least 21 consecutive days, this suggests that the group treatments were often administered for too long a period in this study. Therefore, an important step towards reducing AMU is to comply with the recommended daily dose and treatment duration.²³

Peaks in frequency of treatment were observed during weeks 1, 4 and 9 of the rearing period. As argued by Sjölund et al.⁸, these results strongly suggest that it is common practice in many pig farms to apply treatments to entire batches of pigs at strategic

timepoints when pigs are judged most likely to become infected, i.e. at birth and castration (week 1), at weaning (week 4) and at the start of the finishing period (week 9). Moreover, in accordance with Sjölund et al.,⁸ results indicated that higher AMU at a young age was associated with higher use in older pigs. Possible reasons for these associations are the attitude of the farmer,^{24,25} the overall higher disease pressure in some farms⁸ and the higher disease susceptibility following the effect of AMU at a young age on the bacterial composition.²⁶ Therefore, reducing AMU should already start at a young age (sucklers), even if the total amount used at this stage is still limited, as this AMU seems to affect AMU in the consecutive rearing periods (weaners and finishers).¹⁰ In a recent study the restriction of prophylactic use in weaners by removing antimicrobials from the feed had minimal effects on health and welfare indicators.²⁷

The majority of antimicrobial group treatments were administered orally, although considerable differences at the country level were observed. Only in country C were the majority of the treatments applied parenterally (59%), while only 7% of the treatments were parenteral in country H. It has been argued that the predominant route of administration also influences the average AMU levels.⁸ Oral administration of antimicrobials is likely more easily applied to entire batches compared with parenteral treatments and therefore have a larger impact on AMU, while parenteral administrations allow for more flexible and targeted treatment of diseased animals. Furthermore, in the present study, oral treatments were generally applied for longer compared with parenteral treatments. Finally, studies showed that for some antimicrobials

and bacteria, oral administration of antimicrobials selected for AMR to a greater extent compared with parenteral treatments,^{28,29} while others observed no significant difference between administration routes with respect to resistance.³⁰ More research has been suggested to elucidate the potential effect of the administration route on resistance.³¹ Another factor towards more responsible AMU could therefore be the consideration of parenteral treatment of individuals rather than oral treatment of an entire batch, whenever possible.¹⁰

In this study, colistin (polymyxins) was frequently used in weaned piglets. However, polymyxins have recently been added to the list of HPCIA^s³² and the use of such antimicrobials in food-producing animals is discouraged.¹⁰ Also, macrolides, the fourth most frequently used class in this study, belong to the HPCIA for human medicine list, according to WHO, when considering action on AMU in food-producing animals.^{10,32} The use of other HPCIA^s such as third- and fourth-generation cephalosporins and fluoroquinolones was relatively low. WHO recommends that HPCIA^s should only be used for treatment of refractory infections in food-producing animals, based on veterinary diagnosis and bacterial cultivation and susceptibility results,¹⁰ whereas the European Parliament has asked in a recent resolution for a ban on the veterinary use of antimicrobials that are critical for preventing or treating life-threatening infections in humans.³³

The fact that no group treatments were reported in 11.7% of the farms, for the studied batch, showed that it is possible to rear pigs without systematic use of antimicrobials. Furthermore, there was at least one farm in each participating country that did not report group treatments in either sucklers, weaners or finishers during the observed rearing period. However, the large variation in quantitative and qualitative AMU between farms shows that there is still room for improvement towards responsible AMU. For instance, as mentioned above, one could reduce AMU at strategic timepoints in combination with increased biosecurity^{34,35} and restrict the use of HPCIA^s. Further research is needed to explore what the main drivers for AMU are.

Acknowledgements

We thank all researchers who visited the farms to collect the data and samples (other than those listed below as EFFORT consortium members): K. L. Nielsen, B. E. Hoelstad, P. Ryt-Hansen, C. Nielsen, M. Brunon, B. Gawlik, E. Guyrova, N. Cholakov, T. Ivanova, M. GherPELLI and V. Carfora. The farmers and their herd veterinarians are acknowledged for their participation in the study.

Members of the EFFORT consortium

J. Wagenaar (UUVM), H. Graveland (UUVM), D. Heederik (UUVM), H. Schmitt (UUVM), R. Luiken (UUVM), L. Van Gompel (UUVM), D. Mevius (UUVM), A. van Essen (WBVR), B. Gonzalez-Zorn (UCM), G. Moyano (UCM), P. Sanders (ANSES), C. Chauvin (ANSES), J. David (ANSES), A. Battisti (IZSLT), A. Caprioli (IZSLT), J. Dewulf (UGent), T. Blaha (TIHO), K. Wadepohl (TIHO), M. Brandt (TIHO), F. Aarestrup (DTU), T. Hald (DTU), S. Duarte (DTU), D. Wasyl (NVRI), D. Krasucka (NVRI), B. Biernacki (NVRI), J. Szumilo (NVRI), H. Daskalov (NDRVI), H. Saatkamp (BEC) and K. Stärk (SAFOSO).

Funding

This work was supported by the by the European Commission under the Food, Agriculture and Fisheries, and Biotechnologies theme of the 7th

Framework Programme for Research and Technological Development (Grant Agreement FP7-KBBE-2013-7-613754-EFFORT).

Transparency declarations

None to declare.

Supplementary data

Appendixes A to D, Figure S1 and Tables S1 to S8 are available as Supplementary data at JAC Online.

References

- World Health Organization (WHO). *Global Action Plan on Antimicrobial Resistance*. 2015. http://www.wpro.who.int/entity/drug_resistance/resources/global_action_plan_eng.pdf.
- World Organisation for Animal Health (OIE). *Final Report—83rd General Session*. 2015. http://www.oie.int/fileadmin/Home/eng/About_us/docs/pdf/Session/A_FR_2015_public.pdf.
- Food and Agriculture Organization (FAO). *Report of the Conference of FAO—Thirty-Ninth Session*. 2015. <http://www.fao.org/3/a-mo153e.pdf>.
- da Costa PM, Loureiro L, Matos AJF. Transfer of multidrug-resistant bacteria between intermingled ecological niches: the interface between humans, animals and the environment. *Int J Environ Res Public Health* 2013; **10**: 278–94.
- Tang KL, Caffrey NP, Nóbrega DB *et al*. Restricting the use of antibiotics in food-producing animals and its associations with antibiotic resistance in food-producing animals and human beings: a systematic review and meta-analysis. *Lancet Planet Health* 2017; **1**: e316–27.
- Chantziaras I, Boyen F, Callens B *et al*. Correlation between veterinary antimicrobial use and antimicrobial resistance in food-producing animals: a report on seven countries. *J Antimicrob Chemother* 2014; **69**: 827–34.
- Joint Interagency Antimicrobial Consumption and Resistance Analysis (JIACRA). ECDC/EFSA/EMA second joint report on the integrated analysis of the consumption of antimicrobial agents and occurrence of antimicrobial resistance in bacteria from humans and food-producing animals. *EFSA J* 2017; **15**: e04872.
- Sjölund M, Postma M, Colineau L *et al*. Quantitative and qualitative antimicrobial usage patterns in farrow-to-finish pig herds in Belgium, France, Germany and Sweden. *Prev Vet Med* 2016; **130**: 41–50.
- World Organisation for Animal Health (OIE). *Terrestrial Animal Health Code, Chapter 6.8: Monitoring of the Quantities and Usage Patterns of Antimicrobial Agents Used in Food-Producing Animals*. 2016. http://www.oie.int/fileadmin/Home/eng/Health_standards/tahc/current/chapitre_antibio_monitoring.pdf.
- World Health Organization (WHO). *WHO Guidelines on Use of Medically Important Antimicrobials in Food-Producing Animals*. 2017. <http://apps.who.int/iris/bitstream/handle/10665/258970/9789241550130-eng.pdf?sequence=1>.
- European Medicines Agency (EMA). *Sales of Veterinary Antimicrobial Agents in 30 European Countries in 2015*. 2017. http://www.ema.europa.eu/docs/en_GB/document_library/Report/2017/10/WC500236750.pdf.
- The Danish Integrated Antimicrobial Resistance Monitoring and Research Programme (DANMAP). *DANMAP 2013—Use of Antimicrobial Agents and Occurrence of Antimicrobial Resistance in Bacteria from Food Animals, Food and Humans in Denmark*. 2014. <https://www.danmap.org/~media/Projekt%20sites/Danmap/DANMAP%20reports/DANMAP%202013/DANMAP%202013.ashx>.
- Monitoring of Antimicrobial Resistance and Antibiotic Usage in Animals in the Netherlands (MARAN). *Monitoring of Antimicrobial Resistance and*

- Antibiotic Usage in Animals in the Netherlands in 2016*. 2017. https://www.wur.nl/upload_mm/b/0/1/74ce6009-b112-428d-aeb7-99b95063aab6_Maran%20report%202017.pdf.
- 14** Filippitzi ME, Callens B, Pardon B et al. Antimicrobial use in pigs, broilers and veal calves in Belgium. *Vlaams Diergeneesk Tijdschr* 2014; **83**: 215–24.
- 15** Collineau L, Belloc C, Stärk KDC et al. Guidance on the selection of appropriate indicators for quantification of antimicrobial usage in humans and animals. *Zoonoses Public Health* 2017; **64**: 165–84.
- 16** Munk P, Knudsen BE, Lukjanenko O et al. Abundance and diversity of the faecal resistome in slaughter pigs and broilers in nine European countries. *Nat Microbiol* 2018; **3**: 898–908.
- 17** Callens B, Persoons D, Maes D et al. Prophylactic and metaphylactic antimicrobial use in Belgian fattening pig herds. *Prev Vet Med* 2012; **106**: 53–62.
- 18** Timmerman T, Dewulf J, Catry B et al. Quantification and evaluation of antimicrobial drug use in group treatments for fattening pigs in Belgium. *Prev Vet Med* 2006; **74**: 251–63.
- 19** European Medicines Agency (EMA). *Defined Daily Doses for Animals (DDDvet) and Defined Course Doses for Animals (DCDvet)*. 2016. http://www.ema.europa.eu/docs/en_GB/document_library/Other/2016/04/WC500205410.pdf.
- 20** Postma M, Sjölund M, Collineau L et al. Assigning defined daily doses animal: a European multi-country experience for antimicrobial products authorized for usage in pigs. *J Antimicrob Chemother* 2015; **70**: 294–302.
- 21** Regula G, Torriani K, Gassner B et al. Prescription patterns of antimicrobials in veterinary practices in Switzerland. *J Antimicrob Chemother* 2009; **63**: 805–11.
- 22** Network on Quantification of Veterinary Antimicrobial Usage at Herd Level and Analysis, Communication and Benchmarking to Improve Responsible Usage (AACTING). *Guidelines for Collection, Analysis and Reporting of Farm-Level AMU, in the Scope of Antimicrobial Stewardship*. 2018. http://www.aacting.org/swfiles/files/AACTING_Guidelines_V1.1_2018.03.23_39.pdf.
- 23** European Commission Notice. Guidelines for the prudent use of antimicrobials in veterinary medicine. *Official Journal of the European Union* 2015; 2015/C 299/04. https://ec.europa.eu/health/sites/health/files/antimicrobial_resistance/docs/2015_prudent_use_guidelines_en.pdf.
- 24** Visschers VHM, Backhans A, Collineau L et al. Perceptions of antimicrobial usage, antimicrobial resistance and policy measures to reduce antimicrobial usage in convenient samples of Belgian, French, German, Swedish and Swiss pig farmers. *Prev Vet Med* 2015; **119**: 10–20.
- 25** Kramer T, Jansen LE, Lipman LJA et al. Farmers' knowledge and expectations of antimicrobial use and resistance are strongly related to usage in Dutch livestock sectors. *Prev Vet Med* 2017; **147**: 142–8.
- 26** Callens B, Faes C, Maes D et al. Presence of antimicrobial resistance and antimicrobial use in sows are risk factors for antimicrobial resistance in their offspring. *Microb Drug Resist* 2015; **21**: 50–8.
- 27** Diana A, Manzanilla EG, Calderón Díaz JA et al. Do weaner pigs need in-feed antibiotics to ensure good health and welfare? *PLoS One* 2017; **12**: e0185622.
- 28** Chantziaras I, Smet A, Haesebrouck F et al. Studying the effect of administration route and treatment dose on the selection of enrofloxacin resistance in commensal *Escherichia coli* in broilers. *J Antimicrob Chemother* 2017; **72**: 1991–2001.
- 29** Wiuff C, Lykkesfeldt J, Svendsen O et al. The effects of oral and intramuscular administration and dose escalation of enrofloxacin on the selection of quinolone resistance among *Salmonella* and coliforms in pigs. *Res Vet Sci* 2003; **75**: 185–93.
- 30** Græsbøll K, Damborg P, Møllerup A et al. Effect of tetracycline dose and treatment mode on selection of resistant coliform bacteria in nursery pigs. *Appl Environ Microbiol* 2017; **83**: e00538–17.
- 31** Burow E, Simoneit C, Tenhagen BA et al. Oral antimicrobials increase antimicrobial resistance in porcine *E. coli*—a systematic review. *Prev Vet Med* 2014; **113**: 364–75.
- 32** World Health Organization (WHO). *Critically Important Antimicrobials for Human Medicine—5th Revision* 2016. 2017. <http://apps.who.int/iris/bitstream/handle/10665/255027/9789241512220-eng.pdf?sequence=1>.
- 33** European Parliament. *Resolution of 13 September 2018 on a European One Health Action Plan against Antimicrobial Resistance*. 2018. P8_TA-PROV(2018)0354. <http://www.europarl.europa.eu/sides/getDoc.do?pubRef=-//EP//NONSGML+TA+P8-TA-2018-0354+0+DOC+PDF+V0//EN>.
- 34** Collineau L, Rojo-Gimeno C, Léger A et al. Herd-specific interventions to reduce antimicrobial usage in pig production without jeopardising technical and economic performance. *Prev Vet Med* 2017; **144**: 167–78.
- 35** Postma M, Vanderhaeghen W, Sarrazin S et al. Reducing antimicrobial usage in pig production without jeopardizing production parameters. *Zoonoses Public Health* 2017; **64**: 63–74.