


Adherence to the Mediterranean diet and lymphoma risk in the European Prospective Investigation into Cancer and Nutrition

Marta Solans^{1,2,3}, Yolanda Benavente^{1,4}, Marc Saez^{1,2}, Antonio Agudo⁵, Sabine Naudin⁶, Fatemeh Saberi Hosnijeh^{7,8}, Hwayoung Noh⁶, Heinz Freisling⁶, Pietro Ferrari⁶, Caroline Besson^{9,10,11}, Yahya Mahamat-Saleh^{10,11}, Marie-Christine Boutron-Ruault^{10,11}, Tilman Kühn¹², Rudolf Kaaks¹², Heiner Boeing¹³, Cristina Lasheras¹⁴, Miguel Rodríguez-Barranco^{1,15}, Pilar Amiano^{1,16}, Jose Maria Huerta^{1,17}, Aurelio Barricarte^{1,18,19}, Julie A. Schmidt²⁰, Paolo Vineis²¹, Elio Riboli²², Antonia Trichopoulos^{23,24}, Christina Bamia^{23,24†}, Eleni Peppas^{23,24}, Giovanna Masala²⁵, Claudia Agnoli²⁶, Rosario Tumino²⁷, Carlotta Sacerdote²⁸, Salvatore Panico²⁹, Guri Skeie³⁰, Elisabete Weiderpass^{30,31,32,33}, Mats Jerkeman³⁴, Ulrika Ericson³⁴, Florentin Späth³⁵, Lena Maria Nilsson³⁶, Christina C Dahm³⁷, Kim Overvad³⁷, Anne Katrine Bolvig³⁸, Anne Tjønneland^{38,39}, Silvia de Sanjose^{1,4,40}, Genevieve Buckland⁴¹, Roel Vermeulen⁷, Alexandra Nieters⁴² and Delphine Casabonne ^{1,4}

¹Centro de Investigación Biomédica en Red: Epidemiología y Salud Pública (CIBERESP), Madrid, Spain

²Research Group on Statistics, Econometrics and Health (GRECS), University of Girona, Girona, Spain

³Epidemiology Unit and Girona Cancer Registry, Catalan Institute of Oncology, Girona, Spain

⁴Unit of Molecular and Genetic Epidemiology in Infections and Cancer, Catalan Institute of Oncology (ICO-IDIBELL), Barcelona, Spain

⁵Unit of Nutrition and Cancer, Cancer Epidemiology Research Program, Catalan Institute of Oncology (ICO-IDIBELL), Barcelona, Spain

⁶Nutritional Methodology and Biostatistics Group, International Agency for Research on Cancer (IARC-WHO), Lyon, France

⁷Institute for Risk Assessment Sciences, Division of Environmental Epidemiology, Utrecht University, Utrecht, The Netherlands

Key words: lymphoma, Mediterranean diet, Europe, prospective studies, risk

Abbreviations: arMED: adapted relative Mediterranean diet; BMI: body mass index; CI: confidence interval; CLL/SLL: chronic lymphocytic leukaemia/small lymphocytic lymphoma; DLBCL: diffuse large B-cell lymphoma; EPIC: European Prospective Investigation into Cancer and Nutrition; FL: follicular lymphoma; HL: Hodgkin lymphoma; HR: hazard ratio; MD: Mediterranean diet; MM/PCN: multiple myeloma/plasma cell neoplasm; NHL: Non-Hodgkin lymphoma; NOS: not otherwise specified; WCRF/AICR: World Cancer Research Fund/American Institute for Cancer Research. Additional Supporting Information may be found in the online version of this article.

[†]Deceased.

Grant sponsor: European Commission (DG-SANCO); **Grant sponsor:** International Agency for Research on Cancer; **Grant sponsor:** Danish Cancer Society (Denmark); **Grant sponsor:** Ligue Contre le Cancer; **Grant sponsor:** Institut Gustave Roussy; **Grant sponsor:** Mutuelle Generale de l'Education Nationale; **Grant sponsor:** Institut National de la Sante et de la Recherche Medicale (INSERM) (France); **Grant sponsor:** German Cancer Aid; **Grant sponsor:** German Cancer Research Center (DKFZ); **Grant sponsor:** Federal Ministry of Education and Research (BMBF); **Grant sponsor:** Deutsche Krebshilfe; **Grant sponsor:** Deutsches Krebsforschungszentrum and Federal Ministry of Education and Research (Germany); **Grant sponsor:** the Hellenic Health Foundation (Greece); **Grant sponsor:** Associazione Italiana per la Ricerca sul Cancro-AIRC-Italy and National Research Council (Italy); **Grant sponsor:** Dutch Ministry of Public Health, Welfare and Sports (VWS); **Grant sponsor:** Netherlands Cancer Registry (NKR); **Grant sponsor:** LK Research Funds; **Grant sponsor:** Dutch Prevention Funds; **Grant sponsor:** Dutch ZON (Zorg Onderzoek Nederland); **Grant sponsor:** World Cancer Research Fund (WCRF); **Grant sponsor:** Statistics Netherlands (The Netherlands); **Grant number:** ERC2009-AdG 232997; **Grant sponsors:** Nordforsk, Nordic Centre of Excellence programme on Food, Nutrition and Health (Norway); **Grant sponsor:** German Federal Ministry of Education and Research; **Grant number:** BMBF 01EO1303; **Grant sponsor:** Spanish Ministry of Economy and Competitiveness - Carlos III Institute of Health cofunded by FEDER funds/ European Regional Development Fund (ERDF) - a way to build Europe; **Grant numbers:** PI13/00061 (to Granada), PI13/01162 (to EPIC-Murcia, Regional Governments of Andalusia, Asturias, Basque Country, Murcia and Navarra), PI17/01280 and PI14/01219 (to Barcelona); **Grant sponsor:** Centro de Investigación Biomédica en Red: Epidemiología y Salud Pública (CIBERESP, Spain); **Grant sponsor:** Agència de Gestió d'Ajuts Universitaris i de Recerca (AGAUR, CERCA Programme / Generalitat de Catalunya for institutional support); **Grant number:** (2017SGR1085); **Grant sponsor:** ISCIII RETIC; **Grant number:** RD06/0020 (Spain); **Grant sponsor:** Swedish Cancer Society; **Grant sponsor:** Swedish Research Council and County Councils of Skåne and Vasterbotten (Sweden); **Grant sponsor:** Cancer Research UK; **Grant number:** 14136 (to EPIC-Norfolk), C570/A16491 and C8221/A19170 (to EPIC-Oxford); **Grant sponsor:** Medical Research Council; **Grant number:** 1000143 (to EPIC-Norfolk), MR/M012190/1 (to EPIC-Oxford, UK)

DOI: 10.1002/ijc.32091

History: Received 19 Sep 2018; Accepted 21 Nov 2018; Online 26 Dec 2018

Correspondence to: Delphine Casabonne, Unit of Infections and Cancer, Cancer Epidemiology Research Programme, IDIBELL, Catalan Institute of Oncology, Av. Gran Via de l'Hospitalet 199-203 08908 L'Hospitalet de Llobregat, Barcelona, Spain, E-mail: dcasabonne@iconcologia.net

- ⁸Immunology Department, Erasmus University Medical Center, Rotterdam, The Netherlands
- ⁹Department of Hematology and Oncology, Hospital of Versailles, Le Chesnay, France
- ¹⁰CESP, Fac. de médecine - Univ. Paris-Sud, Fac. de médecine - UVSQ, INSERM, Université Paris-Saclay, Villejuif, France
- ¹¹Gustave Roussy, Villejuif, France
- ¹²Division of Cancer Epidemiology, German Cancer Research Center (DKFZ), Heidelberg, Germany
- ¹³Department of Epidemiology, German Institute of Human Nutrition (DIFE) Postdam-Rehbrücke, Nuthetal, Germany
- ¹⁴Department of Functional Biology, Oviedo University, Oviedo, Spain
- ¹⁵Escuela Andaluza de Salud Pública, Instituto de Investigación Biosanitaria IBS GRANADA, Granada, Spain
- ¹⁶Public Health Division of Gipuzkoa, Regional Government of the Basque Country, Donostia, Spain
- ¹⁷Department of Epidemiology, Murcia Regional Health Council, IMIB-Arrixaca, Murcia, Spain
- ¹⁸Navarra Public Health Institute, Pamplona, Spain
- ¹⁹IdiSNA Navarra Institute for Health Research, Pamplona, Spain
- ²⁰Cancer Epidemiology Unit, Nuffield Department of Population Health, University of Oxford, Oxford, United Kingdom
- ²¹Department of Epidemiology and Biostatistics, MRC-HPA Centre for Environment and Health, School of Public Health, Imperial College London, London, United Kingdom
- ²²School of Public Health, Imperial College London, London, United Kingdom
- ²³Hellenic Health Foundation, Athens, Greece
- ²⁴WHO Collaborating Center for Nutrition and Health, Unit of Nutritional Epidemiology and Nutrition in Public Health, Department of Hygiene, Epidemiology and Medical Statistics, School of Medicine, National and Kapodistrian University of Athens, Athens, Greece
- ²⁵Cancer Risk Factors and Life-Style Epidemiology Unit, Institute for Cancer Research, Prevention and Clinical Network – ISPRO, Florence, Italy
- ²⁶Epidemiology and Prevention Unit, Fondazione IRCCS Istituto Nazionale dei Tumori, Milan, Italy
- ²⁷Cancer Registry and Histopathology Department, "Civic - M. P. Arezzo" Hospital, ASP Ragusa, Ragusa, Italy
- ²⁸Unit of Cancer Epidemiology, Città della Salute e della Scienza University-Hospital and Center for Cancer Prevention (CPO), Turin, Italy
- ²⁹Dipartimento di Medicina Clinica e Chirurgia, Federico II University, Naples, Italy
- ³⁰Department of Community Medicine, University of Tromsø, The Arctic University of Norway, Tromsø, Norway
- ³¹Department of Research, Cancer Registry of Norway, Institute of Population-Based Cancer Research, Oslo, Norway
- ³²Department of Medical Epidemiology and Biostatistics, Karolinska Institutet, Stockholm, Sweden
- ³³Genetic Epidemiology Group, Folkhälsan Research Center, Faculty of Medicine, Helsinki University, Helsinki, Finland
- ³⁴Department of Clinical Sciences in Malmö, Lund University, Malmö, Sweden
- ³⁵Department of Radiation Sciences, Oncology, Umeå University, Umeå, Sweden
- ³⁶Public Health and Clinical Medicine, Nutritional Research and Arctic Research Centre, Umeå University, Umeå, Sweden
- ³⁷Section for Epidemiology, Department of Public Health, Aarhus University, Aarhus, Denmark
- ³⁸Danish Cancer Society Research Center, Copenhagen, Denmark
- ³⁹Department of Public Health, University of Copenhagen, Copenhagen, Denmark
- ⁴⁰Reproductive Health, PATH, Seattle, WA
- ⁴¹NIHR Bristol Biomedical Research Centre Nutrition Theme, Bristol, United Kingdom
- ⁴²Center for Chronic Immunodeficiency (CCI), Medical Center - University of Freiburg, Faculty of Medicine, University of Freiburg, Freiburg, Germany

There is a growing evidence of the protective role of the Mediterranean diet (MD) on cancer. However, no prospective study has yet investigated its influence on lymphoma. We evaluated the association between adherence to the MD and risk of lymphoma and its subtypes in the European Prospective Investigation into Cancer and Nutrition (EPIC) study. The analysis included 476,160 participants, recruited from 10 European countries between 1991 and 2001. Adherence to the MD was estimated through the adapted relative MD (arMED) score excluding alcohol. Cox proportional hazards regression models were used while adjusting for potential confounders. During an average follow-up of 13.9 years, 3,136 lymphomas (135 Hodgkin lymphoma [HL], 2,606 non-HL and 395 lymphoma not otherwise specified) were identified. Overall, a 1-unit increase in the arMED score was associated with a 2% lower risk of lymphoma (95% CI: 0.97; 1.00, p-trend = 0.03) while a statistically nonsignificant inverse association between a high *versus* low arMED score and risk of lymphoma was observed (hazard ratio [HR]: 0.91 [95% CI 0.80; 1.03], p-trend = 0.12). Analyses by lymphoma subtype did not reveal any statistically significant associations. Albeit with small numbers of cases ($N = 135$), a suggestive inverse association was found for HL (HR 1-unit increase = 0.93 [95% CI: 0.86; 1.01], p-trend = 0.07). However, the study may have lacked statistical power to detect small effect sizes for lymphoma subtype. Our findings suggest that an increasing arMED score was inversely related to the risk of overall lymphoma in EPIC but not by subtypes. Further large prospective studies are warranted to confirm these findings.

What's new?

Known risk factors explain only a small proportion of lymphoma cases. Several studies have pointed out the potential role of dietary factors on lymphoma risk, but evidence is still inconclusive. Here, using data from the European Prospective Investigation into Cancer and Nutrition study, the authors found for the first time that adherence to a Mediterranean diet was modestly associated with a reduced risk of overall lymphoma. Further studies are needed to confirm these findings.

Introduction

Lymphomas are a heterogeneous group of malignancies particularly prevalent in Western countries. Although some lymphoid neoplasms have been consistently linked to certain infections and severe immunosuppression, their etiology remains elusive, and evidence from epidemiologic studies increasingly points to etiologic heterogeneity among subtypes.¹

An increase in the incidence of lymphoma has been observed in many regions during the last decades,² and a change in lifestyle might be one possible explanation for this pattern. However, there is limited evidence regarding extrinsic-risk factors, particularly diet, and lymphoma risk.^{3,4} In the recently released Third Expert Report on “Diet, Nutrition, Physical Activity, and Cancer” by the World Cancer Research Fund/American Institute for Cancer Research (WCRF/AICR),⁵ no conclusive associations for specific dietary factors and hematological malignancies were reported. Studies from the European Prospective Investigation into Cancer and Nutrition (EPIC) study have neither shown consistent associations between consumption of meat and dairy⁶ nor vegetables⁷ and overall lymphoma risk, although some statistically significant associations were seen for several lymphoma subtypes.

There is evidence that the Mediterranean diet (MD) has a protective role on risk of overall⁸ and specific types of cancer,⁹ such as breast,¹⁰ colorectal¹¹ or gastric cancers.¹² However, epidemiological research into the effect of a MD pattern on lymphoma remains limited. To our knowledge, no study has yet evaluated the influence of validated *a priori* MD scores on lymphoma risk, while studies on *a posteriori* healthy-like dietary patterns have yielded inconsistent results for overall lymphoma as well as its subtypes.^{13–15}

The aim of our study is to investigate the association between adherence to the MD and lymphoma risk within the EPIC population. The EPIC study provides the opportunity to examine this relation in a prospective design and within a European population with a wide spectrum of dietary habits.

Material and Methods**Study population**

EPIC is a large prospective cohort study designed to investigate the relationship among diet, lifestyle, environmental factors and cancer. The rationale, full methods and study design

have been described previously.^{16,17} In brief, 521,324 subjects, mostly aged 30 to 70 years, were recruited between 1992 and 2000 in 23 centers from 10 European countries (Denmark, France, Germany, Greece, the Netherlands, Italy, Norway, United Kingdom, Spain and Sweden). The ethical review boards from the International Agency for Research on Cancer and all local participating centers approved the study, and all participants gave their informed consent.

Of the 521,324 EPIC cohort participants, we excluded prevalent cancer cases ($n = 25,184$), subjects with missing follow-up information ($n = 4,148$), with incomplete/no dietary information ($n = 6,259$), or those in the highest and lowest 1% of the distribution for the ratio of energy intake to estimate energy requirement ($n = 9,573$). Therefore, the current analysis was based on 476,160 subjects among whom 3,136 incident lymphoma cases occurred.

Data collection

Validated country-specific questionnaires were used to record the usual diet during the previous year,^{17,18} namely through quantitative or semiquantitative food frequency questionnaires (FFQs; administered through a personal interview or self-administered), although few countries used semiquantitative FFQs combined with a food record. Lifestyle questionnaires were used to obtain information on sociodemographic characteristics, physical activity, reproductive history, use of oral contraceptives and hormone replacement therapy, medical history and alcohol and tobacco consumption. Anthropometric measures were also ascertained at recruitment.

Exposure assessment—arMED

The level of adherence to the MD was assessed using the adapted relative MD (arMED) score,¹⁰ which excludes alcoholic beverages as they have been inversely associated with several lymphoma subtypes.¹⁹ The scoring system, adapted from the original index designed by Trichopoulou *et al.*,²⁰ has been detailed previously.¹² In brief, the arMED is a 16-point linear score that incorporates eight key dietary components: six components presumed to reflect the MD (fruit [including nuts and seeds], vegetables, legumes, fish [including seafood], olive oil and cereals) and two components consumed in low quantity in the MD (dairy products and meat). Intake of each component was calculated as a function of energy density (g/day/1,000 kcal) and divided

into tertiles (estimated using the overall study population). A score of 0 to 2 was assigned for the first, second, and third tertile of intakes for the components presumed to fit the MD, while the scoring was inverted for the components presumed to not fit the MD (giving a lower score for higher intakes). The scoring for olive oil was adapted owing to the low consumption of non-Mediterranean countries, by assigning 0 to nonconsumers, 1 for subjects below the median and 2 for subjects equal or above this median (the median was calculated using the overall study population and considering only consumers). The points were summed to define the arMED score, that ranged from 0 to 16 (from the lowest to the highest adherence), and were in turn divided into three categories: low (0–5), medium (6–9) and high (10–16) as described previously.¹⁰

Follow-up and outcome assessment

Incident lymphoma cancer cases were identified by population cancer registries for Denmark, Italy, the Netherlands, Norway, Spain, Sweden and the United Kingdom. A combination of methods was used in France, Germany and Greece, including cancer and pathology registries, health insurance records, and active follow-up contacting participants or their next-of-kin. Mortality data were also obtained from regional or national mortality registries. The follow-up period was defined from the age at recruitment to the age at first cancer diagnosis, death or last complete follow-up, depending on which occurred first. Censoring dates for the last complete follow-up ranged from June 2008 to December 2013, depending on the EPIC center.

Initially, the diagnosis of lymphoma cases was based on the second revision of the International Classification of Diseases for Oncology (ICD-O-2). Later, all cases were reclassified into the ICD-O-3, using a conversion program available on the web site of the Surveillance Epidemiology and End Results program (<http://seer.cancer.gov/tools/conversion/ICD02-3manual.pdf>) and involving a pathology expert and experts from the EPIC centers. Because not all ICD-O-2 diagnostics can be translated unequivocally into the current classification, we left the respective lymphomas unclassified (not otherwise specified “NOS”) when further detailed specification failed. Finally, the Inter-Lymph Pathology Working Group classification, which is based in the 2008 WHO classification, was used to categorize lymphoma histologic subtypes.¹

In the current analysis, the following groups were considered: Hodgkin lymphoma (HL) and non-HL (NHL); within NHL, mature B-cell lymphoma and mature T/NK-cell lymphoma; and among mature B-cell lymphoma, the following entities: diffuse large B-cell lymphoma (DLBCL), follicular lymphoma (FL), chronic lymphocytic leukemia (including small lymphocytic lymphoma), multiple myeloma/plasma cell neoplasm and other B-cell lymphoma (i.e., those cases in which the B-cell lymphoma subtype is unknown or does not

fall within the above mentioned subtypes). Other entities were not considered due to small numbers (Table 1).

Statistical analysis

Cox proportional hazard models were used to estimate the hazard ratio (HR) and 95% confidence interval (CI) of the association between the arMED score and lymphoma risk. The arMED score was analyzed both as a continuous variable (per 1-unit increase) and as a categorical variable (low, medium and high level of adherence). In addition, Cox models were fitted with the arMED ordinal variable as continuous to test for linear trend for comparison with published literature on solid cancers. Two models with two levels of adjustment were used: a basic model, stratified by center, sex and age at recruitment (in 1-year categories), and a multivariable model, further adjusted for body mass index (BMI; <25, 25–30 and ≥ 30 kg/m²), total energy intake (continuous, kcal/day), educational level (no formal education, primary school, secondary school, technical or professional training, university and unknown [3.6%]), height (continuous, cm), physical activity level based on the Cambridge physical activity index (inactive, moderately inactive, moderately active, active and unknown [1.9%]), smoking status (never, former, current and unknown [2.0%]) and alcohol intake at recruitment (continuous, g/day). We tested for interaction by age, sex, alcohol intake and smoking by including a cross-product term along with the arMED score (continuous) in the multivariable Cox model. The statistical significance of the cross-product term was evaluated using likelihood ratio test.

Sensitivity analyses were performed by repeating main Cox analyses (i) including alcohol in the score computation, (ii) censoring participants and excluding cases with less than 2 years of follow-up ($n = 259$ cases), (iii) excluding participants without complete data ($n = 226$ cases) and (iv) restricting HL analysis to classical HL cases. Schoenfeld residuals were used to ensure that the proportional hazard assumption was met in all models. Two-sided p values were reported with statistical significance set at $p < 0.05$. All analyses were performed by using STATA statistical software, version 14 (Stata Corporation, College Station, TX).

Results

During an average follow-up of 13.9 years, 3,136 lymphoma cases (2,606 NHL, 135 HL and 395 lymphoma NOS) were diagnosed. A detailed distribution of cases by lymphoma subtype and country is displayed in Table 1. As expected, the highest levels of the arMED score were found in Mediterranean regions (i.e. Greece, Spain and Italy), while the lowest scores were found in Sweden, the Netherlands and Denmark. Baseline characteristics of the study participants according to category of the arMED score are shown in Table 2. In general, participants with a higher arMED score were more likely to be women, slightly younger, never smokers and physically inactive and to

Table 1. Distribution of lymphoma cases in the EPIC study

	Total cohort	Person years	Lymphoma subtypes				Overall	NHL subtypes ¹			Mature B-cell subtypes					
			NHL	HL	NOS	Mature B-cell		Mature T / NK-cell	DLBCL	FL	CLL/SLL	MM/PCN	Other B-cell	arMED mean (SD)		
Denmark	55,014	815,096.8	538	29	64	506	23	121	78	118	123	66	5.9 (2.4)			
France	67,403	869,362.5	216	11	1	205	8	40	44	44	45	32	8.5 (2.4)			
Germany	48,557	504,479.0	190	13	28	170	12	30	20	39	55	26	6.4 (2.1)			
Greece	26,048	281,283.6	44	3	15	38	2	3	3	13	15	4	11.8 (1.7)			
Italy	44,545	630,951.3	241	15	42	218	11	38	33	44	73	30	10.1 (2.1)			
Norway	33,975	452,171.1	147	5	11	129	14	26	31	26	24	22	7.7 (2.0)			
Spain	39,989	637,947.4	211	14	16	194	10	35	27	51	51	30	10.4 (2.2)			
Sweden	48,674	801,130.2	381	13	123	344	20	57	48	74	132	33	4.6 (2.1)			
The Netherlands	36,539	524,670.7	186	7	8	172	10	43	26	41	43	19	5.6 (2.1)			
United Kingdom	75,416	1,122,765	452	25	87	426	20	95	71	87	115	58	8.6 (2.5)			
Total	476,160	6,639,857.5	2,606	135	395	2,402	130	488	381	537	676	320	7.8 (3.0)			

Abbreviations: arMED, adapted relative Mediterranean diet; CLL/SLL, chronic lymphocytic leukemia/small lymphocytic lymphoma; DLBCL, diffuse large B-cell lymphoma; FL, follicular lymphoma; HL, Hodgkin lymphoma; MM/PCN, multiple myeloma/plasma cell neoplasm; NHL, non-Hodgkin lymphoma; NOS, not otherwise specified; Other B-cell (those cases for which the mature B-cell NHL subtype is unknown or does not fall within the more common subtypes); SD, standard deviation.

¹There were 37 precursor NHL and 37 individuals with NHL without B- or T-cell information.

Table 2. Baseline characteristics of participants in the EPIC study according to adherence to the arMED score

	Total cohort	arMED score		
		Low (0–5) (mean 3.9)	Medium (6–9) (mean 7.5)	High (10–16) (mean 11.4)
Total cohort, <i>n</i>	476,160	116,128	214,649	145,383
Sex (%)				
Men	29.9	45.1	25.0	24.9
Women	70.1	54.9	75.0	75.1
Age at recruitment (mean [SD], years)	51.2 (9.9)	52.0 (10.0)	51.4 (9.5)	50.3 (10.4)
Energy intake (mean [SD], kcal/day)	2,075.1 (619.2)	2,178.5 (648.0)	2,039.8 (604.1)	2,044.6 (608.6)
Alcohol intake (median [25th–75th percentile], g/day)	5.3 (0.9; 14.9)	6.2 (1.4; 17.0)	5.6 (1.1; 15.2)	4.2 (0.4; 13.0)
BMI (mean [SD], kg/m ²)	25.4 (4.3)	25.6 (4.1)	25.1 (4.1)	25.8 (4.5)
Height (mean [SD], cm)	166.0 (8.9)	169.6 (9.2)	165.9 (8.5)	163.2 (8.4)
Smoking status (%)				
Never	49.0	41.5	49.2	54.5
Former	26.6	27.7	27.7	24.3
Current	22.4	29.6	20.8	19.0
Unknown	2.0	1.24	2.3	2.3
Physical activity (%)				
Inactive	21.0	17.6	18.1	27.8
Moderately inactive	32.9	31.7	33.6	32.9
Moderately active	26.4	25.2	28.4	24.3
Active	17.9	22.3	17.8	14.5
Unknown	1.85	3.2	2.0	0.5
Educational level (%)				
None	4.4	0.5	2.4	10.5
Primary school	25.6	30.2	22.4	26.6
Technical/professional school	22.2	30.5	23.8	13.3
Secondary school	20.4	17.1	21.9	20.9
University	23.8	20.0	25.2	24.8
Unknown	3.6	1.7	4.3	3.9

Abbreviations: arMED, adapted relative Mediterranean dietary score; BMI, body mass index; SD, standard deviation.

have a higher educational level and lower alcohol intake compared to those with a low arMED score.

Table 3 shows basic and multivariable HR estimates for category of arMED score associated with lymphoma risk, overall and by subgroups. Overall, a 1-unit increase in the arMED score was associated with a 2% lower risk of lymphoma (95% CI: 0.97; 1.00, *p*-trend = 0.03) while a statistically nonsignificant inverse association between a high *versus* low arMED score and risk of lymphoma was observed (HR: 0.91 [95% CI 0.80; 1.03], *p*-trend = 0.12). Exclusion of lymphoma NOS cases (*N*= 395) did not modify the observed association with lymphoma risk (HR for a 1-unit increase in the arMED score: 0.98 [95% CI: 0.96 to 1.00], *p*-value = 0.04, data not shown). However, albeit with smaller numbers of cases, the lowest HR was observed for HL (HR_{high vs. low} = 0.64 [95% CI: 0.34; 1.19], *p*-trend = 0.16; HR_{1-unit increase} = 0.93 [95% CI: 0.86; 1.01], *p*-trend = 0.07). After restriction to classical HL

(*n* = 127), although the results were still not statistically significant, the inverse association seemed to be strengthened (HR_{high vs. low} = 0.57 [95% CI: 0.30; 1.09], *p*-trend = 0.09; HR_{1-unit increase} = 0.76 [95% CI: 0.55; 1.05], *p*-trend = 0.09; data not shown).

No significant modifications in the association between lymphoma or its subtypes and the arMED score were observed for age, sex and alcohol intake and smoking (Supporting Information Table S1). Similarly, no significant differences were observed among countries (Supporting Information Fig. S1).

In sensitivity analyses, including alcohol in the scoring did not affect the estimates for most of the lymphoma subtypes, although statistically significant associations were found for HL (HR_{1-unit increase} = 0.93 [95% CI: 0.86; 0.99], *p*-trend = 0.04) and DLBCL (HR_{1-unit increase} = 0.96 [95% CI: 0.92; 0.99], *p*-trend = 0.02) (Supporting Information Table S2). Moreover, excluding the first 2 years of follow-up and those individuals

with no information on adjustment variables from the analyses did not materially alter the results (data not shown).

Discussion

This is the first prospective study to investigate the association between a Mediterranean dietary score and risk of lymphoma and its subtypes. Our findings suggest that a higher adherence to the MD is modestly associated with a lower risk of lymphoma.

Current evidence on the adherence to MD and the etiology of lymphoma is scarce. To our knowledge, only one case-control study has evaluated the association of an *a priori* MD score and lymphoma, yielding no significant association for overall cases and only reporting an inverse association for DLBCL.²¹ However, those results require cautious interpretation due to the retrospective study design, small sample size (322 cases for the particular subgroup analysis) and the use of a nonvalidated adaptation of the arMED score, which did not include olive oil, white meat or dairy products. Other studies have extracted healthy-like dietary patterns from their population using data-driven analyses,^{13–15} but also failed in finding patterns with all the MD features (e.g., none of them included olive oil). With the exception of the prospective study of Erber *et al.*¹⁴ who reported an inverse association between a pattern rich in vegetables and fruits and NHL among Caucasian women, none of the other case-control studies found any associations with overall HL¹⁵ or NHL.¹³ However, given the heterogeneity in types of foods eaten within these patterns, the range and absolute amounts of food intakes and cut-offs used to define adherence, direct comparison of study results should be made with caution.

Similarly, there is a lack of consistency for associations between questionnaire-derived dietary components and lymphoma risk. In the 2007 report by the WCRF/AICR,²² the panel did not make any judgments regarding the causality of associations between specific dietary factors and lymphoid neoplasms. However, several suggestive associations were pointed out: (i) vegetables, fruits and alcoholic beverages were associated with decreased risk of lymphoma, (ii) meat, total fat and body fatness with increased risk of lymphoma and (iii) dairy products with increased risk of NHL. Although numerous studies have subsequently emerged, most of them targeted NHL patients, were mainly case-control studies and did not show consistent associations. Indeed, in the recently released third report of the WCRF/AICR, no additional information has been provided for hematological neoplasms.⁵ Recent meta-analyses have shed light on this relationship: (i) two meta-analyses found inverse relationships of vegetables and vegetables and fruit (combined) intake with NHL,^{23,24} (ii) another on foods of animal origin (including red, processed and white meat, fish and seafood, dairy products and eggs) reported positive associations with red meat and dairy intake and NHL²⁵, while (iii) while cohort studies on specific micronutrients point to null associations between supplemented

vitamins A, C and E, total vitamin D intake, as well as dietary lycopene intake and risk of NHL.²⁶ Overall, our results suggest that, more than specific dietary components is the combined effect of a range of nutrients along with the putative biological interactions that take place between them that may be mediating the modest influence of adherence to the MD on lymphoma risk.

In an exploratory analysis, we observed strong inverse associations with HL albeit statistically nonsignificant. Given the small number of HL ($N = 135$), we may have lacked statistical power to detect significant associations within this subgroup. Previous studies of diet and HL risk are scarcer and limited in power.^{6,7,15,21,27–34} Although no consistent associations have been reported for single-food items, the first study on dietary patterns and HL provided some insight.¹⁵ The authors found a suggestive inverse association between a diet characterized by high intake of fruit and low-fat dairy products and mixed cellularity HL. In addition, positive associations were reported between Western-like patterns (rich in meat or desserts and sweets) and specific HL entities and age groups. Together, these and our results suggest that HL might be a lymphoma prone to be influenced by dietary patterns. Thus, further studies with prospective design and with histological subtype-specific analyses, feasible though pooling data from consortium studies, are warranted.

The original MD score included alcohol scored dichotomous variable¹²: 2 points were assigned for moderate consumers (5–25 g/day for women and 10–50 g/day for men) and 0 points for those above and below the sex-specific range, owing its beneficial effects if consumed in moderation. Convincing evidence suggests that alcohol increases the risk of several carcinomas (e.g., mouth, pharynx and larynx, esophagus, liver, colorectal, breast and stomach)⁵ and indeed, the WCRF/AICR recommendations currently promote lowering alcohol consumption.⁵ However, accumulating evidence showed a moderate inverse association between increasing alcohol intake and NHL, especially on DLBCL and FL.¹⁹ Thus, for the current analyses, alcohol was not included in the score, and models were adjusted and further stratified by alcohol intake. In sensitivity analyses, when including alcohol in the score, statistically significant associations were found for HL and DLBCL. The influence of alcohol in lymphomagenesis remains largely unknown, and further studies are needed to clarify its role and possible interaction with dietary factors.

Certain dietary features of the MD and their potentially anticarcinogenic mechanisms make the association with lymphoma plausible from a mechanistic point of view. The abundance of plant-based foods in the MD provides a diet rich in flavonoids, carotenoids and vitamin C or E, whose important antioxidant properties can neutralize free radicals or prevent DNA damage.^{22,35} Indeed, total antioxidant intake has been inversely associated with lymphoma.^{36,37} Moreover, it presents a high monounsaturated to saturated fatty acid ratio, and its

Table 3. Association between adherence to the arMED score and risk of lymphoma and its subtypes in the EPIC study

	arMED score			<i>p</i> -value trend ¹	One-unit increase	<i>p</i> value ²
	Low	Medium	High			
Lymphoma, <i>n</i>	1,016	1,371	749			
HR ³ (95% CI)	Ref	0.92 (0.84; 1.01)	0.90 (0.79; 1.02)	0.08	0.98 (0.97; 1.00)	0.02
HR ⁴ (95% CI)	Ref	0.93 (0.85; 1.02)	0.91 (0.80; 1.03)	0.12	0.98 (0.97; 1.00)	0.03
HL, <i>n</i>	39	65	31			
HR ³ (95% CI)	Ref	0.96 (0.61; 1.51)	0.64 (0.34; 1.18)	0.15	0.93 (0.86; 1.00)	0.07
HR ⁴ (95% CI)	Ref	0.97 (0.61; 1.53)	0.64 (0.34; 1.19)	0.16	0.93 (0.86; 1.01)	0.07
NHL, <i>n</i>	835	1,128	643			
HR ³ (95% CI)	Ref	0.90 (0.81; 0.99)	0.93 (0.81; 1.06)	0.23	0.98 (0.97; 1.00)	0.06
HR ⁴ (95% CI)	Ref	0.90 (0.82; 1.00)	0.94 (0.82; 1.08)	0.31	0.98 (0.97; 1.00)	0.10
NHL subtypes ⁵						
Mature T/NK-cell, <i>n</i>	47	54	29			
HR ³ (95% CI)	Ref	0.73 (0.47; 1.13)	0.76 (0.42; 1.40)	0.31	0.99 (0.91; 1.07)	0.72
HR ⁴ (95% CI)	Ref	0.72 (0.47; 1.14)	0.78 (0.42; 1.44)	0.35	0.99 (0.91; 1.07)	0.80
Mature B-cell, <i>n</i>	766	1,043	593			
HR ³ (95% CI)	Ref	0.91 (0.82; 1.01)	0.94 (0.82; 1.08)	0.31	0.98 (0.97; 1.00)	0.08
HR ⁴ (95% CI)	Ref	0.91 (0.82; 1.01)	0.95 (0.82; 1.10)	0.40	0.98 (0.97; 1.00)	0.11
DLBCL, <i>n</i>	147	245	96			
HR ³ (95% CI)	Ref	1.09 (0.87; 1.37)	0.83 (0.60; 1.14)	0.35	0.96 (0.92; 1.00)	0.07
HR ⁴ (95% CI)	Ref	1.10 (0.88; 1.39)	0.84 (0.61; 1.17)	0.43	0.96 (0.93; 1.01)	0.09
FL, <i>n</i>	111	154	116			
HR ³ (95% CI)	Ref	0.84 (0.64; 1.11)	1.25 (0.90; 1.76)	0.19	1.02 (0.97; 1.07)	0.45
HR ⁴ (95% CI)	Ref	0.84 (0.64; 1.11)	1.27 (0.90; 1.79)	0.17	1.02 (0.97; 1.07)	0.43
CLL/SLL, <i>n</i>	182	213	142			
HR ³ (95% CI)	Ref	0.80 (0.64; 0.99)	0.95 (0.70; 1.27)	0.54	0.99 (0.96; 1.03)	0.75
HR ⁴ (95% CI)	Ref	0.78 (0.63; 0.98)	0.92 (0.68; 1.25)	0.45	0.99 (0.95; 1.03)	0.64
MM/PCN, <i>n</i>	226	283	167			
HR ³ (95% CI)	Ref	0.93 (0.77; 1.14)	0.98 (0.75; 1.28)	0.82	0.98 (0.95; 1.02)	0.40
HR ⁴ (95% CI)	Ref	0.95 (0.78; 1.16)	1.01 (0.77; 1.34)	0.98	0.99 (0.95; 1.03)	0.55
Other B-cell, <i>n</i>	100	148	72			
HR ³ (95% CI)	Ref	0.85 (0.64; 1.13)	0.67 (0.45; 1.00)	0.05	0.96 (0.91; 1.00)	0.08
HR ⁴ (95% CI)	Ref	0.85 (0.64; 1.14)	0.68 (0.46; 1.01)	0.06	0.96 (0.91; 1.00)	0.09

arMED, adapted relative Mediterranean dietary score; HR, hazard ratio; CI, confidence interval; SD, standard deviation; NHL, non-Hodgkin lymphoma; HL, Hodgkin lymphoma; DLBCL, diffuse large B-cell lymphoma; FL, follicular lymphoma; CLL/SLL, chronic lymphocytic leukemia/small lymphocytic lymphoma; MM/PCN, multiple myeloma/ plasma cell neoplasm; Other B-cell (those cases for which the mature B NHL subtype is unknown or does not fall within the more common subtypes).

In bold: *p* value < 0.05.

¹*p* value of Cox proportional model fitted with the arMED ordinal variable as continuous to test for lineal trend.

²*p* value of Cox proportional model fitted with the arMED continuous variable.

³Basic model: Cox proportional hazard model stratified by age (in 1-year categories), center and sex.

⁴Multivariate model: Cox proportional hazard model stratified by age (in 1-year categories), center and sex, and further adjusted for BMI, total energy intake, educational level, height, physical activity, smoking status and alcohol intake.

⁵NHL subtypes, excluding 37 precursor NHL and 37 individuals with NHL without B- or T-cell information.

believed that circulating fatty acids may influence lymphoma risk by modulating inflammation or lymphocyte membrane stability.³⁸ In addition, several studies have consistently linked chronic inflammation or autoimmune conditions with lymphomagenesis^{3,39} and reported associations between plasma levels of cytokines, or other inflammatory markers, and lymphoma.^{40–42} Recent studies are supporting the inflammatory potential of diet⁴³; in particular, for lymphoid neoplasms,

positive associations have been recently reported between a proinflammatory dietary score and NHL.⁴⁴ Thus, the MD's favorable fatty acid profile, as well as a high intake of fiber, vitamins and flavonoids, may be relevant owing its properties.

Interestingly, for the years 2000–2002, the incidence of total lymphoid malignances, in particular HL, was higher in Southern Europe (Italy, Malta and Spain) in comparison with

other European regions.⁴⁵ We are not aware of studies that have attempted to correlate known risk factors for lymphoid neoplasms with regional variations in incidence. Further research on other dietary patterns (e.g., adherence to a Western-like or a proinflammatory diet) or the nutrition transition toward non-Mediterranean dietary patterns in many Mediterranean countries⁴⁶ is warranted to elucidate these incidence patterns.

Limitations of our study should be considered when interpreting the results, including potential measurement errors derived from dietary questionnaires, which could lead to systematic and random errors when estimating adherence to the MD. Although our adjustment for total energy intake would partly remove some of these errors,^{47,48} we cannot rule out that they have affected risk estimates. In addition, we were unable to take into account any possible changes in dietary and lifestyle habits over time. In particular, cases might have modified their diet during the early prediagnostic period of the disease, although sensitivity analyses excluding incident cases diagnosed in the first 2 years of follow-up did not alter the association. Moreover, we lacked data on other potential confounders (e.g., occupational exposures or pesticide use) and, despite adjusting for all known lymphoma's risk factors, residual confounding by other unmeasured or unknown exposure cannot be dismissed. In addition, given the number of comparisons performed, we cannot exclude chance findings. Moreover, despite the large number of enrolled subjects at baseline, the number of observed incident cases of some lymphoma subtype was low (e.g., 135 HL). Therefore, the study might not have sufficient power to detect significant associations within those subgroups. Finally, the arMED score has also limitations, as similar weight is given to each component and the foods within them, assuming that they have the same effects on health. However, according to current evidence, these groupings did not include dietary components that have distinct effects on lymphoma risk. In addition, the EPIC study

included participants from both Mediterranean and non-Mediterranean regions, which may have distinct dietary intakes, food sources and sociodemographic, anthropometric and lifestyle characteristics that cannot be considered when using cohort-wide tertiles to construct the score. However, several studies have found similar associations' when using study-wide or country-specific cut-offs,^{8,49} and no interactions were found by country in our study.

The strengths of our study include its prospective design, long follow-up and large sample size, which allowed us to carry out analyses by lymphoma subentities. In addition, its multicentric European design allowed the inclusion of a geographically diverse population, covering a wide range of dietary patterns and lifestyle habits. Finally, we assessed adherence to the MD with a widely used score in cancer epidemiology, which directly includes olive oil intake, a key feature of this dietary pattern. Moreover, it uses tertiles of intake as cut-offs instead of the frequently used medians to give a better distribution of the subjects with different intakes.

In summary, this is the first prospective study to examine the association between the MD and lymphoma risk. Our findings suggest that an increasing arMED score, reflecting adherence to the MD, was modestly inversely associated with the risk of overall lymphoma. Further studies are needed to confirm these findings.

Availability of Data and Materials

For information on how to submit an application for gaining access to EPIC data and/or biospecimens, please follow the instructions at <http://epic.iarc.fr/access/index.php>.

Acknowledgements

We thank all participants of the EPIC study and Bertrand Hémon at IARC for their valuable work and technical support with the EPIC database.

References

1. Turner JJ, Morton LM, Linet MS, et al. Inter-Lymph hierarchical classification of lymphoid neoplasms for epidemiologic research based on the WHO classification (2008): update and future directions. *Blood* 2010;116:e90–8.
2. Fitzmaurice C, Allen C, Barber RM, et al. Global, regional, and National Cancer Incidence, mortality, years of life lost, years lived with disability, and disability-adjusted life-years for 32 cancer groups, 1990 to 2015: a systematic analysis for the global burden of disease study. *JAMA Oncol* 2017;3:524–48.
3. Morton LM, Slager SL, Cerhan JR, et al. Etiologic heterogeneity among non-Hodgkin lymphoma subtypes: the InterLymph non-Hodgkin lymphoma subtypes project. *J Natl Cancer Inst Monogr* 2014;2014:130–44.
4. Roman E, Smith AG. Epidemiology of lymphomas. *Histopathology* 2011;58:4–14.
5. World Cancer Research Fund/American Institute for Cancer Research. Diet, nutrition, physical activity, and cancer: a global perspective. Continuous update project expert report 2018. 2018. Available at dietandcancerreport.org
6. Rohrmann S, Linseisen J, Jakobsen MU, et al. Consumption of meat and dairy and lymphoma risk in the European prospective investigation into cancer and nutrition. *Int J Cancer* 2011;128:623–34.
7. Rohrmann S, Becker N, Linseisen J, et al. Fruit and vegetable consumption and lymphoma risk in the European prospective investigation into cancer and nutrition (EPIC). *Cancer Causes Control* 2007;18:537–49.
8. Couto E, Boffetta P, Lagiou P, et al. Mediterranean dietary pattern and cancer risk in the EPIC cohort. *Br J Cancer* 2011;104:1493–9.
9. Schwingshackl L, Schwedhelm C, Galbete C, et al. Adherence to Mediterranean diet and risk of cancer: an updated systematic review and meta-analysis. *Nutrients* 2017;9:1063.
10. Buckland G, Travier N, Cottet V, et al. Adherence to the Mediterranean diet and risk of breast cancer in the European prospective investigation into cancer and nutrition cohort study. *Int J Cancer* 2013;132:2918–27.
11. Bamia C, Lagiou P, Buckland G, et al. Mediterranean diet and colorectal cancer risk: results from a European cohort. *Eur J Epidemiol* 2013;28:317–28.
12. Buckland G, Agudo A, Lujan L, et al. Adherence to a Mediterranean diet and risk of gastric adenocarcinoma within the European prospective investigation into cancer and nutrition (EPIC) cohort study. *Am J Clin Nutr* 2010;91:381–90.
13. Ollberding NJ, Aschebrook-Kilfoy B, Caces DBD, et al. Dietary patterns and the risk of non-Hodgkin lymphoma. *Public Health Nutr* 2014;17:1531–7.

14. Erber E, Maskarinec G, Gill JK, et al. Dietary patterns and the risk of non-Hodgkin lymphoma: the multiethnic cohort. *Leuk Lymphoma* 2009;50:1269–75.
15. Epstein MM, Chang ET, Zhang Y, et al. Dietary pattern and risk of Hodgkin lymphoma in a population-based case-control study. *Am J Epidemiol* 2015;182:405–16.
16. Riboli E, Kaaks R. The EPIC project: rationale and study design. European prospective investigation into cancer and nutrition. *Int J Epidemiol* 1997;26(suppl 1):S6–14.
17. Riboli E, Hunt KJ, Slimani TN, et al. European prospective investigation into cancer and nutrition (EPIC): study populations and data collection. *Public Health Nutr* 2002;5:1113–24.
18. Margetts BM, Pietinen P. European prospective investigation into cancer and nutrition: validity studies on dietary assessment methods. *Int J Epidemiol* 1997;26(Suppl 1):S1–5.
19. Psaltopoulou T, Sergentanis TN, Ntanasis-Stathopoulos I, et al. Alcohol consumption and risk of hematological malignancies: a meta-analysis of prospective studies. *Int J Cancer* 2018;143:486–95.
20. Trichopoulou A, Costacou T, Bamia C, et al. Adherence to a Mediterranean diet and survival in a Greek population. *N Engl J Med* 2003;348:2599–608.
21. Campagna M, Cocco P, Zucca M, et al. Risk of lymphoma subtypes and dietary habits in a Mediterranean area. *Cancer Epidemiol* 2015;39:1093–8.
22. World Cancer Research Fund / American Institute for Cancer Research. *Food, nutrition, physical activity, and the prevention of cancer: a global perspective*. Washington, DC: AICR, 2007.
23. Chen G-C, Lv D-B, Pang Z, et al. Fruits and vegetables consumption and risk of non-Hodgkin's lymphoma: a meta-analysis of observational studies. *Int J Cancer* 2013;133:190–200.
24. Sergentanis TN, Psaltopoulou T, Ntanasis-Stathopoulos I, et al. Consumption of fruits, vegetables, and risk of hematological malignancies: a systematic review and meta-analysis of prospective studies. *Leuk Lymphoma* 2018;59:434–47.
25. Caini S, Masala G, Gnagnarella P, et al. Food of animal origin and risk of non-Hodgkin lymphoma and multiple myeloma: a review of the literature and meta-analysis. *Crit Rev Oncol Hematol* 2016;100:16–24.
26. Psaltopoulou T, Ntanasis-Stathopoulos I, Tsilimigras DI, et al. Micronutrient intake and risk of hematological malignancies in adults: a systematic review and meta-analysis of cohort studies. *Nutr Cancer* 2018;1–19.
27. Tavani A, Pregnolato A, Negri E, et al. Diet and risk of lymphoid neoplasms and soft tissue sarcomas. *Nutr Cancer* 1997;27:256–60.
28. Middleton B, Byers T, Marshall J, et al. Dietary vitamin A and cancer—a multisite case-control study. *Nutr Cancer* 1986;8:107–16.
29. Gao Y, Li Q, Bassig BA, et al. Subtype of dietary fat in relation to risk of Hodgkin lymphoma: a population-based case-control study in Connecticut and Massachusetts. *Cancer Causes Control* 2013;24:485–94.
30. Chang ET, Canchola AJ, Clarke CA, et al. Dietary phytochemicals and risk of lymphoid malignancies in the California teachers study cohort. *Cancer Causes Control* 2011;22:237–49.
31. La Vecchia C, Chatenoud L, Negri E, et al. Session: whole cereal grains, fibre and human cancer wholegrain cereals and cancer in Italy. *Proc Nutr Soc* 2003;62:45–9.
32. Lim U, Freedman DM, Hollis BW, et al. A prospective investigation of serum 25-hydroxyvitamin D and risk of lymphoid cancers. *Int J Cancer* 2009;124:979–86.
33. Negri E, La Vecchia C, Franceschi S, et al. Vegetable and fruit consumption and cancer risk. *Int J Cancer* 1991;48:350–4.
34. Tavani A, La Vecchia C, Gallus S, et al. Red meat intake and cancer risk: a study in Italy. *Int J Cancer* 2000;86:425–8.
35. Visioli F, Grande S, Bogani P, et al. The role of antioxidants in the Mediterranean diets: focus on cancer. *Eur J Cancer Prev* 2004;13:337–43.
36. Thompson CA, Habermann TM, Wang AH, et al. Antioxidant intake from fruits, vegetables and other sources and risk of non-Hodgkin's lymphoma: the Iowa Women's health study. *Int J Cancer* 2010;126:992–1003.
37. Holtan SG, O'Connor HM, Fredericksen ZS, et al. Food-frequency questionnaire-based estimates of total antioxidant capacity and risk of non-Hodgkin lymphoma. *Int J Cancer* 2012;131:1158–68.
38. Chiu Y-H, Bertrand KA, Zhang S, et al. A prospective analysis of circulating saturated and monounsaturated fatty acids and risk of non-Hodgkin lymphoma. *Int J Cancer* 2018;143:1914–22.
39. Smedby KE, Ponzoni M. The aetiology of B-cell lymphoid malignancies with a focus on chronic inflammation and infections. *J Intern Med* 2017;282:360–70.
40. Saberi Hosnijeh F, Krop EJM, Scoccianti C, et al. Plasma cytokines and future risk of non-Hodgkin lymphoma (NHL): a case-control study nested in the Italian European prospective investigation into cancer and nutrition. *Cancer Epidemiol Biomarkers Prev* 2010;19:1577–84.
41. Purdue MP, Lan Q, Bagni R, et al. Prediagnostic serum levels of cytokines and other immune markers and risk of non-Hodgkin lymphoma. *Cancer Res* 2011;71:4898–907.
42. Gu Y, Shore RE, Arslan AA, et al. Circulating cytokines and risk of B-cell non-Hodgkin lymphoma: a prospective study. *Cancer Causes Control* 2010;21:1323–33.
43. Shivappa N, Steck SE, Hurley TG, et al. Designing and developing a literature-derived, population-based dietary inflammatory index. *Public Health Nutr* 2014;17:1689–96.
44. Shivappa N, Hebert JR, Taborelli M, et al. Dietary inflammatory index and non-Hodgkin lymphoma risk in an Italian case-control study. *Cancer Causes Control* 2017;28:791–9.
45. Sant M, Allemanni C, Tereanu C, et al. Incidence of hematologic malignancies in Europe by morphologic subtype: results of the HAEMACARE project. *Blood* 2010;116:3724–34.
46. Moreno LA, Sarria A, Popkin BM. The nutrition transition in Spain: a European Mediterranean country. *Eur J Clin Nutr* 2002;56:992–1003.
47. Slimani N, Kaaks R, Ferrari P, et al. European prospective investigation into cancer and nutrition (EPIC) calibration study: rationale, design and population characteristics. *Public Health Nutr* 2002;5:1125–45.
48. Ferrari P, Kaaks R, Fahey MT, et al. Within- and between-cohort variation in measured macronutrient intakes, taking account of measurement errors, in the European prospective investigation into cancer and nutrition study. *Am J Epidemiol* 2004;160:814–22.
49. Molina-Montes E, Sanchez M-J, Buckland G, et al. Mediterranean diet and risk of pancreatic cancer in the European prospective investigation into cancer and nutrition cohort. *Br J Cancer* 2017;116:811–20.

24 days of stem cells

Shape the future of stem cell innovation
October 1- November 1, 2019

Join us for 24 Days of Stem Cells; a premiere virtual event featuring the latest advances in stem cell research.

This year's format will feature a new hour of cutting edge content every week day starting October 1st. Attend the sessions that are most relevant to your work - at your convenience and at your pace.

During the 24-day long event, you can:

- Access leading scientific presentations from thought leaders around the world
- Watch live training demonstrations from our stem cell experts
- Download key stem cell tools and resources
- Complete weekly challenges to earn points towards certification and prizes

Register today at
www.24daysofstemcells.com

ThermoFisher
SCIENTIFIC

WILEY