

LETTERS

Early Palaeogene temperature evolution of the southwest Pacific Ocean

Peter K. Bijl¹, Stefan Schouten³, Appy Sluijs¹, Gert-Jan Reichert², James C. Zachos⁴ & Henk Brinkhuis¹

Relative to the present day, meridional temperature gradients in the Early Eocene age (~56–53 Myr ago) were unusually low, with slightly warmer equatorial regions¹ but with much warmer subtropical Arctic² and mid-latitude³ climates. By the end of the Eocene epoch (~34 Myr ago), the first major Antarctic ice sheets had appeared^{4,5}, suggesting that major cooling had taken place. Yet the global transition into this icehouse climate remains poorly constrained, as only a few temperature records are available portraying the Cenozoic climatic evolution of the high southern latitudes. Here we present a uniquely continuous and chronostratigraphically well-calibrated TEX₈₆ record of sea surface temperature (SST) from an ocean sediment core in the East Tasman Plateau (palaeolatitude ~65° S). We show that southwest Pacific SSTs rose above present-day tropical values (to ~34 °C) during the Early Eocene age (~53 Myr ago) and had gradually decreased to about 21 °C by the early Late Eocene age (~36 Myr ago). Our results imply that there was almost no latitudinal SST gradient between subequatorial and subpolar regions during the Early Eocene age (55–50 Myr ago). Thereafter, the latitudinal gradient markedly increased. In theory, if Eocene cooling was largely driven by a decrease in atmospheric greenhouse gas concentration⁶, additional processes are required to explain the relative stability of tropical SSTs given that there was more significant cooling at higher latitudes.

The Palaeogene temperature evolution of the Antarctic margin, particularly the Pacific sector, is still poorly resolved. One difficulty with obtaining relevant records close to the Antarctic continent is the general absence of biogenic carbonate in most marine facies, which hampers traditional $\delta^{18}\text{O}$ and/or Mg/Ca-based reconstructions of the subpolar temperature evolution. In the absence of biogenic carbonates, organic sea-surface-temperature proxies such as the tetraether index of lipids consisting of 86 carbon atoms (TEX₈₆)⁷ and the alkenone unsaturation index (U^{k}_{37})⁸ are required for reconstructing high-latitude climatic evolution^{2,9}.

We apply TEX₈₆ and U^{k}_{37} on a stratigraphically continuous sedimentary section from the southwest Pacific Ocean, drilled by the Ocean Drilling Program (ODP Leg 189 Site 1172, palaeolatitude ~65° S (ref. 10); Fig. 1). A full methodological description is available in Supplementary Information. The record contains an expanded succession of marginal marine sediments from the lower Palaeocene epoch to the upper Eocene (64–36 Myr ago), with tight chronostratigraphic control, including magnetostratigraphy¹¹ (Supplementary Fig. 2). The presence of typical trans-Antarctic organic-walled dinoflagellate cysts in the Tasman region indicates an Antarctic-derived northward-flowing Tasman Current throughout the Palaeogene, which is verified by experiments based on general circulation models¹² (Fig. 1). This Antarctic influence at the East Tasman Plateau (ETP) persisted until at least the early Late Eocene (~35.5 Myr ago), when

deepening of the Tasmanian Gateway lead to a reorganization of the Tasman and proto-Leeuwin ocean currents¹³.

According to the oldest part of the record, TEX₈₆-derived SSTs at the ETP gradually decreased from ~25 °C around 63 Myr ago to a minimum of ~20 °C around 58 Myr ago (Fig. 2a). During the Late Palaeocene and Early Eocene, Tasman SSTs gradually rose to tropical values of ~34 °C during the Early Eocene climatic optimum (EECO)⁶, between 53 and 49 Myr ago (Fig. 2a). A gradual cooling trend throughout the Middle Eocene (starting at the termination of the EECO ~49 Myr ago) arrived at temperatures of ~23 °C ~42 Myr ago, which is still relatively warm. Subsequently, an interruption of the cooling trend occurred at the Middle Eocene climatic optimum (MECO; ~40 Myr ago)¹⁴, followed by a relatively rapid SST decrease to ~21 °C in the early Late Eocene (Fig. 2a). The late Middle and Late Eocene TEX₈₆-based SSTs are supported by U^{k}_{37} SST estimates derived from the same samples (Supplementary Fig. 3). Both SST estimates also compare well with those for other Late Eocene (Southern Ocean) sites⁹. Unfortunately, sediments from the ETP older than the MECO did not contain alkenones for U^{k}_{37} SST reconstructions.

The Middle Eocene SSTs correspond closely to those from sections in New Zealand^{15,16}, according to records based on TEX₈₆ (Fig. 2a), Mg/Ca and $\delta^{18}\text{O}$, indicating regional consistency of our reconstructed SSTs. Also, trends in our Tasman SST record are remarkably similar to those in the global stack of benthic foraminiferal oxygen isotopes⁶ (Fig. 2b), which we updated and augmented with recently published




Figure 1 | Site location and surface currents. Palaeogeographic reconstruction for the South Pacific Ocean at Early–Middle Eocene times. Surface circulation¹² indicates the Antarctic-derived Tasman Current (TC) over the East Tasman Plateau. Palaeogeographic charts obtained from the Ocean Drilling Stratigraphic Network (ODSN); after ref. 26. The dashed red arrow around New Zealand indicates potential mixing of low-latitude surface waters (from the East Australian Current) with the TC.

¹Palaeoecology, Institute of Environmental Biology, Faculty of Science, Laboratory of Palaeobotany and Palynology, ²Department of Geochemistry, Faculty of Geosciences, Utrecht University, Budapestlaan 4, 3584 CD Utrecht, The Netherlands. ³Department of Marine Organic Biogeochemistry, NIOZ Royal Netherlands Institute of Sea Research, PO Box 59, 1790 AB Den Burg, Texel, The Netherlands. ⁴Earth and Planetary Sciences Department and Institute of Marine Sciences, University of California, Santa Cruz, 1156 High St, Santa Cruz, California 95064, USA.




Figure 2 | Palaeogene deep-sea and sea surface temperatures. **a**, TEX₈₆ SST reconstructions from ODP Site 1172, New Zealand^{15,16} and Tanzania¹ (all according to the same calibration; see Supplementary Information). The black wiggly lines are short (~100 kyr)²⁷ and longer hiatuses at Site 1172. **b**, Global stack of benthic foraminiferal oxygen isotopes (grey data;

Supplementary Information). The temperature scale assumes ice-free conditions ($\delta^{18}\text{O}_{\text{SMOW}} = -1.2\text{‰}$, where $\delta^{18}\text{O}_{\text{SMOW}} = (^{18}\text{O}/^{16}\text{O})_{\text{sample}} / (^{18}\text{O}/^{16}\text{O})_{\text{SMOW}} - 1$; SMOW, standard mean ocean water), and indicates deep-sea temperatures. The black solid line reflects a five-point running average. PETM, Palaeocene–Eocene thermal maximum.

data (Supplementary Information). This correspondence between the two records (Supplementary Fig. 4) indicates that the regional SSTs co-varied with the SSTs where 'global' deep water was sourced. It has previously been suggested that the Southern Ocean was the main region of deep-water formation during the Palaeogene¹⁷.

In contrast, absolute SST estimates from the Tasman region are much higher than those inferred from the benthic foraminiferal oxygen isotopes (Fig. 2). Part of this discrepancy might be due to seasonality, with TEX₈₆ being slightly skewed towards summer temperatures and benthic foraminiferal $\delta^{18}\text{O}$ towards winter temperatures (Supplementary Information). Another possibility is that deep-water formation occurred in areas that were cooler than the Tasman sector. SST reconstructions based on bivalve-shell oxygen isotopes from Seymour Island on the Antarctic shelf, for example, yield much lower SSTs¹⁸. It is possible that the Antarctic margin was more susceptible to winter cooling than the open ocean, or that portions of the coast along the Southern Ocean gyres were somehow isolated from the southern edges of the Southern Ocean gyres. Another possibility is that the aragonite bivalve shells integrate temperature over a greater portion of the year. Regardless, the large SST difference between the Weddell Sea and the ETP would suggest a relatively steep gradient within a few degrees of latitude. Antarctica, being a polar continent, would most likely have experienced extremes in temperature, in particular having cool winters. Such conditions might have been recorded in the bivalves from the Weddell Sea but not in the more distal ETP. In turn, deep-water formation might have been restricted to the Antarctic shelf areas, such as the Weddell Sea.

Planktonic foraminiferal $\delta^{18}\text{O}$ analyses from equatorial regions previously indicated that Palaeogene low-latitude SSTs were the same, or even lower, than those of today¹⁹, a problem that puzzled palaeoclimate scientists for decades. The oxygen isotopic composition of planktonic foraminiferal tests in porous carbonate-rich pelagic facies were later found to be partially altered owing to recrystallization primarily during early diagenesis^{20,21}. In contrast, carbonate-poor and clay-rich facies typically found on the continental margins contain calcite shells without major diagenetic overprint¹. For the Eocene, such well-preserved planktonic foraminifera indicate near-equatorial SSTs that were greater than those of the present day, and agree with TEX₈₆-derived SSTs^{1,21}.

Another observation from well-preserved foraminifera and TEX₈₆ is that (sub)equatorial SSTs were remarkably stable throughout the Eocene¹ (Fig. 2). Stable low-latitude SSTs concomitant with high-latitude Eocene cooling thus suggests that there were increasing SST gradients during the Eocene. Although SST trends are often reconstructed using multi-proxy studies, the difference in absolute SSTs between various proxy reconstructions can be considerably large^{9,15,22}, even when measured on the same sediments. Despite the fact that multi-proxy approaches are generally encouraged in palaeoclimate studies, exclusion of such inter-proxy biases in latitudinal gradient reconstructions requires single-proxy SST records from around the world. Traditional calcite-based SST reconstructions are less suitable for this because calcite is only sparsely available in high-latitude sediments. The organic TEX₈₆ and U^k₃₇ SST proxies, however, can be used independently of latitude and are, hence, suitable for

single-proxy SST gradient reconstructions. Moreover, they do not require critical assumptions about ancient sea-water chemistry, unlike $\delta^{18}\text{O}$ and Mg/Ca .

We compiled Eocene TEX_{86} and U^{K}_{37} SST reconstructions from a suite of sedimentary records from localities worldwide and noted increased Middle Eocene latitudinal SST gradients in both hemispheres (Fig. 3), relative to the Early Eocene. These SST gradients are in general agreement with those found for terrestrial mean annual temperatures, based on Early–Middle Eocene fossil leaves²³. Adding the bivalve-based SST reconstructions from Seymour Island¹⁸ to our organic proxy data suggests a strong gradient between 60° and 70° S, which contrasts with the small gradient between 60° S and the Equator (Fig. 3). A part of this large Southern Ocean SST gradient might be due to biases between organic and calcite proxies. A large part, however, may realistically reflect the influence of the cool Antarctic interior, which cooled the Antarctic shelf. In contrast to the continental South Pole, the Arctic region is an oceanic basin. Instead of amplifying the seasonal cycle, the Arctic Ocean probably moderated seasonal extremes in the northern high-latitude greenhouse. Hence, Palaeogene latitudinal temperature gradients, like those of today, would have exhibited a high degree of asymmetry between the two hemispheres.

It has been suggested that the general warmth that characterized early Palaeogene climates was forced by high atmospheric greenhouse gas concentrations⁶. Concomitantly, the absence of polar ice sheets eliminated ice–albedo feedbacks in the Palaeogene greenhouse. The Middle–Late Eocene global cooling has been related to long-term atmospheric CO_2 decline, eventually resulting in the onset of major Antarctic glaciation around the Eocene/Oligocene boundary⁶. Our results imply that meridional temperature gradients markedly increased together with deep-sea cooling (Fig. 2)⁶. Although high latitudes cooled, tropical temperatures seem to have remained fairly stable throughout the Eocene (Fig. 3)¹. This observation raises questions concerning the precise role of decreasing atmospheric greenhouse gas concentrations in cooling the Eocene poles, as in theory²⁴ they should have cooled tropical regions as well. The role of potential




Figure 3 | Early and Middle Eocene latitudinal SST gradients. Bivalve-shell $\delta^{18}\text{O}$ (triangles), TEX_{86} (squares) and U^{K}_{37} (diamonds) SST reconstructions for the Early (orange) and mid-Middle (blue) Eocene. Data are from Seymour Island¹⁸ (a), the East Tasman Plateau (b), Deep Sea Drilling Project (DSDP) Site 277⁹ (c), New Zealand^{15,16} (d), DSDP Site 511⁹ (e), ODP Site 1090⁹ (f), Tanzania¹ (g), ODP Site 925⁹ (h), New Jersey³ (j, k; circle represents peak PETM SSTs³), ODP Site 336⁹ (m), ODP Site 913⁹ (n) and the Arctic Ocean^{2,28,29} (p) (Supplementary Fig. 1). Error bars indicate the range of variation. Gradients represent second-order polynomials, excluding bivalve-shell data. Black and dashed lines represent the present-day zonally averaged latitudinal temperature gradient³⁰ and age-specific deep-sea temperatures, respectively (Fig. 2b, ref. 6).

high-latitude climate feedbacks involving, for example, differences in cloud/water vapour distribution²⁵ might have been much more instrumental in the Middle Eocene climatic deterioration than previously thought. Another potential positive-feedback mechanism for high-latitude cooling would be ice–albedo feedback. However, the presence of substantial Middle Eocene continental ice is still equivocal given the general warmth and overall absence of conclusive physical evidence.

Received 31 March; accepted 6 August 2009.

- Pearson, P. N. *et al.* Stable tropical climate through the Eocene epoch. *Geology* **35**, 211–214 (2007).
- Sluijs, A. *et al.* Arctic late Paleocene–early Eocene paleoenvironments with special emphasis on the Paleocene–Eocene thermal maximum (Lomonosov Ridge, Integrated Ocean Drilling Program Expedition 302). *Paleoceanography* **23**, doi:10.1029/2007PA001495 (2008).
- Sluijs, A. *et al.* Environmental precursors to rapid light carbon injection at the Palaeocene/Eocene boundary. *Nature* **450**, 1218–1221 (2007).
- Zachos, J. C., Breza, J. R. & Wise, S. W. Jr. Early Oligocene ice sheet expansion on Antarctica: stable isotope and sedimentological evidence from Kerguelen Plateau, Southern Indian Ocean. *Geology* **20**, 569–573 (1992).
- Barker, P. F., Dieckmann, B. & Escutia, C. Onset of Cenozoic Antarctic glaciation. *Deep-Sea Res. II* **54**, 2293–2307 (2007).
- Zachos, J. C., Dickens, G. R. & Zeebe, R. E. An early Cenozoic perspective on greenhouse warming and carbon-cycle dynamics. *Nature* **451**, 279–283 (2008).
- Schouten, S., Hopmans, E. C., Schefuß, E. & Sinninghe Damsté, J. S. Distributional variations in marine crenarchaeotal membrane lipids: a new tool for reconstructing ancient sea water temperatures? *Earth Planet. Sci. Lett.* **204**, 265–274 (2002).
- Müller, P. J., Kirst, G., Rohland, G., von Storch, I. & Rosell-Melé, A. Calibration of the alkenone paleotemperature index U^{K}_{37} based on core-tops from the eastern South Atlantic and the global ocean (60°N–60°S). *Geochim. Cosmochim. Acta* **62**, 1757–1772 (1998).
- Liu, Z. *et al.* Global cooling during the Eocene–Oligocene climate transition. *Science* **323**, 1187–1190 (2009).
- Exon, N., Kennett, J. P. & Malone, M. (eds) *The Cenozoic Southern Ocean: Tectonics, Sedimentation, and Climate Change between Australia and Antarctica* (Geophys. Monogr. Ser. 151, American Geophysical Union, 2004).
- Stickley, C. E. *et al.* in *Proc. Ocean Drilling Program, Scientific Results* (eds Exon, N. F., Kennett, J. P. & Malone, M. J.) 1–57 (2004).
- Huber, M. *et al.* Eocene circulation of the Southern Ocean: was Antarctica kept warm by subtropical waters? *Paleoceanography* **19**, doi:10.1029/2004PA001014 (2004).
- Stickley, C. E. *et al.* Timing and nature of the deepening of the Tasmanian Gateway. *Paleoceanography* **19**, doi:10.1029/2004PA001022 (2004).
- Bohaty, S. M. & Zachos, J. C. Significant Southern Ocean warming event in the late Middle Eocene. *Geology* **31**, 1017–1020 (2003).
- Hollis, C. J. *et al.* Tropical sea temperatures in the high latitude South Pacific during the Eocene. *Geology* **37**, 99–102 (2009).
- Burgess, C. E. *et al.* Middle Eocene climate cyclicity in the southern Pacific: implications for global ice volume. *Geology* **36**, 651–654 (2008).
- Thomas, D. J., Bralower, T. J. & Jones, C. E. Neodymium isotopic reconstruction of the Late Paleocene–Early Eocene thermohaline circulation. *Earth Planet. Sci. Lett.* **209**, 309–322 (2003).
- Ivany, L. C. *et al.* Eocene climate record of a high southern latitude continental shelf: Seymour Island, Antarctica. *Geol. Soc. Am. Bull.* **120**, 659–678 (2008).
- Barron, E. J. Eocene equator-to-pole surface ocean temperatures: a significant climate problem? *Paleoceanography* **2**, 729–739 (1987).
- Schrag, D. P., DePaolo, D. J. & Richter, F. M. Reconstructing past sea surface temperatures: correcting for diagenesis of bulk marine carbonate. *Geochim. Cosmochim. Acta* **59**, 2265–2278 (1995).
- Pearson, P. N. *et al.* Warm tropical sea surface temperatures in the Late Cretaceous and Eocene epochs. *Nature* **413**, 481–487 (2001).
- Huber, M. A hotter greenhouse? *Science* **321**, 353–354 (2008).
- Greenwood, D. R. & Wing, S. L. Eocene continental climates and latitudinal temperature gradients. *Geology* **23**, 1044–1048 (1995).
- Huber, M. & Sloan, L. C. Heat transport, deep waters, and thermal gradients: coupled simulation of an Eocene ‘greenhouse’ climate. *Geophys. Res. Lett.* **28**, 3481–3484 (2001).
- Abbot, D. S. & Tziperman, E. Sea ice, high-latitude convection, and equable climates. *Geophys. Res. Lett.* **35**, doi:10.1029/2007GL032286 (2008).
- Hay, W. W. *et al.* Alternative global Cretaceous paleogeography. *Spec. Pap. Geol. Soc. Am.* **332**, 1–47 (1999).
- Röhl, U. *et al.* in *The Cenozoic Southern Ocean: Tectonics, Sedimentation, and Climate Change between Australia and Antarctica* (eds Exon, N., Kennett, J. P. & Malone, M.) 127–151 (Geophys. Monogr. Ser. 151, American Geophysical Union, 2004).
- Sangiorgi, F. *et al.* Cyclicity in the middle Eocene central Arctic Ocean sediment record: orbital forcing and environmental response. *Paleoceanography* **23**, doi:10.1029/2007PA001487 (2008).

29. Weller, P. & Stein, R. Paleogene biomarker records from the central Arctic Ocean (Integrated Ocean Drilling Program Expedition 302): organic carbon sources, anoxia, and sea surface temperature. *Paleoceanography* **23**, doi:10.1029/2007PA001472 (2008).
30. Shea, D. J., Trenberth, K. E. & Reynolds, R. W. A global monthly sea surface temperature climatology. *J. Clim.* **5**, 987–1001 (1992).

Supplementary Information is linked to the online version of the paper at www.nature.com/nature.

Acknowledgements Funding for this research was provided by Utrecht University, the Netherlands Organisation for Scientific Research (VICI grant to S.S.; VENI grant to A.S.) and the LPP Foundation. This research used samples and data provided by the Ocean Drilling Program (ODP). The ODP was sponsored by the US National Science

Foundation and participating countries under the management of Joint Oceanographic Institutions, Inc. G. Nobbe, E. van Bentum, E. Speelman, J. Ossebaar, A. Mets and E. Hopmans are thanked for technical support. We acknowledge C. J. Hollis, P. N. Pearson and P. F. Sexton for providing published data. A. J. P. (Sander) Houben, P. N. Pearson and M. Huber are thanked for critical comments.

Author Contributions P.K.B., S.S., H.B. and A.S. designed the research, P.K.B. and S.S. performed the organic geochemical analyses; P.K.B. updated the age model for ODP Site 1172 and performed the data compilations. All authors contributed to interpreting the data and writing the paper.

Author Information Reprints and permissions information is available at www.nature.com/reprints. Correspondence and requests for materials should be addressed to P.K.B. (p.k.bijl@uu.nl).