



ELSEVIER

Contents lists available at [ScienceDirect](https://www.sciencedirect.com)

Comparative Biochemistry and Physiology, Part A

journal homepage: www.elsevier.com/locate/cbpa



Corrigendum

Corrigendum to “Termination of puberty in out-of-season male Atlantic salmon smolts” [Comparative Biochemistry and Physiology Part A: Molecular & Integrative Physiology (2019) 60–66]



Thomas W.K. Fraser^{a,*}, Per Gunnar Fjellidal^a, Rüdiger W. Schulz^{b,c}, Birgitta Norberg^d, Tom J. Hansen^a

^a Reproduction and Developmental Biology Group, Institute of Marine research (IMR), Matre Aquaculture Research Station, 5984 Matredal, Norway

^b Reproductive Biology Group, Division Developmental Biology, Institute of Biodynamics and Biocomplexity, Department of Biology, Faculty of Science, University of Utrecht, 3584 CH, Utrecht, The Netherlands

^c Reproduction and Developmental Biology Group, Institute of Marine Research, P.O. Box 1870, Nordnes, 5817 Bergen, Norway

^d Reproduction and Developmental Biology Group, Institute of Marine research (IMR), Austevoll Research Station, 5392 Storebø, Norway

The authors regret that there is a mistake in Fig. 1B, the “common” water temperature has been plotted on the wrong axis, so appears 2 °C lower than it actually was.

The authors would like to apologise for any inconvenience caused.

DOI of original article: <https://doi.org/10.1016/j.cbpa.2019.03.011>

* Corresponding author.

E-mail address: thomas.fraser@hi.no (T.W.K. Fraser).

<https://doi.org/10.1016/j.cbpa.2019.06.015>

Available online 01 July 2019

1095-6433/ © 2019 Elsevier Inc. All rights reserved.




Fig. 1. Temperature and photoperiod from experiments one and two. (A) Experiment one. The photoperiod was simulated natural (60°N) between October and December 2016 and natural from mid-January 2017 onwards. The grey dots indicate the water temperature from the tank inflow. The timing of vaccination is indicated by a dotted line, whereas the dashed line indicates the timing of seawater transfer. (B) Experiment two. Two water temperatures were used between October and December 2016, the 17°C treatment in orange, and the 12°C treatment in blue. The photoperiod was natural from December 2016 onwards. The grey dots indicate the water temperature from the inflow when in tanks, and the average seawater temperature from 3–5 m depth when in sea-cages. The grey dotted line indicates the timing of vaccination, whereas the blue and orange lines indicate the date of seawater transfer, for the 12 and 17°C reared fish, respectively.