

Available online at www.sciencedirect.com

ScienceDirect

Journal homepage: www.elsevier.com/locate/cortex

Editorial

Prism adaptation: From rehabilitation to neural bases, and back

Yves Rossetti ^{a,*}, Shigeru Kitazawa ^b and Tanja Nijboer ^{c,d}

^a Inserm, CNRS, Université de Lyon, Hospices Civils de Lyon, ImpAct, Bron, France

^b Graduate school of Frontier Biosciences, Osaka University, Osaka, Japan

^c Experimental Psychology, Utrecht University, 3584 CS Utrecht, The Netherlands

^d Center of Excellence for Rehabilitation Medicine, Brain Center Rudolf Magnus, University Medical Center Utrecht, Utrecht University and De Hoogstraat Rehabilitation, 3583 TM Utrecht, The Netherlands

Prism adaptation rehabilitation for spatial cognition disorders has been explored since 1998 and a wave of prism adaptation research grew in clinical and basic science fields. A special issue of *Cortex* is devoted to this topic, which illustrates how translational research does not always work from workbench to bedside. Over the last 20 years prism adaptation has spread backward from the rehabilitation of unilateral neglect (McIntosh, Rossetti, & Milner, 2002; Rode et al., 2006a; Jacquin-Courtois, Rode, Pisella, Boisson, & Rossetti, 2008, 2010) to the exploration of the link between sensorimotor adaptation and healthy attentional processes (Colent, Pisella, Bernieri, Rode, & Rossetti, 2000; Michel et al., 2003a; Michel, 2006, 2016), body schema and reference frames (Girardi, McIntosh, Michel, Vallar, & Rossetti, 2004; Michel, Rossetti, Rode, & Tilikete, 2003b) and high level cognitive processes (e.g., Jacquin-Courtois et al., 2010; Rossetti et al., 2004). And parallel explorations led to the clinical use of prism adaptation to improve chronic pain (e.g., Christophe et al., 2016; Sumitani et al., 2007) or constructional disorders (Rode, Klos, Courtois-Jacquin, Rossetti, & Pisella, 2006b).

Uncovering the promising therapeutic usage of prism adaptation has given rise to numerous explorations in patients as well as clinical trials (Frassinetti, Angeli, Meneghello, Avanzi, & Làdavvas, 2002; Shiraishi, Yamakawa, Itou, Muraki, & Asada, 2008; Serino, Barbiani, Rinaldesi, & Làdavvas, 2009; Mizuno et al., 2011; Rode et al., 2015; Ten Brink, Visser-Meily, & Nijboer, 2015, 2017). It also boosted the need for understanding their mechanism of action at both the sensori-motor level (Harris 1963; Gaveau et al., 2018, 2014; Hatada, Miall, & Rossetti, 2006; Inoue, Uchimura, &

Kitazawa, 2016; Michel, Pisella, Prablanc, Rode, & Rossetti, 2007; Nijboer et al. 2008; O'Shea et al., 2017, 2014; Uchimura & Kitazawa, 2013) and their expansion to the cognitive level (e.g., Nijboer, Nys, van der Smagt, van der Stigchel, & Dijkerman, 2011; Rossetti et al., 2004; Rossetti et al., 2015). Despite of this blossoming rise of interest for prism adaptation, we lack empirical evidence and consensus on several important issues. First we need to better understand the basic mechanisms of adaptation and how (conscious vs unconscious, which threshold) and which errors (dynamic movement feedback, terminal errors) can feed back into the adaptive system (Gaveau et al; 2018; Hanajima et al. 2015). Second, the diverging results of clinical trials points to the need to clarify which patient may respond to which adaptation protocol and whether all patients may benefit from prism adaptation. Adaptation protocols used in the clinic vary in term of optical shift generated by prisms (from 5 deg to 20 dioptries), in terms of daily (1 or 2) and weekly dosage (from 1 to 10), total duration (1 day to several weeks) and number of adaptation sessions as well as in terms of activities performed during prism exposure (boring laboratory pointing task to ecological games of activities). Crucially they also vary in terms of sensori-motor adaptation assessment (none, proprioceptive shift or visual-proprioceptive realignment). A flourishing methodological arsenal has been recently invoked to explore the mechanisms of prism adaptation: PET, fMRI, TMS, paired-pulse TMS, tDCS, magnetic resonance spectroscopy and EEG have been used in combination with behavioural variables to identify elementary processes and neural circuits solicited by prism

* Corresponding author.

E-mail address: Yves.rossetti@inserm.fr (Y. Rossetti).

<https://doi.org/10.1016/j.cortex.2019.01.002>

0010-9452/© 2019 Published by Elsevier Ltd.

adaptation. The initial claim that prism adaptation may restore the interhemispheric balance in neglect patients has not received consensual support from these experimental studies and further and more systematic explorations are needed to offer the opportunity to produce a comprehensive view of the rich network that seems to be affected by prism adaptation. The emergence of integrated models of prism adaptation (Inoue et al., 2016; Petitot, O'Reilly, & O'Shea, 2018) is especially expected for optimal clinical applications of this deceptively simple technique.

The Cortex special issue aims at collecting up-to-date theoretical and empirical contributions to blaze the future trails of prism adaptation research. Future contributions in the field should address the following main issues: clarifying terminology, exploring the precise physiological mechanisms of sensori-motor adaptation and their neural network implementation, the peculiar expansion of prism adaptation to spatial cognition and attentional parameters. These explorations and modelling of the available corpus of data should help understanding the pathophysiology of therapeutic mechanisms of prism adaptation far beyond the sensorimotor and reference frame level (Jeannerod & Rossetti, 1993), to predict which patients should respond to this technique (e.g., Goedert, Chen, Foundas, & Barrett, 2018), and to envisage which other pathological conditions may benefit from prism adaptation (Jacquin-Courtois et al., 2013).

Terminology is a key tool for the mind to grasp empirical observations and to elaborate realistic models of the underlying mechanisms. Over the last century the terminology used to describe prism adaptation methods, observations and concepts has remained loosely defined and fluctuating. Attempts to clarify this terminology (e.g., Kornheiser 1976; Welch 1986; Redding, Rossetti, & Wallace, 2005; Bastian, 2008) has remained insufficiently effective and more efforts are needed to develop a consensual glossary in the field. A more detailed dissection of prism adaptation methodology and a more precise terminology will be required in order to outline conceptual and methodological issues and accurately dissect components and consequences of prism adaptation.

Exploring the precise mechanisms of prism adaptation has become a growing issue for several reasons. First clinical applications of prism adaptation have raised the need for a better understanding of this deceptively simple intervention (e.g., Barrett, Goedert, & Basso, 2012; Bastian, 2008; Milner & McIntosh, 2005; Redding et al., 2005; Striemer & Danckert, 2010). Second, new technologies have also begun to address the mechanisms of sensori-motor learning and adaptation (e.g., Darainy, Vahdat, & Ostry, 2013; Hertzfeld et al., 2014), although the boundary between these two mechanisms has remained insufficiently outlined. In-depth analysis of error signals and behavioural consequences of prism exposure allow for refined understanding of this type of sensori-motor adaptation (e.g., Gaveau et al., 2018, 2014; O'Shea et al., 2014; Uchimura & Kitazawa, 2013; Inoue et al. 2015).

Specifying the neural networks involved in prism adaptation is not only necessary to improve our knowledge about brain plasticity (e.g., Bastian, 2008; Pisella, Rode, Farnè, Tilikete,

& Rossetti, 2006) but also aims at better predicting the individual patient's responsiveness to prism adaptation (e.g., Crottaz-Herbette et al., 2017; Goedert et al., 2014, 2018; Ronchi, Rossi, Calzolari, Bolognini, & Vallar, 2018; Ten Brink et al., 2017). Recent neuronal recording (e.g., Inoue et al., 2016; Inoue & Kitazawa, 2018), brain stimulation (Panico, Sagliano, Grossi, & Trojano, 2016, 2017, 2018; Schintu et al., 2016; O'Shea et al., 2018), electrophysiological recordings (Martín-Arévalo et al., 2016) have brought more original insight and more innovative views of prism adaptation circuitry (Inoue & Kitazawa, 2018) than traditional functional brain imaging (e.g., Luauté, Halligan, Rode, Jacquin-Courtois, & Boisson, 2006; Chapman et al., 2010; Danckert, ...) did by confirming lesion studies (e.g., Weiner, Hallett, & Funkenstein, 1983; Pisella et al. 2004, 2005; Hanajima et al. 2015). Innovative functional imaging paradigms and new investigations of functional connectivity should provide fresh insights into the mechanisms of prism adaptation and its expansion properties (e.g., Luauté et al., 2009; Crottaz-Herbette et al., 2017; Tissièrè et al., 2018).


One major challenge about prism adaptation has remained the extraordinary expansion of an apparently basic sensori-motor task to high level cognition (Jacquin-Courtois et al., 2013; Rossetti et al., 2015). Prism adaptation had long remained conceived of a sensori-motor task that could stimulate sensori-motor adaptation objectified by purely sensori-motor testing. Parameters used to describe prism adaptation have therefore been restricted to the error reduction curve (visual feedback available during prism exposure, i.e., closed-loop) and compensatory after-effects (tested without visual feedback after removal of prisms, i. open loop). The total shift (post minus pre measures of open-loop pointing to an untrained visual target) was defined as the global expression of adaptation: a measure of the visual-proprioceptive realignment (Welch, Choe, & Heinrich, 1974; Redding and Wallace book, Redding et al., 2005). Visual and proprioceptive shifts have been additionally used to disentangle between visual and proprioceptive contribution to the total realignment and the only window on prism adaptation has remained sensori-motor. Improvement of spatial cognition impairments seen in spatial neglect (Rode, Rossetti, Li, & Boisson, 1998; Rode et al. 2001; Rossetti et al., 1998) stood in sharp contrast with this long-standing sensori-motor tradition, and this contrast was further emphasised by the discovery that healthy subjects are also affected beyond the sensori-motor level (Colent et al., 2000). Even though it is a very general idea that cognition is grounded on sensorimotor interactions in both the field of developmental psychology (Piaget, 1962) and the more recent field of embodied cognition, prism adaptation offered the possibility to overtly alter cognition following a few minutes of a simple pointing task, opening a window onto this mysterious link between motor interactions with the environment and mental representations. A first wave of studies attempted to delineate in which domains and to what extent such sensorimotor to cognitive expansion of adaptation applies (e.g., Michel et al. 2001; 2003; Girardi et al., 2004; Striemer, Sablatnig, & Danckert, 2006; Fortis, Goedert, & Barrett, 2011). A second, ongoing wave of investigation has begun to explore

the mechanisms of this expansion (Crottaz-Herbette et al., 2017, 2014; Martín-Arévalo et al., 2016; Tsujimoto et al. 2018) and much progress in our scientific conception of embodiment is expected from this endeavour.

At the clinical level three main important issues have to be addressed. First, we have to explore and define how to obtain the most intense, the most generalizable and the most durable effects (e.g., Hatada et al., 2006; O'Shea et al., 2014). Past studies have been trying to define the optimal exposure and optimal regime of prism adaptation (Fortis, Ronchi, Calzolari, Gallucci, & Vallar, 2013, 2018, 2010; Facchin, Bultitude, Mornati, Peverelli, & Daini, 2018; Frassinetti et al., 2002; Goedert, Zhang, & Barrett, 2015; Luauté et al., 2012; Rode et al., 2015; Vaes et al., 2018) and their efficacy in daily life activities (Champod, Frank, Taylor, & Eskes, 2018; Jacquin-

& McIntosh, 2005; Newport & Schenk, 2012; Rode, Fourtassi, Pagliari, Pisella, & Rossetti, 2017; Rossetti et al., 2015; Striemer & Danckert, 2010).

This special issue of Cortex is devoted at paving the ground for the future explorations of the relationship between sensori-motor adaptation and spatial cognition's physiology, patho-physiology and therapeutico-physiology. This first virtual special issue of Cortex will be gathering the most advanced theoretical and methodological, basic empirical and clinical contributions ranging from case reports to large group studies and from refined behavioural to neural approaches, with the aim of contributing to a field that has begun to considerably renew our approaches and understanding of the dynamical interactions between action and cognition.



Number of publications indexed in WoS (ophthalmology excluded):

Courtois et al., 2008; Mizuno et al., 2011; Watanabe & Amimoto, 2010). Accordingly, reviews have been emphasising the need for designing more systematic studies and increasing evidence levels (Barrett et al., 2012; Champod et al., 2018; Fasotti & van Kessel, 2013; Jacquin-Courtois et al., 2013; Luauté et al., 2006; Rode et al., 2006). However recent combinations of techniques have extensively renewed this question (e.g., Guinet et al., 2013; Lådavas et al., 2015; Luauté et al., 2018; O'Shea et al., 2017) and much development is expected in the near future on this point. Second, it is of prime importance to be able to predict whether an individual patient is likely to benefit from prism adaptation, and whether this method should be used alone or in combination with other techniques. Third, further exploration are likely to uncover other domains of application of prism adaptation. So far the available evidence that it can improve spatial neglect (Rossetti et al., 1998), postural control in left hemiparesis (Tilikete et al., 2001) and neglect (Nijboer et al. 2014), spatial dysgraphia (Rode et al., 2006a), constructional deficits (Rode et al., 2006b) or complex regional pain syndrome (Christophe 2016; Sumitani et al., 2007) does not allow to provide a clear and integrated framework to define its application fields. Last, and most perspective gaining is the idea that therapeutical interventions such as prism adaptation may backward shed light on the pathophysiology of the relevant conditions. Beneficial effects of prism adaptation and identification of affected clinical and psychophysical parameters have begun to stimulate new conceptions of the neurophysiological and maladaptive mechanisms leading to impairments (e.g., Milner

REFERENCES

- Barrett, A. M., Goedert, K. M., & Basso, J. C. (2012). Prism adaptation for spatial neglect after stroke: Translational practice gaps. *Nature Reviews. Neurology*, 8(10), 567–577.
- Bastian, A. J. (2008). Understanding sensorimotor adaptation and learning for rehabilitation. *Current Opinion in Neurology*, 21(6), 628–633. <https://doi.org/10.1097/WCO.0b013e328315a293>.
- Champod, A. S., Frank, R. C., Taylor, K., & Eskes, G. A. (2018). The effects of prism adaptation on daily life activities in patients with visuospatial neglect: A systematic review. *Neuropsychological Rehabilitation*, 28(4), 491–514.
- Chapman, H. L., Eramudugolla, R., Gavrilescu, M., Strudwick, M. W., Loftus, A., Cunnington, R., et al. (2010). Neural mechanisms underlying spatial realignment during adaptation to optical wedge prisms. *Neuropsychologia*, 48(9), 2595–2601.
- Christophe, L., Chabanat, E., Delporte, L., Revol, P., Volckmann, P., Jacquin-Courtois, S., et al. (2016). Prisms to shift pain away: Pathophysiological and therapeutic exploration of CRPS with prism adaptation. *Neural Plasticity*, 1–21.
- Colent, C., Pisella, L., Bernieri, C., Rode, G., & Rossetti, Y. (2000). Cognitive bias induced by visuo-motor adaptation to prisms: A simulation of unilateral neglect in normal individuals? *Neuroreport*, 11(9), 1899–1902.
- Crottaz-Herbette, S., Fornari, E., & Clarke, S. (2014). Prismatic adaptation changes visuospatial representation in the inferior parietal lobule. *The Journal of Neuroscience: the Official Journal of the Society for Neuroscience*, 34(35), 11803–11811.
- Crottaz-Herbette, S., Fornari, E., Nottter, M. P., Bindschadler, C., Manzoni, L., & Clarke, S. (2017). Reshaping the brain after

- stroke: The effect of prismatic adaptation in patients with right brain damage. *Neuropsychologia*, 104, 54–63.
- Crottaz-Herbette, S., Fornari, E., Tissieres, I., & Clarke, S. (2017). A brief exposure to leftward prismatic adaptation enhances the representation of the ipsilateral, right visual field in the right inferior parietal lobule. *eNeuro*, 27(5), 4.
- Darainy, M., Vahdat, S., & Ostry, D. J. (2013). Perceptual learning in sensorimotor adaptation. *Journal of Neurophysiology*, 110(9), 2152–2162.
- Danckert, J., Ferber, S., & Goodale, M. A. (2008 Oct). Direct effects of prismatic lenses on visuomotor control: an event-related functional MRI study. *Eur J Neurosci*, 28(8), 1696–1704. <https://doi.org/10.1111/j.1460-9568.2008.06460.x>.
- Facchin, A., Bultitude, J. H., Mornati, G., Peverelli, M., & Daini, R. (2018). A comparison of prism adaptation with terminal versus concurrent exposure on sensorimotor changes and spatial neglect. *Neuropsychological rehabilitation*, 1–28. <https://doi.org/10.1080/09602011.2018.1484374>.
- Fasotti, L., & van Kessel, M. (2013). Novel insights in the rehabilitation of neglect. *Frontiers in Human Neuroscience*, 7, 780.
- Fortis, P., Goedert, K. M., & Barrett, A. M. (2011). Prism adaptation differently affects motor-intentional and perceptual-attentional biases in healthy individuals. *Neuropsychologia*, 49(9), 2718–2727.
- Fortis, P., Maravita, A., Gallucci, M., Ronchi, R., Grassi, E., Senna, I., et al. (2010). Rehabilitating patients with left spatial neglect by prism exposure during a visuomotor activity. *Neuropsychology*, 24(6), 681–697.
- Fortis, P., Ronchi, R., Calzolari, E., Gallucci, M., & Vallar, G. (2013). Exploring the effects of ecological activities during exposure to optical prisms in healthy individuals. *Frontiers in Human Neuroscience*, 7, 29.
- Fortis, P., Ronchi, R., Velardo, V., Calzolari, E., Banco, E., Algeri, L., et al. (2018). A home-based prism adaptation training for neglect patients. *Cortex*. <https://doi.org/10.1016/j.cortex.2018.09.001>. pii: S0010-9452(18)30288-0.
- Frassinetti, F., Angeli, V., Meneghello, F., Avanzi, S., & Làdavas, E. (2002). Long-lasting amelioration of visuospatial neglect by prism adaptation. *Brain*, 125, 608–623.
- Gaveau, V., Prablanc, C., Laurent, D., Rossetti, Y., & Priot, A. E. (2014). Visuomotor adaptation needs a validation of prediction error by feedback error. *Frontiers in Human Neuroscience*, 8(880), 1–15.
- Gaveau, V., Priot, A. E., Pisella, L., Havé, L., Prablanc, C., & Rossetti, Y. (2018). Paradoxical adaptation of successful movements: The crucial role of internal error signals. *Consciousness and Cognition*, 64, 135–145.
- Girardi, M., McIntosh, R. D., Michel, C., Vallar, G., & Rossetti, Y. (2004). Sensorimotor effects on central space representation: Prism adaptation influences haptic and visual representations in normal subjects. *Neuropsychologia*, 42(11), 1477–1487.
- Goedert, K. M., Chen, P., Boston, R. C., Foundas, A. L., & Barrett, A. M. (2014). Presence of motor-intentional aiming deficit predicts functional improvement of spatial neglect with prism adaptation. *Neurorehabilitation and Neural Repair*, 28(5), 483–493.
- Goedert, K. M., Chen, P., Foundas, A. L., & Barrett, A. M. (2018). Frontal lesions predict response to prism adaptation treatment in spatial neglect: A randomised controlled study. *Neuropsychological Rehabilitation*, 1–22.
- Goedert, K. M., Zhang, J. Y., & Barrett, A. M. (2015). Prism adaptation and spatial neglect: The need for dose-finding studies. *Frontiers in Human Neuroscience*, 9, 243.
- Guinet, M., & Michel, C. (2013). Prism adaptation and neck muscle vibration in healthy individuals: Are two methods better than one? *Neuroscience*, 254, 443–451.
- Hanajima, R., Shadmehr, R., Ohminami, S., Tsutsumi, R., Shirota, Y., Shimizu, T., et al. (2015). Modulation of error-sensitivity during a prism adaptation task in people with cerebellar degeneration. *Journal of Neurophysiology*, 114(4), 2460–2471.
- Harris, C. S. (1963). Adaptation to displaced vision: Visual, motor, or proprioceptive change? *Science*, 140(3568), 812–813.
- Hatada, Y., Miall, R. C., & Rossetti, Y. (2006). Two waves of a long-lasting after-effect of prism adaptation measured over 7 days. *Experimental Brain Research*, 169(3), 417–426.
- Herzfeld, D. J., Pastor, D., Haith, A. M., Rossetti, Y., Shadmehr, R., & O'Shea, J. (2014). Contributions of the cerebellum and the motor cortex to acquisition and retention of motor memories. *Neuroimage*, 98, 147–158.
- Inoue, M., & Kitazawa, S. (2018). Motor error in parietal area 5 and target error in area 7 drive distinctive adaptation in reaching. *Current Biology*: CB, 28(14), 2250–2262.
- Inoue, M., Uchimura, M., Karibe, A., O'Shea, J., Rossetti, Y., & Kitazawa, S. (2015). Three timescales in prism adaptation. *Journal of Neurophysiology*, 113(1), 328–338.
- Inoue, M., Uchimura, M., & Kitazawa, S. (2016). Error signals in motor cortices drive adaptation in reaching. *Neuron*, 90(5), 1114–1126.
- Jacquin-Courtois, S., O'Shea, J., Luauté, J., Pisella, L., Revol, P., Mizuno, K., et al. (2013). Rehabilitation of spatial neglect by prism adaptation: A peculiar expansion of sensorimotor after-effects to spatial cognition. *Neuroscience and Biobehavioral Reviews*, 37(4), 594–609.
- Jacquin-Courtois, S., Rode, G., Pavani, F., O'Shea, J., Giard, M. H., Boisson, D., et al. (2010). Effect of prism adaptation on left dichotic listening deficit in neglect patients: Glasses to hear better? *Brain*, 133(3), 895–908.
- Jacquin-Courtois, S., Rode, G., Pisella, L., Boisson, D., & Rossetti, Y. (2008). Wheel-chair driving improvement following visuo-manual prism adaptation. *Cortex*, 44(1), 90–96.
- Jeannerod, M., & Rossetti, Y. (1993). Visuomotor coordination as a dissociable visual function: Experimental and clinical evidence. *Baillieres Clin Neurol*, 2(2), 439–460.
- Kornheiser, A. S. (1976). Adaptation to laterally displaced vision: A review. *Psychological Bulletin*, 83, 783–816.
- Làdavas, E., Giulietti, S., Avenanti, A., Bertini, C., Lorenzini, E., Quinquinio, C., et al. (2015). a-tDCS on the ipsilesional parietal cortex boosts the effects of prism adaptation treatment in neglect. *Restorative Neurology and Neuroscience*, 33(5), 647–662.
- Luauté, J., Halligan, P., Rode, G., Jacquin-Courtois, S., & Boisson, D. (2006). Prism adaptation first among equals in alleviating left neglect: A review. *Restorative Neurology and Neuroscience*, 24(4–6), 409–418.
- Luauté, J., Jacquin-Courtois, S., O'Shea, J., Christophe, L., Rode, G., Boisson, D., et al. (2012). Left-deviating prism adaptation in left neglect patient: Reflexions on a negative result. *Neural Plasticity*, 718604.
- Luauté, J., Schwartz, S., Rossetti, Y., Spiridon, M., Rode, G., Boisson, D., et al. (2009). Dynamic changes in brain activity during prism adaptation. *Journal of Neuroscience*, 29(1), 169–178.
- Luauté, J., Villeneuve, L., Roux, A., Nash, S., Bar, J. Y., Chabanat, E., et al. (2018). Adding methylphenidate to prism-adaptation improves outcome in neglect patients. A randomized clinical trial. *Cortex*, 106, 288–298.
- Martín-Arévalo, E., Laube, I., Koun, E., Farné, A., Reilly, K. T., & Pisella, L. (2016). Prism adaptation alters electrophysiological markers of attentional processes in the healthy brain. *The Journal of Neuroscience: the Official Journal of the Society for Neuroscience*, 36(3), 1019–1030.
- McIntosh, R. D., Rossetti, Y., & Milner, A. D. (2002). Prism adaptation improves chronic visual and haptic neglect: A single case study. *Cortex*, 38(3), 309–320.
- Michel, C. (2006). Simulating unilateral neglect in normals: Myth or reality? *Restorative Neurology and Neuroscience*, 24(4–6), 419–430.
- Michel, C. (2016). Beyond the sensorimotor plasticity: Cognitive expansion of prism adaptation in healthy individuals. *Frontiers in Psychology*, 6, 1979.

- Michel, C., Pisella, L., Halligan, P. W., Luauté, J., Rode, G., Boisson, D., et al. (2003a). Simulating unilateral neglect in normals using prism adaptation: Implications for theory. *Neuropsychologia*, 41(1), 25–39.
- Michel, C., Pisella, L., Prablanc, C., Rode, G., & Rossetti, Y. (2007). Enhancing visuomotor adaptation by reducing error signals: Single-step (aware) versus multiple-step (unaware) exposure to wedge prisms. *Journal of Cognitive Neuroscience*, 19(2), 341–350.
- Michel, C., Rossetti, Y., Rode, G., & Tilikete, C. (2003b). After-effects of visuo-manual adaptation to prisms on body posture in normal subjects. *Experimental Brain Research*, 148(2), 219–226.
- Milner, A. D., & McIntosh, R. D. (2005). The neurological basis of visual neglect. *Current Opinion in Neurology*, 18(6), 748–753.
- Mizuno, K., Tsuji, T., Takebayashi, T., Fujiwara, T., Hase, K., & Liu, M. (2011). Prism adaptation therapy enhances rehabilitation of stroke patients with unilateral spatial neglect: A randomized, controlled trial. *Neurorehabilitation and Neural Repair*, 25(8), 711–720.
- Newport, R., & Schenk, T. (2012). Prisms and neglect: What have we learned? *Neuropsychologia*, 50(6), 1080–1091.
- Nijboer, T. C., McIntosh, R. D., Nys, G. M., Dijkerman, H. C., & Milner, A. D. (2008). Prism adaptation improves voluntary but not automatic orienting in neglect. *Neuroreport*, 19(3), 293–298.
- Nijboer, T. C., Nys, G. M., van der Smagt, M. J., van der Stigchel, S., & Dijkerman, H. C. (2011). Repetitive long-term prism adaptation permanently improves the detection of contralesional visual stimuli in a patient with chronic neglect. *Cortex*, 47(6), 734–740.
- O'Shea, J., Gaveau, V., Kandel, M., Koga, K., Susami, K., Prablanc, C., et al. (2014). Kinematic markers dissociate error correction from sensorimotor realignment during prism adaptation. *Neuropsychologia*, 55, 15–24.
- O'Shea, J., Revol, P., Cousijn, H., Near, J., Petitet, P., Jacquin-Courtois, S., et al. (2017). Induced sensorimotor cortex plasticity remediates chronic treatment-resistant visual neglect. *eLife*, 6, e26602.
- Panico, F., Jacquin-Courtois, S., Di Marco, J., Perrin, C., Trojano, L., & Rossetti, Y. (2017). tDCS reactivation of dormant adaptation circuits. *Cortex*, 94, 196–199.
- Panico, F., Sgaliano, L., Grossi, D., & Trojano, L. (2016). Cerebellar cathodal tDCS interferes with recalibration and spatial realignment during prism adaptation procedure in healthy subjects. *Brain and Cognition*, 105, 1–8.
- Panico, F., Sgaliano, L., Nozzolillo, C., Trojano, L., & Rossetti, Y. (2018). Cerebellar contribution to spatial realignment: A tDCS study during multiple-step prism adaptation. *Neuropsychologia*, 112, 58–65.
- Petitet, P., O'Reilly, J. X., & O'Shea, J. (2018). Towards a neuro-computational account of prism adaptation. *Neuropsychologia*, 115, 188–203.
- Piaget, J. (1962). The stages of the intellectual development of the child. *Bulletin of the Menninger Clinic*, 26, 120–128.
- Pisella, L., Michel, C., Gréa, H., Tilikete, C., Vighetto, A., & Rossetti, Y. (2004). Preserved prism adaptation in bilateral optic ataxia: Strategic versus adaptive reaction to prisms. *Experimental Brain Research*, 156(4), 399–408.
- Pisella, L., Rode, G., Farnè, A., Tilikete, C., & Rossetti, Y. (2006). Prism adaptation in the rehabilitation of patients with visuo-spatial cognitive disorders. *Current Opinion in Neurology*, 19(6), 534.
- Pisella, L., Rossetti, Y., Michel, C., Rode, G., Boisson, D., Pélisson, D., et al. (2005). Ipsidirectional impairment of prism adaptation after unilateral lesion of anterior cerebellum. *Neurology*, 65(1), 150–152.
- Redding, G. M., Rossetti, Y., & Wallace, B. (2005). Applications of prism adaptation: A tutorial in theory and method. *Neuroscience and Biobehavioral Reviews*, 29(3), 431–444.
- Rode, G., Fourtassi, M., Pagliari, C., Pisella, L., & Rossetti, Y. (2017). Complexity vs. unity in unilateral spatial neglect. *Revue Neurologique*, 173(7–8), 440–450.
- Rode, G., Klos, T., Courtois-Jacquin, S., Rossetti, Y., & Pisella, L. (2006b). Neglect and prism adaptation: A new therapeutic tool for spatial cognition disorders. *Restorative Neurology and Neuroscience*, 24(4–6), 347–356.
- Rode, G., Lacour, S., Jacquin-Courtois, S., Pisella, L., Michel, C., Revol, P., et al. (2015). Long-term sensorimotor and therapeutical effects of a mild regime of prism adaptation in spatial neglect. A double-blind RCT essay. *Annals of Physical and Rehabilitation Medicine*, 58(2), 40–53.
- Rode, G., Pisella, L., Marsal, L., Mercier, S., Rossetti, Y., & Boisson, D. (2006a). Prism adaptation improves spatial dysgraphia following right brain damage. *Neuropsychologia*, 44(12), 2487–2493.
- Rode, G., Rossetti, Y., & Boisson, D. (2001). Prism adaptation improves representational neglect. *Neuropsychologia*, 39(11), 1250–1254.
- Rode, G., Rossetti, Y., Li, L., & Boisson, D. (1998). Improvement of mental imagery after prism exposure in neglect: A case study. *Behavioural Neurology*, 11(4), 251–258.
- Ronchi, R., Rossi, I., Calzolari, E., Bolognini, N., & Vallar, G. (2018). Exploring prism exposure after hemispheric damage: Reduced aftereffects following left-sided lesions. *Cortex*. <https://doi.org/10.1016/j.cortex.2018.10.014>. [Epub ahead of print].
- Rossetti, Y., Jacquin-Courtois, S., Calabria, M., Michel, C., Gallagher, S., Honore, J., et al. (2015). Testing cognition and rehabilitation in unilateral neglect with wedge prism adaptation: Multiple interplays between sensorimotor adaptation and spatial cognition. In C. Kansaku, & Birbaumer (Eds.), *Clinical systems neuroscience*. Japan: Springer.
- Rossetti, Y., Jacquin-Courtois, S., Rode, G., Ota, H., Michel, C., & Boisson, D. (2004). Does action make the link between number and space representation? Visuo-manual adaptation improves number bisection in unilateral neglect. *Psychological Science*, 15(6), 426–430.
- Rossetti, Y., Rode, G., Pisella, L., Farnè, A., Li, L., Boisson, D., et al. (1998). Prism adaptation to a rightward optical deviation rehabilitates left hemispatial neglect. *Nature*, 395(6698), 166–169.
- Schintu, S., Martín-Arévalo, E., Vesia, M., Rossetti, Y., Salemme, R., Pisella, L., et al. (2016). Paired-pulse parietal-motor stimulation differentially modulates corticospinal excitability across hemispheres when combined with prism adaptation. *Neural Plasticity*, 5716179.
- Serino, A., Barbiani, M., Rinaldesi, M. L., & Làdavas, E. (2009). Effectiveness of prism adaptation in neglect rehabilitation: A controlled trial study. *Stroke; a Journal of Cerebral Circulation*, 40(4), 1392–1398.
- Shiraishi, H., Yamakawa, Y., Itou, A., Muraki, T., & Asada, T. (2008). Long-term effects of prism adaptation on chronic neglect after stroke. *Neurorehabilitation*, 23(2), 137–151.
- Striemer, C. L., & Danckert, J. A. (2010). Through a prism darkly: Re-evaluating prisms and neglect. *Trends Cogn Sci*, 14(7), 308–316.
- Striemer, C., Sablatnig, J., & Danckert, J. (2006). Differential influences of prism adaptation on reflexive and voluntary covert attention. *Journal of the International Neuropsychological Society: JINS*, 12(3), 337–349.
- Sumitani, M., Rossetti, Y., Shibata, M., Matsuda, Y., Sakaue, G., Inoue, T., et al. (2007). Prism adaptation to optical deviation alleviates pathologic pain. *Neurology*, 68(2), 128–133.
- Ten Brink, A. F., Visser-Meily, J. M. A., & Nijboer, T. C. W. (2015). Study protocol of 'prism adaptation in rehabilitation': A randomized controlled trial in stroke patients with neglect. *BMC Neurology*, 15, 5.
- Ten Brink, A. F., Visser-Meily, J. M. A., Schut, M. J., Kouwenhoven, M., Eijsackers, A., & Nijboer, T. C. W. (2017). Prism adaptation in rehabilitation? No additional effects of prism adaptation on neglect recovery in the subacute phase

- post-stroke - a randomised controlled trial. *Neurorehabilitation and Neural Repair*, 31(12), 1017–1028.
- Tilikete, C., Rode, G., Rossetti, Y., Pichon, J., Li, L., & Boisson, D. (2001). Prism adaptation to rightward optical deviation improves postural imbalance in left-hemiparetic patients. *Current Biology*, 11(7), 524–528.
- Tsujimoto, K., Mizuno, K., Nishida, D., Tahara, M., Yamada, E., Shindo, S., et al. (2018). Prism adaptation changes resting-state functional connectivity in the dorsal stream of visual attention networks in healthy adults: A fMRI study. *Cortex*. <https://doi.org/10.1016/j.cortex.2018.10.018>. [Epub ahead of print].
- Uchimura, M., & Kitazawa, S. (2013). Cancelling prism adaptation by a shift of background: A novel utility of allocentric coordinates for extracting motor errors. *The Journal of Neuroscience: the Official Journal of the Society for Neuroscience*, 33(17), 7595–7602.
- Vaes, N., Nys, G., Lafosse, C., Dereymaeker, L., Oostra, K., Hemelsoet, D., et al. (2018). Rehabilitation of visuospatial neglect by prism adaptation: Effects of a mild treatment regime. A randomised controlled trial. *Neuropsychological Rehabilitation*, 28(6), 899–918.
- Watanabe, S., & Amimoto, K. (2010). Generalization of prism adaptation for wheelchair driving task in patients with unilateral spatial neglect. *Archives of Physical Medicine and Rehabilitation*, 91(3), 443–447.
- Weiner, M. J., Hallett, M., & Funkenstein, H. H. (1983). Adaptation to lateral displacement of vision in patients with lesions of the central nervous system. *Neurology*, 33(6), 766–772.
- Welch, R. B., Choe, C. S., & Heinrich, D. R. (1974). Evidence for a three-component model of prism adaptation. *Journal of Experimental Psychology*, 103(4), 700–705.