

Nivolumab for Relapsed/Refractory Diffuse Large B-Cell Lymphoma in Patients Ineligible for or Having Failed Autologous Transplantation: A Single-Arm, Phase II Study

Stephen M. Ansell, MD, PhD¹; Monique C. Minnema, MD, PhD²; Peter Johnson, MD³; John M. Timmerman, MD⁴; Philippe Armand, MD, PhD⁵; Margaret A. Shipp, MD⁵; Scott J. Rodig, MD, PhD^{5,6}; Azra H. Ligon, PhD⁶; Margaretha G.M. Roemer, PhD⁵; Nishitha Reddy, MD⁷; Jonathon B. Cohen, MD⁸; Sarit Assouline, MD⁹; Michelle Poon, MD¹⁰; Manish Sharma, MD¹¹; Kazunobu Kato, MD, PhD¹¹; Selda Samakoglu, MD, PhD¹¹; Anne Sumbul, MS¹¹; and Andrew Grigg, MD¹²

PURPOSE Treatment options are limited for patients with relapsed/refractory diffuse large B-cell lymphoma (DLBCL). Tumor cells can exploit the programmed death-1 checkpoint pathway to evade immune surveillance. In the current study, we evaluated the efficacy and safety of programmed death-1 blockade by nivolumab in patients with relapsed/refractory DLBCL.

METHODS In this phase II, open-label study, patients with relapsed/refractory DLBCL who were ineligible for autologous hematopoietic cell transplantation (auto-HCT) or who had experienced failure with auto-HCT received nivolumab 3 mg/kg every 2 weeks. We assessed the efficacy and safety of nivolumab as well as genetic alterations of 9p24.1.

RESULTS Among 121 treated patients, patients in the auto-HCT–failed cohort (n = 87) received a median of four nivolumab doses and a median of three doses were administered to those in the auto-HCT–ineligible cohort (n = 34). At a median follow-up of 9 months in the auto-HCT–failed cohort and 6 months in the auto-HCT–ineligible cohort, independently assessed objective response rates were 10% and 3%, and median durations of response were 11 and 8 months, respectively. Median progression-free survival and overall survival were 1.9 and 12.2 months in the auto-HCT–failed cohort and 1.4 and 5.8 months in the auto-HCT–ineligible cohort respectively. All three patients with complete remission—3% of the auto-HCT–failed cohort—had durable response (11 or more, 14 or more, and 17 months). Treatment-related grade 3 and 4 adverse events were reported in 24% of patients. The most common were neutropenia (4%), thrombocytopenia (3%), and increased lipase (3%). Of all evaluable samples for 9p24.1 analysis, 16% exhibited low-level copy gain and 3% had amplification.

CONCLUSION Nivolumab monotherapy is associated with a favorable safety profile but a low overall response rate among patients with DLBCL who are ineligible for auto-HCT or who experienced failure with auto-HCT. Genetic alterations of 9p24.1 are infrequent in DLBCL.

J Clin Oncol 37:481-489. © 2019 by American Society of Clinical Oncology

INTRODUCTION

Diffuse large B-cell lymphoma (DLBCL) is the most common subtype of non-Hodgkin lymphoma worldwide.¹ With standard front-line therapy, approximately two thirds of adult patients achieve long-term remission and others who experience chemosensitive relapse may benefit from autologous hematopoietic cell transplantation (auto-HCT).^{2,3} However, patients with refractory DLBCL or those who are unsuitable for or who have experienced relapse after auto-HCT have limited treatment options,⁴ with a median survival of only 6 to 10 months from progression.^{3,5} Early results of chimeric antigen receptor T-cell therapy in small numbers of selected patients with refractory DLBCL

are promising, but the technology is costly and long-term data are not available.⁶ Accessible treatments that provide durable responses and improved outcomes are needed in this setting.

Programmed death-1 (PD-1) and its ligands PD-L1/PD-L2 are immune checkpoints that, in healthy populations, downregulate immune response and are crucial for maintaining self-tolerance and preventing autoimmunity.^{7,8} Genes that encode PD-L1/PD-L2 are located on chromosome 9p24.1.⁹ Genetic alterations at the locus lead to ligand overexpression, which is common in Hodgkin lymphoma (HL) but are yet to be fully characterized in DLBCL.¹⁰⁻¹² Overexpression of PD-L1 by tumor cells and on tumor-infiltrating

ASSOCIATED CONTENT

Appendix

Data Supplement

Author affiliations and support information (if applicable) appear at the end of this article.

Accepted on November 13, 2018 and published at [jco.org](https://doi.org/10.1200/JCO.18.00766) on January 8, 2019; DOI <https://doi.org/10.1200/JCO.18.00766>

The views expressed in this article are the authors' own and not an official position of Bristol-Myers Squibb or their respective institutions.

Clinical trials information: NCT02038933.

nonmalignant cells in the tumor microenvironment has the potential to interact with PD-1–expressing T cells and B cells, which results in the inhibition of antitumor immune response.^{7,8} Increased PD-L1 expression has been found in certain defined subtypes of large B-cell lymphoma (LBCL), such as primary mediastinal B-cell lymphoma and Epstein-Barr virus (EBV)–positive and select non–germinal center cell DLBCLs, and is associated with inferior overall survival (OS).^{8,10,12,13} Thus, blockade of the PD-1/PD-L1 pathway may have the potential to exert antitumor effects in certain subsets of DLBCL.

Nivolumab is a fully human immunoglobulin G4 anti–PD-1 monoclonal antibody that blocks tumor cell signaling via the PD-1 pathway, which releases T cells from the inhibitory effects of tumor cells and restores T-cell–mediated anti-tumor immune responses.¹⁴ In clinical trials, nivolumab monotherapy has demonstrated activity in solid tumors, including melanoma, non–small-cell lung cancer, renal cell cancer, and bladder cancer, among others,^{15–18} and in relapsed/refractory classic HL (cHL).^{19,20} In a phase I dose-escalation study of nivolumab monotherapy in patients with relapsed/refractory hematologic malignancies, four of 11 patients with DLBCL demonstrated objective responses.²¹ These preliminary results led to this phase II study, which evaluated the efficacy and safety of nivolumab monotherapy in patients with relapsed/refractory DLBCL after auto-HCT or who were not candidates for auto-HCT.

METHODS

Study Design and Patients

This was a multicenter, single-arm, open-label, phase II study conducted in accordance with Good Clinical Practice and the Declaration of Helsinki and approved by the institutional review board and independent ethics committee. All patients provided written informed consent before trial enrollment.

Eligibility criteria included age 18 years or older and Eastern Cooperative Oncology Group performance status of 0 or 1. Patients had de novo DLBCL or transformed lymphoma (confirmed by biopsy before initiation of the study drug) that had either relapsed after high-dose conditioning chemotherapy and auto-HCT or was relapsed/refractory after two or more prior multiagent chemotherapy regimens if auto-HCT ineligible. Key exclusion criteria included prior therapy with any an antibody or drug specifically targeting T-cell costimulation or checkpoint pathways, prior allogeneic HCT, known CNS lymphoma, and history of interstitial lung disease.

Treatment

Patients received nivolumab 3 mg/kg intravenously over 60 minutes every 2 weeks until disease progression, unacceptable toxicity, or withdrawal from the study. Dose delays were permitted for drug-related adverse events

(AEs), but dose escalations or reductions were not. Treatment beyond investigator-assessed disease progression was permitted in patients who did not experience rapid disease progression and who had investigator-determined clinical benefit from nivolumab and stable performance status.

Assessments

Efficacy. Patients were evaluated for tumor response according to the 2007 International Working Group response criteria for malignant lymphoma²² using spiral computed tomography/magnetic resonance imaging. Evaluation was performed at baseline, beginning at week 9 and continuing every 8 weeks through the first 8 months of treatment, every 12 weeks during months 9 to 24, and then every 6 months thereafter until disease progression or until the patient initiated a preparative regimen for allogeneic or auto-HCT. Tumor assessments via fluorodeoxyglucose–positron emission tomography were performed at baseline and were required to confirm complete remission (CR). Baseline and all subsequent scans were submitted to an independent radiology review committee (IRC) for assessment.

Safety. Safety evaluations included assessment of AEs, clinical laboratory tests, and physical examination with assessment of Eastern Cooperative Oncology Group performance status. Local laboratory assessments were performed within 72 hours before dosing. After discontinuation from the study, safety evaluations were scheduled at the first follow-up visit (35 days from the last dose) and the second follow-up (80 days from the first follow-up visit). Patients were then observed every 3 months for ongoing treatment-related AEs and survival.

Biomarkers. In patients with archival tumor biopsies, 9p24.1 genetic alterations were evaluated using a fluorescence in situ hybridization (FISH) assay. Probes encompassed *CD274* (*PD-L1*) or *PDCD1LG2* (*PD-L2*) and included a centromeric control. Copy number alterations were defined as previously described^{11,23} on the basis of the target:control signal ratio in 50 analyzed tumor cells per DLBCL. Nuclei with a target:control signal ratio of 3 or more:1 were defined as coamplified for *PD-L1* and *PD-L2*, and those with a signal ratio of more than 1:1 but less than 3:1 were classified as having relative copy gain. Nuclei with a signal ratio of 1:1 were defined as either polysomic if more than two copies per probe or disomic if there were exactly two copies of the target and control probes.¹¹ For each patient, we noted the percentage and magnitude of 9p24.1 amplification, copy gain, polysomy, and disomy. Patients were classified by the highest observed level of 9p24.1 genetic alteration. Those with 9p24.1 copy gain lacked amplification and those with 9p polysomy lacked 9p24.1 copy gain or amplification. Chromosomal rearrangements that involved *PD-L1* or *PD-L2* were also detected with this assay.

TABLE 1. Baseline Characteristics and Demographics

Characteristic	Auto-HCT Failed (n = 87)	Auto-HCT Ineligible (n = 34)
Age		
Median, years (range)	62 (24-75)	68 (28-86)
< 65	56 (64)	11 (32)
≥ 65	31 (36)	23 (68)
≥ 75	1 (1)	10 (29)
Male sex	56 (64)	21 (62)
Race		
White	71 (82)	31 (91)
ECOG performance status		
0	35 (40)	9 (26)
1	49 (56)	25 (74)
2*	2 (2)	0
3*	1 (1)	0
Prior lines of therapy		
Median (range)	3 (1-11)	3 (1-7)
≥ 4	21 (24)	10 (29)
Refractory to most recent line of therapy	24 (28)	20 (59)
Transformed disease	5 (6)	1 (3)
Time since diagnosis, median, years (range)	3.3 (< 1-28)	1.5 (< 1-11)
Disease stage at diagnosis		
I	8 (9)	1 (3)
II	12 (14)	6 (18)
III	23 (26)	8 (24)
IV	41 (47)	19 (56)
Not reported	3 (3)	0

NOTE. Data presented as No. (%) unless otherwise indicated. Totals may not equal 100% as a result of rounding.

Abbreviations: auto-HCT, autologous hematopoietic cell transplantation; ECOG, Eastern Cooperative Oncology Group.

*Protocol deviation.

We performed dual immunohistochemical staining of PD-L1 (clone 405.9A11)²⁴ and PAX5 (24/Pax-5; BD Biosciences, San Jose, CA) to delineate PD-L1 expression on tumor biopsies, as previously described.^{19,23} H-score (0 to 300) was calculated by multiplying the percentage of malignant (PAX5^{dim+}) cells with PD-L1-positive staining and the average intensity of staining (0 to 3 or more on 50 malignant cells).

Cell-of-origin (COO) assay by gene expression profiling was performed at Bristol-Myers Squibb by targeted expression directly from formalin-fixed, paraffin-embedded sections as previously described.²⁵ The HTG EdgeSeq DLBCL Cell of Origin Assay comprised probes that targeted 93 genes commonly assessed in lymphomas. Cell Of Origin was

assigned using HTG proprietary software (HTG Molecular Diagnostics, Tucson, AZ).

In situ hybridization was performed for EBV-encoded RNA-1 (EBER-1) using formalin-fixed, paraffin-embedded tissue samples. We tested RNA preservation using a nonspecific positive control, and background was assessed with a negative control. Scoring was performed for EBV/EBER if RNA was preserved and score was greater than background. Scores were EBV/EBER positive nuclear staining of any intensity greater than background in tumor cells.

Outcomes

The primary end point was objective response rate (ORR), defined as either partial remission (PR) or CR, as assessed by IRC using the 2007 revised International Working Group criteria. Secondary objectives included duration of response (DOR) on the basis of IRC assessments, IRC-assessed CR rate and duration of CR, IRC-assessed PR rate and duration of PR (DOPR), IRC-assessed progression-free survival (PFS), and investigator-assessed ORR. OS, safety and tolerability, and biomarker assessment were among additional exploratory end points.

Statistical Analyses

Planned sample size was approximately 120 treated patients, divided into two treatment groups on the basis of prior auto-HCT failure (n = 90) or auto-HCT ineligibility (n = 30). The primary analysis population for safety and efficacy was all patients who had received one or more dose of nivolumab. The protocol had an interim analysis that allowed for discontinuation of the study as a result of a lack of efficacy; however, accrual was rapid and the protocol allowed continued enrollment while the interim analysis was performed. This resulted in complete accrual before the interim analysis was concluded. For the auto-HCT-failed cohort, the null hypothesis that the true ORR was 20% or less would be rejected if 25 or more responses were observed in 90 treated patients. For the auto-HCT-ineligible cohort, the sample size of 30 treated patients was determined such that if the observed number of patients with objective response was 10, the true ORR was larger than 17% with 95% confidence.

IRC-assessed ORR, CR, and PR rates, duration of CR, DOPR, and investigator-assessed ORR were summarized for both cohorts separately by binomial response rates and their corresponding two-sided 95% CIs. IRC-assessed DOR, PFS, and OS were analyzed by cohort using the Kaplan-Meier product-limit method with median values and two-sided CI. Survival rate at 6 months was estimated using Kaplan-Meier values from the survival curve. AEs were coded using National Cancer Institute Common Terminology Criteria for Adverse Events, version 4.0.²⁶ All on-study AEs, serious AEs (SAEs), and treatment-related AEs and SAEs were tabulated according to worst grade per National Cancer Institute Common Terminology Criteria for

Adverse Events, version 4.0, by system organ class and Medical Dictionary for Regulatory Activities preferred term.

RESULTS

Patient Baseline Characteristics and Disposition

Patients were enrolled between March 2014 and August 2015 across 45 sites globally (Appendix Fig A1, online only). Of 121 patients treated, 87 had relapsed/refractory disease after auto-HCT and 34 were auto-HCT ineligible. Baseline characteristics of the two groups are listed in Table 1. In general, patients in the auto-HCT–failed group were younger and had better baseline performance status and longer time since diagnosis than did patients who were ineligible for auto-HCT. Patients had received a median of three prior systemic therapies in each cohort. In the auto-HCT–failed cohort, 40 patients (46%) did not receive any additional lines of systemic or radiation therapy between auto-HCT and the initiation of nivolumab. In the auto-HCT–failed group, 40% of patients had germinal center B-cell–like (GCB) subtype and 32% had non-GCB subtype disease, and in the auto-HCT–ineligible group, 56% had GCB and 18% non-GCB subtype disease. Data were missing in the remainder. All but one patient who had evaluable samples were EBV negative (70 of 121 patients), one patient was EBV positive, and the remainder (n = 50) had either nonevaluable data, mixed results (more than one sample), or missing data.

Median number of doses of nivolumab received were four (range, one to 44 doses) in the auto-HCT–failed cohort and

three (range, one to 22 doses) in the auto-HCT–ineligible cohort. At the time of analysis, seven patients (8%) in the auto-HCT–failed group were still on treatment, for whom the median duration of treatment was 14 months from the first study dose. No patients in the auto-HCT–ineligible cohort were still on treatment. Disease progression was the primary reason for discontinuing treatment (83%).

Efficacy

At a median follow-up of 9 months (range, 0.1 to 25 months) in the auto-HCT–failed cohort and 6 months (0.2 to 24 months) in the auto-HCT–ineligible cohort, IRC-assessed ORRs were 10% and 3%, respectively (Table 2). Nearly one third of patients (31%) in the auto-HCT–failed cohort had IRC-assessed stable disease or better response (Fig 1A). Of the nine responders in the auto-HCT–failed cohort, CR occurred in three, with durable response and DOR of 11 or more months, 14 or more months, and 17 months at data cutoff. Of these three patients who achieved CR, two were still on treatment and one developed myelodysplastic syndrome unrelated to study drug and subsequently died. The other six patients achieved PR, with a median DOPR of 7 months (95% CI, 3 to 11 months). The one patient with IRC-assessed PR in the auto-HCT–ineligible group had a DOR of 8.3 months. Patients who achieved CR or PR had early responses, with a median time to response of 1.9 months (Fig 1C). Responses were observed in both the GCB subtype (CR in one and PR in four patients) and non-GCB subtype (CR in one and PR in three patients). Among responders, eight were EBV negative, one was not evaluable, and one had missing data (Appendix Table A1, online only). One patient with PR had

TABLE 2. Response to Nivolumab Treatment

Treatment Response	Auto-HCT Failed (n = 87)		Auto-HCT Ineligible (n = 34)	
	IRC	INV	IRC	INV
Overall response rate,* No. (%)	9 (10)	16 (18)	1 (3)	1 (3)
Nonresponders†	78 (90)	71 (82)	33 (97)	33 (97)
Best overall response, No. (%)				
Complete remission	3 (3)	4 (5)	0	0
Partial remission	6 (7)	12 (14)	1 (3)	1 (3)
Stable disease	18 (21)	12 (14)	3 (9)	1 (3)
Progressive disease	41 (47)	43 (49)	24 (71)	26 (76)
Unable to determine‡	19 (22)	16 (18)	6 (18)	6 (18)
Duration of response, months				
Median (95% CI)	11.4 (3 to 17)	7.4 (2 to 14)	8 (NE to NE)	8 (NE to NE)
Minimum, maximum	2§, 17	2, 19§	8, 8	8, 8

Abbreviations: auto-HCT, autologous hematopoietic cell transplantation; INV, investigator; IRC, independent radiology review committee; NE, not estimable.

*Complete remission or partial remission.

†Nonresponders included patients who had stable disease, progressive disease, or whose status was undetermined.

‡Patients who died or who went off study before the first radiographic assessment.

§Censored value with ongoing response.




FIG 1. Change in target lesion burden per independent radiology review committee by best overall response in all response-evaluable patients. Best reduction in target lesion in patients (A) who experienced failure with autologous hematopoietic cell transplantation (auto-HCT) and (B) who were ineligible for auto-HCT. Tumor burden change over time in patients ($n = 9$) of the auto-HCT–failed cohort who had (C) complete remission (CR) or partial remission (PR) and (D) stable disease (SD; $n = 15$). Horizontal reference line indicates 50% reduction consistent with a response per revised 2007 International Working Group criteria. Response-evaluable patients had assessments at baseline and at one or more postbaseline time point.

transformed lymphoma. Among the seven patients who were still on treatment, two had CR, two PR, and three stable disease. Nineteen patients in the auto-HCT–failed cohort and five patients in the auto-HCT–ineligible cohort were treated with nivolumab beyond disease progression—none achieved response after progression.

Median PFS was 1.9 months in the auto-HCT–failed cohort and 1.4 months in the auto-HCT–ineligible cohort, with a 6-month IRC-assessed PFS rate of 19.1% and 5.2%, respectively (Fig 2A). Median OS was 12.2 months in the auto-HCT–failed cohort and 5.8 months in the

auto-HCT–ineligible cohort, and 6-month OS rate was 67% and 47%, respectively (Fig 2B).

Safety

Grade 3 or 4 AEs were reported in 62% of all treated patients, of which 24% were deemed to be treatment related; none had a frequency of more than 5% (Table 3). The most common treatment-related AEs (TRAEs) were nausea (17%), fatigue (17%), and diarrhea (10%)—these were generally not severe (majority grade 1 to 2). Treatment-related SAEs were reported in 14 patients (12%).




FIG 2. (A) Progression-free survival (PFS) and (B) overall survival (OS) as assessed by independent radiology review committee. Symbols represent censored observations. auto-HCT, autologous hematopoietic cell transplantation.

Four patients (3%) had TRAEs as the primary cause of discontinuation of treatment. TRAEs that contributed to treatment discontinuation included neutropenia, thrombocytopenia, diarrhea, pancreatitis, lipase increase, and dermatitis psoriasiform.

Rates of immune-mediated AEs were low in all treated patients with extended follow-up. The most common grade 3 to 4 immune-mediated AEs were nephritis and renal dysfunction (4%), hepatitis and hepatic dysfunction (3%), diarrhea (3%), and rash (2%).

Overall, 78 patients died. The primary reason for death was disease progression in 39 (45%) and 29 (85%) patients in the auto-HCT–failed and auto-HCT–ineligible cohorts, respectively. No death was attributable to study drug toxicity.

FISH and Immunohistochemistry

In total, 74 of 112 patients had evaluable tumor biopsy specimens for 9p24.1 FISH; 34% of DLBCL specimens were disomic and 45% were polysomic at *PD-L1/PD-L2*. *PD-L1/PD-L2* polysomy in these DLBCLs was low level, largely only three copies in these tumors. Only 16% of DLBCLs exhibited copy gain, which was also low level in the majority of cases (three copies in nine of 13 cases). An additional 3% of tumors had amplification, including *PD-L2*–selective amplification in one patient. Patients were classified by the highest-level 9p24.1 alterations (Figs 3A and 3B), as previously described.^{11,23} DLBCLs with 9p24.1 amplification had additional tumor cells with copy gain (0% to 86%) and/or disomy (4% to 8%). Tumors with relative copy gain had additional cells with polysomy (2% to 58%) and/or disomy (2% to 82%), and DLBCLs that were categorized as polysomic had additional disomic tumor cells (14% to 98%; Fig 3B). The percentage of residual disomic cells was lowest in DLBCLs with amplification, intermediate in tumors with copy gain, and highest in tumors with polysomy ($P < .001$; Fig 3C), which is consistent with an ordered spectrum of 9p24.1 genetic alterations.

In this series of DLBCLs with infrequent low-level 9p24.1 alterations, membranous PD-L1 expression was only detected in four (9%) of 46 evaluable cases. One of the three patients who achieved CR had high-level 9p24.1 amplification, whereas the other two had normal 9p24.1 copy number or an unavailable biopsy specimen. Of the seven patients who achieved PR, five had available biopsy specimens and either normal 9p24.1 copy numbers (one patient), low-level polysomy (three patients), or copy gain (one patient). The five evaluable patients with PR had no detectable PD-L1 expression on tumor cells (Appendix Table A1).

TABLE 3. Drug-Related Adverse Events With an Incidence of $\geq 5\%$

Adverse Event	All Treated Patients (N = 121)	
	Any Grade	Grade 3 and 4
Any adverse event	118 (98)	75 (62)
Drug-related adverse event	75 (62)	29 (24)
Nausea	21 (17)	1 (< 1)
Fatigue	20 (17)	2 (2)
Diarrhea	12 (10)	1 (< 1)
Neutropenia	9 (7)	5 (4)
Thrombocytopenia	7 (6)	4 (3)
Decreased appetite	7 (6)	1 (< 1)
Lipase increased	6 (5)	4 (3)
Rash	6 (5)	1 (< 1)
Pyrexia	6 (5)	1 (< 1)

NOTE. Data presented as No. (%).

improvement of responses in the limited number of patients who had been treated beyond disease progression.

The study protocol included a prespecified stopping rule for futility but, on the basis of the safety profile of nivolumab and promising efficacy results observed in the phase I study, allowed enrollment to continue while the interim analysis result was pending. As a result of rapid accrual and delays related to IRC, the planned enrollment was reached before the futility analysis was concluded. Nevertheless, the majority of patients received only a short duration of nivolumab treatment and were allowed to switch to other cancer therapies upon discontinuation of the study treatment; there was no new safety signal observed. In the future, we shall ensure that study procedures are closely monitored to avoid potential delays in the timely completion of the interim analysis.

Previously reported data suggest that PD-L1 expression is associated with poor prognosis in patients with DLBCL⁸; however, the prevalence of PD-L1 expression on DLBCL tumor cells seemed to be low (11%; with a 30% threshold) in that series,⁸ as well as in a EBV-negative DLBCL series (11%; with a 5% threshold).¹⁰ The 9p24.1 FISH analyses on a large number of DLBCL trial cases in this study indicate that the incidence and magnitude of 9p24.1 alterations are significantly lower in DLBCL than in cHL, as is the level of PD-L1 expression^{20,23} (H-score; Appendix Fig A2, online only). These observations are consistent with the observed differences in the efficacy of PD-1 blockade in

DLBCL and cHL and provide a possible explanation for why only a small proportion of patients with relapsed/refractory DLBCL benefit from nivolumab monotherapy. As a result of the low number of responders, there were insufficient data to accurately assess double-hit or c-Myc association with responses in this study.

Whereas the single-agent activity of PD-1 blockade was low in this unselected series of patients with DLBCL, it may be useful to evaluate the approach in select LBCL subtypes with increased PD-L1 expression, such as primary mediastinal B-cell lymphoma, EBV-positive LBCL, and T-cell/histiocyte-rich LBCL.¹⁰ Recent reports suggest that combination treatment strategies, such as chimeric antigen receptor T cells and nivolumab²⁸ and PD-1 inhibition with concomitant radiotherapy,²⁹ might also be considered.

In conclusion, the results of this phase II study indicate that nivolumab monotherapy has a low response rate but may provide benefit in a small number of patients with DLBCL who have experienced failure with auto-HCT or who are ineligible for auto-HCT and is associated with a favorable safety profile. Additional research on the influence of biologic factors, including the tumor microenvironment, is warranted to characterize whether a subset of patients with DLBCL may be more likely to respond to nivolumab-based treatment, if combination therapy can improve the therapeutic activity of PD-1 blockade, or if earlier use of these agents before intensive immunoablative chemotherapy may be more effective.

AFFILIATIONS

¹Mayo Clinic, Rochester, MN

²University Medical Center Utrecht Cancer Center, Utrecht, The Netherlands, on behalf of the Lunenburg Lymphoma Phase I/II Consortium–HOVON/LLPC

³University of Southampton, Southampton, United Kingdom

⁴University of California, Los Angeles, Los Angeles, CA

⁵Dana-Farber Cancer Institute, Boston, MA

⁶Brigham and Women's Hospital, Boston, MA

⁷Vanderbilt University, Nashville, TN

⁸Emory University, Atlanta, GA

⁹Jewish General Hospital, Montreal, Quebec, Canada

¹⁰National University Cancer Institute, Singapore, Singapore

¹¹Bristol-Myers Squibb, Princeton, NJ

¹²Austin Hospital and Olivia Newton-John Cancer Research Institute, Melbourne, VIC, Australia

CORRESPONDING AUTHOR

Stephen M. Ansell, MD, PhD, Mayo Clinic, 200 First St SW, Rochester, MN 55905; e-mail: ansell.stephen@mayo.edu.

SUPPORT

Supported by Bristol-Myers Squibb, which also funded medical writing support. Also supported by the Harold and Virginia Lash Foundation (P.A.); US National Institutes of Health Grant No. R01CA161026 and the Miller Fund (M.A.S.); the Center for Immuno-Oncology of the Dana-Farber Cancer Institute (S.J.R.); and the American Society of Hematology and Lymphoma Research Foundation (388017; J.B.C.).

AUTHORS' DISCLOSURES OF POTENTIAL CONFLICTS OF INTEREST AND DATA AVAILABILITY STATEMENT

Disclosures provided by the authors and data availability statement (if applicable) are available with this article at DOI <https://doi.org/10.1200/JCO.18.00766>.

AUTHOR CONTRIBUTIONS

Conception and design: Stephen M. Ansell, Peter Johnson, Philippe Armand, Margaret A. Shipp, Scott J. Rodig, Kazunobu Kato

Administrative support: Nishitha Reddy

Provision of study materials or patients: Monique C. Minnema, Peter Johnson, Nishitha Reddy, Jonathon B. Cohen, Sarit Assouline, Michelle Poon, Selda Samakoglu

Collection and assembly of data: Stephen M. Ansell, Monique C. Minnema, Peter Johnson, John M. Timmerman, Philippe Armand, Margaret A. Shipp, Scott J. Rodig, Margaretha G.M. Roemer, Nishitha Reddy, Jonathon B. Cohen, Sarit Assouline, Michelle Poon, Manish Sharma, Andrew Grigg

Data analysis and interpretation: Stephen M. Ansell, Monique C. Minnema, Peter Johnson, John M. Timmerman, Margaret A. Shipp, Scott J. Rodig, Azra H. Ligon, Margaretha G.M. Roemer, Nishitha Reddy, Sarit Assouline, Manish Sharma, Kazunobu Kato, Selda Samakoglu, Anne Sumbul, Andrew Grigg

Manuscript writing: All authors

Final approval of manuscript: All authors

Accountable for all aspects of the work: All authors

ACKNOWLEDGMENT

The authors thank all coinvestigators and the patients and families who participated in the trial. Writing assistance was provided by Janice Zhou,

PhD, at Caudex, under the direction of the authors, and was funded by Bristol-Myers Squibb.

REFERENCES

1. International Agency for Research on Cancer: World Cancer Report 2014: Diffuse Large B-cell Lymphoma. Geneva, Switzerland, WHO Press, 2014
2. Friedberg JW: Relapsed/refractory diffuse large B-cell lymphoma. *Hematology (Am Soc Hematol Educ Program)* 2011:498-505, 2011
3. Crump M, Neelapu SS, Farooq U, et al: Outcomes in refractory diffuse large B-cell lymphoma: Results from the international SCHOLAR-1 study. *Blood* 130:1800-1808, 2017
4. National Comprehensive Cancer Network: NCCN Clinical Practice Guidelines in Oncology: Non-Hodgkin's lymphomas, version 4.2014. <https://www.nccn.org/about/nhl.pdf>
5. Nagle SJ, Woo K, Schuster SJ, et al: Outcomes of patients with relapsed/refractory diffuse large B-cell lymphoma with progression of lymphoma after autologous stem cell transplantation in the rituximab era. *Am J Hematol* 88:890-894, 2013
6. Neelapu SS, Locke FL, Bartlett NL, et al: Axicabtagene ciloleucel CAR T-cell therapy in refractory large B-cell lymphoma. *N Engl J Med* 377:2531-2544, 2017
7. Freeman GJ, Long AJ, Iwai Y, et al: Engagement of the PD-1 immunoinhibitory receptor by a novel B7 family member leads to negative regulation of lymphocyte activation. *J Exp Med* 192:1027-1034, 2000
8. Kiyasu J, Miyoshi H, Hirata A, et al: Expression of programmed cell death ligand 1 is associated with poor overall survival in patients with diffuse large B-cell lymphoma. *Blood* 126:2193-2201, 2015
9. Green MR, Monti S, Rodig SJ, et al: Integrative analysis reveals selective 9p24.1 amplification, increased PD-1 ligand expression, and further induction via JAK2 in nodular sclerosing Hodgkin lymphoma and primary mediastinal large B-cell lymphoma. *Blood* 116:3268-3277, 2010
10. Chen BJ, Chapuy B, Ouyang J, et al: PD-L1 expression is characteristic of a subset of aggressive B-cell lymphomas and virus-associated malignancies. *Clin Cancer Res* 19:3462-3473, 2013
11. Roemer MG, Advani RH, Ligon AH, et al: PD-L1 and PD-L2 genetic alterations define classical Hodgkin lymphoma and predict outcome. *J Clin Oncol* 34:2690-2697, 2016
12. Lenz G, Wright GW, Emre NC, et al: Molecular subtypes of diffuse large B-cell lymphoma arise by distinct genetic pathways. *Proc Natl Acad Sci USA* 105:13520-13525, 2008
13. Andorsky DJ, Yamada RE, Said J, et al: Programmed death ligand 1 is expressed by non-Hodgkin lymphomas and inhibits the activity of tumor-associated T cells. *Clin Cancer Res* 17:4232-4244, 2011
14. Bristol-Myers Squibb: Nivolumab (Opdivo) prescribing information. https://packageinserts.bms.com/pi/pi_opdivo.pdf
15. Weber JS, D'Angelo SP, Minor D, et al: Nivolumab versus chemotherapy in patients with advanced melanoma who progressed after anti-CTLA-4 treatment (CheckMate 037): A randomised, controlled, open-label, phase 3 trial. *Lancet Oncol* 16:375-384, 2015
16. Borghaei H, Paz-Ares L, Horn L, et al: Nivolumab versus docetaxel in advanced nonsquamous non-small-cell lung cancer. *N Engl J Med* 373:1627-1639, 2015
17. Motzer RJ, Escudier B, McDermott DF, et al: Nivolumab versus everolimus in advanced renal-cell carcinoma. *N Engl J Med* 373:1803-1813, 2015
18. Sharma P, Callahan MK, Bono P, et al: Nivolumab monotherapy in recurrent metastatic urothelial carcinoma (CheckMate 032): A multicentre, open-label, two-stage, multi-arm, phase 1/2 trial. *Lancet Oncol* 17:1590-1598, 2016
19. Ansell SM, Lesokhin AM, Borrello I, et al: PD-1 blockade with nivolumab in relapsed or refractory Hodgkin's lymphoma. *N Engl J Med* 372:311-319, 2015
20. Younes A, Santoro A, Shipp M, et al: Nivolumab for classical Hodgkin's lymphoma after failure of both autologous stem-cell transplantation and brentuximab vedotin: A multicentre, multicohort, single-arm phase 2 trial. *Lancet Oncol* 17:1283-1294, 2016
21. Lesokhin AM, Ansell SM, Armand P, et al: Nivolumab in patients with relapsed or refractory hematologic malignancy: Preliminary results of a phase 1b study. *J Clin Oncol* 34:2698-2704, 2016
22. Cheson BD, Pfistner B, Juweid ME, et al: Revised response criteria for malignant lymphoma. *J Clin Oncol* 25:579-586, 2007
23. Roemer MGM, Redd RA, Cader FZ, et al: Major histocompatibility complex class II and programmed death ligand 1 expression predict outcome after programmed death 1 blockade in classic Hodgkin lymphoma. *J Clin Oncol* 36:942-950, 2018
24. Mahoney KM, Sun H, Liao X, et al: PD-L1 antibodies to its cytoplasmic domain most clearly delineate cell membranes in immunohistochemical staining of tumor cells. *Cancer Immunol Res* 3:1308-1315, 2015
25. Xu-Monette ZY, Thompson D, LaFleur B, et al: Diffuse large B-cell lymphoma cell of origin determination from formalin-fixed paraffin-embedded tissues. *Blood* 126:5052, 2015 (suppl)
26. US National Cancer Institute: Common Terminology Criteria for Adverse Events (CTCAE), version 4.0. Bethesda, MD, National Institutes of Health, 2009
27. Topalian SL, Hodi FS, Brahmer JR, et al: Safety, activity, and immune correlates of anti-PD-1 antibody in cancer. *N Engl J Med* 366:2443-2454, 2012
28. Hill BT, Roberts ZJ, Rossi JM, et al: Marked re-expansion of chimeric antigen receptor (CAR) T cells and tumor regression following nivolumab treatment in a patient treated with axicabtagene ciloleucel (axi-cel; KTE-C19) for refractory diffuse large B cell lymphoma (DLBCL). *Blood* 130:2825, 2017 (suppl)
29. Wight JC, Hawkes EA, Berlangieri SU, et al: An abscopal effect may augment PD-1 inhibition in refractory classical Hodgkin lymphoma. *Leuk Lymphoma* [10.1080/10428194.2018.1452217](https://doi.org/10.1080/10428194.2018.1452217) [epub ahead of print on March 23, 2018]



AUTHORS' DISCLOSURES OF POTENTIAL CONFLICTS OF INTEREST**Nivolumab for Relapsed/Refractory Diffuse Large B-Cell Lymphoma in Patients Ineligible for or Having Failed Autologous Transplantation: A Single-Arm, Phase II Study**

The following represents disclosure information provided by authors of this manuscript. All relationships are considered compensated. Relationships are self-held unless noted. I = Immediate Family Member, Inst = My Institution. Relationships may not relate to the subject matter of this manuscript. For more information about ASCO's conflict of interest policy, please refer to www.asco.org/rwc or ascopubs.org/jco/site/ffc.

Stephen M. Ansell

Honoraria: WebMD, Research to Practice

Research Funding: Bristol-Myers Squibb (Inst), Seattle Genetics (Inst), Affimed Therapeutics (Inst), Regeneron (Inst), Pfizer (Inst), LAM Therapeutics (Inst), Trillium Therapeutics (Inst)

Monique C. Minnema

Consulting or Advisory Role: Servier, Takeda, Amgen, Janssen-Cilag

Research Funding: Celgene

Travel, Accommodations, Expenses: Roche, Amgen

Peter Johnson

Honoraria: Takeda, Bristol-Myers Squibb, Novartis, Celgene, Kite Pharma, Genmab, Incyte

Consulting or Advisory Role: Janssen Pharmaceuticals, Epizyme, Boehringer Ingelheim

Research Funding: Epizyme (Inst), Janssen Pharmaceuticals (Inst)

Patents, Royalties, Other Intellectual Property: Combined use of Fc gamma RIIB (CD32b) and CD20, specific antibodies, WO Patent, PCT/GB2011/051572; EU11760819.0

Travel, Accommodations, Expenses: Zenyaku Kogyo

John M. Timmerman

Stock and Other Ownership Interests: Biomarín

Honoraria: Kite Pharma, Gilead Sciences

Consulting or Advisory Role: Celgene, Seattle Genetics, Genmab

Research Funding: Bristol-Myers Squibb, ImmunGene, Kite Pharma, Merck

Travel, Accommodations, Expenses: Bristol-Myers Squibb

Phillippe Armand

Consulting or Advisory Role: Bristol-Myers Squibb, Merck, Pfizer, Affimed Therapeutics, Adaptive Biotechnologies

Research Funding: Bristol-Myers Squibb (Inst), Merck (Inst), Affimed Therapeutics (Inst), Adaptive Biotechnologies (Inst), Genentech (Inst), Tensha Therapeutics (Inst), Otsuka (Inst), Sigma-Tau (Inst)

Travel, Accommodations, Expenses: Genmab

Margaret A. Shipp

Honoraria: Bristol-Myers Squibb, AstraZeneca

Consulting or Advisory Role: Bristol-Myers Squibb

Research Funding: Bristol-Myers Squibb (Inst), Bayer (Inst), Merck (Inst)

Scott J. Rodig

Honoraria: PerkinElmer, Bristol-Myers Squibb

Consulting or Advisory Role: AstraZeneca, PerkinElmer

Research Funding: Bristol-Myers Squibb, Merck, Affimed Therapeutics, Kite Pharma

Patents, Royalties, Other Intellectual Property: Patent pending for use of anti-galectin1 antibodies for diagnostic use

Travel, Accommodations, Expenses: Roche, Bristol-Myers Squibb

Azra H. Ligon

Leadership: Travera (I)

Stock and Other Ownership Interests: Travera (I)

Consulting or Advisory Role: Travera (I)

Nishitha Reddy

Consulting or Advisory Role: Celgene, AbbVie, Bristol-Myers Squibb, Adaptive Biotechnologies

Speakers' Bureau: Gilead Sciences

Research Funding: Bristol-Myers Squibb (Inst)

Jonathon B. Cohen

Consulting or Advisory Role: Pharmacyclics, Celgene, Millennium

Pharmaceuticals, Seattle Genetics, Novartis, Infinity Pharmaceuticals, AbbVie

Research Funding: Bristol-Myers Squibb, Janssen Pharmaceuticals, Novartis, Takeda

Sarit Assouline

Stock and Other Ownership Interests: Knight Pharmaceuticals

Honoraria: Janssen Oncology, Pfizer

Speakers' Bureau: Pfizer, Janssen Oncology, Bristol-Myers Squibb

Research Funding: Roche Canada, Takeda, Epizyme, Gilead Sciences, Astex Pharmaceuticals, Janssen Pharmaceuticals

Travel, Accommodations, Expenses: Roche Canada

Michelle Poon

Consulting or Advisory Role: Takeda, Janssen Pharmaceuticals, AbbVie

Travel, Accommodations, Expenses: Roche

Manish Sharma

Employment: Bristol-Myers Squibb

Stock and Other Ownership Interests: Bristol-Myers Squibb

Kazunobu Kato

Employment: Bristol-Myers Squibb, Celgene

Stock and Other Ownership Interests: Bristol-Myers Squibb, Celgene

Anne Sumbul

Employment: Bristol-Myers Squibb

Andrew Grigg

Honoraria: Gilead Sciences, MSD Oncology

Consulting or Advisory Role: Gilead Sciences, Janssen-Cilag, Roche, Takeda, MSD Oncology

Expert Testimony: MSD Oncology

Travel, Accommodations, Expenses: Roche

No other potential conflicts of interest were reported

APPENDIX




FIG A1. Flow diagram.




FIG A2. Comparison of the incidence and magnitude of 9p24.1 alterations in diffuse large B-cell lymphoma (DLBCL; this study) and classic Hodgkin lymphoma (cHL).²³ (A) Prevalence and (B) spectrum of 9p24.1 alterations in DLBCL. (C) Percentage of tumor cells with residual 9p24.1 disomy in DLBCLs classified by 9p24.1 genetic categories. (D) Prevalence and (E) spectrum of 9p24.1 alterations in cHL.²³ (F) Percentage of tumor cells with residual 9p24.1 disomy in cHLs classified by 9p24.1 genetic categories.²³ As noted, DLBCLs have a much lower incidence of 9p24.1 copy gain and amplification than cHLs. In addition, DLBCLs with low-level 9p24.1 alterations have a higher percentage of residual disomy than cHLs. Ampl, amplification; Coampl, coamplification; PD-L2, programmed death ligand 2; Rearr, rearrangement. [Reprinted with permission. © 2018 American Society of Clinical Oncology.]

TABLE A1. Biomarker Data for Patients With Clinical Response (CR and PR)

Cohort and BOR	PFS, Months	DOR, Months	9p24.1 Alteration			% PD-L1 Positive		COO Status	EBV Status
			Alteration	% Cells	Copies, No.	Malignant Cells	Nonmalignant Cells		
HCT failed									
CR	19	17.1	NA			NA	NA	GCB	NA
CR	15.7*	13.8*	Coamplification	96	6 - >15	NA	NA	Non GCB	Negative
CR	13.1*	11.3*	Disomy			0	10 (cytoplasmic)	NA	Negative
PR	13.1	11.4	Polysomy	58	3-4	0	40 (membrane)	GCB	Negative
PR	8.3	6.6	Polysomy	6	3	NA	NA	Non GCB	Negative
PR	11.8	6.3	Polysomy	16	3	NA	NA	GCB	NA
PR	5.6	3.7	Cogain	28	3-4	0	40 (cytoplasmic/membrane)	Non GCB	Negative
PR	4.4	2.5	NA/disomy†			0	10 (cytoplasmic)/30 (membrane)†	GCB	Negative
PR	10*	2.4*	NA			0	50 (cytoplasmic/membrane)/50 (membrane)†	GCB	Negative
HCT ineligible									
PR	10.2	8.3	NA			0	70 (cytoplasmic)	Non GCB	Negative

Abbreviations: BOR, best of response; COO, cell of origin; CR, complete remission; DOR, duration of response; EBV, Epstein-Barr virus; GCB, germinal center B cell; HCT, hematopoietic cell transplantation; NA, not available; PD-L1, programmed death ligand 1; PFS, progression-free survival; PR, partial remission.

*Ongoing response.

†Two samples from the same patient were analyzed.