

Biomarker evidence for nitrogen-fixing cyanobacterial blooms in a brackish surface layer in the Nile River plume during sapropel deposition

Nicole J. Bale^{1*}, Rick Hennekam², Ellen C. Hopmans¹, Denise Dorhout¹, Gert-Jan Reichart^{2,3}, Marcel van der Meer¹, Tracy A. Villareal⁴, Jaap S. Sinninghe Damsté^{1,3} and Stefan Schouten^{1,3}

¹Department of Marine Microbiology and Biogeochemistry, NIOZ Royal Institute for Sea Research and Utrecht University, 1790 AB Den Burg, Netherlands

²Department of Ocean Systems, NIOZ Royal Institute for Sea Research and Utrecht University, 1790 AB Den Burg, Netherlands

³Department of Earth Sciences, Faculty of Geosciences, Utrecht University, P.O. Box 80.121, 3508 TA Utrecht, Netherlands

⁴Marine Science Institute, The University of Texas at Austin, Port Aransas, Texas 78373, USA

ABSTRACT

Sapropels are organic-rich sediment layers deposited in the eastern Mediterranean Sea during precession minima, resulting from an increase in export productivity and/or preservation. Increased freshwater delivery from the African continent resulted in stratification, causing deep-water anoxia, while nutrient input stimulated productivity, presumably at the deep chlorophyll maximum. Previous studies have suggested that during sapropel deposition, nitrogen fixation was widespread in the highly stratified surface waters, and that cyanobacteria symbiotic with diatoms (diatom-diazotroph associations, DDAs) were responsible. Here we analyzed sapropel S5 sediments for heterocyst glycolipids (HGs) from three locations in the eastern Mediterranean. HG biomarkers can differentiate between those heterocystous cyanobacteria that are free living (found predominately in freshwater or brackish environments) and those that are from DDAs (found in marine settings). In our primary core, from a location which would have been influenced by the Nile River outflow, we detected a HG with a pentose (C₅) head group specific for DDAs. However, HGs with a hexose (C₆) head group, specific to free-living cyanobacteria, were present in substantially (up to 60×) higher concentration. These data suggest that at our study location, free-living cyanobacteria were the dominant diazotrophs, rather than DDAs. The C₆ HGs increased substantially at the onset of sapropel S5 deposition, suggesting that substantial seasonal cyanobacterial blooms were associated with a brackish surface layer flowing from the Nile into the eastern Mediterranean. Two additional S5 sapropels were analyzed, one also from the Nile delta region and one from the region between Libya and southwestern Crete. Overall, comparison of the HG distribution in the three S5 sapropels provides evidence that all three locations were initially influenced by surface salinities that were sufficiently low to support free-living heterocystous cyanobacteria. While free-living heterocystous cyanobacteria continued to outnumber DDAs during sapropel deposition at the two Nile-influenced sites, DDAs, indicators of persistent marine salinities, were the dominant diazotrophs in the upper part of the sapropel at the more westerly site. These results indicate that N₂ fixation by free-living cyanobacteria offers an important additional mechanism to stimulate productivity in regions with strong river discharge during sapropel deposition.

INTRODUCTION

The dark-colored, organic-rich sapropels found in the eastern Mediterranean Sea are associated with a stratification-linked reduction in ventilation resulting from freshwater outflow

during precession minima, concurrent with increased organic-matter export from the photic zone. The increase in productivity is, for sapropel S5 and probably most other sapropels, linked to a highly productive deep chlorophyll maximum (DCM) (Rohling et al., 2015). Increased organic-matter flux into the poorly ventilated

bottom waters and subsequent respiration resulted in anoxia development and sapropel formation (Rossignol-Strick et al., 1982; Rossignol-Strick, 1985; Rohling et al., 2015). Isotope, biomarker, and microfossil studies have suggested that the nutrient-sparse surface waters would have been favorable for cyanobacterial nitrogen fixation and, in particular, diatom-cyanobacteria symbioses (Kemp et al., 1999; Sachs and Repeta, 1999; Bauersachs et al., 2010) termed diatom-diazotroph associations (DDAs).

Heterocyst glycolipids (HGs) are biomarkers for cyanobacteria that use heterocysts to carry out N₂ fixation (Nichols and Wood, 1968; Gambacorta et al., 1995; Bauersachs et al., 2009a). The sugar moiety of HGs found in free-living heterocystous cyanobacteria is typically hexose (hereafter C₆), while the HGs associated with endosymbiotic heterocystous cyanobacteria in DDAs contain a pentose moiety (hereafter C₅) (Schouten et al., 2013; Bale et al., 2015). C₆ HGs have been applied as specific paleo-biomarkers for the presence of N₂-fixing cyanobacteria in marine and lacustrine geological records (Bauersachs et al., 2010; Sollai et al., 2017). Previously, Bauersachs et al. (2010) found that C₆ HGs were abundant in the S5 sapropel from a Nile-influenced location in the eastern Mediterranean Sea and suggested that they were likely derived from DDAs. However, the subsequent discovery that all DDAs studied to date produce C₅ HGs (Schouten et al., 2013; Bale et al., 2015), not C₆ HGs, casts uncertainty on this explanation. Furthermore, the analytical method of Bauersachs et al. (2010) did not cover the detection of C₅ HGs. Here we comprehensively analyzed

*E-mail: nicole.bale@nioz.nl

the C₅ and C₆ HGs in a sediment core from the eastern Mediterranean Sea containing a well-preserved sapropel S5 (ca. 128.3–121.5 ka; Grant et al., 2016), as well as two other S5 sapropels also located in the eastern Mediterranean Sea, in order to unravel the primary mechanism for N₂ fixation by cyanobacteria during this interval.

METHODS

A piston core and suspended particulate matter (SPM) were obtained during Netherlands Earth System Science Centre (NESSC) research cruise 64PE406 onboard the R/V *Pelagia* in the eastern Mediterranean in January 2016. A 920.5-cm-long piston core was collected at cruise station 1 (core 64PE406-E1, 33°18.14898'N, 33°23.71998'E, water depth 1760 m; larger red filled circle in Fig. 1) in the southeastern Levantine Sea. McLane *in situ* pumps (McLane Laboratories Inc., Falmouth, UK) were used to collect SPM from the water column for lipid analysis (see the GSA Data Repository¹ for details). Two other S5 sapropels from core MS66PC (2004 MIMES [Multiscale Investigations of Eastern Mediterranean Seep Systems] cruise; Bauersachs et al., 2010) and Ocean Drilling Program (ODP) core 160-971C-2H-3 (Pearce et al., 1998; Kemp et al., 1999) were also analyzed for their HG lipid content. For details of sample extraction and lipid analysis, see the Data Repository. The core chronology of the sapropel S5 interval (for details, see the

Data Repository) was obtained by correlating the Ba variability to that of published sapropel boundaries (Grant et al., 2016) and, for intervals prior to S5, to other Ba excursions (Ziegler et al., 2010). Sedimentary bulk elemental composition measurements by X-ray fluorescence (XRF) core scanning were performed with trace-metal settings (Hennekam et al., 2019); for details, see the Data Repository.

RESULTS AND DISCUSSION

We examined the HG distribution in 30 × 1 cm sediment slices from a depth interval including sapropel S5 (as evidenced from total organic carbon [TOC] and XRF elemental ratios; Fig. 2). We detected (1) a single C₅ HG, 1-(O-ribose)-3,27,29-triacontanetriol (C₅ HG₃₀ triol); and (2) three C₆ HGs, 1-(O-hexose)-3,25-hexacosanediol (C₆ HG₂₆ diol), 1-(O-hexose)-3,27-octacosanediol (C₆ HG₂₈ diol), and 1-(O-hexose)-3,27,29-tricontanetriol (C₆ HG₃₀ triol) (see Fig. 2 for structures). We found a notably higher concentration of C₆ HGs (maxima between 20 and 31 μg [g TOC]⁻¹) relative to the C₅ HG (maximum 1.8 μg [g TOC]⁻¹). The concentrations of all HGs (normalized to TOC) were higher in the sapropel (Figs. 2B–2E).

The occurrence of C₅ HG₃₀ triol prior to and during the onset of S5 (Fig. 2B) suggests that cyanobacteria belonging to DDAs, specifically *Richelia intracellularis* (Schouten et al., 2013; Bale et al., 2015, 2018), were present in the euphotic zone at our study site before the deposition of the S5 sapropel, and then either they increased in number or their HG was more efficiently preserved during S5 deposition. The presence of C₅ HG₃₀ triol agrees with the high numbers of the DDA host diatoms, in particular *Hemiaulus hauckii*, that have been reported in sapropel S5 (Kemp et al., 1999) at ODP site 971 south of Crete. Kemp et al. (1999) concluded that increased freshwater-induced stratification

of surface waters, although still highly marine, led to conditions that were ideally suited for DDAs. In the modern-day eastern Mediterranean, DDAs containing *R. intracellularis* have been detected in low levels throughout the year with a maximum in autumn (Zeev et al., 2008). We also detected C₅ HG₃₀ triol in modern-day SPM collected in the winter at the site where piston core 64PE406-E1 was recovered (Table DR1 in the Data Repository).

A quite different trend was observed for the C₆ HGs than for the C₅ HG: all three C₆ HGs increased substantially at the onset of S5 deposition (Figs. 2C–2E). The three C₆ HGs detected have been found in varying proportions both in cultures of the order Nostocales (Bauersachs et al., 2009a; Wörmer et al., 2012) and in freshwater and brackish environments (Bauersachs et al., 2009b, 2010, 2011, 2013; Wörmer et al., 2012; Bühring et al., 2014; Bale et al., 2015, 2016). Based on comparison of the sapropel S5 C₆ HG distribution with literature data (cf. Table DR1), we suggest that during sapropel deposition there was an initial peak in the genus *Nodularia* (C₆ HG₂₆ diol; Fig. 2C), followed by dominance in the genus *Aphanizomenon* (C₆ HG₂₈ diol and C₆ HG₃₀ triol; Figs. 2D and 2E). To date, the C₆ HGs detected in this study have not been detected in any DDAs (see the Data Repository for further discussion).

The presence of free-living Nostocales in eastern Mediterranean sapropel layers is unexpected, as these freshwater and/or brackish cyanobacteria have not been reported in the modern eastern Mediterranean, where the surface salinity is >35 PSU and no C₆ HGs were detected in the recovered SPM (Table DR1). It is not likely that the HGs detected were delivered to the sediment by river as the contributions of terrestrial biomarker lipids into sapropels are very low (Bosch et al., 1998; Menzel et al., 2003). Moreover, the Ti/Al ratio, indicative of the relative contribution of aeolian (Ti) versus Nile-derived fluvial (Al) sediment input (Lourens et al., 2001), shows the highest Nile discharge well after the peak in C₆ HGs (Fig. 2I). Hence, we find no support for the hypothesis that C₆ HGs were produced on land, in lakes or rivers, and delivered to the sediment record via rivers. Reports of free-living heterocystous cyanobacteria in present-day marine settings are very limited: planktonic *Nostoc* has been described in the Indian Ocean and the Red and Mediterranean Seas (Taylor, 1966), and *Anabaena gerdii* has been reported to occur in the western Pacific Ocean, Arabian Sea, and Arafura Sea (Carpenter and Janson, 2001). Tuo et al. (2017) reported free-living *Richelia* in the western North Pacific. In contrast, blooms of free-living cyanobacteria are common in brackish environments. For example, there are extensive summer blooms of the free-living Nostocales species *Nodularia*, *Aphanizomenon*, and *Dolichospermum* in the

¹GSA Data Repository item 2019375, details of experimental methods used, additional discussion of diatom-diazotroph associations, Table DR1 (heterocyst glycolipid compositions cited in the text and data from literature), Table DR2 relative abundance of C5 heterocyst glycolipids and C6 heterocyst glycolipids as cited in the literature), and Figure DR1 (records used for age model construction), is available online at <http://www.geosociety.org/datarepository/2019/>, or on request from editing@geosociety.org.




Figure 1. Map of area where core 64PE406-E1 was recovered, eastern Mediterranean Sea. Recovery position of other cores referenced in text are also indicated: Ocean Drilling Program (ODP) Site 160-971. (Kemp et al., 1999; Sachs and Repeta, 1999) and core site MS66PC (Bauersachs et al., 2010). Stations for suspended particulate matter collection are marked with red circles (including 64PE406-E1 recovery station). Present-day location of river Nile is shown in blue, while approximation of extant river-wadi system of Libyan Basin is shown in green (adapted from Rohling et al., 2002).




Figure 2. Depth profile of eastern Mediterranean piston core 64PE406-E1 (33°18.14898'N, 33°23.71998'E, water depth 1760 m; NESSC research cruise 64PE406 onboard the R/V *Pelagia*). Horizontal gray bar shows deposition of sapropel. **A–E:** Organic analyses of total organic carbon (TOC) (**A**) and concentration ($\mu\text{g [g TOC]}^{-1}$) of heterocyst glycolipids (HGs) C_5 HG₃₀ triol (**B**), C_6 HG₂₆ diol (**C**), C_6 HG₂₈ diol (**D**), and C_6 HG₃₀ triol (**E**). Structures of HGs are shown on respective graphs. Stereochemistry of C_5 HG₃₀ triol is as determined by Schouten et al. (2013). **F–I:** Multivariate log-ratio calibrated X-ray fluorescence scan data in concentration (ppm, black lines) and as ratios (gray lines) for Ba and Ba/Al (**F**), Mo and Mo/Al (**G**), U and U/Al (**H**), and Ti/Al (**I**). Concentrations derived by inductively coupled plasma–mass spectrometry data are overlain (gray circles).

upper 20 m of the Baltic Sea (Hajdu et al., 2007; Ploug, 2008; Celepli et al., 2017), where surface salinity is on average 6–8 PSU (Gustafsson and Westman, 2002). Hence, we propose that the abundant C_6 HGs in the S5 sapropel at this location indicate the presence of free-living heterocystous cyanobacteria in a brackish surface layer, likely resulting from the freshwater Nile river plume in the eastern Mediterranean at the onset of S5 deposition.

The presence of a surface brackish layer is seemingly inconsistent with the simultaneous presence of DDAs, albeit their HGs occur in much lower concentrations (Fig. 2B). DDAs would also be expected to inhabit the uppermost 40 m (as seen in the analogous Amazon River plume; Foster et al., 2007). DDAs are unimportant at salinities <30 PSU (Foster et al., 2007, 2009; Subramaniam et al., 2008; Bombar et al., 2011). We postulate that the presence of low levels of the C_5 HG alongside the dominant C_6 HGs is due to variation in monsoonal river flow and/or plume location causing seasonal and/or yearly oscillations between marine and brackish water

at the surface and thus an alternation between Nostocales and DDA-dominated cyanobacterial populations. Furthermore, in the modern-day Baltic Sea, cyanobacterial blooms are not observed as a pervasive cover, but in surface patches (Stal et al., 2003; Ploug, 2008). This can explain how, despite the cyanobacterial bloom, sufficient light penetrated the surface to support the high level of productivity that occurred at the DCM during sapropel formation (Rohling and Gieskes, 1989). Our results indicate that the low-salinity surface cyanobacterial blooms started to establish at the onset of suboxic to euxinic conditions (indicated by U and Mo concentrations and Mo/Al and U/Al ratios [Figs. 2G and 2H]; Tribouillard et al., 2006) and were occurring periodically for ~4.5 k.y. The occurrence of these blooms peaked and then diminished well before the end of the sapropel deposition (Figs. 2C–2E), possibly due to a change in the nutrient availability in the surface waters or a reduced input of freshwater, leading to a system which did not support N_2 fixation. Indeed, the TOC of the sediments was still increasing toward a

maximum (of ~9% at 357–339 cm depth; ca. 126–122 kyr B.P.; Fig. 2A), in tune with increasing export of productivity (high Ba/Al ratios; Fig. 2F) from a potential DCM, while the concentration of all HGs was already decreasing (Figs. 2C–2E).

The presence of a lowered-salinity surface layer in the eastern Mediterranean due to enhanced freshwater influx is well established (Rossignol-Strick et al., 1982; Rossignol-Strick, 1985; Rohling et al., 2004; Rohling, 2007; van der Meer et al., 2007; Rodríguez-Sanz et al., 2017), although the extent and intensity of this surface freshening is uncertain. In order to constrain the geographical spread of the bloom, we also examined the HG distribution in the S5 sapropels recovered in cores MS66PC (Bauer-sachs et al., 2010) and ODP 160-971C-2H-3 (Pearce et al., 1998; Kemp et al., 1999) (see Fig. 1). Cores 64PE406-E1 and MS66PC are both from the Nile delta region, and hence their sites were likely influenced by freshwater delivery from the Nile. Conversely, the ODP 160-971 core site is in the region between Libya

and southwest Crete, which currently is out of reach of any modern-day river systems but has been shown to have been influenced by monsoon flooding during the previous interglacial maximum, thought to have been delivered via extant river systems of northern Libya (Rohling et al., 2002). These additional locations also revealed a high C_6 HG concentration in the lower part of the sapropel (Table DR1), consistent with the expected high input of freshwater at this time. At the site closest to the mouth of the Nile (MS66PC), the relative abundance of C_6 HG was high ($97\% \pm 3\%$ of total HGs), as it was at our site (64PE406-E1) slightly further from the Nile plume ($97\% \pm 2\%$). The ODP 160-971C-2H-3 core had a lower relative abundance of C_6 HG ($78\% \pm 10\%$) consistent with a lower freshwater input. The upper half of the sapropel was also characterized by a moderate or high relative abundance of C_6 HGs in the MS66PC and 64PE406-E1 cores (on average 54% and 87%, respectively). Conversely, the upper part of S5 in the ODP 160-971C-2H-3 core contained a high percent of C_5 HG (96%). Interestingly, the percent of the DDA-forming diatom *H. hauckii* has also been reported to be significantly higher in the upper half of S5 in the ODP 160-971C-2H-3 core (Pearce et al., 1998; Kemp et al., 1999), further confirming the C_5 HG₃₀ triol is a marker for DDAs.

Overall, comparison of the HG distribution in the three S5 sapropels provides evidence that all three locations were initially influenced by surface salinities that were sufficiently low to support free-living heterocystous cyanobacteria. While free-living heterocystous cyanobacteria continued to outnumber DDAs during sapropel deposition at the two Nile-influenced sites, DDAs, indicators of persistent marine salinities, were the dominant diazotrophs in the upper part of the sapropel at the ODP 160-971 site.

ACKNOWLEDGMENTS

We thank the captain and crew of the R/V *Pelagia* for their support during cruise 64PE406, J.P. Dirksen for assistance with sample collection and processing, and P. Laan and W. Boer for the inductively coupled plasma-mass spectrometry measurements. We thank C. Yildiz, T. Bauersachs, and Z. Erdem for assistance acquiring samples, sample extraction, and total organic carbon analysis. We also thank three anonymous reviewers who greatly improved a previous version of this manuscript. This research used samples provided by the Ocean Drilling Program (ODP). ODP is sponsored by the U.S. National Science Foundation and participating countries under management of Joint Oceanographic Institutions, Inc. The work of Bale is supported by the Netherlands Organisation for Scientific Research (NWO) through grant 822.01.017 awarded to Schouten, and funding from the European Research Council under the European Union's Horizon 2020 research and innovation program, grant agreement 694569 awarded to Sinninghe Damsté. Hennekam is supported through NWO grant ALW OP.2015.113 awarded to Reichart. Fieldwork was carried out under the program of the

Netherlands Earth System Science Centre, financially supported by the Ministry of Education, Culture and Science, grant 024.002.001.

REFERENCES CITED

- Bale, N.J., Hopmans, E.C., Zell, C., Sobrinho, R.L., Kim, J.-H., Sinninghe Damsté, J.S., Villareal, T.A., and Schouten, S., 2015, Long chain glycolipids with pentose head groups as biomarkers for marine endosymbiotic heterocystous cyanobacteria: *Organic Geochemistry*, v. 81, p. 1–7, <https://doi.org/10.1016/j.orgchem.2015.01.004>.
- Bale, N.J., Hopmans, E.C., Schoon, P.L., de Kluijver, A., Downing, J.A., Middelburg, J.J., Sinninghe Damsté, J.S., and Schouten, S., 2016, Impact of trophic state on the distribution of intact polar lipids in surface waters of lakes: *Limnology and Oceanography*, v. 61, p. 1065–1077, <https://doi.org/10.1002/lno.10274>.
- Bale, N.J., Villareal, T.A., Hopmans, E.C., Brussaard, C.P.D., Besseling, M., Dorhout, D., Sinninghe Damsté, J.S., and Schouten, S., 2018, C_5 glycolipids of heterocystous cyanobacteria track symbiont abundance in the diatom *Hemiaulus hauckii* across the tropical North Atlantic: *Biogeosciences*, v. 15, p. 1229–1241, <https://doi.org/10.5194/bg-15-1229-2018>.
- Bauersachs, T., Compaoré, J., Hopmans, E.C., Stal, L.J., Schouten, S., and Sinninghe Damsté, J.S., 2009a, Distribution of heterocyst glycolipids in cyanobacteria: *Phytochemistry*, v. 70, p. 2034–2039, <https://doi.org/10.1016/j.phytochem.2009.08.014>.
- Bauersachs, T., Hopmans, E.C., Compaoré, J., Stal, L.J., Schouten, S., and Sinninghe Damsté, J.S., 2009b, Rapid analysis of long-chain glycolipids in heterocystous cyanobacteria using high-performance liquid chromatography coupled to electrospray ionization tandem mass spectrometry: *Rapid Communications in Mass Spectrometry*, v. 23, p. 1387–1394, <https://doi.org/10.1002/rcm.4009>.
- Bauersachs, T., Speelman, E.N., Hopmans, E.C., Reichart, G.-J., Schouten, S., and Sinninghe Damsté, J.S., 2010, Fossilized glycolipids reveal past oceanic N_2 fixation by heterocystous cyanobacteria: *Proceedings of the National Academy of Sciences of the United States of America*, v. 107, p. 19,190–19,194, <https://doi.org/10.1073/pnas.1007526107>.
- Bauersachs, T., Compaoré, J., Severin, I., Hopmans, E.C., Schouten, S., Stal, L.J., and Sinninghe Damsté, J.S., 2011, Diazotrophic microbial community of coastal microbial mats of the southern North Sea: *Geobiology*, v. 9, p. 349–359, <https://doi.org/10.1111/j.1472-4669.2011.00280.x>.
- Bauersachs, T., Miller, S.R., van der Meer, M.T.J., Hopmans, E.C., Schouten, S., and Sinninghe Damsté, J.S., 2013, Distribution of long chain heterocyst glycolipids in cultures of the thermophilic cyanobacterium *Mastigoeladus laminosus* and a hot spring microbial mat: *Organic Geochemistry*, v. 56, p. 19–24, <https://doi.org/10.1016/j.orgchem.2012.11.013>.
- Bombar, D., Moisaner, P.H., Dippner, J.W., Foster, R.A., Voss, M., Karfeld, B., and Zehr, J.P., 2011, Distribution of diazotrophic microorganisms and *nifH* gene expression in the Mekong River plume during intermonsoon: *Marine Ecology Progress Series*, v. 424, p. 39–52, <https://doi.org/10.3354/meps08976>.
- Bosch, H.-J., Sinninghe Damsté, J.S., and de Leeuw, J.W., 1998, Molecular palaeontology of eastern Mediterranean sapropels: Evidence for photic zone euxinia, in Robertson, A.H.F., et al., eds., *Proceedings of the Ocean Drilling Program, Scientific Results, Volume 160*: College Station, Texas, Ocean Drilling Program, p. 285–295, <https://doi.org/10.2973/odp.proc.sr.160.066.1998>.
- Bühning, S.I., Kamp, A., Wörmer, L., Ho, S., and Hinrichs, K.-U., 2014, Functional structure of laminated microbial sediments from a supratidal sandy beach of the German Wadden Sea (St. Peter-Ording): *Journal of Sea Research*, v. 85, p. 463–473, <https://doi.org/10.1016/j.seares.2013.08.001>.
- Carpenter, E.J., and Janson, S., 2001, *Anabaena gerdii* sp. nov., a new planktonic filamentous cyanobacterium from the South Pacific Ocean and Arabian Sea: *Phycologia*, v. 40, p. 105–110.
- Celepli, N., Sundh, J., Ekman, M., Dupont, C.L., Yooseph, S., Bergman, B., and Ininbergs, K., 2017, Meta-omic analyses of Baltic Sea cyanobacteria: Diversity, community structure and salt acclimation: *Environmental Microbiology*, v. 19, p. 673–686, <https://doi.org/10.1111/1462-2920.13592>.
- Foster, R.A., Subramaniam, A., Mahaffey, C., Carpenter, E.J., Capone, D.G., and Zehr, J.P., 2007, Influence of the Amazon River plume on distributions of free-living and symbiotic cyanobacteria in the western tropical north Atlantic Ocean: *Limnology and Oceanography*, v. 52, p. 517–532, <https://doi.org/10.4319/lo.2007.52.2.0517>.
- Foster, R.A., Subramaniam, A., and Zehr, J.P., 2009, Distribution and activity of diazotrophs in the Eastern Equatorial Atlantic: *Environmental Microbiology*, v. 11, p. 741–750, <https://doi.org/10.1111/j.1462-2920.2008.01796.x>.
- Gambacorta, A., Soriente, A., Trincone, A., and Sodano, G., 1995, Biosynthesis of the heterocyst glycolipids in the cyanobacterium *Anabaena cylindrica*: *Phytochemistry*, v. 39, p. 771–774, [https://doi.org/10.1016/0031-9422\(95\)00007-T](https://doi.org/10.1016/0031-9422(95)00007-T).
- Grant, K.M., Grimm, R., Mikolajewicz, U., Marino, G., Ziegler, M., and Rohling, E.J., 2016, The timing of Mediterranean sapropel deposition relative to insolation, sea-level and African monsoon changes: *Quaternary Science Reviews*, v. 140, p. 125–141, <https://doi.org/10.1016/j.quascirev.2016.03.026>.
- Gustafsson, B.G., and Westman, P., 2002, On the causes for salinity variations in the Baltic Sea during the last 8500 years: *Paleoceanography*, v. 17, p. 12–12–14, <https://doi.org/10.1029/2000PA000572>.
- Hajdu, S., Höglander, H., and Larsson, U., 2007, Phytoplankton vertical distributions and composition in Baltic Sea cyanobacterial blooms: *Harmful Algae*, v. 6, p. 189–205, <https://doi.org/10.1016/j.hal.2006.07.006>.
- Hennekam, R., Sweere, T., Tjallingii, R., de Lange, G.J., and Reichart, G.-J., 2019, Trace metal analysis of sediment cores using a novel X-ray fluorescence core scanning method: *Quaternary International*, v. 514, p. 55–67, <https://doi.org/10.1016/j.quaint.2018.10.018>.
- Kemp, A.E.S., Pearce, R.B., Koizumi, I., Pike, J., and Rance, S.J., 1999, The role of mat-forming diatoms in the formation of Mediterranean sapropels: *Nature*, v. 398, p. 57–61, <https://doi.org/10.1038/18001>.
- Lourens, L.J., Wehausen, R., and Brumsack, H.J., 2001, Geological constraints on tidal dissipation and dynamical ellipticity of the Earth over the past three million years: *Nature*, v. 409, p. 1029–1033, <https://doi.org/10.1038/35059062>.
- Menzel, D., van Bergen, P.F., Schouten, S., and Sinninghe Damsté, J.S., 2003, Reconstruction of changes in export productivity during Pliocene sapropel deposition: A biomarker approach: *Palaeogeography, Palaeoclimatology, Palaeoecology*, v. 190, p. 273–287, [https://doi.org/10.1016/S0031-0182\(02\)00610-7](https://doi.org/10.1016/S0031-0182(02)00610-7).

- Nichols, B.W., and Wood, B.J.B., 1968, New glycolipid specific to nitrogen-fixing blue-green algae: *Nature*, v. 217, p. 767–768, <https://doi.org/10.1038/217767a0>.
- Pearce, R.B., Kemp, A.E.S., Koizumi, I., Pike, J., Cramp, A., and Rowland, S.J., 1998, A lamina-scale, SEM-based study of a late Quaternary diatom-ooze sapropel from the Mediterranean Ridge, Site 971, in Robertson, A.H.F., et al., eds., *Proceedings of the Ocean Drilling Program Scientific Results, Volume 160*: College Station, Texas, Ocean Drilling Program, p. 333–348, <https://doi.org/10.2973/odp.proc.sr.160.016.1998>.
- Ploug, H., 2008, Cyanobacterial surface blooms formed by *Aphanizomenon* sp. and *Nodularia spumigena* in the Baltic Sea: Small-scale fluxes, pH, and oxygen microenvironments: *Limnology and Oceanography*, v. 53, p. 914–921, <https://doi.org/10.4319/lo.2008.53.3.0914>.
- Rodríguez-Sanz, L., Bernasconi, S.M., Marino, G., Heslop, D., Müller, I.A., Fernandez, A., Grant, K.M., and Rohling, E.J., 2017, Penultimate deglacial warming across the Mediterranean Sea revealed by clumped isotopes in foraminifera: *Scientific Reports*, v. 7, 16572, <https://doi.org/10.1038/s41598-017-16528-6>.
- Rohling, E.J., 2007, Progress in paleosalinity: Overview and presentation of a new approach: *Paleoceanography*, v. 22, PA3215, <https://doi.org/10.1029/2007PA001437>.
- Rohling, E.J., and Gieskes, W.W.C., 1989, Late Quaternary changes in Mediterranean intermediate water density and formation rate: *Paleoceanography*, v. 4, p. 531–545, <https://doi.org/10.1029/PA004i005p00531>.
- Rohling, E.J., et al., 2002, African monsoon variability during the previous interglacial maximum: *Earth and Planetary Science Letters*, v. 202, p. 61–75, [https://doi.org/10.1016/S0012-821X\(02\)00775-6](https://doi.org/10.1016/S0012-821X(02)00775-6).
- Rohling, E.J., et al., 2004, Reconstructing past planktic foraminiferal habitats using stable isotope data: A case history for Mediterranean sapropel S5: *Marine Micropaleontology*, v. 50, p. 89–123, [https://doi.org/10.1016/S0377-8398\(03\)00068-9](https://doi.org/10.1016/S0377-8398(03)00068-9).
- Rohling, E.J., Marino, G., and Grant, K.M., 2015, Mediterranean climate and oceanography, and the periodic development of anoxic events (sapropels): *Earth-Science Reviews*, v. 143, p. 62–97, <https://doi.org/10.1016/j.earscirev.2015.01.008>.
- Rossignol-Strick, M., Nesteroff, W., Olive, P., and Vergnaud-Grazzini, C., 1982, After the deluge: Mediterranean stagnation and sapropel formation: *Nature*, v. 295, p. 105–110, <https://doi.org/10.1038/295105a0>.
- Rossignol-Strick, M., 1985, Mediterranean Quaternary sapropels, an immediate response of the African monsoon to variation of insolation: *Palaeogeography, Palaeoclimatology, Palaeoecology*, v. 49, p. 237–263, [https://doi.org/10.1016/0031-0182\(85\)90056-2](https://doi.org/10.1016/0031-0182(85)90056-2).
- Sachs, J.P., and Repeta, D.J., 1999, Oligotrophy and nitrogen fixation during eastern Mediterranean sapropel events: *Science*, v. 286, p. 2485–2488, <https://doi.org/10.1126/science.286.5449.2485>.
- Schouten, S., Villareal, T.A., Hopmans, E.C., Mets, A., Swanson, K.M., and Sinninghe Damsté, J.S., 2013, Endosymbiotic heterocystous cyanobacteria synthesize different heterocyst glycolipids than free-living heterocystous cyanobacteria: *Phytochemistry*, v. 85, p. 115–121, <https://doi.org/10.1016/j.phytochem.2012.09.002>.
- Sollai, M., Hopmans, E.C., Bale, N.J., Mets, A., Warden, L., Moros, M., and Sinninghe Damsté, J.S., 2017, The Holocene sedimentary record of cyanobacterial glycolipids in the Baltic Sea: An evaluation of their application as tracers of past nitrogen fixation: *Biogeosciences*, v. 14, p. 5789–5804, <https://doi.org/10.5194/bg-14-5789-2017>.
- Stal, L.J., Albertano, P., Bergman, B., von Bröckel, K., Gallon, J.R., Hayes, P.K., Sivonen, K., and Walsby, A.E., 2003, BASIC: Baltic Sea cyanobacteria. An investigation of the structure and dynamics of water blooms of cyanobacteria in the Baltic Sea—Responses to a changing environment: *Continental Shelf Research*, v. 23, p. 1695–1714, <https://doi.org/10.1016/j.csr.2003.06.001>.
- Subramaniam, A., et al., 2008, Amazon River enhances diazotrophy and carbon sequestration in the tropical North Atlantic Ocean: *Proceedings of the National Academy of Sciences of the United States of America*, v. 105, p. 10,460–10,465, <https://doi.org/10.1073/pnas.0710279105>.
- Taylor, F.J.R., 1966, Phytoplankton of the south western Indian Ocean: *Nova Hedwigia*, v. 12, p. 433–476.
- Tribouillard, N., Algeo, T.J., Lyons, T., and Riboulleau, A., 2006, Trace metals as paleoredox and paleoproductivity proxies: An update: *Chemical Geology*, v. 232, p. 12–32, <https://doi.org/10.1016/j.chemgeo.2006.02.012>.
- Tuo, S.-H., Lee Chen, Y.-L., Chen, H.-Y., and Chen, T.-Y., 2017, Free-living heterocystous cyanobacteria in the tropical marginal seas of the western North Pacific: *Journal of Plankton Research*, v. 39, p. 404–422, <https://doi.org/10.1093/plankt/fbx023>.
- van der Meer, M.T.J., Baas, M., Rijpstra, W.I.C., Marino, G., Rohling, E.J., Sinninghe Damsté, J.S., and Schouten, S., 2007, Hydrogen isotopic compositions of long-chain alkenones record freshwater flooding of the Eastern Mediterranean at the onset of sapropel deposition: *Earth and Planetary Science Letters*, v. 262, p. 594–600, <https://doi.org/10.1016/j.epsl.2007.08.014>.
- Wörmer, L., Cirés, S., Velázquez, D., Quesada, A., and Hinrichs, K.-U., 2012, Cyanobacterial heterocyst glycolipids in cultures and environmental samples: Diversity and biomarker potential: *Limnology and Oceanography*, v. 57, p. 1775–1788, <https://doi.org/10.4319/lo.2012.57.6.1775>.
- Zeev, E.B., Yogeve, T., Man-Aharonovich, D., Kress, N., Herut, B., Béjà, O., and Berman-Frank, I., 2008, Seasonal dynamics of the endosymbiotic, nitrogen-fixing cyanobacterium *Richelia intracellularis* in the eastern Mediterranean Sea: *The ISME Journal*, v. 2, p. 911–923, <https://doi.org/10.1038/ismej.2008.56>.
- Ziegler, M., Tuenter, E., and Lourens, L.J., 2010, The precession phase of the boreal summer monsoon as viewed from the eastern Mediterranean (ODP Site 968): *Quaternary Science Reviews*, v. 29, p. 1481–1490, <https://doi.org/10.1016/j.quascirev.2010.03.011>.

Printed in USA