

Structural completeness in propositional logics of dependence

Rosalie Iemhoff and Fan Yang*

Department of Philosophy

Utrecht University

Utrecht, the Netherlands

June 2015

Abstract

In this paper we prove that three of the main propositional logics of dependence (including propositional dependence logic and inquisitive logic), none of which is structural, are structurally complete with respect to a class of substitutions under which the logics are closed. We obtain an analogous result with respect to stable substitutions, for the negative variants of some well-known intermediate logics, which are intermediate theories that are closely related to inquisitive logic.

Keywords: structural completeness, admissible rules, dependence logic, inquisitive logic, intermediate logic

1 Introduction

In recent years there have appeared many results on admissible rules in logics. The diversity of the results show that the properties of admissibility vary from logic to logic, and the complexity of some of the results show that describing these rules is not always an easy matter. The admissible rules of a logic are the rules under which the logic is closed, meaning that one could add them to the logic without obtaining new theorems. Since adding a derivable rule to a logic cannot alter what can be derived, derivable rules are always admissible, thus showing that the notion of admissibility is a natural extension of the notion of derivability.

For *structural logics*, which means logics that are closed under uniform substitution, a rule is admissible if every substitution that unifies the premiss, unifies the conclusion, where a substitution σ unifies a formula φ in a logic if $\sigma\varphi$ is derivable in the logic. Until now, most logics for which the admissibility relation has been studied are structural. Main examples are classical and intuitionistic propositional logic and certain modal logics such as K, K4, and S4. Except for classical logic, all these logics

*Support by the Netherlands Organisation for Scientific Research under grant 639.032.918 is gratefully acknowledged.

have nonderivable admissible rules and their admissibility relations are decidable and have concise axiomatizations [6, 13, 15, 21, 22]. In recent years, admissibility has been studied for a plethora of other logics as well. However, logics that are not structural, have received less attention. In order to obtain a meaningful notion of admissibility for such a logic one first has to isolate a set of substitutions, as large as one thinks possible, under which the logic is closed. Admissibility can then be studied with respect to this class of substitutions.

In this paper we show that three of the main propositional logics of dependence, none of which is structural, are structurally complete with respect to the class of *flat* substitutions. We obtain an analogous result, but then with respect to *stable* substitutions, for the negative variants of some well-known intermediate logics, which are intermediate theories that are closely related to one of the logics of dependence. As a byproduct we develop an extension of the usual logics of dependence in which the use of negation and the dependence atom is not restricted to propositional variables, but to the much larger class of flat formulas instead.

We think the interest in these results lies in the fact that logics of dependence, to be described below, are versatile and widely applicable nonclassical logics. And knowing that many nonclassical logics have nontrivial admissible rules, establishing that in these logics all rules that are admissible (with respect to flat substitutions) are derivable, provides a useful insight in the logics. Moreover, these results provide one of the first examples of natural nonstructural logics for which admissibility is studied. A paper in which various admissibility relations of nonstructural logics, the same logics that we treat in Theorem 5.5, have been studied is [20], but the results are different from the ones obtained here.

Dependence logic is a new logical formalism that characterizes the notion of “dependence” in social and natural sciences. First-order dependence logic was introduced by Väänänen [24] as a development of *Henkin quantifier* [8] and *independence-friendly logic* [10]. Recently, propositional dependence logic (PD) was studied and axiomatized in [23, 27]. With a different motivation, Ciardelli and Roelofsen [4] introduced and axiomatized *propositional inquisitive logic* (InqL), which can be regarded as a natural variant of propositional dependence logic. Both PD and InqL are fragments of *propositional downward closed team logic* (PT), which was studied in [27] and essentially also in [3]. Dependency relations are characterized in these propositional logics of dependence by a new type of atoms $\equiv(\vec{p}, q)$, called *dependence atoms*. Intuitively, the atom specifies that *the proposition q depends completely on the propositions \vec{p}* . The semantics of these logics is called *team semantics*, introduced by Hodges [11, 12]. The basic idea of this new semantics is that properties of dependence cannot be manifested in *single* valuations, therefore unlike the case of classical propositional logic, formulas in propositional logics of dependence are evaluated on *sets* of valuations (called *teams*) instead.

The three logics PD, InqL and PT are of particular interest, because they are all *expressively complete*, in the sense that they characterize all downward closed nonempty collections of teams. As a result of the feature of team semantics, the sets of theorems of these logics are closed under flat substitutions, but not closed under *uniform substitution*. As mentioned above, in this paper we prove that the three logics are structurally

complete with respect to flat substitutions.

In the study of admissible rules there is a technical detail that needs to be addressed. In PD and PT, negation and the dependence operator can only be applied to atoms. Therefore, the only substitutions under which these logics are closed are *renamings*, substitutions that replace atoms by atoms. However, these logics can be conservatively extended to logics that are closed under flat substitutions. These extensions, **PD** and **PT**, are closed under flat substitutions, and for these logics, as well as for InqL, the notion of admissibility with respect to flat substitutions is shown to be equal to derivability (Theorem 5.4).

There is a close connection between inquisitive logic and certain intermediate logics. The set of theorems of the former equals the negative variant of Kreisel-Putnam logic (KP), which is equal to the negative variant of Medvedev logic (ML). It is open whether KP is structurally complete, whereas ML is known to be structurally complete but not hereditarily structurally complete. An interesting corollary we obtain in this paper is that the negative variants of both ML and KP are hereditarily structurally complete with respect to negative substitutions.

2 Logics of dependence

2.1 Syntax and semantics

We first define the language of propositional downward closed team logic. All of the logics of dependence we consider in the paper are fragments of propositional downward closed team logic.

Definition 2.1. Fix a set Prop of propositional variables and denote its elements by p, q, r, \dots (possibly with subscripts). A sequence p_1, \dots, p_k of propositional variables will be denoted by \vec{p} . Well-formed formulas of *propositional downward closed team logic* (PT) are given by the following grammar:

$$\varphi ::= p \mid \neg p \mid \perp \mid \top \mid =(\vec{p}, q) \mid \varphi \wedge \varphi \mid \varphi \otimes \varphi \mid \varphi \vee \varphi \mid \varphi \rightarrow \varphi.$$

We call the formulas p , $\neg p$, \perp and \top *propositional atoms*. The formula $=(\vec{p}, q)$ is called a *dependence atom*, and it shall be read as “ q depends on \vec{p} ”. The connective \otimes is called *tensor* (disjunction), and the connectives \vee and \rightarrow are called *intuitionistic disjunction* and *intuitionistic implication*, respectively. The formula $\varphi \rightarrow \perp$ is abbreviated as $\neg\varphi$, and the team semantics to be given guarantees that the formula $\neg p$ and $p \rightarrow \perp$ are semantically equivalent.

Fragments of PT formed by certain sets of atoms and connectives in the standard way are called (*propositional*) *logics of dependence*. The following table defines the language of the logics of dependence we consider in this paper.

Logic	Atoms	Connectives
Propositional dependence logic (PD)	$p, \neg p, \perp, \top, =(\vec{p}, q)$	\wedge, \otimes
Propositional inquisitive logic (InqL)	p, \perp, \top	$\wedge, \vee, \rightarrow$

Given any of the three logics $L \in \{\text{PD}, \text{InqL}, \text{PT}\}$, let \mathcal{L}_L denote the language of L . We say that a formula φ is in \mathcal{L}_L if all symbols in φ belong to \mathcal{L}_L . Clearly, $\mathcal{L}_{\text{InqL}}$ is the same as the language of intuitionistic propositional logic or intermediate logics. We will discuss the connection between InqL and intermediate logics in the sequel. Note that formulas in \mathcal{L}_{PD} are assumed to be in (*strict*) *negation normal form*, in the sense that negation is allowed only in front of propositional variables and dependence atoms can not be negated. We will revisit the issue about negation in Section 3.1.

For the semantics, propositional logics of dependence adopt *team semantics*. A *team* is a set of valuations, i.e., a set of functions $v : \text{Prop} \rightarrow \{0, 1\}$, where Prop denotes the set of all propositional variables.

Definition 2.2. We inductively define the notion of a formula φ in \mathcal{L}_{PT} being *true* on a team X , denoted by $X \models \varphi$, as follows:

- $X \models p$ iff for all $v \in X$, $v(p) = 1$;
- $X \models \neg p$ iff for all $v \in X$, $v(p) = 0$;
- $X \models \perp$ iff $X = \emptyset$;
- $X \models \top$ for all teams X ;
- $X \models =(\vec{p}, q)$ iff for any $v, v' \in X$, $v(\vec{p}) = v'(\vec{p})$ implies $v(q) = v'(q)$;
- $X \models \varphi \wedge \psi$ iff $X \models \varphi$ and $X \models \psi$;
- $X \models \varphi \otimes \psi$ iff there exist teams $Y, Z \subseteq X$ with $X = Y \cup Z$ such that $Y \models \varphi$ and $Z \models \psi$;
- $X \models \varphi \vee \psi$ iff $X \models \varphi$ or $X \models \psi$;
- $X \models \varphi \rightarrow \psi$ iff for any team $Y \subseteq X$, $Y \models \varphi$ implies $Y \models \psi$.

We say that a formula φ is *valid*, denoted by $\models \varphi$, if $X \models \varphi$ holds for all teams X . For a finite set Γ of formulas, we write $\Gamma \models \varphi$ and say that φ is a *logical consequence* of Γ if $X \models \bigwedge \Gamma$ implies $X \models \varphi$ for any team X . We write $\varphi \models \psi$ for $\{\varphi\} \models \psi$. If $\varphi \models \psi$ and $\psi \models \varphi$, then we write $\varphi \equiv \psi$ and say that φ and ψ are *semantically equivalent*. Two logics of dependence L_1 and L_2 are said to have the *same expressive power* if for every L_1 -formula φ , $\varphi \equiv \psi$ for some L_2 -formula ψ , and vice versa.

The logics of dependence mentioned above are defined as follows. Since in this paper we consider the logics from a semantical point of view, using the team semantics, we define their finitary consequence relations semantically.

Definition 2.3 (Consequence relations for logics of dependence). For a logic $L \in \{\text{PD}, \text{InqL}, \text{PT}\}$, a formula φ and a finite set of formulas Γ , $\Gamma \vdash_L \varphi$ if and only if φ and all formulas in Γ are in \mathcal{L}_L and $\Gamma \models \varphi$. A formula φ is *valid in L* , or a *theorem* of L , if $\vdash_L \varphi$, which is short for $\emptyset \vdash_L \varphi$. Thus theorems of PD and InqL are the restrictions of the theorems of PT to \mathcal{L}_{PD} and $\mathcal{L}_{\text{InqL}}$, respectively.

Because of the semantical definition of the consequence relation \vdash_{\perp} , soundness and completeness with respect to the team semantics trivially holds. We will see, however, that there do exist genuine syntactic characterizations of propositional logics of dependence, as given in Theorem 2.7 and the comments thereafter. Since in this paper the methods are purely semantical, the semantically defined consequence relations suffice for our aims.

We write $\varphi(p_1, \dots, p_n)$ if the propositional variables occurring in φ are among p_1, \dots, p_n . Given a set V of propositional variables, a *valuation on V* is a function $v : V \rightarrow \{0, 1\}$, and a *team on V* is a set of valuations on V .

Theorem 2.4. Let $\varphi(p_1, \dots, p_n)$ be a formula and Γ a set of formulas in \mathcal{L}_{PT} , and X and Y two teams. Then the following holds.

(Locality) If $\{v \upharpoonright \{p_1, \dots, p_n\} : v \in X\} = \{v \upharpoonright \{p_1, \dots, p_n\} : v \in Y\}$, then

$$X \models \varphi \iff Y \models \varphi.$$

(Downward Closure Property) If $X \models \varphi$ and $Y \subseteq X$, then $Y \models \varphi$.

(Empty Team Property) $\emptyset \models \varphi$.

(Deduction Theorem) $\Gamma, \varphi \models \psi$ if and only if $\Gamma \models \varphi \rightarrow \psi$.

(Compactness Theorem) If $\Gamma \models \varphi$, then there exists a finite set $\Delta \subseteq \Gamma$ such that $\Delta \models \varphi$.

Given a formula φ and a finite set $\{\varphi_i \mid i \in I\}$ of formulas we introduce a meta-symbol \sqcup and use $\varphi \sqcup_{i \in I} \varphi_i$ as an abbreviation for the statement: For all teams X : $X \models \varphi_i$ implies $X \models \varphi$ for all $i \in I$, and $X \models \varphi$ implies $X \models \varphi_i$ for some $i \in I$.

Theorem 2.5 (Disjunction property). Let φ be a formula and $\{\varphi_i \mid i \in I\}$ a finite set of formulas in \mathcal{L}_{\perp} . If $\varphi \sqcup_{i \in I} \varphi_i$ and $\models \varphi$, then $\models \varphi_i$ for some $i \in I$.

Proof. Let Prop be the set of all propositional variables. Since $\models \varphi$, for the team $X = \{0, 1\}^{\text{Prop}}$, we have $X \models \varphi$. It follows from $\varphi \sqcup_{i \in I} \varphi_i$ that $X \models \varphi_i$ for some $i \in I$. Noting that every team Y is a subset of X , by the downward closure property we obtain that $Y \models \varphi_i$, which implies $\models \varphi_i$. \square \square

A formula of PT is said to be *classical* if it does not contain any dependence atoms or intuitionistic disjunction. Classical formulas φ of PT are *flat*, that is,

$$X \models \varphi \iff \forall v \in X, \{v\} \models \varphi$$

holds for all teams X . The following lemma shows that classical tautologies of PT are exactly the tautologies of classical propositional logic (CPC).

Lemma 2.6. For any classical formula φ in \mathcal{L}_{PT} , identifying tensor disjunction with classical disjunction of CPC, we have that $\models_{\text{CPC}} \varphi \iff \models_{\text{PT}} \varphi$.

Proof. An easy inductive proof shows that $v \models_{\text{CPC}} \varphi \iff \{v\} \models_{\text{PT}} \varphi$ for all valuations v and all classical formulas φ . \square \square

Having the same syntax as intuitionistic logic, the logic InqL has a close relationship with intermediate logics between ND and ML. In [4], a Hilbert-style deduction system for InqL is given. The axioms of this system will play a role in this paper, so we present the system in detail as follows.

Theorem 2.7 (see [4]). The following Hilbert-style deduction system of InqL is sound and strongly complete:

Axioms:

- all substitution instances of IPC axioms
- $\neg\neg p \rightarrow p$ for all $p \in \text{Prop}$
- all substitution instances of ND_k for all $k \in \mathbb{N}$:

$$(\text{ND}_k) \quad (\neg\varphi \rightarrow \bigvee_{1 \leq i \leq k} \neg\psi_i) \rightarrow \bigvee_{1 \leq i \leq k} (\neg\varphi \rightarrow \neg\psi_i).$$

Rule:

$$\text{Modus Ponens: } \frac{\varphi \rightarrow \psi \quad \psi}{\varphi} \text{ (MP)}$$

Remark 2.8. InqL extended with dependence atoms is called *propositional intuitionistic dependence logic* (PID) in the literature (see e.g., [26, 27]). As noted in [26, 27], PID and InqL have the same expressive power, as dependence atoms are definable in InqL:

$$\equiv(p_1, \dots, p_n, q) \equiv (p_1 \vee \neg p_1) \wedge \dots \wedge (p_n \vee \neg p_n) \rightarrow (q \vee \neg q). \quad (1)$$

Adding an axiom that corresponds to the above equivalence to the deduction system of InqL, one obtains a complete axiomatization for PID. For simplicity, we will not discuss the logic PID in this paper, but we remark that results obtained in this paper can be easily generalized to PID.

The logic PD was first axiomatized by a natural deduction system in [26, 27], and a Hilbert-style axiomatization and a labelled tableau calculus for PD can be found in [23]. Based on these, a natural deduction system for the fragment of PT without dependence atoms was given in [3]. Adding to the deduction system in [3] obvious rules for dependence atom that correspond to the equivalence in (1), one easily obtains a complete natural deduction system for full PT. Interested readers are referred to the literature given for the exact definitions of the deduction systems. Throughout this paper, we take for granted the strong completeness theorem for these logics.

It is important to note that the deduction systems for PD, InqL and PT do *not* admit *uniform substitution*. Here substitutions, a crucial notion in this paper, are defined as follows. The definition is sufficiently general to apply to both propositional logics of dependence and intermediate logics that we consider later in the paper.

Definition 2.9 (Substitution). A *substitution* of a propositional logic or theory L is a mapping σ from the set of all formulas in \mathcal{L}_L to the set of all formulas in \mathcal{L}_L , that commutes with the connectives and atoms.

Definition 2.10. Let $\vdash_{\mathcal{L}}$ be a consequence relation of a logic or theory \mathcal{L} . A substitution σ is called a $\vdash_{\mathcal{L}}$ -substitution if $\vdash_{\mathcal{L}}$ is closed under σ , i.e., for all formulas φ, ψ in $\mathcal{L}_{\mathcal{L}}$,

$$\varphi \vdash_{\mathcal{L}} \psi \implies \sigma(\varphi) \vdash_{\mathcal{L}} \sigma(\psi).$$

If $\vdash_{\mathcal{L}}$ is closed under all substitutions, then we say that $\vdash_{\mathcal{L}}$ is *structural*.

The consequence relations of the logics PD, InqL and PT are not structural, because, for example, $p \otimes p \vdash_{\text{PD}} p$ and $\vdash_{\text{InqL}} \neg\neg p \rightarrow p$, but $\models(p) \otimes \models(p) \not\vdash_{\text{PD}} \models(p)$ and $\not\vdash_{\text{InqL}} \neg\neg(p \vee \neg p) \rightarrow p \vee \neg p$.

2.2 Normal forms

In this section, we recall from [4] and [27] the disjunctive normal forms for formulas of PD, InqL and PT. These normal forms, reminiscent of the disjunctive normal form in classical logic, play an important role in the main proofs of this paper and are defined as follows.

Fix $V = \{p_1, \dots, p_n\}$. Let X be a team on V . For each of the logics PD, InqL and PT, we define a formula Θ_X as follows:

$$\Theta_X := \begin{cases} \bigotimes_{v \in X} (p_1^{v(p_1)} \wedge \dots \wedge p_n^{v(p_n)}) & \text{for PD} \\ \neg\neg \bigvee_{v \in X} (p_1^{v(p_1)} \wedge \dots \wedge p_n^{v(p_n)}) & \text{for InqL, PT} \end{cases} \quad (2) \quad (3)$$

where $p^1 := p$ and $p^0 := \neg p$ and we stipulate that $\Theta_{\emptyset} := \perp$. The reader can verify readily that the two formulas in (2) and (3) are semantically equivalent. This is why we decide to be sloppy here and use the same notation Θ_X to stand for two syntactically different formulas. We tacitly assume that Θ_X is given by (2) in the context of PD and by (3) in the context of InqL. For PT we could as well have chosen (2) as the definition of Θ_X , as both defining formulas belong to \mathcal{L}_{PT} and are equivalent.

With respect to the domain V , the formula Θ_X defines the family of subteams of the team X , as stated in the following lemma, whose proof is left to the reader or see [27].

Lemma 2.11. Let X and Y be teams on V . For the logics PD, InqL and PT, we have $Y \models \Theta_X \iff Y \subseteq X$.

The set $\llbracket \varphi \rrbracket = \{X \subseteq \{0, 1\}^V : X \models \varphi\}$ is nonempty (as $\emptyset \in \llbracket \varphi \rrbracket$) and *downward closed*, i.e., $Y \subseteq X \in \llbracket \varphi \rrbracket$ implies $Y \in \llbracket \varphi \rrbracket$. We say that a propositional logic \mathcal{L} of dependence is *expressively complete*, if every nonempty downward closed collection \mathcal{K} of teams on V is definable by a formula φ in $\mathcal{L}_{\mathcal{L}}$, i.e., $\mathcal{K} = \llbracket \varphi \rrbracket$.

Theorem 2.12 (see [4, 27]). (i) All of the logics PT, PD and InqL are expressively complete and have the same expressive power.

(ii) **(Normal Forms)** Let $\varphi(p_1, \dots, p_n)$ be a formula in \mathcal{L}_{PT} or \mathcal{L}_{PD} or $\mathcal{L}_{\text{InqL}}$. There exists a finite collection $\{X_i \mid i \in I\}$ of teams on V such that $\varphi \equiv \bigvee_{i \in I} \Theta_{X_i}$. In particular, $\varphi \equiv \bigvee_{i \in I} \Theta_{X_i}$ holds for PT and InqL.

Proof. We only give a proof sketch. For (i), let \mathcal{K} be a nonempty downward closed collection of teams on V . The formula $\bigvee_{X \in \mathcal{K}} \Theta_X$ in \mathcal{L}_{PT} or $\mathcal{L}_{\text{InqL}}$ satisfies $\mathcal{K} = \llbracket \bigvee_{X \in \mathcal{K}} \Theta_X \rrbracket$ by Lemma 2.11. The proof for the logic PD follows from a different argument; we refer the reader to [27] for details.

For every formula φ , the set $\llbracket \varphi \rrbracket$ is nonempty and downward closed. Thus item (ii) follows from the proof of item (i). \square \square

2.3 Intermediate logics

There is a close relationship between logics of dependence and intermediate theories (i.e., theories between intuitionistic and classical propositional logic), as first formulated in [4]. Here we describe this connection, and in the sections on projectivity and admissibility we will treat dependence logics and intermediate theories side by side.

An *intermediate theory* is a set L of formulas closed under modus ponens such that $\text{IPC} \subseteq L \subseteq \text{CPC}$. An *intermediate logic* is an intermediate theory closed under uniform substitution. The intermediate logics that are most relevant in this paper are Maksimova’s logic ND, Kreisel-Putnam logic KP and Medvedev’s logic ML (“the logic of finite problems”). It is well-known that $\text{ND} \subseteq \text{KP} \subseteq \text{ML}$, and ML is the maximal intermediate logic extending ND that has the disjunction property.

We call a substitution σ *stable* in a logic L that has implication and negation in its language, if $\sigma(p)$ is *stable* in L , i.e., $\vdash_L \sigma(p) \leftrightarrow \neg \neg \sigma(p)$, for all $p \in \text{Prop}$. It is easy to verify that the substitution $(\cdot)^\neg$, defined as $p^\neg = \neg p$ for all $p \in \text{Prop}$, is a stable substitution in all intermediate logics. For any intermediate logic L , define its *negative variant* L^\neg as

$$L^\neg = \{\varphi \mid \varphi^\neg \in L\}.$$

Lemma 2.13 (see [4]). Let L be an intermediate logic.

- (i) L^\neg is the smallest intermediate theory that contains L and $\neg \neg p \rightarrow p$ for every $p \in \text{Prop}$.
- (ii) The consequence relation \vdash_{L^\neg} of L^\neg is closed under stable substitutions.
- (iii) If L has the disjunction property, then so does L^\neg .

Lemma 2.14. Let L be an intermediate logic such that $\text{ND} \subseteq L$. Every formula is provably equivalent to a formula of the form $\bigvee_{i \in I} \neg \varphi_i$ in L^\neg .

Proof. The lemma follows essentially from [4]. Each formula $\neg \varphi_i$ is a Θ_X formula as defined in (3) for some set X of valuations, and the proof makes essential use of the axioms of ND and Lemma 2.13(i). \square \square

It was shown in [4] that the negative variants of all of the intermediate logics between ND and ML (including KP) are identical. Propositional inquisitive logic InqL is the negative variant of such logics. We state this and other properties of InqL in the following theorem.

Theorem 2.15 (see [4]). **(i)** For any intermediate logic L such that $ND \subseteq L \subseteq ML$, we have $\text{Inq}L = L^\neg$.

(ii) $\text{Inq}L$ has the disjunction property and its consequence relation $\vdash_{\text{Inq}L}$ is closed under stable substitutions.

There are many intermediate logics, including ND and KP , for which not much is known about their admissible rules. In Theorem 5.5 we show that the negative fragment of intermediate logics between ND and ML is structurally complete with respect to stable substitutions. Although we cannot immediately draw conclusions from this about the admissibility in the original logics, we hope that our results can be of help in the understanding of admissibility in these logics some day.

3 Extensions of the logics and substitutions

3.1 Extensions of the logics

For intermediate logics and $\text{Inq}L$, all possible substitutions are well-defined, meaning that given a formula and a substitution in the language of the logic, applying that substitution to the formula results in a formula in that language. However, for the other logics of dependence that we consider in this paper (i.e., PD and PT), substitution is *not* well-defined. A counter example is the formulas $\text{=}(p_1, \dots, p_n, q)$ and $\neg p$, for which the substitution instances $\text{=}(\sigma p_1, \dots, \sigma p_n, \sigma q)$ and $\neg \sigma(p)$ only belong to \mathcal{L}_{PD} or/and \mathcal{L}_{PT} if σ maps every propositional variable to a propositional variable.

For the study of admissibility one has to isolate the (or a meaningful) set of *well-defined* substitutions under which a consequence relation of a logic is *closed*. For this purpose, in this section we expand the languages of the logics PD and PT so as to force flat substitutions to be well-defined, and we will show in the next section that these extensions are closed under flat substitutions.

Definition 3.1. The following grammars define well-formed formulas of the extended logics of dependence.

- The *extended propositional downward closed team logic (PT)*:

$$\varphi ::= p \mid \perp \mid \top \mid \text{=}(\vec{\varphi}, \varphi) \mid \neg \varphi \mid \varphi \wedge \varphi \mid \varphi \otimes \varphi \mid \varphi \vee \varphi \mid \varphi \rightarrow \varphi.$$

- The *extended propositional dependence logic (PD)*:

$$\varphi ::= p \mid \perp \mid \top \mid \text{=}(\vec{\alpha}, \beta) \mid \neg \varphi \mid \varphi \wedge \varphi \mid \varphi \otimes \varphi,$$

where $\vec{\alpha}, \beta$ are flat formulas.

The extended logics have arbitrary negations as well-formed formulas. This negation will be defined as the intuitionistic negation that is conservative over the restricted negation in the original logics. The intuitionistic negation in the team semantics context can be traced back to the negation $\sim\downarrow$ considered by Hodges [11, 12], and a recent

work [17] considered this negation also in the context of modal dependence logic. The extension **PT** has dependence atoms with arbitrary arguments, while in the extension **PD** we only allow dependence atoms with flat arguments. The restriction for **PD** is made for technical simplicity that we discuss in the sequel, but as we consider flat substitutions only, this limitation does not affect the generality of the results in this paper. Generalized dependence atoms with flat arguments are also studied in the context of modal dependence logic, see [5, 7].

Below we define the semantics of the new formulas. We first treat **PT** and then **PD**.

Definition 3.2. Let $\varphi_1, \dots, \varphi_n, \psi$ be arbitrary formulas of **PT**. Define

- (a) $X \models (\varphi_1, \dots, \varphi_n, \psi)$ iff $X \models \bigwedge_{i=1}^n (\varphi_i \vee (\varphi_i \rightarrow \perp)) \rightarrow (\psi \vee (\psi \rightarrow \perp))$;¹
- (b) $X \models \neg\varphi$ iff $X \models \varphi \rightarrow \perp$ iff $\{v\} \not\models \varphi$ for all $v \in X$.

In order for these definitions to be well-defined they have to agree with previously defined notions. For the dependence atom the observation (1) suffices. For negation, it suffices for **PT** that $\neg\varphi$ has been defined as a shorthand for $\varphi \rightarrow \perp$, thus the semantics for negation as given in item (b) coincides with that in this logic.

We turn to **PD**. To define the semantics of the new formulas we need the following equivalence relation between valuations. Given a sequence $\vec{\varphi} = \varphi_1 \dots \varphi_n$ of formulas, define an equivalence relation $\sim_{\vec{\varphi}}$ on the set of all valuations as follows:

$$u \sim_{\vec{\varphi}} v \quad \text{iff} \quad \forall 1 \leq i \leq n (\{u\} \models \varphi_i \Leftrightarrow \{v\} \models \varphi_i).$$

Definition 3.3. Define

- (a) for flat formulas $\alpha_1, \dots, \alpha_k, \beta$ of **PD**,

$$X \models (\vec{\alpha}, \beta) \equiv_{\text{df}} \forall v, v' \in X (v \sim_{\vec{\alpha}} v' \implies v \sim_{\beta} v'); \quad (4)$$

- (b) full negation in **PD** as $X \models \neg\varphi$ iff $\{v\} \not\models \varphi$ for all $v \in X$.

We have to show that the notions defined in Definition 3.3 are extensions of the corresponding notions for **PD**, and also special case of those of **PT**. Obviously for the formula (\vec{p}, q) , the semantics given in item (a) coincides with the semantics given in Definition 2.2, and we leave it to the reader to check that it also coincides with Definition 3.2(a).

The negation defined in item (b) deserves more comments. It is straightforward from the definition that $\neg\varphi$ is always flat, and the so defined negation being the intuitionistic negation coincides with that of **PT**. In the literature of first-order dependence logic, negation is usually treated only *syntactically*, in the sense that a negated formula $\neg\varphi$ is defined to have the same semantics as the unique formula φ^\sim in *negation normal form* obtained by exhaustively applying the De Morgan's laws and some other syntactic

¹The authors would like to thank Ivano Ciardelli for suggesting this definition, see also [3].

rewrite rules. The corresponding syntactic rewrite rules for propositional dependence logic are as follows:

$$\begin{array}{lll}
p^\sim & \mapsto & \neg p \\
(\neg p)^\sim & \mapsto & p \\
=(\vec{p}, \psi)^\sim & \mapsto & \perp \\
\top^\sim & \mapsto & \perp \\
\perp^\sim & \mapsto & \top \\
(\varphi \wedge \psi)^\sim & \mapsto & \varphi^\sim \otimes \psi^\sim \\
(\varphi \otimes \psi)^\sim & \mapsto & \varphi^\sim \wedge \psi^\sim
\end{array} \tag{5}$$

It is easy to see that the syntactic rewrite procedure for a negated formula $\neg\varphi$ of **PD** defined as above always terminates on a unique dependence atom-free formula φ^\sim in negation normal form in \mathcal{L}_{PD} .

When applying the syntactic negation, special attention needs to be paid to double negations of dependence atoms, i.e., formulas of the form $\neg\neg=(\vec{a}, b)$, where the variables \vec{a}, b are first-order or propositional. Following Hintikka's game-theoretic perspective of logic (see, e.g., [9]), the negation in logics of dependence is usually treated as a connective for which the two players in the corresponding semantic game swap their roles. This way $\neg\neg=(\vec{a}, b)$ should have the same meaning as $=(\vec{a}, b)$. However, this reading is not consistent with the syntactic rewrite rules as in (5). To avoid ambiguity, most literature of logics of dependence does not allow double negation to occur in front of dependence atoms. In this paper, in the extended logic **PD** we do include double negated dependence atoms as well-formed formulas, but as we do not take the game-theoretic approach to propositional logics of dependence, the semantics of double negated dependence atoms is computed simply according to Definition 3.3(b), namely, $\neg\neg=(\vec{p}, q)$ is always semantically equivalent to \top (noting that $=(\vec{p}, q)$ is always true on singleton teams). Given such interpretation of the double negated dependence atoms, the negation defined in Definition 3.3(b) coincides with the syntactic negation given by the rewrite rules in (5), as we will show in the next lemma. However, on the other hand, in the context of first-order dependence logic, regardless of how double negated dependence atoms are treated, the negation defined as in Definition 3.3(b) does *not* coincide with the syntactic negation given by the rewrite rules; rather, it corresponds to the defined connective $\sim\downarrow$ in Hodges [11, 12]. For instance, the reader who is familiar with the semantics of first-order dependence logic can easily verify that $M \not\models_{\{s\}} \forall x=(x)$ holds for all assignments s on all models M , assuming that the domain of a model has at least two elements. Thus by Definition 3.3(b) $M \models_X \neg\forall x=(x)$ for all teams X on all models M , namely $\neg\forall x=(x) \equiv \top$. However, by the syntactic rewrite rules, $(\forall x=(x))^\sim = \exists x=(x)^\sim = \exists x\perp$.

Lemma 3.4. For any formula φ in \mathcal{L}_{PD} , we have $\neg\varphi \equiv \varphi^\sim$.

Proof. We prove by induction on φ that $X \models \neg\varphi \iff X \models \varphi^\sim$ for all teams X .

The case $\varphi = p$ or \perp or \top is easy. If $\varphi = \neg p$, then $\varphi^\sim = p$ and we have that $X \models \neg\neg p \iff \forall v \in X : \{v\} \not\models \neg p \iff \forall v \in X : \{v\} \models p \iff X \models p$.

If $\varphi = =(\vec{p}, q)$, then $\varphi^\sim = \perp$ and we have that $X \models \neg=(\vec{p}, q) \iff \forall v \in X : \{v\} \not\models =(\vec{p}, q) \iff X = \emptyset \iff X \models \perp$.

If $\varphi = \psi \wedge \chi$, then $\varphi^\sim = \psi^\sim \otimes \chi^\sim$ and we have that

$$\begin{aligned}
& X \models \neg(\psi \wedge \chi) \\
& \iff \forall v \in X : \{v\} \not\models \psi \wedge \chi \\
& \iff \exists Y, Z \subseteq X \text{ s.t. } Y \cup Z = X, (\forall v \in Y : \{v\} \not\models \psi) \text{ and } (\forall u \in Z : \{u\} \not\models \chi) \\
& \iff \exists Y, Z \subseteq X \text{ s.t. } Y \cup Z = X, Y \models \neg\psi \text{ and } Z \models \neg\chi \\
& \iff \exists Y, Z \subseteq X \text{ s.t. } Y \cup Z = X, Y \models \psi^\sim \text{ and } Z \models \chi^\sim \text{ (by the induction hypothesis)} \\
& \iff X \models \psi^\sim \otimes \chi^\sim.
\end{aligned}$$

If $\varphi = \psi \otimes \chi$, then $\varphi^\sim = \psi^\sim \wedge \chi^\sim$ and we have by the induction hypothesis that

$$\begin{aligned}
X \models \neg(\psi \otimes \chi) & \iff \forall v \in X : \{v\} \not\models \psi \otimes \chi \\
& \iff \forall v \in X : \{v\} \not\models \psi \text{ and } \{v\} \not\models \chi \\
& \iff X \models \neg\psi \text{ and } X \models \neg\chi \\
& \iff X \models \psi^\sim \wedge \chi^\sim \text{ (by the induction hypothesis)}.
\end{aligned}$$

□

□

It is evident from Definition 3.3(b) that the full negation of **PD** is a *semantic connective*. A k -ary connective $*$ is called a *semantic connective*, if

$$[\varphi_1 \equiv \psi_1, \dots, \varphi_k \equiv \psi_k] \implies *(\varphi_1, \dots, \varphi_k) \equiv *(\psi_1, \dots, \psi_k).$$

Lemma 3.4 states that the semantic negation of **PD** defined in Definition 3.3(b) and the syntactic negation given by (5) coincide. It is worth emphasizing that in contrast to **PD** and other familiar logics with negation, the syntactic negation of first-order dependence logic is *not* a semantic connective (regardless of how double negated dependence atoms are treated), as shown by Burgess [1] and Väänänen and Kontinen [16]. For an illustration, $\forall x = (x) \equiv \forall x \forall y (x = y)$, whereas, by the syntactic rewrite rules, $(\forall x = (x))^\sim = \exists x \perp \not\equiv \exists x \exists y (x \neq y) = (\forall x \forall y (x = y))^\sim$.

The logics **PD** and **PT** are expressively complete, therefore their extensions have the same expressive power as the original ones. Thus it is straightforward to verify that Theorem 2.4 and Theorem 2.12 hold also for the extended logics **PD** and **PT**. One can easily extend the deduction systems of the original logics by adding characterization rules for the negation and generalized dependence atoms and prove the sound and completeness theorems for the extensions. To characterize the negation, to the deduction systems of **PT** and **PD** one adds the obvious rules that characterize the equivalence between $\neg\varphi$ and $\varphi \rightarrow \perp$, and the obvious rules that characterize the rewrite rules in (5), respectively. To characterize the generalized dependence atoms, to the deduction system of **PT** one adds obvious rules that correspond to the equivalence in Definition 3.2(a). For **PD**, following the idea in [27] one generalizes the rules for dependence atoms in the deduction system of **PD** according to the equivalence in Definition 3.2(a) in an obvious way. To prove the completeness theorem for such obtained system of **PD**, one observes that whenever $\varphi_1, \dots, \varphi_n, \psi$ are flat,

$$= (\varphi_1, \dots, \varphi_n, \psi) \equiv \bigvee_{f \in \{0,1\}^X} \bigotimes_{v \in X} (\varphi_1^{v(\varphi_1)} \wedge \dots \wedge \varphi_n^{v(\varphi_n)} \wedge \psi^{f(v)}) \quad (6)$$

holds, where $X = \{0, 1\}^{\{\varphi_1, \dots, \varphi_n\}}$, $\varphi_i^1 = \varphi_i$ and $\varphi_i^0 = \neg\varphi_i$, and modifies the definition of a *realization* of a generalised dependence atom accordingly. Note that if the arguments $\varphi_1, \dots, \varphi_n, \psi$ of a generalized dependence atom are not assumed to be flat, Equation (6) will no longer hold, and we do not see at this moment how to obtain a complete axiomatization of the extended logic also in the general case. But since the notion of admissibility we study in this paper concerns *theoremhood* of our logics only, and we intensionally defined the consequence relations of our logics in a semantic manner (see Definition 2.3), this obstacle in the axiomatization of the extended logic is not essential for the main results of this paper. In view of this, for the sake of simplicity in **PD** we only allow generalized dependence atoms with flat arguments.

3.2 Closure under flat substitutions

The consequence relations of the logics **PD**, **lnqL**, and **PT** are not structural. In this section we prove, however, that the consequence relations of these logics are closed under *flat substitutions*, i.e., substitutions σ that map propositional variables to flat formulas. To this end, we define the following translation on teams. For any valuation v and any substitution σ , define a valuation v_σ as

$$v_\sigma(p) = \begin{cases} 1 & \text{if } \{v\} \models \sigma(p) \\ 0 & \text{if } \{v\} \not\models \sigma(p) \end{cases}$$

For any team X , we define $X_\sigma = \{v_\sigma \mid v \in X\}$. Given a team $Y \subseteq X_\sigma$, let Y_X^σ denote the set $\{v \in X \mid v_\sigma \in Y\}$. Clearly $Y_X^\sigma \subseteq X$ and $(Y_X^\sigma)_\sigma = Y$.

Lemma 3.5. Let $L \in \{\mathbf{PD}, \mathbf{lnqL}, \mathbf{PT}\}$. For all formulas φ and all flat substitutions σ in \mathcal{L}_L ,

$$X \models \sigma(\varphi) \iff X_\sigma \models \varphi.$$

Proof. We prove this lemma for all three logics at the same time by induction on the complexity of φ , where we use the following complexity $c(\varphi)$ on formulas in $\mathcal{L}_{\mathbf{PT}}$. The use of the complicated clause for the dependence atoms will become clear in the proof below.

$$\begin{aligned} c(p) &= 0 && p \text{ a propositional variable} \\ c(\perp) &= 0 \\ c(\top) &= 0 \\ c(\neg\varphi) &= c(\varphi) + 1 \\ c(\varphi \circ \psi) &= c(\varphi) + c(\psi) + 1 && \circ \in \{\wedge, \rightarrow, \otimes\} \\ c(=\vec{\varphi}, \psi) &= (\sum_{i=1}^n (2c(\varphi_i) + 4)) + 2c(\psi) + 4 && \text{where } \vec{\varphi} = \varphi_1, \dots, \varphi_n \end{aligned}$$

The cases $\varphi = \perp$ and $\varphi = \top$ are trivial. Since $\sigma(p)$ is flat, the following equivalences hold:

$$X \models \sigma(p) \iff \forall v \in X (\{v\} \models \sigma(p)) \iff \forall v_\sigma \in X_\sigma (\{v_\sigma\} \models p) \iff X_\sigma \models p.$$

Thus the case $\varphi = p$ is proved.

Case $\varphi = \Rightarrow(\vec{\theta}, \psi)$. For **PT**, we know from Definition 3.2(a) that φ is semantically equivalent to a formula in its language whose subformulas are of lower complexity, thus this case is reduced to the other cases. However for **PD**, the equivalent formula given by Definition 3.2(a) is not in its language, neither is the equivalent formula given by Equation (6). Since PD is expressively complete, there indeed exists a formula φ' in the language of PD that is equivalent to φ . However, this translation is not done in a compositional manner, nor in an inductive manner (see Theorem 2.12). We therefore cannot reduce this case to the other cases for **PD**, as the reduction would assume

$$\varphi \equiv \varphi' \implies \sigma(\varphi) \equiv \sigma(\varphi'),$$

a fact that we establish only in Theorem 3.7. To avoid such a circular argument, we now proceed to prove this case for **PD** directly, using the equivalent semantics given in Definition 3.2(a) and assuming that $\vec{\theta}$ and ψ are flat.

For the direction “ \implies ”, assume $X \models \Rightarrow(\sigma(\vec{\theta}), \sigma(\psi))$ and $Y \models \bigwedge_{i=1}^n (\theta_i \vee \neg\theta_i)$ for some $Y \subseteq X_\sigma$. As $(Y_X^\sigma)_\sigma = Y$ and $c(\bigwedge_{i=1}^n (\theta_i \vee \neg\theta_i)) < \sum_{i=1}^n (2c(\theta_i) + 4) < c(\Rightarrow(\vec{\theta}, \chi))$, by the induction hypothesis, we obtain that $Y_X^\sigma \models \bigwedge_{i=1}^n (\sigma(\theta_i) \vee \neg\sigma(\theta_i))$. Clearly $Y_X^\sigma \subseteq X$, thus the assumption implies that $Y_X^\sigma \models \sigma(\psi) \vee \neg\sigma(\psi)$, which by the induction hypothesis again gives the desired $(Y_X^\sigma)_\sigma \models \psi \vee \neg\psi$, because $c(\psi \vee \neg\psi) = 2c(\psi) + 3 < c(\Rightarrow(\vec{\theta}, \psi))$. The other direction “ \impliedby ” is symmetric, using $Y_\sigma \subseteq X_\sigma$ for any X, Y with $Y \subseteq X$.

The cases that $\varphi = \psi \wedge \chi$ and $\varphi = \psi \vee \chi$ follow immediately from the induction hypothesis.

Case $\varphi = \psi \otimes \chi$. We first prove the direction “ \implies ”. Assume $X \models \sigma(\varphi)$ and consider $Y, Z \subseteq X$ such that $X = Y \cup Z$ and $Y \models \sigma(\psi)$ and $Z \models \sigma(\chi)$. Using that $Y_\sigma \cup Z_\sigma = X_\sigma$, this implies $X_\sigma \models \psi \otimes \chi$ by the induction hypothesis.

For the direction “ \impliedby ”, assume $X_\sigma \models \varphi$ and consider $Y, Z \subseteq X_\sigma$ such that $X_\sigma = Y \cup Z$, $Y \models \psi$ and $Z \models \chi$. Thus $Y_X^\sigma \models \sigma(\psi)$ and $Z_X^\sigma \models \sigma(\chi)$ by the induction hypothesis. Since $X = Y_X^\sigma \cup Z_X^\sigma$, this implies $X \models \sigma(\psi) \otimes \sigma(\chi)$, as required.

Case $\varphi = \psi \rightarrow \chi$. We first prove the direction “ \implies ”. Assume $X \models \sigma(\varphi)$ and consider $Y \subseteq X_\sigma$ such that $Y \models \psi$. As $(Y_X^\sigma)_\sigma = Y$, $Y_X^\sigma \models \sigma(\psi)$ follows by the induction hypothesis. And as $Y_X^\sigma \subseteq X$, this implies $Y_X^\sigma \models \sigma(\chi)$. Hence $Y \models \chi$ by the induction hypothesis, as required. The direction “ \impliedby ” is similar.

Case $\varphi = \neg\psi$. It follows from the induction hypothesis that $X \models \neg\sigma(\psi) \iff \forall v \in X : \{v\} \not\models \sigma(\psi) \iff \forall v \in X : \{v_\sigma\} \not\models \psi \iff X_\sigma \models \neg\psi$. \square \square

Lemma 3.6. The set of flat formulas in $\mathcal{L}_{\mathbf{PT}}$ is closed under flat substitutions, i.e., whenever φ is a flat formula and σ is a flat substitution, $\sigma(\varphi)$ is flat too.

Proof. Suppose X is a team such that for all $v \in X$, $\{v\} \models \sigma(\varphi)$. To show that $X \models \sigma(\varphi)$, by Lemma 3.5, it suffices to show that $X_\sigma \models \varphi$. As φ is flat, we therefore have to show that $\{v_\sigma\} \models \varphi$ for all $s \in X$. Again by Lemma 3.5 it suffices to show that $\{v\} \models \sigma(\varphi)$ for all $v \in X$. But that is what we assumed, so we are done. \square \square

As a consequence of the above lemma, for every generalized dependence atom $\Rightarrow(\varphi_1, \dots, \varphi_n, \psi)$ in $\mathcal{L}_{\mathbf{PD}}$, where $\varphi_1, \dots, \varphi_n, \psi$ are flat formulas, the resulting formula $\Rightarrow(\sigma(\varphi_1), \dots, \sigma(\varphi_n), \psi)$

under an arbitrary flat substitution σ is still a well-formed formula in $\mathcal{L}_{\mathbf{PD}}$. This shows that flat substitutions are well-defined in \mathbf{PD} .

Theorem 3.7. The consequence relations of \mathbf{PD} , \mathbf{InqL} , and \mathbf{PT} are closed under flat substitutions. In particular, we have for all flat substitutions σ that $\varphi \equiv \psi$ implies $\sigma(\varphi) \equiv \sigma(\psi)$.

Proof. By the definition of the consequence relations, it suffices to prove that for all formulas φ and ψ , $\varphi \models \psi$ implies $\sigma(\varphi) \models \sigma(\psi)$ for all flat substitutions σ .

Assume $\varphi \models \psi$. For any team X , any flat substitution σ ,

$$\begin{aligned} X \models \sigma(\varphi) &\implies X_\sigma \models \varphi \quad (\text{by Lemma 3.5}) \\ &\implies X_\sigma \models \psi \quad (\text{by the assumption}) \\ &\implies X \models \sigma(\psi) \quad (\text{by Lemma 3.5}). \end{aligned}$$

Hence $\sigma(\varphi) \models \sigma(\psi)$. \square \square

4 Flat formulas and projective formulas

Having proved that our logics are closed under flat substitutions we work towards the proof of our main results by showing that flatness in these logics is nothing but projectivity, a key notion in the study of admissible rules.

The formulas Θ_X that play a crucial role in the normal form of formulas in $\mathcal{L}_{\mathbf{PT}}$ defined in Section 2.2 turn out to be of particular interest. They actually serve as a syntactic characterization of flat formulas, as the following lemma shows.

Lemma 4.1. Let $\varphi(p_1, \dots, p_n)$ be a consistent formula in $\mathcal{L}_{\mathbf{PT}}$. The following are equivalent.

- (i) φ is flat
- (ii) $\varphi \equiv \Theta_X$ for some nonempty team X on $\{p_1, \dots, p_n\}$
- (iii) $\varphi \equiv \neg\neg\varphi$
- (iv) $\varphi \models \varphi \otimes \neg\varphi$

Proof. (ii) \implies (i) and (iii) \implies (i) follow from the fact that negated formulas are flat, and (i) \implies (iii) follows immediately from the definition of negation.

(i) \implies (ii): In view of Lemma 2.11 and Theorem 2.12, without loss of generality, we may assume that $\varphi(p_1, \dots, p_n) = \bigvee_{i=1}^k \Theta_{X_i}$, where $\{X_1, \dots, X_k\}$ is a collection of some nonempty maximal (with respect to set inclusion) teams on $\{p_1, \dots, p_n\}$. Suppose φ is flat and $k > 1$. For each $1 \leq i < k$, pick $v_i \in X_i \setminus X_{i+1}$ and pick $v_k \in X_k \setminus X_1$. The maximality of the X_i 's guarantees that such v_i 's exist. Since $\{v_i\} \subseteq X_i$ and $\{v_1, \dots, v_k\} \not\subseteq X_i$ for all $1 \leq i < k$, by Lemma 2.11, $\{v_i\} \models \Theta_{X_i}$ and $\{v_1, \dots, v_k\} \not\models \Theta_{X_i}$ for all $1 \leq i < k$, thereby $\{v_i\} \models \varphi$ for all $1 \leq i < k$ whereas $\{v_1, \dots, v_k\} \not\models \varphi$. Hence we conclude that $k = 1$ and $\varphi = \Theta_{X_1}$, as required.

(i) \Rightarrow (iv): If φ is flat, to show (iv), it suffices to show $\{v\} \models \varphi \otimes \neg\varphi$, i.e., $\{v\} \models \varphi$ or $\{v\} \models \neg\varphi$, for all valuations v . But this is obvious.

(iv) \Rightarrow (i): Suppose $\{v\} \models \varphi$ for all valuations v in a team X . Then $Y \not\models \neg\varphi$ for all nonempty $Y \subseteq X$. Now, if $\models \varphi \otimes \neg\varphi$, then we must have that $X \models \varphi$, which shows that φ is flat. \square \square

Since some of the logics we consider in this paper do not have implication in the language, and none of them is closed under uniform substitution, we modify the usual definition of projective formula.

Definition 4.2 (Projective formula). Let L be a logic, and S a set of L -substitutions. A formula φ in \mathcal{L}_L is said to be S -projective in L if there exists $\sigma \in S$ such that

- (a) $\vdash_L \sigma(\varphi)$,
- (b) $\varphi, \sigma(p) \vdash_L p$ and $\varphi, p \vdash_L \sigma(p)$ for all propositional variables p .

Such substitutions are called S -projective unifiers for φ in L .

Because of the Deduction Theorem (Theorem 2.4) of those of our logics that have built-in implication, the notion of projectivity can in those cases be formulated purely in terms of theoremhood. A standard inductive proof shows that the condition in Definition 4.2(b) implies that $\varphi, \sigma(\psi) \vdash_L \psi$ and $\varphi, \psi \vdash_L \sigma(\psi)$ hold for all formulas φ and ψ of our logics.

The proof of the following lemma uses what is known as *Prucnal's trick*, which consists of a method to prove projectivity via a connection between valuations and substitutions.

Lemma 4.3. Let $L \in \{\text{InqL}, \text{PT}\}$ and X a nonempty set of teams on a finite set of propositional variables. The formula Θ_X in \mathcal{L}_L (defined by Equation (3)) is \mathcal{F} -projective in L , where \mathcal{F} is the class of all flat substitutions.

Proof. Put $\varphi = \Theta_X$ and pick $v \in X$. View φ as a formula of CPC. Clearly we have $v(p_1^{v(p_1)} \wedge \dots \wedge p_n^{v(p_n)}) = 1$, and thereby $v(\varphi) = 1$. Define a substitution σ_v^φ as follows:

$$\sigma_v^\varphi(p) = \begin{cases} \varphi \wedge p & \text{if } v(p) = 0 \\ \varphi \rightarrow p & \text{if } v(p) = 1 \end{cases} \quad (7)$$

Put $\sigma = \sigma_v^\varphi$. Clearly, $\sigma(p)$ (in both cases) is classical, thus flat.

By a standard inductive argument, one proves that

$$\vdash_{\text{CPC}} \sigma(\psi) \iff v(\psi) = 1 \quad (8)$$

for all subformulas ψ of φ . Now, as $v(\varphi) = 1$, we obtain $\vdash_{\text{CPC}} \sigma(\varphi)$. Since φ is a classical formula, by Lemma 2.6 we derive $\vdash_L \sigma(\varphi)$. Moreover, it follows from the definition of σ that $\vdash_L \varphi \rightarrow (\sigma(p) \leftrightarrow p)$ holds for all $p \in \text{Prop}$. Hence we conclude that φ is \mathcal{F} -projective in L . \square \square

It is known that negated formulas $\neg\varphi$ are projective in every intermediate logic L . This follows, for example, from Ghilardi's characterization in [6]. Here we prove that the same holds for the negative variants of intermediate logics and that the projective unifiers involved are moreover stable.

Lemma 4.4. Let L be an intermediate logic. Every consistent formula $\neg\varphi$ is \mathcal{ST} -projective in L^\neg , where \mathcal{ST} is the class of all stable substitutions.

Proof. Take a valuation v such that $v(\neg\varphi) = 1$. Define a substitution $\sigma_v^{\neg\varphi}$ for $\neg\varphi$ in exactly the same way as in (7) of the preceding lemma. Put $\sigma = \sigma_v^{\neg\varphi}$. The definition of σ guarantees that $\vdash_L \varphi \rightarrow (\sigma(p) \leftrightarrow p)$ holds for all $p \in \text{Prop}$. By (8) and Glivenko's Theorem (see e.g. Theorem 2.47 in [2]), we obtain that $\vdash_L \neg\sigma(\varphi)$. Hence we have proved that $\neg\varphi$ is projective in L . Now, by Lemma 2.13 $L \subseteq L^\neg$, thus $\neg\varphi$ is projective also in L^\neg .

It remains to check that the σ defined as above is a stable substitution in L^\neg , i.e., $\vdash_{L^\neg} \sigma(p) \leftrightarrow \neg\neg\sigma(p)$ for all $p \in \text{Prop}$. If $v(p) = 0$, then by the definition, we have that $\sigma(p) = \neg\varphi \wedge p$. Since $\vdash_{L^\neg} \neg\neg p \rightarrow p$ (by Lemma 2.13), we have that

$$\neg\neg\sigma(p) = \neg\neg(\neg\varphi \wedge p) \dashv\vdash \neg\neg\neg\varphi \wedge \neg\neg p \dashv\vdash \neg\varphi \wedge p = \sigma(p),^2$$

as required. If $v(p) = 1$, then by the definition we have that $\sigma(p) = \neg\varphi \rightarrow p$. Since $\vdash_{L^\neg} \neg\neg p \rightarrow p$, we have that

$$\neg\neg\sigma(p) = \neg\neg(\neg\varphi \rightarrow p) \dashv\vdash \neg\neg\neg\varphi \rightarrow \neg\neg p \dashv\vdash \neg\varphi \rightarrow p = \sigma(p),$$

as required. □ □

Lemma 4.5. For any nonempty team X on a set $\{p_1, \dots, p_n\}$ of propositional variables, the formula Θ_X in $\mathcal{L}_{\mathbf{PD}}$ (defined by Equation (2)) is \mathcal{F} -projective in \mathbf{PD} .

Proof. This lemma is proved using a similar argument to that of Lemma 4.3 too. Put $\varphi = \Theta_X$. Take an arbitrary $v \in X$. Clearly, $v(\varphi) = 1$ when φ is viewed as a formula of CPC (hereafter in the proof, we identify the tensor disjunction \otimes with the classical disjunction). Define a substitution σ_v^φ as follows:

$$\sigma_v^\varphi(p) = \begin{cases} \varphi \wedge p & \text{if } v(p) = 0 \\ \neg\varphi \vee p & \text{if } v(p) = 1 \end{cases}$$

Put $\sigma = \sigma_v^\varphi$. Clearly, the formula $\sigma(p)$ (in both cases) is classical, thus flat.

As in the proof of Lemma 4.3, we have that (8) holds for all subformulas ψ of φ , thus $\vdash_{\text{CPC}} \sigma(\varphi)$. Now, since the formula $\sigma(\varphi)$ is classical, we obtain by Lemma 2.6 that $\vdash_{\mathbf{PD}} \sigma(\varphi)$.

It remains to show that $\varphi, \sigma(p) \vdash_L p$ and $\varphi, p \vdash_L \sigma(p)$ for all $p \in \text{Prop}$. If $v(p) = 0$, then clearly $\varphi, \varphi \wedge p \vdash_{\mathbf{PD}} p$ and $\varphi, p \vdash_{\mathbf{PD}} \varphi \wedge p$. If $v(p) = 1$, to see that $\varphi, \neg\varphi \otimes p \vdash_{\mathbf{PD}} p$, if $X \models \varphi \wedge (\neg\varphi \otimes p)$, then for all $v \in X$, we have that $\{v\} \models \varphi \wedge (\neg\varphi \otimes p)$, which implies that $\{v\} \models p$, thereby $X \models p$, as required. That $\varphi, p \vdash_{\mathbf{PD}} \neg\varphi \otimes p$ follows from the fact that $p \vdash_{\mathbf{PD}} \neg\varphi \otimes p$. □ □

²The notation $\varphi \dashv\vdash \psi$ is short for $\varphi \vdash \psi$ and $\psi \vdash \varphi$.

Lemma 4.6. Let $L \in \{\mathbf{PD}, \text{InqL}, \mathbf{PT}\}$, and φ a consistent formula in \mathcal{L}_L . The following are equivalent:

- (i) $\varphi \Vdash \Theta_X$ for some nonempty n -team X
- (ii) φ is flat
- (iii) φ is \mathcal{F} -projective in L

Proof. (ii) \iff (i) \implies (iii) follows from Lemmas 4.1, 4.3 and 4.5. Now, we show that (iii) \implies (i). Suppose φ is \mathcal{F} -projective in L and σ is an \mathcal{F} -projective unifier for φ . Then $\vdash_L \sigma(\varphi)$, which implies $\models \sigma(\varphi)$. By Theorem 2.5, this implies that there exists $1 \leq i \leq k$ such that $\models \sigma(\Theta_{X_i})$. Since Θ_{X_i} is in \mathcal{L}_L and thus so is $\sigma(\Theta_{X_i})$, $\vdash_L \sigma(\Theta_{X_i})$ follows. On the other hand, we also have that $\varphi, \sigma(\Theta_{X_i}) \vdash_L \Theta_{X_i}$. It then follows that $\varphi \vdash_L \Theta_{X_i}$. Hence $\varphi \equiv \Theta_{X_i}$, which gives $\varphi \Vdash \Theta_{X_i}$. \square \square

Lemma 4.7. Let L be an intermediate logic such that $\text{ND} \subseteq L$ and φ a consistent formula in \mathcal{L}_L . The following are equivalent:

- (i) $\vdash_{L^\neg} \varphi \leftrightarrow \neg\neg\varphi$
- (ii) φ is \mathcal{ST} -projective in L^\neg

Proof. (i) \implies (ii) follows from Lemma 4.4. For (ii) \implies (i), by Theorem 2.15, in L^\neg we have that $\varphi \Vdash \bigvee_{i \in I} \neg\varphi_i$ for some formulas $\{\neg\varphi_i \mid i \in I\}$. By a similar argument to that in the proof of “(iii) \implies (i)” of Lemma 4.6, we obtain in L^\neg that $\varphi \Vdash \neg\varphi_i$ for some $i \in I$, which implies that $\vdash_{L^\neg} \varphi \leftrightarrow \neg\neg\varphi$. \square \square

5 Structural completeness of the logics

In this section we prove the main results of our paper, namely that the three propositional logics of dependence \mathbf{PD} , InqL , and \mathbf{PT} are \mathcal{F} -structurally complete and that the negative variants of logics extending ND are hereditarily \mathcal{ST} -structurally complete. In both cases the proof of the fact is based on the existence, for every formula φ , of certain \mathcal{S} -projective formulas φ_i such that $\varphi \sqcup_{i \in I} \varphi_i$, where in the first case \mathcal{S} consists of all flat substitutions and in the second case of all stable ones. As mentioned in Remark 2.8, it is not hard to prove by the same methods that also the logic \mathbf{PID} is \mathcal{F} -structurally complete, where \mathbf{PID} is an extension of propositional intuitionistic dependence logic PID in the same manner as \mathbf{PT} is an extension of PT .

Definition 5.1. Let L be a logic, and \mathcal{S} a set of \vdash_L -substitutions. A rule φ/ψ of L is said to be \mathcal{S} -admissible, in symbols $\varphi \vdash_{\mathcal{S}} \psi$, if for all $\sigma \in \mathcal{S}$, $\vdash_L \sigma(\varphi) \implies \vdash_L \sigma(\psi)$.

In case \mathcal{S} is the set of all substitutions, we write \vdash for $\vdash_{\mathcal{S}}$, and such a rule is called an *admissible rule*.

Definition 5.2. A logic L is said to be \mathcal{S} -structurally complete if every \mathcal{S} -admissible rule of L is derivable in L , i.e., $\varphi \vdash_{\mathcal{S}} \psi \iff \varphi \vdash_L \psi$. In case \mathcal{S} is the set of all substitutions and L is \mathcal{S} -structurally complete, we say that L is *structurally complete*.

Informally, a rule is admissible in a logic L if its addition to the logic does not change the theorems that are derivable. Clearly, if \mathcal{S} is a set of L -substitutions, then $\varphi \vdash_L \psi \implies \varphi \sim_{\mathcal{S}}^L \psi$ for all formulas φ and ψ in \mathcal{L}_L . In particular, by Theorem 3.7, all derivable rules of **PD** and **InqL** are \mathcal{F} -admissible in the logics. A logic that is \mathcal{S} -structurally complete has no nontrivial \mathcal{S} -admissible rules: all such rules are derivable in the logic. Classical logic is structurally complete, but intuitionistic logic is not, as neither are many other intermediate logics. The well-known example showing that intuitionistic logic is not structurally complete uses *Harrop's Rule*:

$$\varphi \rightarrow \psi \vee \theta \vdash_{\text{IPC}} (\varphi \rightarrow \psi) \vee (\varphi \rightarrow \theta) \text{ but } \varphi \rightarrow \psi \vee \theta \not\vdash_{\text{IPC}} (\varphi \rightarrow \psi) \vee (\varphi \rightarrow \theta).$$

Recall the definition of \sqcup just below Lemma 2.4: $\varphi \sqcup_i \varphi_i$ holds if and only if $\varphi_i \models \varphi$ for all i , and for all teams X : $X \models \varphi$ implies $X \models \varphi_i$ for some i .

Lemma 5.3. For any L which is an intermediate theory or one of **PD**, **InqL** or **PT**, and any set \mathcal{S} of L -substitutions, if for every consistent formula φ in \mathcal{L}_L there exists a finite set $\{\varphi_i \mid i \in I\}$ of \mathcal{S} -projective formulas in \mathcal{L}_L such that $\varphi \sqcup_{i \in I} \varphi_i$, then L is \mathcal{S} -structurally complete.

Proof. We show that every \mathcal{S} -admissible rule $\varphi \sim_{\mathcal{S}}^L \psi$ of L is derivable, i.e., $\varphi \vdash_L \psi$. If φ is inconsistent, then clearly $\varphi \vdash_L \perp \vdash_L \psi$. Now assume that φ is consistent. By assumption there exists a finite set $\{\varphi_i \mid i \in I\}$ of \mathcal{S} -projective formulas such that $\varphi \sqcup_{i \in I} \varphi_i$. Let $\sigma_i \in \mathcal{S}$ be the projective unifier of φ_i . Thus $\vdash_L \sigma_i(\varphi_i)$. Hence $\vdash_L \sigma_i(\varphi)$ for all $i \in I$. From $\varphi \sim_{\mathcal{S}}^L \psi$ we derive $\vdash_L \sigma_i(\psi)$ for each $i \in I$. Since σ_i is a projective unifier for φ_i , we have that $\varphi_i, \sigma_i(\psi) \vdash_L \psi$. It follows that $\varphi_i \vdash_L \psi$ for each $i \in I$. Therefore $\varphi \vdash_L \psi$. □ □

Theorem 5.4. **PD**, **InqL** and **PT** are \mathcal{F} -structurally complete.

Proof. By Theorem 2.12 for the extended logics, Lemmas 4.6 and 5.3. □ □

Let L be an intermediate theory/logic and \mathcal{S} a set of \vdash_L -substitutions. We say that L is *\mathcal{S} -hereditarily structurally complete* if for any intermediate theory L' such that $L \subseteq L'$ and \mathcal{S} is a set of $\vdash_{L'}$ -substitutions, L' is \mathcal{S} -structurally complete. In case \mathcal{S} is the class of all substitutions of L , then we say that L is *hereditarily structurally complete*. It is known that none of **ND**, **KP** and **ML** are hereditarily structurally complete.

Theorem 5.5. For any intermediate logic L such that $\text{ND} \subseteq L$, its negative variant L^\neg is \mathcal{ST} -hereditarily structurally complete. In particular, ND^\neg , KP^\neg and ML^\neg are \mathcal{ST} -hereditarily structurally complete.

Proof. By Theorem 2.15, Lemmas 4.7 and 5.3. □ □

6 Concluding remarks

We have shown that the three propositional logics of dependence, **PD**, **PT**, **InqL**, are structurally complete with respect to flat substitutions and that the negative variant of

every intermediate logic that is an extension of ND is hereditarily structurally complete with respect to stable substitutions. In particular, ND^\neg , KP^\neg and ML^\neg are ST -hereditarily structurally complete. The reason for this are the strong normal forms that hold in these logics or theories. In this aspect they resemble classical logic, with its disjunctive normal form, that is also hereditarily structurally complete.

Apart from [20] there has not been much research on admissibility on intermediate theories that are not intermediate logics, and for propositional logics of dependence the above results are the first of that kind. Thus, naturally, many questions remain open. We discuss several of them.

Theorem 5.5 states that the negative variants of extensions of ND are hereditarily structurally complete. It follows from results of Maximova and Prucnal that any structurally complete intermediate logic with the disjunction property contains KP and is contained in ML, and in [25], which recaptures these results, it is moreover shown that KP itself is not structurally complete. The same holds for ND, since it is properly contained in KP. One wonders whether the fact that the negative variants of ND and KP are structurally complete could shed some light on admissibility in the original logics.

In this paper the results on admissibility are with respect to sets of substitutions, such as the flat and the stable substitutions. There exist logics for which establishing whether admissibility has certain properties, such as decidability, seems hard. These problems are often considered only for admissibility with respect to all substitutions, but one could start with smaller sets of substitutions, which may be easier to deal with, and although certain properties, such as decidability of admissibility, do not transfer from a smaller set of substitutions to its extensions, understanding a restricted case may still help understanding the general case.

On a more abstract level, there are two definitions of admissibility in the literature that in most instances amount to the same notion. Although intuitively clear, the proper connection between the two is not completely straightforward [14, 19]. It is mostly considered only for admissibility with respect to the set of all substitutions. It would be nice to see whether this connection can be generalized to admissibility with respect to any set of substitutions.

The results obtained in this paper made essential use of the disjunctive normal form of formulas of propositional logics of dependence. It is known from the literature that modal dependence logic and propositional independence logic both have a similar disjunctive normal form [26, 18, 5]. We conjecture that the argument in this paper may apply to these two logics and lead to similar results.

The authors would like to thank Ivano Ciardelli, Dick de Jongh and Jouko Väänänen for useful discussions on the topic of this paper.

References

- [1] Burgess, J.P.: A remark on henkin sentences and their contraries. *Notre Dame Journal of Formal Logic* **44**(3), 185–188 (2003)
- [2] Chagrov, A., Zakharyashev, M.: *Modal Logic*. Oxford University Press, USA (1997)

- [3] Ciardelli, I.: Dependency as Question Entailment. in: H.V.S. Abramsky, J. Kontinen, J. Väänänen (ed.), *Dependence Logic: Theory and Application*, in: *Progress in Computer Science and Applied Logic*, Birkhauser, (2016, to appear)
- [4] Ciardelli, I., Roelofsen, F.: Inquisitive logic. *Journal of Philosophical Logic* **40**(1), 55–94 (2011)
- [5] Ebbing, J., Hella, L., Meier, A., Müller, J.S., Virtema, J., Vollmer, H.: Extended modal dependence logic. In: 20th International Workshop, WoLLIC 2013, Proceedings, *Lecture Notes in Computer Science*, vol. 8071, pp. 126–137. Springer Berlin Heidelberg (2013)
- [6] Ghilardi, S.: Unification in intuitionistic logic. *Journal of Symbolic Logic* **64**, 859–880 (1999)
- [7] Hella, L., Luosto, K., Sano, K., Virtema, J.: The Expressive Power of Modal Dependence Logic. in: *Advances in Modal Logic 10, Invited and Contributed Papers from the Tenth Conference on Advances in Modal Logic*, College Publications, London, 294–312 (2014)
- [8] Henkin, L.: Some remarks on infinitely long formulas. In: *Infinistic Methods, Proceedings Symposium Foundations of Mathematics*, pp. 167–183. Pergamon, Warsaw (1961)
- [9] Hintikka, J.: *The Principles of Mathematics Revisited*. Cambridge University Press (1998)
- [10] Hintikka, J., Sandu, G.: Informational independence as a semantical phenomenon. In: R.H. J. E. Fenstad I. T. Frolov (ed.) *Logic, Methodology and Philosophy of Science*, pp. 571–589. Amsterdam: Elsevier (1989)
- [11] Hodges, W.: Compositional semantics for a language of imperfect information. *Logic Journal of the IGPL* **5**, 539–563 (1997)
- [12] Hodges, W.: Some strange quantifiers. In: J. Mycielski, G. Rozenberg, A. Salomaa (eds.) *Structures in Logic and Computer Science: A Selection of Essays in Honor of A. Ehrenfeucht, Lecture Notes in Computer Science*, vol. 1261, pp. 51–65. London: Springer (1997)
- [13] Iemhoff, R.: On the admissible rules of intuitionistic propositional logic. *Journal of Symbolic Logic* **66**, 281–294 (2001)
- [14] Iemhoff, R.: Consequence relations and admissible rules. Tech. Rep. 314, Department of Philosophy, Utrecht University (2013)
- [15] Jeřábek, E.: Admissible rules of modal logics. *Journal of Logic and Computation* **15**, 411–431 (2005)
- [16] Kontinen, J., Väänänen, J.: A remark on negation in dependence logic. *Notre Dame Journal of Formal Logic* **52**(1), 55–65 (2011)
- [17] Kuusisto, A.: A Double Team Semantics for Generalized Quantifiers logic. *Journal of Logic, Language and Information* **24**(2), 149–191 (2015)
- [18] Lohmann, P., Vollmer, H.: Complexity results for modal dependence logic. *Studia Logica* **101**(2), 343–366 (2013)
- [19] Metcalfe, G.: Admissible rules: From characterizations to applications. In: *Proceedings of WoLLIC 2012, LNCS*, vol. 7456, pp. 56–69. Springer (2012)
- [20] Miglioli, P., Moscato, U., Ornaghi, M., Quazza, S., Usberti, G.: Some results on intermediate constructive logics. *Notre Dame Journal of Formal Logic* **30**(4), 543–562 (1989)
- [21] Rozière, P.: *Regles admissibles en calcul propositionnel intuitionniste*. Ph.D. thesis, Université Paris VII (1992)
- [22] Rybakov, V.: *Admissibility of Logical Inference Rules*. Elsevier (1997)
- [23] Sano, K., Virtema, J.: Axiomatizing Propositional Dependence Logics *Proceedings of the 24th EACSL Annual Conference on Computer Science Logic*, 292–307 (2015)
- [24] Väänänen, J.: *Dependence Logic: A New Approach to Independence Friendly Logic*. Cambridge: Cambridge University Press (2007)
- [25] Wojtylak, P.: On a problem of H. Friedman and its solution by T. Prucnal. *Reports on Mathematical Logic* **38**, 69–86 (2004)
- [26] Yang, F.: On extensions and variants of dependence logic. Ph.D. thesis, University of Helsinki (2014)

- [27] Yang, F., Väänänen, J.: Propositional Logics of Dependence, *Annals of Pure and Applied Logic*, **167**(7), 557-589 (2016)