

# Developing sensor technologies to inform breeding approaches to reduce damaging behaviour in laying hens and pigs: The GroupHouseNet approach

T.B. Rodenburg<sup>1</sup>, J. Bennewitz<sup>2</sup>, E.N. de Haas<sup>1</sup>, L. Košťál<sup>3</sup>, K. Pichová<sup>3</sup>, D. Piette<sup>4</sup>, J. Tetens<sup>5</sup>, B. Visser<sup>6</sup>, B. De Klerk<sup>7</sup>, M. van der Sluis<sup>8,1</sup>, L.E. van der Zande<sup>9</sup>, J. Siegford<sup>10</sup>, M. Toscano<sup>11</sup>, T. Norton<sup>4</sup>, O. Guzhva<sup>12</sup> and E.D. Ellen<sup>8</sup>

<sup>1</sup>Faculty of Veterinary Medicine, Utrecht University, Utrecht, Netherlands; <sup>2</sup>Institute of Animal Science, University of Hohenheim, Hohenheim, Germany; <sup>3</sup>Slovak Academy of Sciences, Institute of Animal Biochemistry and Genetics, Ivanka pri Dunaji, Slovakia; <sup>4</sup>M3-BIORES, KU Leuven, Leuven, Belgium; <sup>5</sup>Functional Breeding Group, Department of Animal Sciences, Georg-August University, Göttingen, Germany; <sup>6</sup>Hendrix Genetics, Boxmeer, Netherlands; <sup>7</sup>Cobb Vantress, Boxmeer, Netherlands; <sup>8</sup>Animal Breeding and Genomics, Wageningen University & Research, Wageningen, Netherlands; <sup>9</sup>Adaptation Physiology Group, Wageningen University & Research Wageningen, Netherlands; <sup>10</sup>Department of Animal Science, Michigan State University, East Lansing, USA; <sup>11</sup>ZTHZ Veterinary Public Health Institute, Animal Welfare Division, University of Bern, Zollikofen, Switzerland; <sup>12</sup>Dept. Biosystems and Technology, Swedish University of Agricultural Sciences, Alnarp, Sweden  
Corresponding author: t.b.rodenburg@uu.nl

## Abstract

The European COST Action GroupHouseNet aims to provide synergy for preventing damaging behaviour in group-housed pigs and laying hens. One area of focus of this network is how genetic and genomic tools can be used to breed animals that are less likely to develop damaging behaviour directed at their pen-mates. Reducing damaging behaviour in large groups is a challenge, because it is difficult to identify and monitor individual animals. With the current developments in sensor technologies and animal breeding, there is the possibility to identify individual animals, monitor individual behaviour, and link this information to the genotype. Using a combination of sensor technologies and genomics enables us to select against damaging behaviour in pigs and laying hens.


**Keywords:** damaging behaviour, genetic selection, automatic tracking

## Introduction

The European COST Action GroupHouseNet ([www.grouphousenet.eu](http://www.grouphousenet.eu)) aims to provide synergy for preventing damaging behaviour in group-housed pigs and laying hens. One area of focus of this network is how genetic and genomic tools can be used to breed animals that are less likely to develop damaging behaviour directed at pen-mates. The behaviours we are focussing on are feather pecking in laying hens (Rodenburg *et al.*, 2013) and tail biting in pigs (Valros *et al.*, 2015). As both animal species are kept in groups, identifying actual performers of this behaviour (peckers and biters) at the individual level remains challenging. At the same time individual tracking is pivotal for breeding approaches. Using traditional behavioural observations is possible, but time consuming and costly. Here, we propose that a combination of sensor technologies and genomic methods should be used as a more feasible strategy to select against damaging behaviour in laying hens and pigs (Rodenburg *et al.*, 2017a). We compare different sensor technologies that can be used to identify individual animals for breeding, discuss the identification of indicator traits using data from sensor technologies, and discuss applications to animal breeding.

## Sensor technology

With the current developments in sensor technologies, breeding for laying hens and pigs that show less damaging behaviour by selecting animals using sensor data might offer solutions to these welfare challenges. We propose using a combination of sensor technology and genomic methods to solve this issue (Figure 1).



**Figure 1.** Overview of the approach to derive individual sensor and genomic data from group-housed pigs and laying hens and then combine this information to develop a genomic profile of individuals with the desired behavioural phenotype

We compare different sensor technologies (like ultra-wideband, RFID, computer vision) that can be used for detection of damaging behaviour or related behavioural traits (proxy measures). In laying hens, using video tracking and computer vision is challenging, given the small size of the animals and the large similarity between animals. When evaluating the sensor technologies used to this point, for laying hens RFID (Richards *et al.*, 2011) and accelerometer-based (Quwaider *et al.*, 2010) approaches seem most promising. Using UWB tracking, a specific type of RFID tracking using active tags on the animals, it was possible to distinguish feather peckers from non-feather peckers based on activity levels (Rodenburg *et al.*, 2017b). In pigs, computer vision is already used to record technical performance and there seems to be potential for expanding this approach to the recording of damaging behaviour (D'Eath *et al.*, 2018; Mittek *et al.*, 2018). Using computer vision, one of the main challenges is to link the correct identity to each individual and to maintain this link between identity and video image throughout the tracking period. Here, a combination between video tracking and passive RFID systems seems promising, as the RFID system can be used to re-assign the correct identity to individual animals. If sensor signatures and genomic fingerprints of individual animals can be combined, this would greatly improve our possibilities to reduce damaging behaviour through genetic selection.

### Linking sensor information to genetic information

We are now at a point where both sensor technology and genomics approaches have the potential to provide a large amount of data at the level of the individual animal, which can be used to understand and selectively breed against damaging behaviours, such as feather pecking (FP) in laying hens. For example, the high and low FP lines (Kjaer, 2017), selected on whether they show high or low FP behaviour, have been characterised in genomic and transcriptomic studies. These studies have added to our knowledge of the mechanisms underlying FP behaviour and can also be used to record genomic profiles of individual birds. Similarly, using sensor technology, we can now record detailed information on hens from the lines, creating an individual behavioural profile, describing a hen's activity, location and proximity to other individuals (Rodenburg *et al.*, 2017b; Rufener *et al.*, 2018). If we use both sensor and genomic technological approaches in a breeding population, we can link the genomic data to the behavioural data, and define the genomic profile of individuals that show the desired behaviour (for example, low or no damaging behaviour). As a prerequisite,

however, we need to determine genetic parameters for each indicator trait derived from sensor data, especially the genetic correlations between indicator and target trait. Once thought impossible, this approach may now be feasible, because breeding companies have begun to genotype their breeding stock routinely and they are also investing in methods for automatic phenotyping. Once the desired genomic profile has been defined, we can test whether selecting for this profile will reduce damaging behaviour by breeding a next generation based on genomic selection and then phenotyping this generation with the same tools that were used to phenotype the parent stock. We feel that a combined sensor and genomics approach has great promise to select against complex behavioural traits that involve multiple individual animals in a group, such as damaging behaviour in pigs and laying hens.

## Conclusions

Reducing damaging behaviour is an important goal for commercial poultry and pig production. The current developments in animal breeding and Precision Livestock Farming offer solutions to reduce damaging behaviour. We argue that a combined sensor and genomics approach has great promise to select against complex behavioural traits, especially when combining sensors like computer vision and RFID tracking.

## Acknowledgements

This article is based upon work from COST Action CA 15134 GroupHouseNet, supported by COST (European Cooperation in Science and Technology). Support was provided to J.M. Siegford as an AgBioResearch supported faculty member of Michigan State University by the National Institute of Food and Agriculture, U.S. Department of Agriculture, Hatch projects #1002990 and #1010765.

## References

- D'Eath, R.B., Jack, M., Futro, A., Talbott, D., Zhu, Q.M., Barclay, D., and Baxter, E.M. (2018). Automatic early warning of tail biting in pigs: 3D cameras can detect lowered tail posture before an outbreak. *PLoS One*, 13: 18.
- Kjaer, J.B. (2017). Divergent selection on home pen locomotor activity in a chicken model: Selection program, genetic parameters and direct response on activity and body weight. *PLoS ONE*, 12, e0182103.
- Mittek, M., Psota, E.T., Carlson, J.D., Pérez, L.C., Schmidt, T., and Mote, B. (2018). Tracking of group-housed pigs using multi-ellipsoid expectation maximisation. *IET Computer Vision*, 12: 121–128.
- Quwaider, M.Q., Daigle, C.L., Biswas, S.K., Siegford, J.M., and Swanson, J.C. (2010). Development of a wireless body-mounted sensor to monitor location and activity of laying hens in a non-cage housing system. *Transactions of the ASABE* 53, 1705–1713.
- Richards, G.J., Wilkins, L.J., Knowles, T.G., Booth, F., Toscano, M.J., Nicol, C.J., and Brown, S.N. (2011). Continuous monitoring of pop hole usage by commercially housed free-range hens throughout the production cycle. *Veterinary Record*, 169: 338-U358.
- Rodenburg, T.B., van Krimpen, M.M., de Jong, I.C., de Haas, E.N., Kops, M.S., Riedstra, B.J., Nordquist, R.E., Wagenaar, J.P., Bestman, M., and Nicol, C.J. (2013). The prevention and control of feather pecking in laying hens: identifying the underlying principles. *World's Poultry Science Journal*, 69: 361–374.
- Rodenburg, T.B., Bennewitz, J., de Haas, E.N., Kostal, L., Pichova, K., Piette, D., Tetens, J., van der Eijk, J.A.J., Visser, B., and Ellen, E.D. (2017a). The use of sensor technology and genomics to breed for laying hens that show less damaging behaviour. In: Berckmans, D., Keita, A. (Eds.), *Precision Livestock Farming '17*, Nantes, France, pp. 532–541.
- Rodenburg, T.B., van der Eijk, J.A.J., van Mil, B., Pichova, K., and de Haas, E.N. (2017b). PhenoLab: automatic recording of location, activity and proximity in group-housed laying hens. In: Berckmans, D., Keita, A. (Eds.), *Precision Livestock Farming '17*, Nantes, France, pp. 269–275.

- Rufener, C., Berezowski, J., Maximiano Sousa, F., Abreu, Y., Asher, L., and Toscano, M.J. (2018). Finding hens in a haystack: Consistency of movement patterns within and across individual laying hens maintained in large groups. *Scientific Reports* 8, 12303.
- Valros, A., Palander, P., Heinonen, M., Munsterhjelm, C., Brunberg, E., Keeling, L., and Piepponen, P. (2015). Evidence for a link between tail biting and central monoamine metabolism in pigs (*Sus scrofa domestica*). *Physiology & Behavior*, 143: 151–157.

# Precision Livestock Farming '19

Papers presented at the  
9<sup>th</sup> European Conference on Precision Livestock Farming  
Cork, Ireland  
26-29 August '19

***Edited by:***

B. O'Brien  
D. Hennessy  
L. Shalloo

