

Identification of New Pathogens in the Intraocular Fluid of Patients With Uveitis

JOLANDA D.F. DE GROOT-MIJNES, LENNEKE DE VISSER, STEPHANIE ZUURVEEN, ROALDY A. MARTINUS, RENÉ VÖLKER, NINETTE H. TEN DAM-VAN LOON, JOKE H. DE BOER, GINA POSTMA, RAOUL J. DE GROOT, ANTON M. VAN LOON, AND ANIKI ROTHOVA

- **PURPOSE:** To determine infectious causes in patients with uveitis of unknown origin by intraocular fluids analysis.
- **DESIGN:** Case-control study.
- **METHODS:** Ocular fluids from 139 patients suspected of infectious uveitis, but negative for herpes simplex virus, varicella-zoster virus, cytomegalovirus, and *Toxoplasma gondii* by polymerase chain reaction and/or antibody analysis in intraocular fluids, were assessed for the presence of 18 viruses and 3 bacteria by real-time polymerase chain reaction (PCR). The ocular fluids from 48 patients with uveitis of known etiology or with cataract were included as controls.
- **RESULTS:** Positive PCR results were found for Epstein-Barr virus, for rubella virus, and for human herpesvirus 6 each in 1 patient and for human parechovirus in 4 patients. Of the human parechovirus-positive patients, 1 was immunocompromised and had panuveitis. The other 3 patients were immunocompetent and had anterior uveitis, all with corneal involvement.
- **CONCLUSIONS:** Human parechovirus might be associated with infectious (kerato)uveitis. (Am J Ophthalmol 2010;150:628–636. © 2010 by Elsevier Inc. All rights reserved.)

UVEITIS CAN BE OF INFECTIOUS OR NONINFECTIOUS origin. Infections are thought to cause approximately 20% to 25% of cases; about 30% are associated with a noninfectious systemic disease. Although for patient management and the efficacy of treatment, the differential diagnosis is crucial, in more than half of the uveitis patients the underlying cause remains unknown.

See Accompanying Editorial on page 595.

Accepted for publication May 19, 2010.

From the Departments of Virology (J.D.F.d.G.-M., L.d.V., S.Z., R.A.M., R.V., A.M.v.L.) and Ophthalmology (J.D.F.d.G.-M., L.d.V., N.H.t.D.-v.L., J.H.d.B., G.P., A.R.), University Medical Center Utrecht, Utrecht, The Netherlands; and the Virology Section (R.J.d.G.), Department of Immunology and Infectious Diseases, Faculty of Veterinary Medicine, Utrecht University, Utrecht, The Netherlands.

Inquiries to Jolanda D.F. de Groot-Mijnes, Department of Virology, University Medical Center Utrecht, Heidelberglaan 100, G04.614, 3584 CX Utrecht, The Netherlands; e-mail: J.D.FdeGroot@umcutrecht.nl

The pathogens most commonly associated with infectious uveitis in immunocompetent patients are *Toxoplasma gondii*, herpes simplex virus (HSV), and varicella-zoster virus (VZV). In recent years, a few other infectious agents have been implicated in the etiology of uveitis, most notably rubella virus and cytomegalovirus (CMV).^{1–7} CMV is currently recognized as the most common cause of uveitis in immunocompromised patients.

In this study we performed an extensive search for infectious agents that cause uveitis but so far have escaped attention. Aqueous humor samples from 139 uveitis patients were analyzed retrospectively in our laboratory by available real-time polymerase chain reaction (PCR) assays for a variety of viruses and bacteria. Included were pathogens previously associated with uveitis (enteroviruses, Epstein-Barr viruses [EBV], human herpesvirus 6 [HHV6], influenza virus, rubella virus, *Mycoplasma pneumoniae*) and those causing conjunctivitis and/or keratitis or encephalitis (adenoviruses, coronaviruses, enteroviruses, influenza virus, parainfluenzaviruses, human parechovirus [HPeV], respiratory syncytial virus, *Chlamydia pneumoniae*, *Chlamydia trachomatis*).^{1,5,8–30} Our findings suggest that human parechovirus may be involved in the pathogenesis of infectious (kerato)uveitis.

METHODS

- **PATIENTS AND SAMPLES:** Ocular fluid samples analyzed in this study were from 629 uveitis patients who visited the ophthalmology clinic of the University Medical Center Utrecht from October 1, 2001 until June 30, 2006 and were suspected of infectious uveitis. The patients were classified using the uveitis nomenclature according to the recommendations of the Standardization of Uveitis Nomenclature working group 2005.³¹ All patients had undergone the uveitis screening consisting of erythrocyte sedimentation rate, red and white blood cell counts, determination of serum angiotensin-converting enzyme levels, serologic tests for syphilis, and chest radiography. Selected patients also underwent serologic testing for *Borrelia burgdorferi*. For all 629

TABLE 1. General Characteristics of Uveitis Patients and Controls for Which the Intraocular Fluids Were Analyzed for the Presence of New Pathogens

	N	Immunocompromised	Gender (M:F)	Mean Age \pm SD
Patients				
Anterior uveitis	49	2 (4%)	29:20	50.8 \pm 16.7
Panuveitis/posterior uveitis	90	8 (9%)	46:44	48.9 \pm 18.3
Controls				
Ocular toxoplasmosis	13	0	7:6	47.2 \pm 15.4
HSV anterior uveitis	10	0	5:5	44.1 \pm 22.1
Rubella virus-associated Fuchs heterochromic uveitis syndrome	14	0	10:4	42.3 \pm 16.5
Cataract	11	0	6:5	71.3 \pm 15.8

F = female; HSV = herpes simplex virus; M = male; SD = standard deviation.

patients aqueous sampling was performed for diagnostic purposes. The samples were stored at -80°C within 5 hours of collection before processing for laboratory analysis. Initial analysis was performed for HSV, VZV, and in the case of posterior uveitis also for *Toxoplasma* and CMV, by PCR and by Goldmann-Witmer coefficient (GWC), to determine intraocular antibody production. Of the 629 patients, 486 were negative for the above mentioned agents. A sufficient amount of ocular fluid remained for this study in 139 of these cases. Forty-nine patients had anterior uveitis (AU) and 90 had posterior uveitis (PU) or panuveitis (Table 1). Of the 49 AU patients, 2 were immunocompromised as a result of immunosuppressive medications (1 for lethal midline granuloma and the other after allogeneic stem cell transplantation for hematologic malignancy). Of the 90 patients with PU and panuveitis, 8 were immunocompromised, 5 of whom had acquired immunodeficiency syndrome and 3 of whom received immunosuppressive drugs (Table 1).

The remainders of ocular fluid samples from patients with PCR- and/or GWC-confirmed infectious uveitis (ocular toxoplasmosis, $n = 13$; HSV anterior uveitis, $n = 10$; rubella virus-associated Fuchs heterochromic uveitis syndrome (FHUS), $n = 14$) and from patients with cataract in the absence of intraocular inflammation ($n = 11$) served as controls.

• **NUCLEIC ACID ISOLATION AND REAL-TIME PCR:** The ocular fluid samples were analyzed for the presence of adenovirus, EBV, HHV6, *Mycoplasma pneumoniae*, *Chlamydia pneumoniae*, and *Chlamydia trachomatis* DNA and of coronaviruses 229E, OC43, and NL63, enteroviruses, human metapneumovirus, influenza A and B virus, parainfluenza virus 1 to 4, HPeV, respiratory syncytial virus A and B, and rubella virus RNA. If not done previously, samples from patients with anterior uveitis were also analyzed for CMV and *Toxoplasma*. DNA and RNA were extracted from 30 μL of ocular fluid using the MagNa Pure LC Total Nucleic Acid isolation kit

(Roche, Mannheim, Germany). To monitor the quality of the extraction and the subsequent amplification procedure, a standard dose of phocine herpesvirus type 1 and encephalomyocarditis virus was added to each sample as an internal control prior to extraction.³²⁻³⁴ Nucleic acid was collected in a volume of 240 μL . For detection of RNA viruses, copyDNA (cDNA) was produced by mixing 40 μL of extracted nucleic acid with 60 μL of reverse transcriptase mix (Taqman, reverse transcription reagents; Applied Biosystems, Foster City, California, USA) and incubating the mixture for 10 minutes at 25°C and 30 minutes at 48°C . The cDNA synthesis reaction was stopped by incubating for 5 minutes at 95°C . Per amplification reaction 10 μL of extracted nucleic acid (for DNA detection) or 10 μL of cDNA (for RNA detection) was used. Real-time PCR assays were performed on an ABI Prism 7700 sequence detection system (Applied Biosystems, Branchburg, New Jersey, USA). For *Chlamydia trachomatis*, 25 μL of extracted nucleic acid was analyzed using the Cobas Amplicor *Chlamydia trachomatis* detection kit according to the instructions of the manufacturer (Roche). All samples were examined once. In case of positive outcomes the real-time PCR reaction was repeated. Two human parechovirus-positive samples were confirmed by nucleic acid sequencing. Samples for which the internal control was inhibited were excluded. The primers and probes used are listed in Table 2.

• **ANTIBODY DETECTION ASSAYS:** Intraocular production of antibody against rubella virus (Goldmann-Witmer coefficient) was assessed as described previously.¹ Serum and intraocular immunoglobulin (Ig)G titers against HHV6 were determined using the Biotrin International Human Herpes Virus 6 IgG immunofluorescence assay (Dublin, Ireland). Serum and intraocular IgG against EBV was determined using the Panbio VCA IgG ELISA (Grenoble, France).

TABLE 2. Primers, Probes, and Sensitivities of the Real-Time Polymerase Chain Reaction Assays Used to Identify New Pathogens in the Intraocular Fluids of Patients With Unidentified Uveitis

Pathogen	Primers/Probes	Sequence 5' to 3'	Sensitivity	References
Adenoviruses	Forward	TTT GAG GTG GA(C/T) CC(A/C) ATG GA TTT GAG GTG GA(C/T) CC(A/C) ATG GA	100 copies/mL	51
	Reverse	AGA A(G/C)G G(G/C)G T(A/G)C GCA GGT A AGA A(G/C)G GTG T(A/G)C GCA GAT A		
	Probe	FAM-ACC ACG TCG AAA ACT TCG AA-MGBNFQ FAM-ACC ACG TCG AAA ACT TCA AA-MGBNFQ FAM-ACA CCG CGG CGT CA-MGBNFQ		
Coronavirus 229E	Forward	CAG TCA AAT GGG CTG ATG CA	ND	53
	Reverse	CAA AGG GCT ATA AAG AGA ATA AGG TAT TCT		
	Probe	FAM-CCC TGA CGA CCA CGT TGT GGT TCA -TAMRA		
Coronavirus NL63	Forward	AAG GGT TTT CCA CAG CTT GCT AAA GGT TTT CCA CAG CTT GCT	ND	53
	Reverse	ATC ACC CAC TTC ATC AGT GCT AAC		
	Probe	FAM-TCA CTA TCA AAG AAT AAC GCA GCC TGA TTA GGA A -TAMRA		
Coronavirus OC43	Forward	CGA TGA GGC TAT TCC GAC TAG GT	ND	53
	Reverse	CCT TCC TGA GCC TTC AAT ATA GTA ACC		
	Probe	FAM-TCC GCC TGG CAC GGT ACT CCC T -TAMRA		
Enteroviruses	Forward	TCC TCC GGC CCC TGA	ND	50
	Reverse	AAT TGT CAC CAT AAG CAG CCA GAT TGT CAC CAT AAG CAG CCA		
	Probe	FAM-CGG AAC CGA CTA CTT TGG GTG ACC GT -TAMRA FAM-CGG AAC CGA CTA CTT TGG GTG TCC GT -TAMRA		
Epstein-Barr virus	Forward	GGA ACC TGG TCA TCC TTT GC	50 copies/mL	42
	Reverse	ACG TGC ATG GAC CGG TTA AT		
	Probe	FAM-CGC AGG CAC TCG TAC TGC TCG CT -TAMRA		
Human herpesvirus 6	Forward	GAA GCA GCA ATC GCA	160 copies/mL	42
	Reverse	ACA CA ATG TAA CTC		
	Probe	FAM-AAC CCG TGC GCC GCT -TAMRA		
Human metapneumovirus	Forward	CAT ATA AGC ATG CTA TAT TAA AAG AGT CTC	ND	49
	Reverse	CCT ATT TCT GCA GCA TAT TTG TAA TCA G		
	Probe	FAM-TG(C/T) AAT GAT GAG GGT GTC ACT GCG GTT G -TAMRA		
Influenza virus A	Forward	AAG ACC AAT CCT GTC ACC TCT GA	ND	52
	Reverse	CAA AGC GTC TAC GCT GCA GTC C		
	Probe	FAM-TTT GTG TTC ACG CTC ACC GTG CC -TAMRA		
Influenza virus B	Forward	AAA TAC GGT GGA TTA AAC AAA AGC AA	ND	52
	Reverse	CCA GCA ATA GCT CCG AAG AAA		
	Probe	FAM- CAC CCA TAT TGG GCA ATT TCC TAT GGC -TAMRA		
Parainfluenza virus 1	Forward	TGA TTT AAA CCC GGT AAT TTC TCA T	ND	51
	Reverse	CCT TGT TCC TGC AGC TAT TAC AGA		
	Probe	FAM- ACG ACA ACA GGA AAT C -TAMRA		
Parainfluenza virus 2	Forward	AGG ACT ATG AAA ACC ATT TAC CTA AGT GA	ND	51
	Reverse	AAG CAA GTC TCA GTT CAG CTA GAT CA		
	Probe	FAM- ATC AAT CGC AAA AGC TGT TCA GTC ACT GCT ATA C -TAMRA		
Parainfluenza virus 3	Forward	TGA TGA AAG ATC AGA TTA TGC AT	ND	51
	Reverse	CCG GGA CAC CCA GTT GTG		
	Probe	FAM-TGG ACC AGG GAT ATA CTA CAA AGG CAA AAT AAT ATT TCT C -TAMRA		

Continued on next page

TABLE 2. Primers, Probes, and Sensitivities of the Real-Time Polymerase Chain Reaction Assays Used to Identify New Pathogens in the Intraocular Fluids of Patients With Unidentified Uveitis (*Continued*)

Pathogen	Primers/Probes	Sequence 5' to 3'	Sensitivity	References
Parainfluenza virus 4	Forward	CAA ATG ATC CAC AGC AAA GAT TC	ND	51
	Reverse	ATG TGG CCT GTA AGG AAA GCA		
	Probe	FAM- GTA TCA TCA TCT GCC AAA TCG GCA ATT AAA CA -TAMRA		
Human parechovirus	Forward	TGC AAA CAC TAG TGG TA(A/T) GGC CC	1 TCID50/mL	Forthcoming
	Reverse 1	TCA GAT CCA TAG TG(C/T) CAC TTG TTA CCT		
	Reverse 2	TCA GAT CCA CAG TGT CTC TTG TTA CCT		
	Probe	FAM- CGA AGG ATG CCC AGA AGG TAC CCG -TAMRA		
Respiratory syncytial virus A	Forward	AGA TCA ACT TCT GTC ATC CAG CAA	ND	54
	Reverse	TTC TGA ACA TCA TAA TTA GGA GTA TCA AT		
	Probe	FAM- CAC CAT CCA ACG GAG CAC AGG AGA T -TAMRA		
Respiratory syncytial virus B	Forward	AAG ATG CAA ATC ATA AAT TCA CAG GA	ND	54
	Reverse	TGA TAT CCA GCA TCT TTA AGT ATC TTT ATA GTG		
	Probe	FAM- TCC CCT TCC TAA CCT GGA CAT AGC ATA TAA CAT ACC T -TAMRA		
Rubella virus	Forward	CAC GCC GCA CGG ACA	1.7 PFU/mL	Forthcoming
	Reverse 1	CAC CGG GAC TG(C/T) TG(A/G) TTG C		
	Reverse 2	CAC CGG GAC TGT TGG TTG C		
	Probe	FAM- AGG TCC CGC CCG AC- MGBNFQ		
<i>Mycoplasma pneumoniae</i>	Forward	GGT CAA TCT GGC GTG CAT CT	50 CCU/mL	34
	Reverse	TGG TAA CTG CCC CAC AAG C		
	Probe	FAM- TCC CCC GTT GAA AAA GTG AGT GGG T -TAMRA		
<i>Chlamydia pneumoniae</i>	Forward	TCC GCA TTG CTC AGC C	4.9 IFU/mL	34
	Reverse	AAA CAA TTT GCA TGA AGT CTG AGA A		
	Probe	FAM-TAA ACT TAA CTG CAT GGA ACC CTT CTT TAC TAG G -TAMRA		

CCU = color-changing units; FAM = 6-carboxyfluorescein fluorescent label; IFU = inclusion-forming units; MGBNFQ = minor groove binding nonfluorescent quencher; ND = not determined; PFU = plaque-forming units; TAMRA = tetramethylrhodamine quencher; TCID50 = tissue culture infectious dose 50.

RESULTS

IN NONE OF THE OCULAR FLUIDS THE INTERNAL CONTROL was inhibited. In the patient samples positive PCR reactions were found for Epstein-Barr virus (n = 1), rubella virus (n = 1), human herpesvirus 6 (n = 1), and human parechovirus (n = 4). The PCR reactions for all other pathogens were negative. All control samples were negative except for 3; 2 toxoplasma chorioretinitis control samples were positive for EBV and 1 sample positive for rubella virus intraocular antibody production also tested PCR-positive for rubella virus RNA.

The patients with uveitis of unknown cause and a positive PCR result for rubella virus, HHV6, and human parechovirus are described below.

• **RUBELLA VIRUS (CASE 1):** A 40-year-old female patient complained of gradual decrease of visual acuity in the right eye (OD). Her medical history included pneumotho-

rax and bilateral pneumonia many years ago, but she had no signs of systemic disease.

The visual acuity of the OD was 20/80. The anterior chamber and vitreous of the OD revealed cells, but no synechiae. There was a subcapsular posterior cataract, fine keratic precipitates (KPs) and vitreous opacities. The retina was normal. The left eye (OS) had full visual acuity and no abnormalities on examination. Uveitis screening results were within normal limits. The clinical diagnosis of Fuchs heterochromic uveitis syndrome was made and a cataract extraction with implantation of an intraocular lens was performed as well as pars plana vitrectomy for vitreous opacities. On examination of the vitreous, there was no evidence for systemic and/or intraocular infection using PCR and GWC for CMV, HSV, VZV, *Toxoplasma*, *Borrelia burgdorferi*, and *Bartonella henselae*. Microbiological cultures were negative and cytologic examination revealed no malignant cells. By PCR, rubella virus was detected in the vitreous fluid. Subsequent antibody analysis for rubella

TABLE 3. Clinical Data of Uveitis Patients With a Human Parechovirus-Positive Polymerase Chain Reaction on Intraocular Fluid

Case	Gender	Age	Immune Status	Location	Unilateral or Bilateral	Cells Ant ^a	Cells Post ^a	Corneal Involvement	Retinitis	Vasculitis	Papillitis
3	M	54	Normal	Anterior	Unilateral	1+	No	Corneal scar	No	No	No
4	M	53	Normal	Anterior	Unilateral	No	No	Corneal infiltrate	No	No	No
5	F	73	Normal	Anterior	Unilateral	No	No	Corneal edema and keratic precipitates	No	No	No
6	M	37	HIV-positive	Panuveitis ^b	Unilateral	3+	2+	Corneal edema and keratic precipitates	Yes	Yes	Yes

Ant = anterior; F = female; M = male; Post = posterior.

^aGrading of cells was performed as recommended by Jabs and associates.³¹

^bThis patient was also diagnosed with neurosyphilis.

virus revealed the presence of intraocular IgG, but the GWC was negative (ratio 2.02). However, in comparison with HSV, VZV, CMV en *Toxoplasma*, intraocular antibody production against rubella virus was elevated.

• **HUMAN HERPESVIRUS-6 (CASE 2):** A 42-year-old man was referred because of decrease in visual acuity of his OD and floaters since 3 months prior. His medical history was not contributory and the patient used no medications. Uveitis screening results were within normal range. Remarkable was the heterochromia of his eyes, present since childhood. There was no serologic evidence for an active infection with CMV, HSV, VZV, *Toxoplasma*, or *Treponema pallidum*. *Borrelia burgdorferi* serum IgG and immunoblot were positive; however, a distinction between a past and an ongoing infection could not be made.

On ocular examination, the visual acuity of the OD was 20/25; the cornea revealed the presence of keratic precipitates, but the anterior chamber was clear. There were no synechiae, but several small noduli were present on the pupillary edge of the iris. Cataract was not observed. Funduscopy of the OD revealed vitreous cells and several peripheral snowballs. The fundoscopic findings were normal. The OS had full visual acuity; however, some peripheral vitreous opacities were observed.

Because of the possible (previous) infection with *Borrelia*, the patient was treated with intravenous ceftriaxone and additionally with periocular steroids, but with no effect. Diagnostic vitrectomy was performed and cytologic and microbiological examinations did not reveal a cause of his uveitis. Vitreous analysis was negative for CMV, HSV, VZV, and *Borrelia*, both by PCR and by GWC. The rubella virus GWC was negative (ratio 2.68), although intraocular IgG was detected and the GWC was elevated in comparison to CMV, HSV, and VZV. Therefore, rubella virus-associated FHUS could not be excluded. Three months after vitrectomy the patient regained full visual acuity, although the keratic precipitates in his OD remained. Retrospectively, the vitreous fluid appeared to be positive for HHV6 by PCR. Immunofluorescence assay demonstrated that the patient was seropositive for HHV6.

• **HUMAN PARECHOVIRUS:CASE 3.** A 54-year-old male patient was referred to our center with anterior uveitis of 2 years' duration in his pseudophakic OS. Twenty-nine years prior, the patient underwent cataract extraction and implantation of an iris-clip lens in his OS because of previous trauma. On ocular examination, the visual acuity of the OS was hand movements (Table 3). A central corneal scar was seen, cells were present in the anterior chamber, and the vitreous was clear. Funduscopy revealed no abnormalities. The OD had full visual acuity and no abnormalities. Both eyes had normal intraocular pressure. The patient had no systemic complaints and used no medications. Uveitis screening results were within normal limits.

Uveitis was clinically attributed to irritation caused by the iris-clip lens, which was therefore surgically removed. A vitreous sample was obtained during surgery. Analysis of the ocular fluid was negative for HSV and VZV, both by PCR and GWC, and for CMV and *Toxoplasma* by PCR, but was retrospectively positive for human parechovirus by PCR. Ocular examination 4 months after removal of the intraocular lens revealed a quiet OS.

Case 4. A 53-year-old male was referred because of persistent keratitis of his OS. It was thought to be caused by HSV, but the patient did not respond to systemic and topical treatments with acyclovir and valacyclovir. His previous medical and ophthalmic histories were unremarkable. Uveitis screening did not reveal any abnormalities.

On ocular examination, the visual acuity of the OS was 20/100. An infiltrate in the upper part of the cornea with epithelial defect and sporadic cells in the anterior chamber were observed (Table 3). Corneal sensitivity was normal. The vitreous was clear and the retinal findings were unremarkable. Intraocular pressure was normal. The OD had full visual acuity and no abnormalities. Aqueous analysis was negative for HSV, VZV, and *Toxoplasma* by both PCR and GWC and for CMV by PCR; however, retrospectively PCR was positive for human parechovirus. The patient was treated with antibiotic eye ointment and the corneal lesion and anterior uveitis slowly became quiet.

Case 5. A 73-year-old female patient with an ophthalmologic history of cataract extraction in both eyes at the age of 70 was referred to our institution because of secondary glaucoma in the OD. The patient had no systemic complaints and used no medications. The OD revealed pupillary seclusion with an intraocular pressure (IOP) of 50 mm Hg. On examination corneal edema and keratic precipitates were noted in the OD (Table 3). The iris revealed atrophic areas. Fluorescein angiography revealed slight optic disc leakage and cystoid macular edema in the OD. The IOP initially normalized with laser iridotomy and local treatment; however, intermittent periods with IOP elevations up to 50 mm Hg were regularly encountered and trabeculectomy was required. The presumed diagnosis included low-grade *Propionibacterium* endophthalmitis and other various causes of hypertensive uveitis (Table 3). Screening examinations were within normal limits as well as serology for *Borrelia* and *Bartonella*. Aqueous sampling was performed and PCR was negative for HSV, VZV, CMV, and *Toxoplasma* and GWC was negative for HSV and VZV. Cultures were negative for *Propionibacterium*. Retrospectively, the patient was found positive for human parechovirus by PCR.

Case 6. A 37-year-old male patient was referred because of panuveitis with a focal chorioretinitis lesion in his OS since 3 months prior. At that time visual acuity in his OS

decreased to finger counting. On examination KPs and cells in the anterior chamber and vitreous were noted with an active lesion located in the periphery of the retina (Table 3). Intraocular pressure was normal. Fluorescein angiography demonstrated optic disc leakage and vasculitis with changes of retinal pigment epithelium in the mid-periphery of the retina. The OD had full visual acuity and no abnormalities. The patient had no systemic complaints and used no medications. The presumed diagnosis of toxoplasma chorioretinitis was made. Extensive screening for panuveitis revealed positive human immunodeficiency virus (HIV) serology, an HIV RNA plasma load of 69 700 copies/mL, 619 CD4 cells/mL, and positive syphilis serology (*Treponema pallidum* hemagglutination [TPHA] >1:2560 and a Venereal Diseases Research Laboratory [VDRL] test result of 1:256). Aqueous analysis was negative for *Treponema pallidum* by PCR and for HSV, VZV, CMV, and *Toxoplasma* both by PCR and GWC. The aqueous was, however, retrospectively positive for human parechovirus by PCR. Although both the aqueous and cerebrospinal fluid analyses were negative for syphilis, the tentative diagnosis of ocular syphilis was made and the patient was treated with systemic penicillin. Antiretroviral treatment was considered not necessary at that time. Ocular inflammation subsided slowly, visual acuity increased to 20/20, and the eye remained quiet during 2 years of follow-up.

DISCUSSION

IN 139 OCULAR FLUID SAMPLES FROM PATIENTS WITH UVEITIS that were analyzed with a panel of 21 real-time PCRs, we found positive results in 7 cases (5%): 1 case with EBV, 1 case with rubella virus, 1 case with human herpesvirus 6, and 4 cases with human parechovirus.

EBV, the causative agent of infectious mononucleosis and several malignancies, has been implicated as a possible cause of uveitis and in primary ocular non-Hodgkin lymphoma of the central nervous system.^{16,26,35} In our study EBV was detected in the ocular fluid of 1 patient with anterior uveitis of unknown cause and in 2 patients with toxoplasma chorioretinitis. IgG analysis in these 3 patients did not show any evidence of intraocular antibody production against EBV. The presence of EBV genome in the eyes of patients with various causes of uveitis was demonstrated previously and was found independent of the clinical diagnosis. The clinical significance of this phenomenon has not yet been established.^{18,36,37} Moreover, as in our study, Ongkosuwito and associates found EBV also in ocular fluids from patients with laboratory-confirmed toxoplasmosis and in ocular fluids of patients without ocular inflammation.¹⁸ Apparently, PCR detection of EBV in ocular fluids should be interpreted with caution, and may in most cases be considered an epiphenomenon, mostly likely attributable to the presence of EBV in B cells present in the inflamed eye. Further studies combining PCR and

intraocular antibody production analysis are required to determine whether EBV is a true cause of intraocular inflammation.

One patient was PCR-positive for rubella virus, the causative agent of German measles and congenital rubella syndrome.³⁸ Rubella virus has been associated with FHUS and FHUS-like uveitis.^{1,5,39,40} This patient was clinically diagnosed with incomplete FHUS, as she did not have iris heterochromia or atrophy.

HHV6 is a beta-herpesvirus and the causative agent of roseola infantum (or exanthema subitum), a childhood disease.⁴¹ In addition, HHV6 is being recognized as an important opportunistic infection following bone marrow and/or stem cell transplantation.⁴² Our HHV6 PCR-positive patient (Case 2) was neither a child nor immunosuppressed. Antibody analysis of the ocular fluid by immunofluorescence assay did not reveal the presence of intraocular IgG against HHV6. However, absence of intraocular antibody production does not necessarily exclude intraocular infection. Previously, we reported that by simultaneous use of PCR and GWC in immunocompetent patients a diagnosis by PCR only was established in 9% of cases.² Moreover, de Boer and associates found that in patients with presumed herpetic anterior uveitis, PCR was more frequently positive than GWC.⁴³ HHV6 has been implicated in ocular inflammation, most notably when the posterior part of the eye was affected.^{14,15,17,19} Our patient had anterior uveitis with heterochromia that had been present since childhood. Heterochromia is classically associated with FHUS but can develop in other viral infections such as HSV or VZV.⁴⁴ The detection of HHV6 in the eye might not be a clinically relevant finding; however, like other (herpes)viruses, HHV6 can reside latently in cells of the lymphoid and myeloid lineage, and it may have entered the inflamed eye via immune cells, similar to HIV and possibly EBV.^{6,18,19,36,45} The role of HHV6 as a cause of anterior uveitis is inconclusive and further studies are required.

Human parechoviruses belong to the genus parechovirus within the family of *Picornaviridae*. They may cause gastroenteritis, encephalitis, and flaccid paralysis in young children, but rarely in adults.^{29,46} Ocular diseases attributable to other picornavirus infections, particularly enteroviruses such as echoviruses 11 and 19 and coxsackieviruses, have been published, but an association between parechoviruses and ocular disease has not been reported yet.⁸⁻¹² In this study the ocular fluids of 4 patients with undiagnosed

unilateral uveitis were PCR-positive for HPeV. Unfortunately, intraocular antibody production could not be established as appropriate serologic assays are not available and there was not enough ocular fluid left to perform viral culture.

One patient (Case 6) with intraocular HPeV was immunocompromised (Table 3). He had been diagnosed with active syphilis, but there were no indications for neurosyphilis or ocular syphilis. However, upon treatment with penicillin his ocular condition improved. As this patient was HIV-positive, multiple uveitis entities may have contributed to ocular disease.

The other 3 patients all had unilateral anterior uveitis with corneal involvement and cells in the anterior chamber, which suggested an ocular viral infection (Table 3). Further research is required to determine if HPeV plays a role in the pathogenesis of infectious (anterior) uveitis. It is surprising to find a virus that is associated with disease in children in the eyes of adults. However, other infections of childhood are known to cause intraocular disease in adults, as is the case for rubella virus.^{1,5,39}

The vast majority (132/139 [95%]) of ocular fluid samples were negative by PCR analysis for multiple viruses and bacteria. There are several explanations for this result. First, the uveitis might be of noninfectious origin. Second, the number of DNA or RNA copies present in the ocular samples may have been below the detection level. This may be attributable to the low amount of input nucleic acid, which is inherent to diagnostic assays with intraocular fluid. Alternatively, the time of sampling may have been such that the causative agent had already been cleared from the eye. In this case, other diagnostic approaches may be more useful, such as the detection of intraocular antibody production by Goldmann-Witmer coefficient calculation. It is known, also for systemic and neurologic viral infections, that a PCR assay is most sensitive early in infection, whereas antibody can be detected over a much longer period of time and thus provide a wider window of detection.^{2,43,47,48} Finally, it may be that other pathogens not covered by our assays were involved in these cases.

Our study addressed multiple infectious causes in patients with undiagnosed uveitis and revealed a possible new cause of infectious uveitis. Further investigations are required to narrow the diagnostic gap in patients with presumed infectious uveitis.

PUBLICATION OF THIS STUDY WAS SUPPORTED BY THE DR. F.P. FISCHER FOUNDATION, AMERSFOORT, THE NETHERLANDS (J.D.F.d.G.-M., L.d.V., and S.J.Z.) and by the Foundation for Dutch Ophthalmic Research (SNOO), Rotterdam, The Netherlands (S.J.Z.). The authors indicate no financial conflict of interest. Involved in design of study (J.D.F.d.G.-M., A.M.v.L., A.R.); conduct of study (L.d.V., S.J.Z., R.A.M., R.V., R.J.d.G.); collection, management, analysis, and interpretation data (J.D.F.d.G.-M., L.d.V., N.H.t.D.-v.L., J.H.d.B., G.P., R.J.d.G., A.M.v.L., A.R.); and preparation, review, or approval of manuscript (J.D.F.d.G.-M., L.d.V., N.H.t.D.-v.L., J.H.d.B., R.J.d.G., A.M.v.L., A.R.). This study was performed according to the tenets of the Declaration of Helsinki and with approval of the Institutional Review Board of the Medical Ethics Committee of the University Medical Center Utrecht, The Netherlands. The authors thank John W.A. Rossen for developing the human parechovirus real-time polymerase chain reaction.

REFERENCES

- de Groot-Mijnes JDF, de Visser L, Rothova A, Schuller M, van Loon AM, Weersink AJ. Rubella virus is associated with fuchs heterochromic iridocyclitis. *Am J Ophthalmol* 2006; 141(1):212–214.
- de Groot-Mijnes JDF, Rothova A, Van Loon AM, et al. Polymerase chain reaction and Goldmann-Witmer coefficient analysis are complimentary for the diagnosis of infectious uveitis. *Am J Ophthalmol* 2006;141(2):313–318.
- de Schryver I, Rozenberg F, Cassoux N, et al. Diagnosis and treatment of cytomegalovirus iridocyclitis without retinal necrosis. *Br J Ophthalmol* 2006;90(7):852–855.
- Markomichelakis NN, Canakis C, Zafirakis P, Marakis T, Mallias I, Theodossiadis G. Cytomegalovirus as a cause of anterior uveitis with sectoral iris atrophy. *Ophthalmology* 2002;109(5):879–882.
- Quentin CD, Reiber H. Fuchs heterochromic cyclitis: rubella virus antibodies and genome in aqueous humor. *Am J Ophthalmol* 2004;138(1):46–54.
- Rothova A, de Boer JH, Ten Dam-van Loon NH, et al. Usefulness of aqueous humor analysis for the diagnosis of posterior uveitis. *Ophthalmology* 2008;115(2):306–311.
- van Bortel LA, van der Lelij A, van der Meer J, Los LL. Cytomegalovirus as a cause of anterior uveitis in immunocompetent patients. *Ophthalmology* 2007;114(7):1358–1362.
- Förster W, Bialasiewicz AA, Busse H. Coxsackievirus B3-associated panuveitis. *Br J Ophthalmol* 1993;77(3):182–183.
- Hirakata K, Oshima T, Azuma N. Chorioretinitis induced by coxsackievirus B4 infection. *Am J Ophthalmol* 1990;109(2): 225–227.
- Kadrmas EF, Buzney SM. Coxsackievirus B4 as a cause of adult chorioretinitis. *Am J Ophthalmol* 1999;127(3):347–349.
- Lashkevich VA, Koroleva GA, Lukashev AN, Denisova EV, Katargina LA. Enterovirus uveitis. *Rev Med Virol* 2004; 14(4):241–254.
- Takeuchi M, Sakai J, Usui M. Coxsackievirus B4 associated uveoretinitis in an adult. *Br J Ophthalmol* 2003;87(4):501–502.
- Yoser SL, Forster DJ, Rao NA. Systemic viral infections and their retinal and choroidal manifestations. *Surv Ophthalmol* 1993;37(5):313–352.
- Maslin J, Bigaillon C, Froussard F, Enouf V, Nicand E. Acute bilateral uveitis associated with an active human herpesvirus-6 infection. *J Infect* 2007;54(4):e237–e240.
- Mechai F, Boutolleau D, Manceron V, et al. Human herpesvirus 6-associated retrobulbar optic neuritis in an HIV-infected patient: response to anti-herpesvirus therapy and long-term outcome. *J Med Virol* 2007;79(7):931–934.
- Mittra RA, Pulido JS, Hanson GA, Kajdacsy-Balla A, Brummitt CF. Primary ocular Epstein-Barr virus-associated non-Hodgkin's lymphoma in a patient with AIDS: a clinicopathologic report. *Retina* 1999;19(1):45–50.
- Moschetti D, Franceschini R, Vaccaro NM, et al. Human herpesvirus-6B active infection associated with relapsing bilateral anterior optic neuritis. *J Clin Virol* 2006;37(4): 244–247.
- Ongkosuwito JV, Van der Lelij A, Bruinenberg M, et al. Increased presence of Epstein-Barr virus DNA in ocular fluid samples from HIV negative immunocompromised patients with uveitis. *Br J Ophthalmol* 1998;82(3):245–251.
- Sugita S, Shimizu N, Kawaguchi T, Akao N, Morio T, Mochizuki M. Identification of human herpesvirus 6 in a patient with severe unilateral panuveitis. *Arch Ophthalmol* 2007;125(10):1426–1427.
- Arnold JJ, McIntosh ED, Martin FJ, Mense MA. A fifty-year follow-up of ocular defects in congenital rubella: late ocular manifestations. *Aust N Z J Ophthalmol* 1994;22(1):1–6.
- Gümbel HO, Lipke KJ, Schafer HG, Cinatl I, Hattenbach LO. Bilateral influenza-triggered panuveitis and subsequent therapy with amantadine and hyperimmunoglobulins. *Ophthalmologie* 2004;101(1):720–724.
- Haller-Schober EM, El-Shabrawi Y. Chlamydial conjunctivitis (in adults), uveitis, and reactive arthritis, including SARA. Sexually acquired reactive arthritis. *Best Pract Res Clin Obstet Gynaecol* 2002;16(16):815–828.
- Numazaki K, Chiba S, Aoki K, Suzuki K, Ohno S. Detection of serum antibodies to Chlamydia pneumoniae in patients with endogenous uveitis and acute conjunctivitis. *Clin Infect Dis* 1997;25(4):928–929.
- Weinstein O, Shneck M, Levy J, Lifshitz T. Bilateral acute anterior uveitis as a presenting symptom of Mycoplasma pneumoniae infection. *Can J Ophthalmol* 2006;41(5):594–595.
- Yashar SS, Yashar B, Epstein E, Viani RM. Uveitis associated with Mycoplasma pneumoniae meningitis. *Acta Ophthalmol Scand* 2001;79(1):100–101.
- Ritterband DC, Friedberg DN. Virus infections of the eye. *Rev Med Virol* 1998;8(4):187–201.
- Athanasu P, Anghelescu S, Predescu E, et al. Rapid detection by immunofluorescence of multiple viral infections in patients with keratitis. *Virologie* 1984;35(2):83–88.
- Fouchier RA, Rimmelzwaan GF, Kuiken T, Osterhaus AD. Newer respiratory virus infections: human metapneumovirus, avian influenza virus, and human coronaviruses. *Curr Opin Infect Dis* 2005;18(2):141–146.
- Benschop KS, Schinkel J, Minnaar RP, et al. Human parechovirus infections in Dutch children and the association between serotype and disease severity. *Clin Infect Dis* 2006;42(2):204–210.
- Fujishima H, Okamoto Y, Saito I, Tsubota K. Respiratory syncytial virus and allergic conjunctivitis. *J Allergy Clin Immunol* 1995;95(3):663–667.
- Jabs DA, Nussenblatt RB, Rosenbaum JT. Standardization of uveitis nomenclature for reporting clinical data. Results of the First International Workshop. *Am J Ophthalmol* 2005; 140(3):509–516.
- Niesters HG. Standardization and quality control in molecular diagnostics. *Expert Rev Mol Diagn* 2001;1(2):129–131.
- Stranska R, Schuurman R, de Vos M, van Loon AM. Routine use of a highly automated and internally controlled real-time PCR assay for the diagnosis of herpes simplex and varicella-zoster virus infections. *J Clin Virol* 2004;30(1):39–44.
- van de Pol AC, Wolfs TF, Jansen NJ, van Loon AM, Rossen JW. Diagnostic value of real-time polymerase chain reaction to detect viruses in young children admitted to the paediatric

- intensive care unit with lower respiratory tract infection. *Crit Care* 2006;10(2):R61.
35. Kutok JL, Wang F. Spectrum of Epstein-Barr virus-associated diseases. *Annu Rev Pathol* 2006;1:375–404.
 36. Matos K, Muccioli C, Belfort Junior R, Rizzo LV. Correlation between clinical diagnosis and PCR analysis of serum, aqueous, and vitreous samples in patients with inflammatory eye disease. *Arq Bras Oftalmol* 2007;70(1):109–114.
 37. Sugita S, Shimizu N, Watanabe K, et al. Use of multiplex PCR and real-time PCR to detect human herpes virus genome in ocular fluids of patients with uveitis. *Br J Ophthalmol* 2008;92(7):928–932.
 38. Banatvala JE, Brown DW. Rubella. *Lancet* 2004;363(9415):1127–1137.
 39. Birnbaum AD, Tessler HH, Schultz KL, et al. Epidemiologic relationship between fuchs heterochromic iridocyclitis and the United States rubella vaccination program. *Am J Ophthalmol* 2007;144(3):424–428.
 40. de Visser L, Braakenburg A, Rothova A, de Boer JH. Rubella virus-associated uveitis: clinical manifestations and visual prognosis. *Am J Ophthalmol* 2008;146(2):292–297.
 41. Levy JA. Three new human herpesviruses (HHV6, 7, and 8). *Lancet* 1997;349(9051):558–563.
 42. de Pagter PJ, Schuurman R, Meijer E, van Baarle D, Sanders EA, Boelens JJ. Human herpesvirus type 6 reactivation after haematopoietic stem cell transplantation. *J Clin Virol* 2008;43(4):361–366.
 43. de Boer JH, Verhagen C, Bruinenberg M, et al. Serologic and polymerase chain reaction analysis of intraocular fluids in the diagnosis of infectious uveitis. *Am J Ophthalmol* 1996;121(6):650–658.
 44. Harper S, Chorich L III, Foster C. Principles of diagnosis and therapy. In: Foster C, Vitale A, eds. *Diagnosis and Treatment of Uveitis*. Philadelphia: W.B. Saunders Company; 2002:79–103.
 45. Mitchell SM, Fox JD, Tedder RS, Gazzard BG, Lightman S. Vitreous fluid sampling and viral genome detection for the diagnosis of viral retinitis in patients with AIDS. *J Med Virol* 1994;43(4):336–340.
 46. Stanway G, Joki-Korpela P, Hyypia T. Human parechoviruses—biology and clinical significance. *Rev Med Virol* 2000;10(1):57–69.
 47. Fomsgaard A, Kirkby N, Jensen IP, Vestergaard BF. Routine diagnosis of herpes simplex virus (HSV) encephalitis by an internal DNA controlled HSV PCR and an IgG-capture assay for intrathecal synthesis of HSV antibodies. *Clin Diagn Virol* 1998;9(1):45–56.
 48. Furuta Y, Fukuda S, Suzuki S, Takasu T, Inuyama Y, Nagashima K. Detection of varicella-zoster virus DNA in patients with acute peripheral facial palsy by the polymerase chain reaction, and its use for early diagnosis of zoster sine herpette. *J Med Virol* 1997;52(3):316–319.
 49. Maertzdorf J, Wang CK, Brown JB, et al. Real-time reverse transcriptase PCR assay for detection of human metapneumoviruses from all known genetic lineages. *J Clin Microbiol* 2004;42(3):981–986.
 50. Nijhuis M, van Maarseveen N, Schuurman R, et al. Rapid and sensitive routine detection of all members of the genus enterovirus in different clinical specimens by real-time PCR. *J Clin Microbiol* 2002;40(10):3666–3670.
 51. van de Pol AC, van Loon AM, Wolfs TF, et al. Increased detection of respiratory syncytial virus, influenza viruses, parainfluenza viruses, and adenoviruses with real-time PCR in samples from patients with respiratory symptoms. *J Clin Microbiol* 2007;45(7):2260–2262.
 52. van Elden LJ, Nijhuis M, Schipper P, Schuurman R, van Loon AM. Simultaneous detection of influenza viruses A and B using real-time quantitative PCR. *J Clin Microbiol* 2001;39(1):196–200.
 53. van Elden LJ, van Loon AM, van Alphen F, et al. Frequent detection of human coronaviruses in clinical specimens from patients with respiratory tract infection by use of a novel real-time reverse-transcriptase polymerase chain reaction. *J Infect Dis* 2004;189(4):652–657.
 54. van Elden LJ, van Loon AM, van der Beek A, et al. Applicability of a real-time quantitative PCR assay for diagnosis of respiratory syncytial virus infection in immunocompromised adults. *J Clin Microbiol* 2003;41(9):4378–4381.