

Reviews of Geophysics

REVIEW ARTICLE

10.1029/2019RG000681

Key Points:

- Titration and charge balance alkalinity differ
- The impact of biogeochemical processes on pH depends on environmental conditions
- Ocean alkalinity budget is balanced when the additional alkalinity input from riverine particulate inorganic carbon and sedimentary sources is included

Supporting Information:

- Supporting Information S1
- Software S1

Correspondence to:

J. J. Middelburg,
j.b.m.middelburg@uu.nl

Citation:

Middelburg, J. J., Soetaert, K., & Hagens, M. (2020). Ocean alkalinity, buffering and biogeochemical processes. *Reviews of Geophysics*, 58, e2019RG000681. <https://doi.org/10.1029/2019RG000681>

Received 7 DEC 2019

Accepted 23 APR 2020

Accepted article online 9 JUN 2020

Ocean Alkalinity, Buffering and Biogeochemical Processes

Jack J. Middelburg¹ , Karline Soetaert² , and Mathilde Hagens³ 

¹Department of Earth Sciences, Geosciences, Utrecht University, Utrecht, The Netherlands, ²Department of Estuarine and Delta Systems, Royal Netherlands Institute for Sea Research (NIOZ Yerseke) and Utrecht University, Yerseke, The Netherlands, ³Soil Chemistry and Chemical Soil Quality, Wageningen University, Wageningen, The Netherlands

Abstract Alkalinity, the excess of proton acceptors over donors, plays a major role in ocean chemistry, in buffering and in calcium carbonate precipitation and dissolution. Understanding alkalinity dynamics is pivotal to quantify ocean carbon dioxide uptake during times of global change. Here we review ocean alkalinity and its role in ocean buffering as well as the biogeochemical processes governing alkalinity and pH in the ocean. We show that it is important to distinguish between measurable titration alkalinity and charge balance alkalinity that is used to quantify calcification and carbonate dissolution and needed to understand the impact of biogeochemical processes on components of the carbon dioxide system. A general treatment of ocean buffering and quantification via sensitivity factors is presented and used to link existing buffer and sensitivity factors. The impact of individual biogeochemical processes on ocean alkalinity and pH is discussed and quantified using these sensitivity factors. Processes governing ocean alkalinity on longer time scales such as carbonate compensation, (reversed) silicate weathering, and anaerobic mineralization are discussed and used to derive a close-to-balance ocean alkalinity budget for the modern ocean.

Plain Language Summary The ocean plays a major role in the global carbon cycle and the storage of anthropogenic carbon dioxide. This key function of the ocean is related to the reaction of dissolved carbon dioxide with water to form bicarbonate (and minor quantities of carbonic acid and carbonate). Alkalinity, the excess of bases, governs the efficiency at which this occurs and provides buffering capacity toward acidification. Here we discuss ocean alkalinity, buffering, and biogeochemical processes and provide quantitative tools that may help to better understand the role of the ocean in carbon cycling during times of global change.

1. Introduction

The ocean plays a major role in controlling atmospheric carbon dioxide and storage of anthropogenic carbon (Gruber et al., 2019). For the last decade, ocean uptake of anthropogenic carbon was 2.5 ± 0.6 Pg C year⁻¹, that is, about 23% of annual anthropogenic carbon emissions due to fossil fuels, cement production, and land-use change (Friedlingstein et al., 2019). The cumulative (1850–2019) total release of anthropogenic carbon was 655 ± 65 Pg C, of which 160 ± 20 Pg C (about 24%) has accumulated in the ocean (Friedlingstein et al., 2019). This crucial role of the ocean in attenuating the increase in atmospheric carbon dioxide, and thus global warming, is related to the large volume (and surface area) of the ocean and the reaction of dissolved carbon dioxide with water to form carbonic acid, a weak acid that dissociates to protons and the conjugated bases bicarbonate and carbonate, which are not directly exchangeable with the atmosphere (Butler, 1982). The redistributions among gaseous and dissolved carbon dioxide, carbonic acid, bicarbonate, and carbonate ions are governed by multiple co-occurring equilibria with the result that approximately 19 out of the 20 molecules of carbon dioxide entering the ocean are converted into bicarbonate and carbonate ions. The total amount of dissolved inorganic carbon (DIC) in the ocean is typically about 200 times that of dissolved carbon dioxide (Middelburg, 2019; Zeebe & Wolf-Gladrow, 2001).

This reequilibration following the principles of le Chatelier (1884) provides resistance to, but does not entirely eliminate, changes in ocean carbon chemistry. Oceanic uptake of anthropogenic carbon dioxide has caused increases in dissolved carbon dioxide and total inorganic carbon concentrations and decreases in carbonate ions and ocean pH, that is, ocean acidification (Gattuso & Hansson, 2011). Ocean acidification has consequences for further ocean carbon dioxide uptake, the precipitation and dissolution of carbonate minerals, and for the functioning and survival of marine organisms (Kroeker et al., 2013). It is therefore important that we understand and are able to quantify the buffering, that is, resistance, of the ocean in

©2020. The Authors.

This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

the changing world of the Anthropocene. Detailed understanding and quantification of how biogeochemical processes impact pH and marine carbon dioxide equilibria is pivotal to predicting the impact of ocean acidification on marine organisms, carbonate mineral precipitation and dissolution, (seasonal) variability in carbonate system parameters and the resilience of various ecosystem functions (Orr et al., 2018). Understanding is also required to use pH observations to infer the intensity and changes in biogeochemical processes and to evaluate the feasibility of ocean engineering options (Gattuso et al., 2018; Renforth & Henderson, 2017; Soetaert et al., 2007).

Although acid–base equilibria of simple solutions are well understood (Butler, 1964, 1982; Morel & Hering, 1993; Stumm & Morgan, 1981), the carbon dioxide system in seawater remains challenging because of the complexity of multiple equilibria (Zeebe & Wolf-Gladrow, 2001). Alkalinity, the excess of bases (proton acceptors) over acids (proton donors) in a solution (a complete definition is provided in section 2), is not only impacted by acid–base additions but also by redox reactions and mineral dissolution and precipitation. Oxidation reactions involving oxygen generally consume alkalinity, while anaerobic processes usually produce alkalinity. Dissolution of minerals is often accompanied by alkalinity generation. Alkalinity is a central concept in our treatment of the oceanic carbon dioxide system, because it is measurable, it remains unchanged with pressure and temperature (i.e., it is conservative), and it is governed by the net effect of multiple chemical equilibria and often needed to solve the mathematical equilibrium problem (Butler, 1982; Stumm & Morgan, 1981). However, there are multiple interpretations, and even definitions, of alkalinity that are not always used in a consistent way. One of the goals of this review is to clarify inconsistencies or sources of confusion, for example, the distinction between titration alkalinity (that can be measured) and charge balance alkalinity (that should be used to interpret biogeochemical processes in nature). Another goal is to discuss approaches to quantify the resistance (buffering) or its inverse, i.e. the sensitivity of the ocean carbon dioxide system and pH to change. While many geochemical and oceanographic studies mention ocean buffering, there are few where buffer and/or sensitivity factors are being used, except for the well-known Revelle factor expressing the sensitivity of $p\text{CO}_2$ to changes in DIC (Bolin & Eriksson, 1959; Revelle & Suess, 1957; Sarmiento & Gruber, 2006; Sundquist et al., 1979). This is surprising as rigorous treatments of buffering have been published a century ago (Koppel & Spiro, 1914; van Slyke, 1922).

Following a treatment of ocean alkalinity (section 2) and sensitivity and buffer factors (section 3), we will discuss the impact of biogeochemical processes on pH and $p\text{CO}_2$ (section 4), heterogeneous buffering, including carbonate compensation (section 5), and factors governing ocean alkalinity, including an alkalinity budget of the ocean (section 6). Basic terminology is explained in the Glossary. The supporting information accompanying this article contains three sections, and the R script used to generate figures and results.

2. Ocean Alkalinity

There is a long history from the eighteenth century observation that seawater is alkaline (Marsigli, 1725) to the modern concept of seawater alkalinity (Dickson, 1981; Zeebe & Wolf-Gladrow, 2001). The term alkalinity was already in use by chemists in the first half of the nineteenth century (e.g., Donovan, 1839) and utilized since in multiple disciplines, including medicine (Andral, 1850) and oceanography (Dittmar, 1884). Dickson (1992) provides an excellent historical account on the alkalinity concept in seawater and showed that it involved both advances in analytical procedures as well as the development of a chemical model for seawater. Rather than recapitulating the historical context, we believe it is instructive to formally distinguish between titration alkalinity, that is, total alkalinity, as defined by Dickson (1981) and the charge-balance alkalinity needed to quantify buffering and pH changes in natural environments. Observational and experimental studies in the ocean are normally based on titration alkalinity, but theoretical, modeling, and geological studies sometimes employ the charge balance approach (e.g., Broecker, 1974; Boudreau, 1996; Turchyn & DePaolo, 2019). The charge-balance alkalinity concept is often used in freshwater systems (with high concentrations of dissolved organic matter) and is also known as the excess negative charge (ENC; Soetaert et al., 2007) and linked to the explicit conservative expression of total alkalinity (Wolf-Gladrow et al., 2007; Zeebe & Wolf-Gladrow, 2001). This difference between titration alkalinity (TA) and charge-balance alkalinity

(CBA) is related to the equations used to solve the chemical equilibrium problem: The TA is based on a proton balance, while CBA is based on a charge balance closure (supporting information S1). Depending on the specific problem at hand and definition of the system, TA and CBA may differ or be identical. The lack of distinction between TA and CBA has caused confusion and discussion.

2.1. Titration Alkalinity

In 1981, Dickson defined the alkalinity (TA) as follows: “*The total alkalinity of a natural water is thus defined as the number of moles of hydrogen ion equivalent to the excess of proton acceptors (bases formed from weak acids with a dissociation constant $K \leq 10^{-4.5}$ and zero ionic strength) over proton donors (acids with $K > 10^{-4.5}$) in one kilogram of sample.*” The definition is stated in gravimetric units to remain independent of the temperature and pressure of the system. Furthermore, Dickson (1981) adopted a pK value of 4.5 as the reference level to distinguish between proton donors (acids with a dissociation constant $pK < 4.5$) and proton acceptors ($pK \geq 4.5$) to continue the common practice to match the reference level with the carbonic acid equivalence point of a titration.

Using this exact definition of alkalinity of Dickson (1981), it is straightforward to calculate the titration alkalinity for any system for which the contributing components are known and characterized in terms of dissociation constants. Figure 1a shows the distribution diagram of acid–base pairs (Bjerrum plot) for the carbonate system in seawater. For the $\text{CO}_2\text{--H}_2\text{O}$ system, at $\text{pH} = 4.5$, carbonic acid is by far the dominant species and used as reference. Referenced to this point, we then arrive at the proton balance, a mass balance for protons (see supporting information S1)

$$\text{H}^+ = \text{HCO}_3^- + 2\text{CO}_3^{2-} + \text{OH}^- \quad (1)$$

with proton donors on the left-hand side and proton acceptors on the right-hand side. The carbonate ion is counted twice because it is two protons below the reference level H_2CO_3 .

The titration alkalinity, that is, excess of proton acceptors over donors with respect to carbonic acid, the reference level, is then defined as follows

$$\text{TA} = \text{HCO}_3^- + 2\text{CO}_3^{2-} + \text{OH}^- - \text{H}^+ \quad (2)$$

Other acid–base systems can be included in the alkalinity expression. To this end, all chemical species in the solution have to be classified either as proton donor or acceptor relative to the zero level of protons for each acid–base system (Figure 1b). Dickson (1981) included fluoride, sulfate, borate, phosphate, silicate, ammonia, and hydrogen sulfide to eventually arrive at

$$\begin{aligned} \text{TA} = & \text{HCO}_3^- + 2\text{CO}_3^{2-} + \text{OH}^- + \text{B}(\text{OH})_4^- + \text{HPO}_4^{2-} + 2\text{PO}_4^{3-} + \text{H}_3\text{SiO}_4^- + \text{NH}_3 + \text{HS}^- + 2\text{S}^{2-} \\ & - \text{H}^+ - \text{HF} - \text{HSO}_4^- - \text{H}_3\text{PO}_4 \end{aligned} \quad (3)$$

This proton balance approach toward alkalinity allows an exact definition of alkalinity. Dickson (1981) focused on the quantification of alkalinity in seawater from titration data and, therefore, did not include the strong acids H_2SO_4 and HNO_3 nor HNO_2 and $\text{H}_2\text{SiO}_4^{2-}$. However, these can easily be included using the same approach (Soetaert et al., 2007; Wolf-Gladrow et al., 2007) and results in

$$\begin{aligned} \text{TA} = & \text{HCO}_3^- + 2\text{CO}_3^{2-} + \text{OH}^- + \text{B}(\text{OH})_4^- + \text{HPO}_4^{2-} + 2\text{PO}_4^{3-} + \text{H}_3\text{SiO}_4^- \\ & + 2\text{H}_2\text{SiO}_4^{2-} + \text{NH}_3 + \text{HS}^- + 2\text{S}^{2-} - \text{H}^+ - \text{HF} - \text{HSO}_4^- - 2\text{H}_2\text{SO}_4 - \text{H}_3\text{PO}_4 - \text{HNO}_2 - \text{HNO}_3 \end{aligned} \quad (4)$$

where H_2SO_4 and HNO_3 are zero for the pH values > 0 .

The titration alkalinity definition of Dickson (1981) is fully consistent with the conservation equation for total hydrogen ions (TOT) of Morel and Hering (1993), which is also based on a proton (mass) balance (supporting information S1). Specifically, $\text{TA} = -\text{TOT}$ when the components chosen are the reference level species at $\text{pH} = 4.5$.




Figure 1. Speciation, proton acceptor levels, and charge as a function of pH. (a) Bjerrum plot showing the distribution of carbonic acid, bicarbonate, and carbonate as a function of pH and the corresponding proton acceptor level (solid black line) and charge (red dashed line); (b) the proton acceptor level for selected acid–base pairs; (c) the charge for selected acid–base pairs. Details of the calculations are presented in the supporting information.

2.2. Charge Balance Alkalinity

Electrolyte solutions, including seawater, obey the electroneutrality condition, that is, the sum of negative and positive charges balances at the macroscale (Boudreau et al., 2004; Soetaert et al., 2007; Wolf-Gladrow et al., 2007). For seawater, we thus have to balance the sum of cation concentrations

$$\text{Na}^+ + 2 \text{Mg}^{2+} + 2 \text{Ca}^{2+} + \text{K}^+ + 2 \text{Sr}^{2+} + (\dots) + \text{NH}_4^+ + \text{H}^+ \quad (5a)$$

with the sum of anion concentrations

$$\begin{aligned} \text{Cl}^- + \text{Br}^- + (\dots) + \text{HCO}_3^- + 2\text{CO}_3^{2-} + \text{OH}^- + \text{B}(\text{OH})_4^- + \text{H}_2\text{PO}_4^- + 2\text{HPO}_4^{2-} + 3\text{PO}_4^{3-} \\ + \text{H}_3\text{SiO}_4^- + 2\text{H}_2\text{SiO}_4^{2-} + \text{HS}^- + 2\text{S}^{2-} + \text{F}^- + \text{HSO}_4^- + 2\text{SO}_4^{2-} + \text{NO}_2^- + \text{NO}_3^- \end{aligned} \quad (5b)$$

where the ellipses (...) stand for additional ions in solution. In charge conservation equations, ions are multiplied with their charge. This equation can be rearranged into a part that is conservative and a part that is not and involves species exchanging protons. Accordingly, when ignoring minor conservative species (...), the sum of strong base cations minus the sum of strong acid anions (i.e., excess positive charge of conservative ions)

$$\text{EPC} = \text{Na}^+ + 2\text{Mg}^{2+} + 2\text{Ca}^{2+} + \text{K}^+ + 2\text{Sr}^{2+} - \text{Cl}^- - \text{Br}^- \quad (6a)$$

should balance the excess negative charge (ENC) of nonconservative ions that are involved in proton exchange reactions

$$\begin{aligned} \text{ENC} = \text{HCO}_3^- + 2\text{CO}_3^{2-} + \text{OH}^- + \text{B}(\text{OH})_4^- + \text{H}_2\text{PO}_4^- + 2\text{HPO}_4^{2-} + 3\text{PO}_4^{3-} + \text{H}_3\text{SiO}_4^- \\ + .2\text{H}_2\text{SiO}_4^{2-} + \text{HS}^- + 2\text{S}^{2-} + \text{F}^- + \text{HSO}_4^- + 2\text{SO}_4^{2-} + \text{NO}_2^- + \text{NO}_3^- - \text{NH}_4^+ - \text{H}^+ \end{aligned} \quad (6b)$$

This excess negative charge (Soetaert et al., 2007) is also known as CBA. Hence, CBA is defined as the sum of nonconservative ions involved in proton exchange reactions that account for the difference between the sum of conservative cations and anions.

CBA (Equation 6b) and TA (Equation 4) are linked via

$$\text{TA} = \text{CBA} + \sum \text{NH}_3 - \sum \text{NO}_3 - \sum \text{NO}_2 - \sum \text{PO}_4 - 2\sum \text{SO}_4 - \sum \text{F} \quad (7)$$

where $\sum \text{NH}_3 = \text{NH}_3 + \text{NH}_4^+$, $\sum \text{NO}_3 = \text{NO}_3^- + \text{HNO}_3$, $\sum \text{NO}_2 = \text{NO}_2^- + \text{HNO}_2$, $\sum \text{PO}_4 = \text{H}_3\text{PO}_4 + \text{H}_2\text{PO}_4^- + \text{HPO}_4^{2-} + \text{PO}_4^{3-}$, $\sum \text{SO}_4 = \text{H}_2\text{SO}_4 + \text{HSO}_4^- + \text{SO}_4^{2-}$ and $\sum \text{F} = \text{HF} + \text{F}^-$ are the total concentrations of ammonia, nitrate, nitrite, phosphate, sulfate, and fluoride, respectively. This difference between titration and charge balance alkalinity is due to the charge of components at the reference pH level of 4.5. At pH 4.5, ammonia is present as ammonium (+1), while nitrate, nitrite, fluoride, and phosphate have an overall charge of -1 and sulfate is present as SO_4^{2-} with charge -2 (Figure 1c). Accordingly, dissolved inorganic carbon does not appear in Equation 7 because it is present as the uncharged carbon dioxide at pH 4.5. In other words, the difference between CBA and TA is caused by components for which the species used as zero proton level are charged (e.g., H_2PO_4^- is the zero-proton level for phosphate; Dickson, 1981).

Wolf-Gladrow et al. (2007) introduced the explicitly conservative expression of total alkalinity (TA_{ec}) that relates the sum of conservative cations and anions (i.e., excess positive charge; Equation 6a) and total concentrations of ammonia, nitrate, nitrite, phosphate, sulfate, and fluoride with titration alkalinity (TA)

$$\begin{aligned} \text{TA}_{\text{ec}} = \text{Na}^+ + 2\text{Mg}^{2+} + 2\text{Ca}^{2+} + \text{K}^+ + 2\text{Sr}^{2+} - \text{Cl}^- - \text{Br}^- + (\dots) \\ + \sum \text{NH}_3 - \sum \text{NO}_3 - \sum \text{NO}_2 - \sum \text{PO}_4 - 2\sum \text{SO}_4 - \sum \text{F} \end{aligned} \quad (8)$$

or alternatively formulated: $\text{TA}_{\text{ec}} = \text{EPC} + \text{TA} - \text{CBA}$.

This explicitly conservative form of alkalinity is equivalent to Dickson's expression (as $\text{EPC}-\text{CBA} = 0$) for titration alkalinity (Equation 4), but each single term is conservative to proton exchange and pressure and temperature changes (Wolf-Gladrow et al., 2007). Since charge balance alkalinity is directly related to the difference between conservative cations and anions, it is evident that CBA, and thus, also TA covary with salinity. Alkalinity and salinity are both affected to the same degree by processes that dilute or concentrate seawater, such as precipitation, evaporation, and melting or formation of ice.

2.3. Alternative Alkalinity and Related Expressions

The titration and charge balance alkalinity expressions (Equations 4 and 6b) are well defined and traceable to the use of a proton or charge balance (supporting information S1), but alternative expressions are often

used. There are a number of reasons for this. One, alkalinity and related concepts are used in multiple disciplines (e.g., chemistry, medicine, environmental engineering, ecology, geology, hydrology, limnology, and oceanography) with their own specific scientific traditions and terminology. Two, although many species are included in the formal definition of alkalinity in seawater, a few of these dominate by far and most others can be ignored as a first-order approximation. Carbonate alkalinity ($CA = HCO_3^- + 2CO_3^{2-}$) typically accounts for >95% of the total alkalinity in the ocean. Many studies (e.g., Broecker & Peng, 1982) use a simple form of alkalinity including only water and carbonate alkalinity terms (Equation 2). In seawater, a slightly more accurate expression is obtained when borate alkalinity is included as well. Zeebe and Wolf-Gladrow (2001) termed this alkalinity for most practical purposes (PA_{ZW-G})

$$PA_{ZW-G} = HCO_3^- + 2CO_3^{2-} + B(OH)_4^- + OH^- - H^+ = CA + \text{borate alkalinity} + \text{water alkalinity}. \quad (9)$$

PA_{ZW-G} is often used interchangeably with TA because it typically represents >99% of total alkalinity in oxygenated ocean surface waters. In anoxic waters and pore waters of marine sediments in which metabolites (ammonia, phosphate, sulfide, and silicate) have accumulated, some of these are then included in the operational definition of alkalinity for that system (Ben-Yaakov, 1973; Boudreau & Canfield, 1993; Hiscock & Millero, 2006). Finally, a major reason for alternative alkalinity expressions relates to application of the measurable property TA to biogeochemical processes that impact CBA because of electroneutrality constraints.

2.3.1. Use of Titration Alkalinity as Proxy for Charge Balance Alkalinity

Charge balance and titration alkalinity differ by the total amounts of nitrite, nitrate, ammonia, phosphate, sulfate, and fluoride (Equation 7; Soetaert et al., 2007; Wolf-Gladrow et al., 2007). Processes such as primary production, organic matter degradation, and nitrification involve these components and thus potentially impact alkalinity. Brewer and Goldman (1976) and Goldman and Brewer (1980) documented increases in alkalinity due to nitrate and phosphate uptake and a decrease in alkalinity due to ammonium uptake (see section 4). These alkalinity changes (ΔTA) due to biological consumption or production processes have to be taken into account when using measured TA values for quantification of calcium carbonate precipitation or dissolution. Specifically, Brewer et al. (1975) introduced the potential alkalinity change ($\Delta P.A.$) as a measure of calcite formation/dissolution

$$\Delta P.A. = \Delta TA + \Delta \sum NO_3 + \Delta \sum PO_4. \quad (10)$$

By comparing Equations 10 and 7, it is clear that potential alkalinity change is a proxy for CBA change based on measured nitrate, phosphate, and TA. However, only changes in nitrate and phosphate due to biological processes should be considered, not those due to physical mixing (Huang et al., 2015). Similarly, Kanamori and Ikegami (1982) identified the need to include nitrate, phosphate, and sulfate when using measured TA for calculating alkalinity changes due to calcium carbonate dynamics. Including nitrate, phosphate, and sulfate contributions in Redfield proportions would result in the following expression for potential alkalinity (P.A.)

$$P.A. = TA + a^* \sum NO_3 \quad (11)$$

where a varies from 1.06 (Chen, 1978) and 1.26 (Kanamori & Ikegami, 1982) to 1.36 (Wolf-Gladrow et al., 2007), depending on the Redfield ratios considered.

The Alk^* tracer (Carter et al., 2014) combines the potential alkalinity (Equation 11 with $a = 1.26$) with salinity normalization to single out the effect of calcium carbonate dynamics on alkalinity. Similarly, Feely et al. (2002) introduced the TA^* tracer which expresses the change in TA due to calcium carbonate dynamics: $TA^* = 0.5 (TA_s - TA_s^0) + 0.63(0.0941 \text{ AOU})$, where TA_s and TA_s^0 are the measured and preformed salinity-normalized TA, respectively, and AOU is the apparent oxygen utilization, introduced to correct for charges generated during organic matter dynamics. Finally, ecologists studying calcification by benthic communities (coral reefs and bivalves) often employ the alkalinity anomaly technique (Chisholm & Gattuso, 1991; Gazeau et al., 2015; Kinsey, 1978), which involves measurement of TA and correcting it with ammonium, nitrate, and phosphate for obtaining calcium carbonate dynamics.

2.3.2. Organic Alkalinity

While oceanographers usually measure TA and introduce empirical corrections to arrive at the CBA needed for quantitative applications (Equations 10 and 11), freshwater scientists studying soft natural waters have to use a charge balance of the major conservative ions because of an important contribution of organic acids (Hemond, 1990). Contrary to the inorganic acid–base species discussed before, dissolved organic compounds comprise a complex, heterogeneous group, which poses a challenge on classifying its acid–base properties and quantifying their contribution to TA (Hu, 2020). The composition and thus acid–base properties of dissolved organic compounds depend on whether these compounds are derived from locally produced organic matter or transported from adjacent terrestrial ecosystems (Leenheer & Croué, 2003). Phytoplankton-derived dissolved organic compounds are found to have two distinct proton binding sites with pK values of 4.4–4.9 and 6.1–6.9, respectively (Ko et al., 2016). In contrast, terrestrially derived organic matter is dominated by humic substances. These comprise a much wider range of proton binding sites, often described by carboxyl and phenolic groups having average pK values of about 3.7 ± 2.4 and about 12.5 ± 1.8 , respectively (Perdue et al., 1984). This wide range in pK values implies that at least part of the dissolved organic compounds, either of autochthonous or allochthonous origin, acts as proton acceptor at pK 4.5, thus contributing to TA despite being absent in Equation 4.

Substantial contributions of organic alkalinity to TA have been found in laboratory incubations (Ko et al., 2016), estuaries (Cai et al., 1998), sediment pore waters (Łukawska-Matuszewska, 2016; Łukawska-Matuszewska et al., 2018), coastal waters receiving high terrestrial inputs, like the Baltic Sea (Hammer et al., 2017; Kuliński et al., 2014) or salt marsh-influenced coastal waters (Song et al., 2020), and ocean waters (Fong & Dickson, 2019). Organic alkalinity is normally assessed by difference, that is, carbonate alkalinity is derived from two out of three other measurable parameters in the $\text{CO}_2\text{-H}_2\text{O}$ system (pH, DIC, or pCO_2), and organic alkalinity is calculated as the difference between TA measured and calculated from the contributions of the inorganic species following Equation 3. This method relies on at least one parameter (pH or pCO_2) which value is affected by the presence of organic compounds and, therefore, does not allow for an exact value of organic alkalinity. Back-titration methods to directly quantify organic alkalinity are used by others (Cai et al., 1998; Hernandez-Ayon et al., 2007; Muller & Bleie, 2008; Yang et al., 2015), showing no clear correlation with organic alkalinity estimated by difference (Song et al., 2020). Alternatively, chemical equilibrium models describing proton binding to humic substances, which are well-known in the freshwater community (Kinniburgh et al., 1999), can be coupled to inorganic carbonate system calculations (Ulfsbo et al., 2015).

2.3.3. Acid Neutralizing Capacity

Although not often used in oceanography, the term acid neutralizing capacity (ANC) is closely linked to titration alkalinity (TA). The ANC of a solution to the carbonic acid equivalent point of a titration is fully consistent with the Dickson (1981) definition of TA (Stumm & Morgan, 1981; Weber & Stumm, 1963). Other equivalence points are termed p-alkalinity (phenolphthalein endpoint of titration, corresponding to the proton balance of Equation 1.13 in supporting information S1) and caustic alkalinity, the reverse of acidity, with the proton balance: $\text{TOTH} = \text{OH}^- - \text{H}^+ - 2\text{H}_2\text{CO}_3 - \text{HCO}_3^-$ (Stumm & Morgan, 1981). Theoretically, one can use any expression for alkalinity to solve the $\text{CO}_2\text{-H}_2\text{O}$ system as long it is properly defined.

Some researchers distinguish between TA and ANC whether water samples are filtered or not, respectively (Asuero & Michałowski, 2011; Michałowski & Asuero, 2012). The chemical model underlying Dickson's TA only includes homogeneous reactions in solution and ignores proton exchange with particles and organisms. This implies that water samples for alkalinity should be filtered before titration because of potential proton exchange with the surface of phytoplankton, bacteria, and inorganic particles and the dissolution of suspended particulate inorganic carbon (Kim et al., 2006), and dedicated filtration methods have been developed (Bockman & Dickson, 2014). However, differences between filtered and unfiltered samples are often negligible (open ocean, Chanson & Millero, 2007; coastal systems, Hagens et al., 2015) but might be substantial in experimental systems with high densities of organisms or particles.

3. Buffering and Sensitivity Factors

Seawater is a solution with multiple weak acids and bases in contact with both the atmosphere and sediments containing minerals that have the potential to react when solution composition or physical

conditions change. Seawater is consequently well buffered, that is, able to resist changes by transferring protons. The response of a chemical equilibrium system to a perturbation follows the principle of le Chatelier. The original statement of Henry Louis le Chatelier (1884) “*Tout système en équilibre chimique stable soumis à l'influence d'une cause extérieure qui tend à faire varier soit sa température, soit sa condensation (pression, concentration, nombre de molécules dans l'unité de volume) dans sa totalité ou seulement dans quelques-unes de ses parties, ne peut éprouver que des modifications intérieures, qui, si elles se produisaient seules, amèneraient un changement de température ou de condensation de signe contraire à celui résultant de la cause extérieure.*” is often rephrased as follows: whenever a system in equilibrium is disturbed by changing the conditions, the positions of the equilibria shift in such a way that the effect of the change will be moderated.

In this section, we discuss the sensitivity and resistance of ocean chemistry to changes. It is instructive to distinguish between homogeneous reactions in solution and heterogeneous buffering involving interactions with particles (e.g., dissolution or precipitation of carbonate minerals modifying alkalinity). Homogeneous buffering takes place nearly instantaneously and is most relevant for quantifying and understanding the impact of biogeochemical processes on pH on short (hour-days) timescales (Egleston et al., 2010; Frankignoulle, 1994; Soetaert et al., 2007). Heterogeneous buffering reactions may involve very long time scales (months to millions of years) and will be discussed in section 5.

3.1. Buffer Capacity Systematics

Although the buffer capacity of seawater and its role in earth system science has been recognized in the first part of the twentieth century (Mitchell & Rakestraw, 1933; Thompson & Bonnar, 1931) and mathematical tools to quantify buffer efficiency have been developed a century ago (Koppel & Spiro, 1914; van Slyke, 1922), quantitative treatments of seawater buffering have historically received little attention, except for the homogeneous Revelle factor (Revelle & Suess, 1957) and the acid–base buffer capacity (van Slyke, 1922; Weber & Stumm, 1963). The acid–base buffer value β was originally defined for biological fluids by Koppel and Spiro (1914) but is commonly attributed to van Slyke (1922)

$$\beta = \frac{-dC_a}{dpH} \quad (12)$$

where C_a is the quantity of acid added to a solution. For seawater, TA is substituted for C_a and partial derivatives are used to indicate that other properties are kept constant during the titration

$$\beta = \frac{\partial TA}{\partial pH} \quad (13)$$

The buffer value β is always positive because every solution resists pH change according to the principle of le Chatelier. It is based on pH rather than proton concentrations because of historical reasons and laboratory procedures.

The Revelle factor (R ; Revelle & Suess, 1957; Bolin & Eriksson, 1959; Sundquist et al., 1979) expresses the sensitivity of pCO_2 to changes in DIC

$$R = \frac{\partial \ln pCO_2}{\partial \ln DIC} = \frac{DIC}{pCO_2} \left(\frac{\partial pCO_2}{\partial DIC} \right) \quad (14)$$

where use is made of the property $\partial \ln x = \frac{1}{x} \partial x$. This Revelle factor is limited to homogeneous systems, because the partial derivatives indicate that other variables such as alkalinity are kept constant. Sundquist and Plummer (1981) extended the homogeneous Revelle factor to allow for changes in alkalinity (e.g., due to calcification/dissolution)

$$R_{tot} = \frac{DIC}{pCO_2} \left(\frac{dpCO_2}{dDIC} \right) = \frac{DIC}{pCO_2} \left[\left(\frac{\partial pCO_2}{\partial DIC} \right)_{TA} + \left(\frac{\partial pCO_2}{\partial TA} \right)_{DIC} \cdot \frac{dTA}{dDIC} \right] \quad (15)$$

In seawater, the term $\left(\frac{\partial pCO_2}{\partial TA} \right)_{DIC}$ is negative, while $\frac{dTA}{dDIC}$ varies from zero (no TA change) to 2 when all changes in DIC are due to calcium carbonate dissolution; heterogeneous buffering thus lowers the Revelle factor (i.e., ocean buffering is larger when carbonate minerals are involved).

The Revelle and acid–base buffer factors are just two out of many ways to quantify the response of seawater to changes and some additional (chemical) buffer factors have been proposed (e.g., Frankignoulle, 1994; Frankignoulle et al., 1994; Egleston et al., 2010; Hagens & Middelburg, 2016a; Table 1). However, there are multiple inconsistencies in terminology and their relationships are not clear (Table 1). To clarify matters and link the various buffer capacities and factors in the literature, we present a systematic treatment involving partial derivatives as sensitivities. While the application of Le Chatelier's principle is straightforward for simple systems, it becomes difficult to predict the response of individual reactions when multiple reactions sharing ions are involved (Fishtik et al., 1995) and a sensitivity analysis is then useful. Sensitivities are also known as chemical buffer factors (Egleston et al., 2010; Frankignoulle, 1994; Soetaert et al., 2007). Sensitivities express the rate of change of output quantities (Y) in terms of input quantities (X), that is, their partial derivatives (Morel et al., 1976; Smith & Missen, 2003). The (first order) sensitivity coefficient (Smith & Missen, 2003) or interaction capacity (Morel et al., 1976) is defined as follows

$$S_{Y,X} = \frac{\partial Y}{\partial X} \quad (16)$$

which can be normalized using logarithms so that the percentage change in output can be directly linked to percentage change in input (interaction intensity values [Morel et al., 1976] or normalized first-order sensitivities [Smith & Missen, 2003])

$$S_{Y,X}^{nor} = \frac{\partial \ln Y}{\partial \ln X} \quad (17)$$

In the case of pH during an acid titration of TA, the sensitivity is the inverse of the well-known acid–base buffer capacity (β ; Equation 13).

$$S_{pH,TA} = \frac{\partial pH}{\partial TA} = \beta^{-1} \quad (18)$$

The use of both sensitivities (=tendency to change), in which the cause of change is in the denominator and the resulting change is in the numerator (e.g., $\frac{\partial pH}{\partial TA}$), and buffering capacities (=resistance to change), which are just the inverse (e.g., $\frac{\partial TA}{\partial pH}$) is one of the reasons for confusion in the literature. Buffer capacity β expresses the ability to resist changes and is normally presented as

$$\beta_{pH} = \left(\frac{\partial pH}{\partial TA} \right)^{-1} \quad (19)$$

(Middelburg, 2019; Morel & Hering, 1993; Stumm & Morgan, 1981). The Revelle factor (R; Equation 14) is a (normalized) sensitivity factor.

Another cause of inconsistencies among studies relates to the use of pH, $\ln[H^+]$, or $[H^+]$. The original buffer factor β is based on pH (Equations 13 and 19), but Egleston et al. (2010) and Hofmann, Middelburg, et al. (2010) presented definitions based on the natural logarithm of proton concentrations and proton concentrations, respectively

$$\beta_{TA} = \left(\frac{\partial \ln H}{\partial TA} \right)^{-1} \text{ and } \beta_H = \left(\frac{\partial H}{\partial TA} \right)^{-1} \quad (20,21)$$

While β_{pH} values are always positive, β_H and β_{TA} are negative. Note that these buffer factors have been defined as inverse of sensitivity factors. The interchangeable and inconsistent use of the terms buffer capacity, intensity, and index for β is another cause of confusion. Analytical chemists favor the term buffer capacity, geochemists prefer buffer intensity and engineers use the term buffer index (Urbansky & Schock, 2000). Others distinguish between buffer intensity for the actual value at a certain pH (the differential) and buffer capacity for the integral over a distinct range (Chiriac & Balea, 1997).

Table 1

Overview of sensitivities of the Ocean Carbon System $\left(\frac{\partial \text{response}}{\partial \text{driver}}\right)$ and Their Relations to Buffering Quantities in the Literature

Driver	Response	Sensitivity factor	Symbol	Name	Reference
TA	pH	$\left(\frac{\partial \text{pH}}{\partial \text{TA}}\right)$	$-\Phi_H$ β_{pH}^{-1}	Chemical buffer factor Inverse of Buffer capacity	Frankignoulle (1994) van Slyke (1922)
	lnH	$\left(\frac{\partial \ln H}{\partial \text{TA}}\right)$	β_{TA}^{-1}		Egleston et al. (2010)
	H	$\left(\frac{\partial H}{\partial \text{TA}}\right)$	$-\beta_H^{-1}$	Negative inverse of Proton concentration buffer factor	Hofmann et al. (2010a & 2010b)
	pCO ₂	$\left(\frac{\partial \text{pCO}_2}{\partial \text{TA}}\right)$	$-\Pi_H$	Buffer factor	Frankignoulle (1994)
	lnpCO ₂	$\left(\frac{\partial \text{pCO}_2}{\partial \text{TA}}\right) \frac{\text{TA}}{\text{pCO}_2}$ or $\left(\frac{\partial \ln \text{pCO}_2}{\partial \ln \text{TA}}\right)$	γ_{TA}	Alkalinity factor	Sarmiento and Gruber (2006)
	lnCO ₂	$\left(\frac{\partial \ln \text{CO}_2}{\partial \text{TA}}\right)$	γ_{TA}^{-1}		Egleston et al. (2010)
	CO ₂	$\frac{1}{K_0} \left(\frac{\partial \text{CO}_2}{\partial \text{TA}}\right)$	$-\Pi_H$		Frankignoulle (1994)
	lnCO ₃ ²⁻	$\left(\frac{\partial \ln \text{CO}_3^{2-}}{\partial \text{TA}}\right)$	ω_{TA}^{-1}		Egleston et al. (2010)
DIC	pH	$\left(\frac{\partial \text{pH}}{\partial \text{DIC}}\right)$	Φ_{CO_2} $\beta_{\text{CO}_2}^{-1}$		Frankignoulle (1994) and Weber & Stumm (1963)
	lnH	$\left(\frac{\partial \ln H}{\partial \text{DIC}}\right)$	β_{DIC}^{-1}		Egleston et al. (2010)
	pCO ₂	$\left(\frac{\partial \text{pCO}_2}{\partial \text{DIC}}\right)$	Π_D		Frankignoulle (1994)
	lnpCO ₂	$\left(\frac{\partial \text{pCO}_2}{\partial \text{DIC}}\right) \frac{\text{DIC}}{\text{pCO}_2}$ or $\left(\frac{\partial \ln \text{pCO}_2}{\partial \ln \text{DIC}}\right)$	R B_{hom} β_D γ_{DIC}	Homogeneous buffer factor or Revelle factor	Bolin and Eriksson (1959), Sundquist et al. (1979), Frankignoulle (1994), and Sarmiento and Gruber (2006)
	lnCO ₂	$\left(\frac{\partial \ln \text{CO}_2}{\partial \text{DIC}}\right)$	γ_{DIC}^{-1}		Egleston et al. (2010)
	TA	$\left(\frac{\partial \text{TA}}{\partial \text{DIC}}\right)$	Q	Isocapnic quotient	Humphreys et al. (2018)
	lnCO ₃ ²⁻	$\left(\frac{\partial \ln \text{CO}_3^{2-}}{\partial \text{DIC}}\right)$	ω_{DIC}^{-1}		Egleston et al. (2010)
pCO ₂	pH	$\left(\frac{\partial \text{pH}}{\partial \text{pCO}_2}\right)$	Φ_D		Frankignoulle (1994)
	lnH	$\left(\frac{\partial \ln H}{\partial \ln \text{pCO}_2}\right)$	H^{-1}		Omta et al. (2010)
	lnCO ₃ ²⁻	$\left(\frac{\partial \text{CO}_3^{2-}}{\partial \text{pCO}_2}\right) \frac{\text{pCO}_2}{\text{CO}_3^{2-}}$ or $\left(\frac{\partial \ln \text{CO}_3^{2-}}{\partial \ln \text{pCO}_2}\right)$	β_C^{-1} O^{-1}		Frankignoulle (1994) and Omta et al. (2010)

Note. Based on the Hagens and Middelburg (2016a) approach, supporting information S2 explicitly links the various sensitivities.

Using a first-order sensitivity approach the response in Y due to changes in the drivers TA , DIC , T , S , and any other property X can be described as follows

$$dY = \left(\frac{\partial Y}{\partial TA} \right) dTA + \left(\frac{\partial Y}{\partial DIC} \right) dDIC + \left(\frac{\partial Y}{\partial T} \right) dT + \left(\frac{\partial Y}{\partial S} \right) dS + \left(\frac{\partial Y}{\partial X} \right) dX \quad (22)$$

where the partial derivatives imply that the other factors are constant. These and other sensitivities have been used and named in the literature, either in this particular, a normalized or similar form (Table 1). Sometimes a different name is used for the same sensitivity or the same name is used for different sensitivities. For instance, Sarmiento and Gruber (2006) define their alkalinity factor (γ_{TA}) as

$$\gamma_{TA} = \left(\frac{\partial pCO_2}{\partial TA} \right) \frac{TA}{pCO_2} = \left(\frac{\partial \ln pCO_2}{\partial \ln TA} \right) \quad (23)$$

while Egleston et al. (2010) use the same symbol for a different sensitivity:

$$\gamma_{TA} = \left(\frac{\partial TA}{\partial \ln CO_2} \right) \quad (24)$$

Similarly, the Revelle factor (R ; Equation 14) is also known as homogeneous buffer factor B_{hom} (Sundquist et al., 1979), β_D (Frankignoulle, 1994), and γ_{DIC} (Sarmiento & Gruber, 2006). Moreover, it relates to DIC/γ_{DIC} using the γ_{DIC} definition of Egleston et al. (2010), which is again different from that of Sarmiento and Gruber (2006). Most of the sensitivities in Table 1 can be explicitly linked (Frankignoulle, 1994; Hagens & Middelburg, 2016a) as documented in supporting information S2. *To prevent further confusion and propagation of inconsistencies, we propose to explicitly add the term sensitivity to factors such as the Revelle sensitivity factor and restrict the use of buffer for its inverse, that is, the resistance to change. Consequently, we discourage the common use of the term buffer or chemical buffer factor for sensitivities.*

3.2. Sensitivity of Seawater

Most of the sensitivities presented above depend nonlinearly on the solution composition. Figure 2 shows the sensitivities of pH toward changes in CBA and DIC and of pCO_2 toward a change in DIC (Revelle sensitivity factor) as a function of pH for average seawater. The carbonic acid system dominates the buffering capacity of seawater and these sensitivities thus show extrema related to the pK_1 (≈ 5.9) and pK_2 (≈ 9) values of carbonic acid in seawater. The sensitivity $\left(\frac{\partial pH}{\partial CBA} \right)$ shows maxima at pH values of about 4.5 and 7.5. The former is the proton reference level chosen at the well-known inflection point of the alkalinity titration (Dickson, 1981), and the latter is consistent with the minor species theorem that it should be half way between pK_1 and pK_2 (Egleston et al., 2010; Morel & Hering, 1993). These sensitivities show minima close to the pK_1 and pK_2 values of carbonic acid in seawater, consistent with the well-established concept that buffers are most efficient close to their pK value (Butler, 1964; Stumm & Morgan, 1981). At pH values of about 7.5, sensitivity toward addition of dissolved inorganic carbon $\left(\frac{\partial pH}{\partial DIC} \right)$, a weak acid, is similar to that of $\left(\frac{\partial pH}{\partial CBA} \right)$, addition of proton acceptors, but for the sign. However, seawater is more sensitive to CBA than to DIC changes below the pK_1 of the carbonic acid system ($pH \approx 5.9$). Conversely, the sensitivity toward DIC changes is larger than that due to CBA changes at $pH > 7.5$. This can be attributed to the number of protons released (Egleston et al., 2010). The Revelle sensitivity factor is very low at pH values below 6 and above 12 and shows maxima at pH values of about 7.5 and 10 and a minimum around the pK_2 of the carbonic acid system ($pH \approx 9$) because of the prominent role of the carbonate ion in buffering the carbon dioxide added (Gattuso & Hansson, 2011)



These and other seawater sensitivities have many applications, ranging from propagating uncertainties in the carbonic acid system (Orr et al., 2018), attributing changes in pCO_2 to temperature, salinity, and other




Figure 2. The sensitivities $dpH/dCBA$ (a), $dpH/dDIC$ (b), and the Revelle sensitivity factor (c) as function of pH. Left column over the entire pH range; right column focuses on the changes from pH 6 to 9. Vertical gray lines indicate maxima/minima. Details of the calculations are presented in the supporting information.

factors (Middelburg, 2019; Sarmiento & Gruber, 2006; Takahashi et al., 1993, 2014), understanding factors governing pH seasonality (Hagens & Middelburg, 2016b), and how these factors will change because of global warming and ocean acidification (Hagens & Middelburg, 2016a). For instance, Richier et al. (2018) showed that the CO_2 sensitivity of phytoplankton correlates with the sensitivity $\left(\frac{\partial pH}{\partial DIC}\right)$ of seawater. This quantification of sensitivities is pivotal to understanding earth system functioning and the magnitude of climate feedbacks during times of global change. For instance, Frankignoulle et al. (1994) showed how the stoichiometry of carbon dioxide release during calcite precipitation would change with increasing atmospheric carbon dioxide levels. Multiple authors have shown that the seasonality of pH and pCO_2 will increase due to elevated sensitivities induced by ocean acidification (Gallego et al., 2018; Hagens & Middelburg, 2016a; Kwiatkowski & Orr, 2018; Riebesell et al., 2009; Schulz & Riebesell, 2013). Seawater sensitivity analysis has also contributed to elucidating interactions among various factors perturbing seawater pH and pCO_2 such as hypoxia (Cai et al., 2011, 2017; Hagens et al., 2015; Hagens & Middelburg, 2016a) and atmospheric deposition (Hagens et al., 2014). Finally, explicit quantification of sensitivities allows estimation of simple projections or retrodictions. For instance, the present-day value for $\left(\frac{\partial pH}{\partial pCO_2}\right)$ is about -0.0011 ($ppmv^{-1}$; Hagens & Middelburg, 2016a). Combining this sensitivity with




Figure 3. TA versus DIC plots (i.e., Deffeyes diagrams) showing the equilibrium pH at free scale (a) and $p\text{CO}_2$ in μatm (b) as contours. Both graphs show vectors for the addition of protons (vertically downward) and dissolution of calcium carbonate (slope 2:1). Notice that the resulting change in pH and $p\text{CO}_2$ for the same amount of calcite dissolved or acid added (same vector) differs because of differences in sensitivity (buffering). For instance, the ΔpH and $\Delta p\text{CO}_2$ for proton additions are -0.074 and $+136 \mu\text{atm}$, respectively, at low buffering (high DIC/TA ratio), and -0.037 and $+20.9 \mu\text{atm}$ at high buffering (low DIC/TA ratio). Similarly, for the calcite dissolution vector, the ΔpH values are 0.022 and 0.013 and the $\Delta p\text{CO}_2$ values are -33.9 and $-5.9 \mu\text{atm}$ for low and high buffering, respectively.

global annual increases in $p\text{CO}_2$ of 1.1 to $2.1 \text{ ppmv year}^{-1}$ for the periods 1964–1975 and 2005–2014 generates annual ocean pH declines of about 0.0012 to 0.0023 units, similar to that observed (Dore et al., 2009). Similarly, using the sensitivity $\left(\frac{\partial \text{DIC}}{\partial p\text{CO}_2}\right)$ of about 0.51 ($\mu\text{M kg}^{-1}/\text{ppmv}$), one would retrodict that the global ocean surface DIC would have increased by about $1 \mu\text{M kg}^{-1} \text{ year}^{-1}$ from 1994 to 2007, consistent with observations by Gruber et al. (2019).

4. Biogeochemical Processes and Inorganic Carbon Dynamics

The marine carbon dioxide system is impacted by many biogeochemical processes: mineral dissolution and precipitation, organic matter production and respiration, and transfers of electrons, that is, redox processes (Middelburg, 2019; Soetaert et al., 2007; Wolf-Gladrow et al., 2007). The impact of biogeochemical processes on pH and $p\text{CO}_2$ is often analyzed graphically in the form of TA versus DIC plots with isolines for pH and $p\text{CO}_2$ (Figure 3). The impact of biogeochemical processes on DIC, TA, or any of its constituents can be represented as a vector (Deffeyes, 1965). For instance, calcium carbonate dissolution results in the release of one unit DIC and two units of TA and the resulting vector on Figure 3 shows that it will cause an increase in pH and decrease in $p\text{CO}_2$. However, for the very same process intensity (vector length and direction), the resulting change in pH and $p\text{CO}_2$ is different because it depends on the initial conditions (i.e., the sensitivity of the system; section 3.1). Changes in pH and $p\text{CO}_2$ are smaller in well buffered water with a high TA: DIC ratio, that is, low sensitivity (Figure 3).

Although this graphical approach is instructive, there is a need to quantify these changes to improve our predictive capabilities. Ben-Yaakov (1973) recognized that a given reaction can change the pH of a solution by changing the total charge or by adding (or removing) an acid or base. Soetaert et al. (2007) elaborated this approach and showed that the effect of a single biogeochemical process on pH can be calculated from the product of the net charge exchanged during a biogeochemical process (Δcharge) and the sensitivity factor of seawater $\left(\frac{\partial \text{pH}}{\partial \text{CBA}}\right)$. Specifically, the instantaneous effect of a single process with intensity I_{process} ($\text{mol m}^{-3} \text{ s}^{-1}$) on pH can be calculated as follows

$$dpH = \Delta\text{charge} \cdot \left(\frac{\partial \text{pH}}{\partial \text{CBA}}\right) \cdot I_{\text{process}} \quad (26)$$

where Δcharge is the sum of the pH independent ΔCBA (Table 2) and the pH dependent charge (i.e., proton transfer) calculated from reaction stoichiometry and charge of acid–base systems (see Soetaert et al., 2007). Since both the sensitivity of seawater $\left(\frac{\partial \text{pH}}{\partial \text{CBA}}\right)$ and Δcharge depend on the pH, this equation shows that the effect of a specific biogeochemical process on pH also depends on pH. Hofmann et al. (2010a) and Middelburg (2019) presented similar approaches, but focused on proton rather than charge transfers and used the sensitivity factors $\left(\frac{\partial \text{H}}{\partial \text{TA}}\right)$ and $\left(\frac{\partial \text{pH}}{\partial \text{TA}}\right)$, respectively.

Table 2

Biogeochemical Processes and Their Impact on Charge Balance Alkalinity (ΔCBA) and Titration Alkalinity (ΔTA) (After Soetaert et al., 2007)

Process	Reaction	ΔCBA	ΔTA
(R1) Aerobic Mineralization	$(CH_2O)(NH_3)_n(H_3PO_4)_p + O_2 \rightarrow CO_2 + n NH_3 + p H_3PO_4 + H_2O$	0	$n - p$
(R2) Denitrification	$(CH_2O)(NH_3)_n(H_3PO_4)_p + 0.8 HNO_3 \rightarrow CO_2 + n NH_3 + p H_3PO_4 + 0.4 N_2 + 1.4 H_2O$	0	$0.8 + n - p$
(R3) Mn-Oxide Reduction	$(CH_2O)(NH_3)_n(H_3PO_4)_p + 2 MnO_2 + 4 H^+ \rightarrow CO_2 + n NH_3 + p H_3PO_4 + 2 Mn^{2+} + 3 H_2O$	4	$n - p + 4$
(R4) Fe-Oxide Reduction	$(CH_2O)(NH_3)_n(H_3PO_4)_p + 2 Fe_2O_3 + 8 H^+ \rightarrow CO_2 + n NH_3 + p H_3PO_4 + 4 Fe^{2+} + 5 H_2O$	8	$n - p + 8$
(R5) Sulfate Reduction	$(CH_2O)(NH_3)_n(H_3PO_4)_p + 0.5 H_2SO_4 \rightarrow CO_2 + n NH_3 + p H_3PO_4 + 0.5 H_2S + H_2O$	0	$n - p + 1$
(R6) Methanogenesis	$(CH_2O)(NH_3)_n(H_3PO_4)_p \rightarrow 0.5 CO_2 + n NH_3 + p H_3PO_4 + 0.5 CH_4 + H_2O$	0	$n - p$
(R7) Nitrification	$NH_3 + 2 O_2 \rightarrow HNO_3 + H_2O$	0	-2
(R8) Annamox	$NH_3 + HNO_2 \rightarrow N_2 + H_2O$	0	0
(R9) Aerobic Oxidation of Methane	$CH_4 + O_2 \rightarrow CO_2 + 2 H_2O$	0	0
(R10) Anaerobic Oxidation of Methane	$CH_4 + H_2SO_4 \rightarrow CO_2 + H_2S + 2 H_2O$	0	2
(R11) Calcite Precipitation	$Ca^{2+} + CO_3^{2-} \rightarrow CaCO_3$	-2	-2
(R12) Primary Production (Nitrate)	$CO_2 + n HNO_3 + p H_3PO_4 + (1 + n) H_2O \rightarrow (CH_2O)(NH_3)_n(H_3PO_4)_p + (1 + 2n) O_2$	0	$p + n$
(R13) Primary Production (Ammonium)	$CO_2 + n NH_3 + p H_3PO_4 + H_2O \rightarrow (CH_2O)(NH_3)_n(H_3PO_4)_p + O_2$	0	$p - n$
(R14) CO_2 Emission to Air	$CO_2 \rightarrow CO_2(g)$	0	0
(R15) Proton Sorption	$H^+ \rightarrow H^+_{-surface}$	1	1
(R15) Ammonium sorption	$NH_4^+ \rightarrow NH_4^+_{-surface}$	1	0

Note. n = N/C ratio of organic matter and p = P/C ratio of organic matter.

Figure 4 shows the impact of denitrification with Redfield organic matter (R2 in Table 2) on the Δ charge and pH. The process involves the production of DIC, $\sum NH_3$, and $\sum PO_4$, and the consumption of $\sum NO_3$. Denitrification does not impact charge balance alkalinity ($\Delta CBA = 0$). At $pH < 4$, nitrate is charged negatively, ammonium positively while DIC and $\sum PO_4$ are present as uncharged carbonic and phosphoric acids, the Δ charge of the reaction is about +0.95. At $pH > 10$, nitrate is still charged negatively, while carbonate and phosphate ions dominate the DIC and $\sum PO_4$ with the consequence that the Δ charge of the reaction is about -1.08 (Figure 4). At a pH of about 7, the Δ charge is zero because the positive charge due to nitrate consumption and ammonium production is compensated by the production of bicarbonate, the major species in the DIC pool at this pH, with contributions from carbonate, dihydrogenphosphate and hydrogenphosphate. The overall effect of denitrification on pH is obtained by multiplying the Δ charge and sensitivity at a specific pH value (Equation 26; Figure 4). Consequently, the pH increases at pH values below about 7 (Δ charge > 0), while it decreases pH at higher pH because the production of bicarbonate, carbonate and phosphate ions outcompetes the consumption of nitrate and production of ammonium (Δ charge < 0). Thus, Δ charge defines the direction of the pH change (i.e., increase or decrease), while the sensitivity determines the magnitude of the response.

Soetaert et al. (2007) presented a full list of processes and how they impact pH over the full pH range; here we focus on aerobic and anaerobic mineralization, primary production, and calcium carbonate production and




Figure 4. The response in pH due to denitrification as a function of pH. (a) The sensitivity factor $dpH/dCBA$ of seawater, (b) the change in charge due to denitrification, and (c) the resulting change in pH as calculated by Equation 26. Details of the calculations are presented in the supporting information.




Figure 5. The impact of biogeochemical processes on pH. (a) The change in pH for aerobic mineralization with and without nitrification. (b) The change in pH due to sulfate reduction and methanogenesis. (c) The change in pH due to primary production based on ammonium or nitrate. (d) The change in pH due to calcification and calcium carbonate dissolution. Note the break at the pH corresponding to calcium carbonate equilibrium (modified from Soetaert et al., 2007). Details of the calculations are presented in the supporting information.

dissolution (Table 2). Figure 5 shows that the overall impact of a biogeochemical process on pH is a highly nonlinear function with multiple processes crossing the zero-pH-change line, implying that the direction (sign of Δcharge) and magnitude (sensitivity times Δcharge) of pH change depend on the initial conditions. For instance, calcium carbonate dissolution always increases the pH but the response depends on the sensitivity factor, that is, on the initial conditions, consistent with the graphical approaches presented above (Figure 3). Aerobic respiration (R1) and methanogenesis (R6) increase pH at pH < 5.2 and 5.6, respectively (Soetaert et al., 2007), because the production of ammonium is not compensated by sufficient production of anions (bicarbonate and phosphate anions). At higher pH, aerobic respiration and methanogenesis decrease pH because of bicarbonate, carbonate, and phosphate ion production (Figure 5). Aerobic respiration accompanied by nitrification (oxidation of ammonium to nitrate, R 7) always results in a pH decrease (Figure 5) because there is no positive charge produced (Soetaert et al., 2007). Similarly, primary production based on nitrate always results in pH increase, while regenerated production based on ammonium results in pH decrease at low pH (ammonium uptake is compensated by proton release for electroneutrality; Soetaert et al., 2007; Wolf-Gladrow et al., 2007). Sulfate reduction (R5) causes a stronger pH increase at low pH values because of sulfate ion consumption and switches to proton production (pH decrease) at a higher pH value. In other words, the impact of sulfate reduction in sediments on pH depends on the initial conditions, that is, whether the sediment has experienced extensive denitrification and metal-oxide reduction before initiation of sulfate reductions (Boudreau & Canfield, 1993; Meister, 2013; Soetaert et al., 2007).

5. Heterogeneous Buffering

While homogeneous buffering due to equilibria in solution is nearly instantaneous and can be quantified using seawater sensitivities discussed in section 3, heterogeneous buffering mechanisms involving particles occur over multiple time scales (Archer et al., 1998; Boudreau et al., 2018; Sarmiento & Gruber, 2006). For instance, proton sorption to surfaces occurs almost instantaneously, while mineral dissolution and

precipitation are kinetically controlled (Lasaga, 1998). Moreover, homogeneous buffering is spatially rather uniform in the ocean because of the limited range and relative uniform distribution of salinity, dissolved inorganic carbon, and alkalinity. Heterogeneous buffering involves particles suspended in the water column, sediments deposited at the seafloor, and benthic and pelagic calcifying organisms. Calcification is dominated by pelagic organisms in the open ocean, while benthic organisms dominate in the coastal domain (Milliman, 1993; Morse et al., 2007; Morse & Mackenzie, 1990). Sediments dominate heterogeneous buffering because of the large size of this reservoir, that is, there are orders of magnitude more particles at the seafloor than suspended in the water column.

Particles suspended in the water column and deposited on the seafloor may contribute to buffering of seawater via dissolution, precipitation, and mineral surface reactions. Sorption of protons to surfaces increases alkalinity (Table 2). Reactions at the surfaces of organic, biogenic carbonate, and detrital silicate particles contribute to instantaneous buffering; alkalinity titrations of unfiltered samples containing phytoplankton and bacteria have shown that particulate matter surfaces neutralize some of the protons added (Kim et al., 2006). However, this heterogeneous buffer capacity is very limited for typical marine suspended matter concentrations. The role of surface reactions in buffering pore-water chemistry is largely unknown although Jahnke and Jahnke (2004) identified the need to consider mineral surface reactions to properly understand sediment pH dynamics.

Dissolution, precipitation, and transformation reactions of sedimentary silicates and biogenic carbonates govern heterogeneous buffering in the ocean. Heterogeneous buffering in the ocean is dominated by carbonate compensation (Berner, 2004; Pytkowicz, 1967; Ridgwell & Zeebe, 2005), and we will focus on this as well (section 5.2), realizing that processes involving silicate minerals, such as reverse weathering and submarine weathering contribute as well, in particular on very long times scales (section 5.1; Isson & Planavsky, 2018; Mackenzie & Garrels, 1966; Sillén, 1967; Wallmann et al., 2008).


5.1. Silicate Reactions

The impact of seawater-rock interactions on alkalinity is often quantified via a charge balance of major cations on the one hand and chloride, sulfate, and alkalinity on the other hand (Antonelli et al., 2017; Turchyn & DePaolo, 2019)

$$\text{Na}^+ + \text{K}^+ + 2 \text{Mg}^{2+} + 2 \text{Ca}^{2+} = \text{Cl}^- + 2 \text{SO}_4^{2-} + \text{alkalinity}, \quad (27)$$

where exchanges between the univalent and divalent cations or between magnesium and calcium are considered of less importance for the balance. High-temperature hydrothermal vents result in the removal of Ca^{2+} and SO_4^{2-} via anhydrite precipitation and of Mg^{2+} via hydroxy-silicate formation (Antonelli et al., 2017). The latter process generates acidity that enhances release of Ca^{2+} from basalt so that charge remains balanced. Most of the calcium released is eventually precipitated as calcium carbonate in the oceanic crust. Overall, submarine weathering results in carbon dioxide consumption and bicarbonate and calcium release (Berner, 2004; Caldeira, 1995; Staudigel et al., 1989).

Reverse weathering refers to the consumption of alkalinity and generation of protons during marine authigenic clay formation. Weathering on the continents results in the formation of cation-depleted clay minerals which after transport and deposition at the seafloor react with major elements in seawater. Reverse weathering can be written in multiple forms, for example, kaolinite to mica transformation (Sillén, 1967),



or cation-poor amorphous Al-silicate to clay (Mackenzie & Garrels, 1966),



The latter reaction involves the transformation of bicarbonate to carbon dioxide via reaction with cation-depleted, acidic Al-silicates. Reverse weathering can be written in multiple forms, but they all lower marine pH and alkalinity, and increase carbon dioxide concentrations (Isson & Planavsky, 2018).

Another submarine weathering process is induced by organic matter degradation in anoxic sediments. This degradation results in the release of carbon dioxide and dissolved organic matter (fulvic and humic acids), and these may cause dissolution of primary silicate minerals and generate high alkalinity levels (Wallmann et al., 2008). Marine weathering contributes to carbon dioxide consumption and alkalinity release and, thus, counteracts reverse weathering processes. Although seafloor weathering and in particular reverse weathering processes may have played a prominent role during past periods with warm, silica-rich ocean waters (e.g., Precambrian), these heterogeneous buffering reactions involving silicates are considered of less importance for the present ocean than those involving carbonate minerals (Isson & Planavsky, 2018; Pytkowicz, 1967; Berner, 2004; see section 6).

5.2. Carbonate Compensation Dynamics

Carbonate compensation refers to the reactions between carbonate minerals and seawater, and it is instructive to distinguish between chemical and biological carbonate compensation. Chemical compensation focuses on the dissolution or preservation of carbonates at the seafloor, while biological compensation centers on the role of precipitation and its dependence on solution chemistry (Boudreau et al., 2018).

Surface oceans waters are supersaturated with respect to most carbonate minerals (Morse & Mackenzie, 1990). Supersaturation must be reduced to undersaturation, at least in the local microenvironment, before carbonate minerals will dissolve and contribute to buffering (but for surface reactions). Carbonate particles settling to the ocean floor will experience pressure increases and temperature decreases that increase solubility of carbonate minerals (Millero, 2007; Morse & Mackenzie, 1990). Moreover, subsurface ocean waters are usually richer in carbon dioxide and lower in carbonate ions because of organic matter degradation (Sarmiento & Gruber, 2006). As a consequence, seawater becomes undersaturated with respect to carbonate minerals at a certain depth and below this saturation depth mineral dissolution rates increase with depth (Ridgwell & Zeebe, 2005). At the carbonate compensation depth, the flux of carbonate particles downward is exactly balanced by the rate of carbonate dissolution (at the seafloor) with the consequence that no carbonate minerals accumulate at steady state (Boudreau, Middelburg, Hoffmann, & Meysman, 2010). The lysocline refers to the depth range between the carbonate saturation and compensation depths (Boudreau et al., 2018). Ocean buffering dynamics is reflected in changes in the depth distribution of the saturation and compensation depths (Boudreau, Middelburg, & Meysman, 2010; Ridgwell & Zeebe, 2005; Sigman et al., 1998). During periods of ocean acidification, saturation and carbonate compensation depths will shallow, causing increases in carbonate mineral dissolution and alkalinity release, counteracting the acidification. Conversely, during ocean alkalinization, saturation and compensation depths will deepen, with the result that carbonate dissolution and alkalinity release diminish (Boudreau et al., 2018; Ridgwell & Zeebe, 2005; Sigman et al., 1998).

Almost all marine carbonate minerals are of biological origin, for example, coccoliths, pteropods, and foraminifera in the open ocean and corals and mollusks in the coastal domain (Milliman, 1993; Morse & Mackenzie, 1990). Calcifying organisms consume alkalinity and any change in their activity due to alteration in environmental conditions (e.g., temperature and ocean acidification) or food web structure (e.g., food resources, predators, or viruses) consequently impacts their role in ocean buffering. This role of calcifiers in carbonate compensation has impact on buffering at multiple timescales (Boudreau et al., 2018; Caldeira & Rampino, 1993). Lower calcification rates because of ocean acidification (Gazeau et al., 2007; Kroeker et al., 2013) or global warming (Hoegh-Guldberg et al., 2007) directly impact alkalinity removal and thus represent a rapidly operating negative feedback mechanism that will be detectable within decades (Schlunegger et al., 2019). Less calcification in the surface layer also implies less export of biogenic calcite and thus less calcite dissolution in the subsurface. Boudreau, Middelburg, Hoffmann, and Meysman (2010), Boudreau et al. (2018) provided an equation to approximate carbonate compensation depth (z_{CCD})

$$z_{CCD} \approx z_{ref} \ln \left(\frac{F_{car} [Ca^{2+}]}{K_{sp} A \beta_{mt}} + \frac{[Ca^{2+}] [CO_3^{2-}]}{K_{sp}} \right) \quad (30)$$

where F_{car} is the export flux of calcium carbonate, A is the surface area of the seafloor, β_{mt} is the mass transfer of solutes across the diffusive boundary layer at the seafloor, K_{sp} is the temperature, salinity, and pressure dependent stoichiometric solubility product, $[Ca^{2+}]$ and $[CO_3^{2-}]$ are the concentrations of

dissolved calcium and carbonate ions, and z_{ref} is a scaling parameter. Lower calcification and export of calcium carbonate F_{car} will thus lead to a shallowing of the carbonate compensation depth on shorter time scales (years) but would cause additional deepening on longer timescales (10^4 years), because of alkalinity accumulation during periods of lower calcification (Boudreau et al., 2018). This additional deepening due to biological carbonate compensation is an alternative to CO_2 -enhanced continental weathering (Ridgwell, 2007; Zeebe et al., 2009) for CaCO_3 overshooting in the geological record (Luo et al., 2016).

6. Processes Governing Alkalinity in the Ocean

The distribution of alkalinity often covaries with salinity. This is logical because oceanographic processes impacting salinity by freshwater addition (such as precipitation, river, and groundwater discharge and ice melting) or removal (e.g., evaporation and ice formation) also impact alkalinity. These processes are most intense in surface waters that are exposed to the atmosphere, cryosphere, and riverine inputs. It is for this reason that alkalinity can be predicted quite well in ocean surface water using salinity and temperature (Lee et al., 2006; Millero et al., 1998). Alkalinity is often normalized to salinity to infer the other processes affecting alkalinity such as calcite production and dissolution (Carter et al., 2014; Feely et al., 2004; Sarmiento & Gruber, 2006). However, salinity normalization might induce biases, because of regional differences in salinity-alkalinity relationships (Friis et al., 2003; Jiang et al., 2014).

Whole ocean alkalinity is largely governed by the balance between ions generated by weathering and removal of ions via the formation of calcium carbonate. Continental weathering on land generates cations that are charge balanced by alkalinity (Berner & Berner, 2012; Mackenzie & Garrels, 1966; Turchyn & DePaolo, 2019). Rivers and groundwater deliver cations and alkalinity to the ocean where calcifiers produce skeletons and remove alkalinity. On million-year timescales, this is usually represented by the Urey-Ebelman reaction (Berner, 2004; Urey, 1952)



which illustrates the net transfer of carbon from the atmosphere to the sedimentary record. Accordingly, over geological timescales, one would expect that riverine delivery of alkalinity to the ocean is balanced by burial of carbonate in marine sediments (Figure 6a).

Estimates of riverine alkalinity delivery range from 26 to 36 Tmol year^{−1} (supporting information S3 and Table 3) and have been derived either from river DIC export or global estimates of CO_2 consumption by chemical weathering of silicate and carbonate rocks (Berner et al., 1983; Gaillardet et al., 1999; Hartmann et al., 2014; Ludwig et al., 1996; Li et al., 2017; Meybeck, 1987; 1998; Suchet et al., 2003). River biogeochemists normally assume that bicarbonate equals the alkalinity and dissolved inorganic carbon (Raymond & Hamilton, 2018; Suchet et al., 2003). Consequently, global estimates of DIC delivery to the ocean can be used as a proxy for alkalinity transfer from weathering to the ocean. Carbonate burial estimates range from 18 to 34 Tmol C year^{−1} (supporting information S3), with clear consensus about carbonate burial in the deep sea of 11–12 Tmol C year^{−1}, while ocean margin contributions vary from 6 to 23 Tmol C year^{−1} (Iglesias-Rodriguez et al., 2002; Milliman, 1993; Milliman & Droxler, 1996; Morse & Mackenzie, 1990; O'Mara & Dunne, 2019; Smith, 2013; Smith & Mackenzie, 2016; Wollast, 1994). Calcium carbonate formation involves the consumption of two moles of alkalinity per mole of carbon (Table 2), indicating an imbalance between alkalinity inputs from chemical weathering (26–36 Tmol year^{−1}) and alkalinity outputs by carbonate burial (54–62 Tmol year^{−1}), the latter based on carbonate (carbon) burial equal to 27–31 Tmol C year^{−1} (Figure 6 and supporting information S3). This imbalance has been identified before based on Ca and HCO_3^- budgets of the ocean (Berner & Berner, 2012).

This imbalance of 18 to 36 Tmol year^{−1} can be explained in three ways. One, the present-day ocean may not be at steady state regarding alkalinity (Milliman, 1993). The inventory of alkalinity in the ocean is about $3.15 \cdot 10^{18}$ mol, based on a total ocean volume of $1.34 \cdot 10^{18} \text{ m}^3$ and a mean ocean alkalinity of about 2.35 mol m^{-3} (Sarmiento & Gruber, 2006). Accordingly, the residence time of alkalinity is about 88–121 ky with respect to a riverine input of 26–36 Tmol year^{−1}. Consequently, the modern ocean alkalinity budget may still be recovering from last glacial sea level drop and related shifts in carbonate burial/exposure on continental shelves. Two, carbonate burial in ocean margin sediments may be lower than consensus




Figure 6. Alkalinity balance of the ocean (fluxes are in Tmol year^{-1}). (a) Traditional alkalinity balance considering river input of DIC and calcium carbonate burial. (b) Revised oceanic alkalinity balance considering additional sources and sinks based on Table 3.

Table 3

Global Alkalinity Balance of the Ocean (Tmol year^{-1})

Alkalinity sources/sinks	Riverine DIC and carbonate burial balance	Complete alkalinity balance
Riverine DIC	32	32
Riverine PIC		21
Submarine Groundwater		1
Submarine Silicate		2.8
Sulfur Burial		4.7
Denitrification		1.5
Organic Matter Burial		3
Total Sources	32	66
Open Ocean Carbonate Burial	23	23
Ocean Margin Carbonate Burial	36	36
Reverse Weathering		1
Total Sinks	59	60
Imbalance	27	-6

Note. Supporting Information S3 provides a detailed documentation for the various terms, including the range.

values (16–20 Tmol C year⁻¹; Milliman & Droxler, 1996; Iglesias-Rodriguez et al., 2002; Smith, 2013; Smith and Mackenzie, 2016; O'Mara & Dunne, 2019). Morse and Mackenzie (1990) (their Figure 5.1) reported a long-term carbonate burial of about 6 Tmol C year⁻¹ in ocean margin sediments. Total alkalinity removal by carbonate burial would then be about 34 Tmol year⁻¹ (2*(6 + 11)) and result in a balanced budget. Van der Ploeg et al. (2019) reported an alkalinity removal via Cenozoic marginal carbonate burial of 14.3 Tmol year⁻¹ by balancing riverine and anaerobic mineralization inputs with marginal and deep-sea carbonate burial. However, there is consensus that modern carbonate burial in ocean margins is about 16–20 Tmol C year⁻¹ (supporting information S3), hence an alkalinity removal of 32–40 Tmol year⁻¹. Three, this simple depiction of riverine alkalinity balancing calcium carbonate burial in marine sediment (Figure 6a) requires revision because of additional alkalinity inputs from land (e.g., riverine particulate inorganic carbon and groundwater alkalinity) or from marine sediments (Figure 6b).

Rivers deliver elements to the ocean not only in dissolved form but also in particulate form (Martin & Meybeck, 1979). If riverine particulate carbonates are unreactive, they will eventually be buried in marine sediments and are then implicitly included in the marine carbonate burial term. Similarly, if riverine particulate carbonates are reactive and dissolve, they represent an alkalinity source and should be added to the input. Irrespective of their fate, transfer of particulate inorganic carbon from rivers to the ocean would reduce the imbalance. There are very few data on the global particulate inorganic carbon input to the ocean. On the basis of a very small data set, Meybeck (1987) reported a PIC flux of 14.2 Tmol C year⁻¹ and this is still used in global carbon assessments (Li et al., 2017). This PIC flux implies an additional alkalinity source of 28.4 Tmol year⁻¹. This is not only of similar magnitude as the alkalinity delivered in dissolved form, but it would also close the modern ocean alkalinity budget (Table 3). This large, so far overlooked, alkalinity input to the ocean is however poorly constrained. Using the global sediment delivery estimate (19 Pg year⁻¹; range 11–27 Pg year⁻¹) of Beusen et al. (2005) and the average of PIC content of US rivers (0.47 wt% C) from Canfield (1997), a lower riverine PIC delivery is estimated (7.4 Tmol C year⁻¹; range 4.3–10.6 Tmol C year⁻¹). However, this would still correspond to an alkalinity flux of almost 15 Tmol year⁻¹. This difference is primarily due to difference in PIC content of suspended particles (0.9 wt% for Meybeck (1982) vs. 0.47 wt% in US river data base; Canfield, 1997). The average of these two uncertain numbers is used for the alkalinity budget (Figure 6b and Table 3).

Part of the freshwater return flow to the ocean occurs via groundwater discharge rather than via rivers (Slomp & Van Cappellen, 2004), and this is an additional alkalinity input. Zhou et al. (2019) reported a global freshwater submarine discharge flux of 489 km³ year⁻¹, which corresponds to about 1.3% of the global river discharge and is significantly lower than the 5% adopted by Slomp and van Cappellen (2004). Combining these fractions with assumptions about the alkalinity of groundwater, one to three times that of river water (Zhang & Planavsky, 2019), we estimate a global fresh groundwater alkalinity input of about 1 Tmol year⁻¹, range 0.4–4.7 Tmol year⁻¹ (supporting information S3).

Besides these additional alkalinity inputs from riverine PIC and submarine groundwater discharge, there is also alkalinity production and consumption in marine sediments due to anaerobic degradation of organic matter, organic matter burial, and reactions involving silicates (Ben-Yaakov, 1973; Berner et al., 1970; Boudreau & Canfield, 1993; Hu & Cai, 2011; Soetaert et al., 2007; Wallmann et al., 2008). Although we understand the impact of individual processes on TA quite well (see sections 4 and 5), the overall effect of these processes on ocean TA is more difficult to assess because of the tight coupling between alkalinity generating and consuming processes within a sediment column. For instance, dissimilatory sulfate reduction and sulfate reduction coupled to anaerobic methane oxidation generate TA (Table 2), but most of the sulfide and ammonium generated during these processes are reoxidized, resulting in alkalinity consumption (Table 2). Consequently, net overall impact of sedimentary sulfate reduction on ocean alkalinity is limited to the small fraction of reduced sulfur that is eventually buried (Gustafsson et al., 2019; Hu & Cai, 2011; Krumins et al., 2013; Wallmann et al., 2008). Moreover, alkalinity released from the sediment in the form of reduced substances (e.g., ammonium and sulfide) that are subsequently oxidized (e.g., nitrification and sulfide oxidation) in the water column does not contribute to the whole ocean alkalinity balance either (Hu & Cai, 2011).

In the context of the global ocean alkalinity balance, anaerobic alkalinity production can conceptually be considered as an anion charge transfer process, as discussed in section 4 (Ben-Yaakov, 1973; Hu &

Cai, 2011). During denitrification (reduction of nitrate to dinitrogen gas), the charge of nitrate is transferred to bicarbonate and thus increases alkalinity. Similarly, the reduced sulfur buried in marine sediments initially entered the ocean as a sulfate ion; this implies a charge transfer to bicarbonate. In contrast, solid phase oxidants such as iron and manganese oxides enter the ocean uncharged and leave the ocean in an uncharged solid form (e.g., FeS_2), with no impact on global ocean alkalinity. Accordingly, it is only the charge transfer from land-derived nitrate and sulfate to bicarbonate produced by anaerobic respiration that matters for the whole ocean alkalinity balance (Hu & Cai, 2011).

Net alkalinity production due to denitrification/anammox is thus about $1.5 \text{ Tmol year}^{-1}$ (Hu & Cai, 2011) based on a river nitrate input of $21 \text{ Tg N year}^{-1}$ (Seitzinger et al., 2006). The net alkalinity production due to sulfate reduction is derived from the burial of sulfur in marine sediments because most of the sulfate produced is reoxidized (Jørgensen, 1977, 1982). Estimates of sulfur burial are based on organic carbon burial and vary from 1.2 (Berner, 1982) to $3.4 \text{ Tmol S year}^{-1}$ (supporting information S3), implying a potential alkalinity source of 2.4 to $6.9 \text{ Tmol year}^{-1}$ to the ocean. These are upper estimates because part of the net alkalinity generated within sediments results in authigenic carbonate formation and does not contribute to ocean alkalinity. Accordingly, the total contribution of anaerobic biogeochemical processes to the ocean alkalinity balance varies between 3.9 and $8.4 \text{ Tmol year}^{-1}$.

Primary production based on new nitrogen (e.g., nitrate) is an alkalinity source, while aerobic respiration accompanied by nitrification represent an alkalinity sink (Table 2). If all organic matter produced were respired there would be no impact on alkalinity, but a small part of the organic matter produced in the photic zone is eventually buried in marine sediments. Based on organic carbon burial estimates of Berner (1982) and Burdige (2007), we estimate a net alkalinity production of 3 Tmol year^{-1} (see supporting information S3).

Quantifying the contribution of sediment silicate reactions to ocean alkalinity is also complicated by multiple alkalinity production and consumption processes: reverse weathering in surface sediments, marine weathering at depth, in particular in the methanogenic zone (Wallmann et al., 2008), and ocean crust weathering (Berner, 2004; Caldeira, 1995; Staudigel et al., 1989). Moreover, the alkalinity generated at depth by mineral weathering results in authigenic carbonate formation. Although alteration of oceanic crust is likely on the order of 2 Tmol year^{-1} , most bicarbonate generated is precipitated as calcite and does not contribute to ocean alkalinity. Reverse weathering is a sink of alkalinity (e.g., Equations . 28 and 29) on the order of about $0.5\text{--}1.5 \text{ Tmol year}^{-1}$ (Isson & Planavsky, 2018) but varies locally depending on the supply of materials (Michalopoulos & Aller, 1995, 2004; Rahman et al., 2016). Wallmann et al. (2008) reported high rates of submarine weathering of 3.3 to $13.3 \text{ Tmol year}^{-1}$. Their estimate is based on the assumption that all carbon dioxide produced during methanogenesis is converted to bicarbonate and that one third is removed by authigenic carbonate formation and two third is released as alkalinity to the ocean. These numbers are likely too high given that these are based on global methane production rates of 5 (Reeburgh et al., 1993) and $20 \text{ Tmol C year}^{-1}$ (Hinrichs & Boetius, 2002). More recent estimates for marine methane formation are $2.8 \text{ Tmol year}^{-1}$ (Egger et al., 2018) and 0.3 to $2.1 \text{ Tmol C year}^{-1}$ (Wallmann et al., 2012). This would lower submarine weathering alkalinity input to $2\text{--}3 \text{ Tmol year}^{-1}$ (supporting information S3).

Figure 6a and Table 3 clearly show that the ocean alkalinity budget based on the balance between riverine alkalinity inputs and carbonate burial at the ocean floor is imbalanced by about $27 \text{ Tmol year}^{-1}$ and provides only part of the story. Additional alkalinity from riverine PIC delivery (about $21 \text{ Tmol year}^{-1}$) and anaerobic mineralization (about $6.2 \text{ Tmol year}^{-1}$) provides the majority of the alkalinity to balance the budget with minor additional inputs from submarine groundwater discharge (about 1 Tmol year^{-1}), organic matter burial (about 3 Tmol year^{-1}), and silicate interactions (about 1 Tmol year^{-1} ; submarine weathering minus reversed weathering).

7. Conclusions

Alkalinity is a central concept in ocean buffering and it is of the utmost importance to understand and quantify its role in carbon dioxide uptake, carbonate mineral formation, and ocean buffering during times of global change. In section 2, we have shown that it is instructive to distinguish between measurable titration

alkalinity (TA) that is based on a proton balance and CBA. This distinction is needed to understand and quantify the impact of biogeochemical processes such as calcification or primary production on alkalinity.

Although much progress has been made in accurate and reproducible measurements of titration alkalinity, quantification of ocean buffering through the use of buffer or sensitivity factors is underexplored, except for the Revelle sensitivity factor (Sundquist et al., 1979) and the acid–base buffer capacity (Weber & Stumm, 1963). This is unfortunate because such a sensitivity analysis is critical to attribute changes in $p\text{CO}_2$ and/or pH to physical (temperature and salinity) and chemical/biological changes (alkalinity and dissolved inorganic carbon). In section 3, we have provided a systematic treatment of these sensitivity factors and identified and resolved inconsistent terminology. Together with the availability of public domain packages such as *seacarb* (Gattuso et al., 2019) and *AquaEnv* (Hofmann, Soetaert, et al. (2010b)) that facilitate their calculation, this provides researchers with the tools to understand and predict changes in ocean chemistry. This can be retrodiction of past or prediction of future pH changes due to ocean acidification or prediction of future pH, $p\text{CO}_2$, or carbonate saturation values upon alkalinity manipulation during geo-engineering (Renforth & Henderson, 2017).

While these sensitivity factors provide a powerful approach to deal with equilibrium reactions, they are less useful when buffering is provided by interactions with slowly reacting solids or by changes in biological processes (e.g., calcification). These heterogeneous buffering reactions are dominated by calcium carbonate formation and dissolution and normally discussed in terms of ocean carbonate compensation dynamics (Boudreau et al., 2018). Traditionally, the focus has been on the role of carbonate mineral dissolution in the water column and sediments, that is, chemical dissolution, which provides long-term buffering (Archer et al., 1998, 2009; Ridgwell & Zeebe, 2005; Sigman et al., 1998). However, ocean acidification or warming also impact calcification (e.g., coral bleaching). This biological carbonate compensation mechanism has implications on the short term (months) for the carbonate compensation depth and on the longer term (kyr) for alkalinity accumulation in the ocean (Boudreau et al., 2018). Homogeneous buffering, chemical and biological carbonate compensation, and weathering feedbacks together govern the long-term fate of anthropogenic carbon dioxide (Archer et al., 1998, 2009; Boudreau, Middelburg, Hoffmann, & Meysman, 2010). The very same processes have also been involved in the recovery of ocean chemistry to carbon perturbations in the past (Boudreau et al., 2018; Ridgwell & Zeebe, 2005).

Ocean alkalinity is controlled by multiple processes operating over multiple time scales complicating the elucidation of the present-day budget (Table 3) as well as reconstructions of alkalinity, carbon, and pH of the past ocean. Nevertheless, some studies (Boudreau et al., 2019; Caves et al., 2016; Tyrrell & Zeebe, 2004; Zeebe & Tyrrell, 2019) have reported alkalinity reconstructions for the last 50–60 million years indicating relatively minor fluctuations, for example, between 1.5 and 3 mM. The Urey-Ebelmen concept implies that riverine alkalinity delivery should be balanced by carbonate burial in the ocean on geological timescales (Figure 6a). The imbalance of the present-day ocean alkalinity budget between riverine DIC input and marine carbonate burial cannot be resolved by including alkalinity delivery by submarine groundwater or within ocean alkalinity production by anaerobic mineralization or submarine weathering. However, the alkalinity budget can readily be balanced by including alkalinity input to the ocean through riverine particulate inorganic carbon delivery (Table 3 and Figure 6b). The global riverine flux of PIC is poorly known, but reasonable estimates (7.5 to 15 Tmol C year^{−1}; corresponding to an alkalinity flux of 15–30 Tmol year^{−1}) indicate that it is similarly sized as the global riverine DIC flux (26–36 Tmol year^{−1}). This not only urges detailed research to better constrain this flux but also implies a reconsideration of past and present ocean alkalinity budgets that ignore physical weathering products delivered to the ocean.

Glossary

- *Acid*: substance donating a proton
- *Base*: substance accepting a proton
- *Conjugated acids/bases*: substances that only differ by one proton
- *strong acid*: acid that donates all protons in natural water
- *weak acid*: acid that partly dissociates to conjugate base and proton in water
- *pH*: $-\log_{10}$ of proton concentration (H^+)
- *pK*: $-\log_{10}$ of equilibrium constant (K)
- *Bjerrum plot*: graph showing the distribution of acids and their conjugated bases as a function of pH

- *alkalinity*: the excess of proton acceptors (over proton donors) in a solution
- *titration alkalinity (TA)*: measurable alkalinity based on a proton balance
- *charge balance alkalinity (CBA)*: alkalinity based on excess of proton exchanging anions over cations, also known as excess negative charge (ENC)
- *buffer*: solution with a mixture of weak acids and conjugated bases that resist changes by transferring protons
- *homogeneous buffer*: solution resisting drastic changes by rearrangement of protons in solution phase only
- *heterogeneous buffer*: a buffer system comprising both a solution and particles
- *buffer capacity*: ability of a solution to resist changes, also known as buffer index or intensity.
- *sensitivity factor*: change in output due to change in input, also known as chemical buffer factor, and the inverse of buffer capacity
- *carbonate compensation*: response of carbonate production and dissolution processes in the ocean upon changes
- *proton balance*: mass balance of protons
- *proton acceptor level*: the number of protons that can be accepted for an acid–base system at a specific pH, the reverse is known as the proton level
- *reference level species*: the major species of an acid–base system present at the reference level (e.g., pH = 4.5)
- *Revelle factor*: a sensitivity factor expressing the change in carbon dioxide to the change in dissolved inorganic carbon

Author Contributions

J. J. M. designed the research and wrote the manuscript with contributions from K. S. and M. H., in particular for sections 2–4. K. S. and M. H. performed the calculations presented in Figures 1–5.

Acknowledgments

No data sets were used in this review, and the R script is provided in the supporting information. The authors declare that they have no conflicts of interest. This research contributes to the Netherlands Earth System Science Centre, financially supported by the Ministry of Education, Culture and Science (OCW). Ton Marcus assisted with drawing of Figure 6. Wei-Jun Cai, David Burdige, and an anonymous referee are thanked for their constructive reviews and Bernie Boudreau, Matthew Humphreys, Olivier Sulpis, and Alastair McDonald for their informal feedback.

References

- Andral, M. (1850). Researches on the acidity and alkalinity of certain liquids of the human body, in the state of health and disease. *Edinburgh Medical and Surgical Journal*, 73(182), 124–132.
- Antonelli, M., Pester, N. J., Brown, S. T., & DePaolo, D. J. (2017). Effect of paleoseawater composition on hydrothermal exchange in midocean ridges. *Proceedings of the National Academy of Sciences of the United States of America*, 114, 12413–12418. <https://doi.org/10.1073/pnas.1709145114>
- Archer, D., Eby, M., Brovkin, V., Ridgwell, A., Cao, L., Mikolajewicz, U., et al. (2009). Atmospheric lifetime of fossil fuel carbon dioxide. *Annual Review of Earth and Planetary Sciences*, 37, 117–134. <https://doi.org/10.1146/annurev.earth.031208.100206>
- Archer, D., Khesghi, H., & Maier-Reimer, E. (1998). Dynamics of fossil fuel CO₂ neutralization by marine CaCO₃. *Global Biogeochemical Cycles*, 12, 259–276. <https://doi.org/10.1029/98gb00744>
- Asuero, A. G., & Michałowski, T. (2011). Comprehensive formulation of titration curves referred to complex acid–base systems and its analytical implications. *Critical Reviews in Analytical Chemistry*, 41, 151–187. <https://doi.org/10.1080/10408347.2011.559440>
- Ben-Yaakov, S. (1973). pH buffering of pore water of recent anoxic marine sediments. *Limnology and Oceanography*, 18. <https://doi.org/10.4319/lm.1973.18.1.0086>
- Berner, E. K., & Berner, R. A. (2012). *Global environment: Water, air, and geochemical cycles*. Princeton, NJ: Princeton University Press.
- Berner, R. A. (1982). Burial of organic carbon and pyrite sulfur in the modern ocean; its geochemical and environmental significance. *American Journal of Science*, 282, 451–473. <https://doi.org/10.2475/ajs.282.4.451>
- Berner, R. A. (2004). *The Phanerozoic carbon cycle: CO₂ and O₂*. Oxford: Oxford University Press.
- Berner, R. A., Lasaga, A. C., & Garrels, R. M. (1983). Carbonate-silicate geochemical cycle and its effect on atmospheric carbon dioxide over the past 100 million years. *American Journal of Science*, 283, 641–683. <https://doi.org/10.2475/ajs.283.7.641>
- Berner, R. A., Scott, M. R., & Thomlinson, C. (1970). Carbonate alkalinity in the pore waters of anoxic marine sediments. *Limnology and Oceanography*, 15, 544–549. <https://doi.org/10.4319/lm.1970.15.4.0544>
- Beusen, A. H. W., Dekkers, A. L. M., Bouwman, A. F., Ludwig, W., & Harrison, J. (2005). Estimation of global river transport of sediments and associated particulate C, N, and P. *Global Biogeochemical Cycles*, 19, GB4S05. <https://doi.org/10.1029/2005GB002453>
- Bockman, E. E., & Dickson, A. G. (2014). A seawater filtration method suitable for total dissolved inorganic carbon and pH analyses. *Limnology and Oceanography: Methods*, 12, 191–195. <https://doi.org/10.4319/lom.2014.12.191>
- Bolin, B., & Eriksson, E. (1959). Changes in the carbon dioxide content of the atmosphere and sea due to fossil fuel combustion. In *The Atmosphere and the Sea in Motion* (pp. 130–141). Woods Hole, MA, USA: Woods Hole Oceanographic Institution.
- Boudreau, B. P., & Canfield, D. E. (1993). A comparison of closed and open-system models for porewater pH and calcite-saturation state. *Geochimica et Cosmochimica Acta*, 57, 317–334. [https://doi.org/10.1016/0016-7037\(93\)90434-x](https://doi.org/10.1016/0016-7037(93)90434-x)
- Boudreau, B. P. (1996). Method-of-lines code for carbon and nutrient diagenesis in aquatic sediments. *Computer & Geosciences*, 22(5), 479–496. [https://doi.org/10.1016/0098-3004\(95\)00115-8](https://doi.org/10.1016/0098-3004(95)00115-8)
- Boudreau, B. P., Meysman, F. J. R., & Middelburg, J. J. (2004). Multicomponent ionic diffusion in porewaters: Coulombic effects revisited. *Earth and Planetary Science Letters*, 222, 653–666. <https://doi.org/10.1016/j.epsl.2004.02.034>
- Boudreau, B. P., Middelburg, J. J., Hoffmann, A. F., & Meysman, F. J. R. (2010). Ongoing transients in carbonate compensation. *Global Biogeochemical Cycles*, 24, GB4010. <https://doi.org/10.1029/2009GB003654>

- Boudreau, B. P., Middelburg, J. J., & Luo, Y. (2018). The role of calcification in carbonate compensation. *Nature Geoscience*, 11, 894–900. <https://doi.org/10.1038/s41561-018-0259-5>
- Boudreau, B. P., Middelburg, J. J., & Meysman, F. J. R. (2010). Carbonate compensation dynamics. *Geophysical Research Letters*, 37, L03603. <https://doi.org/10.1029/2009GL041847>
- Boudreau, B. P., Middelburg, J. J., Sluijs, A., & van der Ploeg, R. (2019). Secular variations in the carbonate chemistry of the oceans over the Cenozoic. *Earth and Planetary Science Letters*, 512, 194–206. <https://doi.org/10.1016/j.epsl.2019.02.004>
- Brewer, P. G., & Goldman, J. C. (1976). Alkalinity changes generated by phytoplankton growth. *Limnology and Oceanography*, 21(1), 108–117. <https://doi.org/10.4319/lo.1976.21.1.0108>
- Brewer, P. G., Wong, G. T. F., Bacon, M. P., & Spencer, D. W. (1975). An oceanic calcium problem? *Earth and Planetary Science Letters*, 26(1), 81–87. [https://doi.org/10.1016/0012-821x\(75\)90179-x](https://doi.org/10.1016/0012-821x(75)90179-x)
- Broecker, W. S. (1974). *Chemical oceanography*. New York City: Harcourt Brace Jovanovich.
- Broecker, W. S., & Peng, T. H. (1982). *Tracers in the sea*. Palisades, New York: Lamont-Doherty Geological Observatory.
- Burdige, D. J. (2007). Preservation of organic matter in marine sediments: Controls, mechanisms, and an imbalance in sediment organic carbon budgets. *Nature Reviews*, 107, 467–485. <https://doi.org/10.1021/cr050347q>
- Butler, J. N. (1964). *Solubility and pH calculations*. Reading Mass: Addison-Wesley Publishing Company Inc.
- Butler, J. N. (1982). *Carbon dioxide equilibria and their applications*. Reading, Mass.: Addison-Wesley.
- Cai, W.-J., Hu, X., Huang, W.-J., Murrell, M. C., Lehrter, J. C., Lohrenz, S. E., et al. (2011). Acidification of subsurface coastal waters enhanced by eutrophication. *Nature Geoscience*, 4, 766–770. <https://doi.org/10.1038/ngeo1297>
- Cai, W.-J., Huang, W.-J., Luther, G. W., Pierrot, D., Li, M., Testa, J., et al. (2017). Redox reactions and weak buffering capacity lead to acidification in the Chesapeake Bay. *Nature Communications*, 8(1), 369. <https://doi.org/10.1038/s41467-017-00417-7>
- Cai, W. J., Wang, Y. C., & Hodson, R. E. (1998). Acid-base properties of dissolved organic matter in the estuarine waters of Georgia, USA. *Geochimica et Cosmochimica Acta*, 62(3), 473–483. [https://doi.org/10.1016/s0016-7037\(97\)00363-3](https://doi.org/10.1016/s0016-7037(97)00363-3)
- Caldeira, K. (1995). Long-term control of atmospheric carbon dioxide; low-temperature seafloor alteration or terrestrial silicate-rock weathering? *American Journal of Science*, 295, 1077–1114. <https://doi.org/10.2475/ajs.295.9.1077>
- Caldeira, K., & Rampino, M. R. (1993). Aftermath of the end-Cretaceous mass extinction: possible biogeochemical stabilization of the carbon cycle and climate. *Paleoceanography*, 8, 515–525. <https://doi.org/10.1029/93PA01163>
- Canfield, D. E. (1997). The geochemistry of river particulates from the continental USA: Major elements. *Geochimica et Cosmochimica Acta*, 61, 3349–3365. [https://doi.org/10.1016/s0016-7037\(97\)00172-5](https://doi.org/10.1016/s0016-7037(97)00172-5)
- Carter, B. R., Toggweiler, J. R., Key, R. M., & Sarmiento, J. L. (2014). Processes determining the marine alkalinity and calcium carbonate saturation state distributions. *Biogeosciences*, 11, 7349–7362. <https://doi.org/10.5194/bg-11-7349-2014>
- Caves, J. K., Jost, A. B., Lau, K. V., & Maher, K. (2016). Cenozoic carbon cycle imbalances and a variable weathering feedback. *Earth and Planetary Science Letters*, 450, 152–163. <https://doi.org/10.1016/j.epsl.2016.06.035>
- Chanson, M., & Millero, F. J. (2007). Effect of filtration on the total alkalinity of open-ocean seawater. *Limnology and Oceanography: Methods*, 5. <https://doi.org/10.4319/lom.2007.5.293>
- Chen, C. T. (1978). Decomposition of calcium carbonate and organic carbon in the deep oceans. *Science*, 201, 735–736. <https://doi.org/10.1126/science.201.4357.735>
- Chiriac, V., & Balea, G. (1997). Buffer index and buffer capacity for a simple buffer solution. *Journal of Chemical Education*, 74, 937–939. <https://doi.org/10.1021/ed074p937>
- Chisholm, J. R. M., & Gattuso, J. P. (1991). Validation of the alkalinity anomaly technique for investigating calcification of photosynthesis in coral reef communities. *Limnology and Oceanography*, 36, 1232–1239. <https://doi.org/10.4319/lo.1991.36.6.1232>
- Deffeyes, K. S. (1965). Carbonate equilibria: A graphic and algebraic approach. *Limnology and Oceanography*, 10, 412–426. <https://doi.org/10.4319/lo.1965.10.3.0412>
- Dickson, A. G. (1981). An exact definition of total alkalinity and a procedure for the estimation of alkalinity and total inorganic carbon from titration data. *Deep Sea Research Part A: Oceanographic Research Papers*, 28, 609–623. [https://doi.org/10.1016/0198-0149\(81\)90121-7](https://doi.org/10.1016/0198-0149(81)90121-7)
- Dickson, A. G. (1992). The development of the alkalinity concept in marine chemistry. *Marine Chemistry*, 40, 49–63. [https://doi.org/10.1016/0304-4203\(92\)90047-e](https://doi.org/10.1016/0304-4203(92)90047-e)
- Dittmar, W. (1884). Report on researches into the composition of ocean water, collected by H.M.S. Challenger, during the years 1873–1876. In J. Murray (Ed.), *Report on the Scientific Results of the Voyage of H.M.S. Challenger, during the years 1873–1876, Physics and Chemistry* (Vol. 1, pp. 1–251). London: H.M.S.O.
- Donovan, M. (1839). *A treatise on chemistry*. UK: Longman.
- Dore, J. E., Lukas, R., Sadler, D. W., Church, M. J., & Karl, D. M. (2009). Physical and biogeochemical modulation of ocean acidification in the central North Pacific. *Proceedings of the National Academy of Sciences of the United States of America*, 106(30), 12,235–12,240. <https://doi.org/10.1073/pnas.0906044106>
- Egger, M., Riedinger, N., Mogollón, J. M., & Jørgensen, B. B. (2018). Global diffusive fluxes of methane in marine sediments. *Nature Geoscience*, 11, 421–425. <https://doi.org/10.1038/s41561-018-0122-8>
- Egleston, E. S., Sabine, C. L., & Morel, F. M. M. (2010). Revelle revisited: Buffer factors that quantify the response of ocean chemistry to changes in DIC and alkalinity. *Global Biogeochemical Cycles*, 24, GB1002. <https://doi.org/10.1029/2008GB003407>
- Feely, R. A., Sabine, C. L., Lee, K., Berelson, W., Kleypas, J., Fabry, V. J., & Millero, F. J. (2004). Impact of anthropogenic CO₂ on the CaCO₃ system in the oceans. *Science*, 305, 362–366. <https://doi.org/10.1126/science.1097329>
- Feely, R. A., Sabine, C. L., Lee, K., Millero, F. J., Lamb, M. F., Greeley, D., et al. (2002). In situ calcium carbonate dissolution in the Pacific Ocean. *Global Biogeochemical Cycles*, 16(4), 1144. <https://doi.org/10.1029/2002GB001866>
- Fishtik, I., Nagypál, I., & Gutman, I. (1995). Sensitivity analysis of multiple chemical equilibria: Sensitivity coefficients and response equilibria. *The Journal of Chemical Physics*, 103(17), 7545–7555. <https://doi.org/10.1063/1.470271>
- Fong, M. B., & Dickson, A. G. (2019). Insight from GO-SHIP hydrography data into the thermodynamic consistency of CO₂ system measurements in seawater. *Marine Chemistry*, 211, 52–63. <https://doi.org/10.1016/j.marchem.2019.03.006>
- Frankignoulle, M. (1994). A complete set of buffer factors for acid/base CO₂ system in seawater. *Journal of Marine Systems*, 5(2), 111–118. [https://doi.org/10.1016/0924-7963\(94\)90026-4](https://doi.org/10.1016/0924-7963(94)90026-4)
- Frankignoulle, M., Canon, C., & Gattuso, J. P. (1994). Marine calcification as a source of carbon dioxide: Positive feedback of increasing atmospheric CO₂. *Limnology and Oceanography*, 39, 458–462. <https://doi.org/10.4319/lo.1994.39.2.0458>
- Friedlingstein, P., Jones, M. W., O'Sullivan, M., Andrew, R. M., Hauck, J., Peters, G. P., et al. (2019). Global carbon budget 2019. *Earth System Science Data*, 11, 1783–1838. <https://doi.org/10.5194/essd-11-1783-2019>

- Friis, K., Körtzinger, A., & Wallace, D. W. R. (2003). The salinity normalization of marine inorganic carbon chemistry data. *Geophysical Research Letters*, 30(2), 1085. <https://doi.org/10.1029/2002GL015898>
- Gaillardet, J., Dupré, B., Louvat, P., & Allègre, C. J. (1999). Global silicate weathering and CO₂ consumption rates deduced from the chemistry of large rivers. *Chemical Geology*, 159, 3–30. [https://doi.org/10.1016/S0009-2541\(99\)00031-5](https://doi.org/10.1016/S0009-2541(99)00031-5)
- Gallego, M. A., Timmermann, A., Friedrich, T., & Zeebe, R. E. (2018). Drivers of future seasonal cycle changes in oceanic pCO₂. *Biogeosciences*, 15, 5315–5327. <https://doi.org/10.5194/bg-15-5315-2018>
- Gattuso, J. P., Epitalon, J. M., Lavigne, H., & Orr, J. C. (2019). Seacarb: seawater carbonate chemistry R package version 3.2.12. <https://CRAN.R-project.org/package=seacarb>
- Gattuso, J.-P., & Hansson, L. (2011). Ocean acidification: Background and history. In J.-P. Gattuso, & L. Hansson (Eds.), Chap. 1 *Ocean Acidification* (pp. 1–20). Oxford: Oxford University Press.
- Gattuso, J.-P., Magnan, A. K., Bopp, L., Cheung, W. W. L., Duarte, C. M., Hinkel, J., et al. (2018). Ocean solutions to address climate change and its effects on marine ecosystems. *Frontiers in Marine Science*, 5, 337. <https://doi.org/10.3389/fmars.2018.00337>
- Gazeau, F., Quiblier, C., Jansen, J. M., Gattuso, J.-P., Middelburg, J. J., & Heip, C. H. R. (2007). Impact of elevated CO₂ on shellfish calcification. *Geophysical Research Letters*, 34, L07603. <https://doi.org/10.1029/2006GL028554>
- Gazeau, F., Urbini, L., Cox, T. E., Alliouane, S., & Gattuso, J.-P. (2015). Comparison of the alkalinity and calcium anomaly techniques to estimate rates of net calcification. *Marine Ecology Progress Series*, 527, 1–12. <https://doi.org/10.3354/meps11287>
- Goldman, J. C., & Brewer, P. G. (1980). Effect of nitrogen source and growth rate on phytoplankton-mediated changes in alkalinity. *Limnology and Oceanography*, 25, 352–357. <https://doi.org/10.4319/lm.1980.25.2.0352>
- Gruber, N., Clement, D., Carter, B. R., Feely, R. A., van Heuven, S., Hoppema, M., et al. (2019). The oceanic sink for anthropogenic CO₂ from 1994 to 2007. *Science*, 363(6432), 1193–1199. <https://doi.org/10.1126/SCIENCE.AAU5153>
- Gustafsson, E., Hagens, M., Sun, X., Reed, D. C., Humborg, C., Slomp, C. P., & Gustafsson, B. G. (2019). Sedimentary alkalinity generation and long-term alkalinity development in the Baltic Sea. *Biogeosciences*, 16, 437–456. <https://doi.org/10.5194/bg-16-437-2019>
- Hagens, M., Hunter, K. A., Liss, P. S., & Middelburg, J. J. (2014). Biogeochemical context impacts seawater pH changes resulting from atmospheric sulfur and nitrogen deposition. *Geophysical Research Letters*, 41, 935–941. <https://doi.org/10.1002/2013GL058796>
- Hagens, M., & Middelburg, J. J. (2016a). Generalised expressions for the response of pH to changes in ocean chemistry. *Geochimica et Cosmochimica Acta*, 187, 334–349. <https://doi.org/10.1016/j.gca.2016.04.012>
- Hagens, M., & Middelburg, J. J. (2016b). Attributing seasonal pH variability in surface ocean waters to governing factors. *Geophysical Research Letters*, 43, 12,528–12,537. <https://doi.org/10.1002/2016GL071719>
- Hagens, M., Slomp, C. P., Meysman, F. J. R., Seitaj, D., Harlay, J., Borges, A. V., & Middelburg, J. J. (2015). Biogeochemical processes and buffering capacity concurrently affect acidification in a seasonally hypoxic coastal marine basin. *Biogeosciences*, 12, 1561–1583. <https://doi.org/10.5194/bg-12-1561-2015>
- Hammer, K., Schneider, B., Kulinski, K., & Schulz-Bull, D. (2017). Acid-base properties of Baltic Sea dissolved organic matter. *Journal of Marine Systems*, 173, 114–121. <https://doi.org/10.1016/j.jmarsys.2017.04.007>
- Hartmann, J., Moosdorf, N., Lauerwald, R., Hinderer, M., & West, A. J. (2014). Global chemical weathering and associated P-release—The role of lithology, temperature and soil properties. *Chemical Geology*, 363, 145–163. <https://doi.org/10.1016/j.chemgeo.2013.10.025>
- Hemond, H. F. (1990). Acid neutralizing capacity, alkalinity, and acid-base status of natural waters containing organic acids. *Environmental Science & Technology*, 24(10), 1486–1489. <https://doi.org/10.1021/es00080a005>
- Hernandez-Ayon, J., Zirino, A., Dickson, A., Camiro-Vargas, T., & Valenzuela, E. (2007). Estimating the contribution of organic bases from microalgae to the titration alkalinity in coastal seawaters. *Limnology and Oceanography: Methods*, 5, 225–232. <https://doi.org/10.4319/lom.2007.5.225>
- Hinrichs, K. H., & Boetius, A. (2002). The anaerobic oxidation of methane: New insights in microbial ecology and biogeochemistry. In G. Wefer, D. Billett, D. Hebbeln, B. B. Jorgensen, M. Schlüter, & T. V. Weering (Eds.), *Ocean Margin Systems* (pp. 457–477). Heidelberg, Germany: Springer.
- Hiscock, W. T., & Millero, F. J. (2006). Alkalinity of the anoxic waters in the western Black Sea. *Deep Sea Research, Part II*, 53, 1787–19.
- Hoegh-Guldberg, O., Mumby, P. J., Hooten, A. J., Steneck, R. S., Greenfield, P., Gomez, E., et al. (2007). Coral reefs under rapid climate change and ocean acidification. *Science*, 318, 1737–1742. <https://doi.org/10.1126/science.1152509>
- Hofmann, A. F., Middelburg, J. J., Soetaert, K., Wolf-Gladrow, D. A., & Meysman, F. J. R. (2010a). Proton cycling, buffering, and reaction stoichiometry in natural waters. *Marine Chemistry*, 121, 246–255. <https://doi.org/10.1016/j.marchem.2010.05.004>
- Hofmann, A. F., Soetaert, K., Middelburg, J. J., & Meysman, F. J. R. (2010b). AquaEnv: An aquatic acid-base modelling environment in R. *Aquatic Geochemistry*, 16, 507–546. <https://doi.org/10.1007/s10498-009-9084-1>
- Hu, X. (2020). Effect of organic alkalinity on seawater buffer capacity: A numerical exploration. *Aquatic Geochemistry*. <https://doi.org/10.1007/s10498-020-09375-x>
- Hu, X., & Cai, W.-J. (2011). An assessment of ocean margin anaerobic processes on oceanic alkalinity budget. *Global Biogeochemical Cycles*, 25, GB3003. <https://doi.org/10.1029/2010GB003859>
- Huang, W.-J., Cai, W.-J., Wang, Y., Hu, X., Chen, B., Lohrenz, S. E., et al. (2015). The response of inorganic carbon distributions and dynamics to upwelling-favorable winds on the northern Gulf of Mexico during summer. *Continental Shelf Research*, 111, 211–222. <https://doi.org/10.1016/j.csr.2015.08.020>
- Humphreys, M. P., Daniels, C. J., Wolf-Gladrow, D. A., Tyrrell, T., & Achterberg, E. P. (2018). On the influence of marine biogeochemical processes over CO₂ exchange between the atmosphere and ocean. *Marine Chemistry*, 199, 1–11. <https://doi.org/10.1016/j.marchem.2017.12.006>
- Iglesias-Rodriguez, M. D., Armstrong, R., Feely, R., Hood, R., Kleypas, J., Sabine, C., & Sarmiento, J. (2002). Progress made in study of ocean's calcium carbonate budget. *Eos, Transactions American Geophysical Union*, 83(34), 365. <https://doi.org/10.1029/2002EO000267>
- Isson, T. T., & Planavsky, N. J. (2018). Reverse weathering as a long-term stabilizer of marine pH and planetary climate. *Nature*, 560, 471–475. <https://doi.org/10.1038/s41586-018-0408-4>
- Jahnke, R. A., & Jahnke, D. B. (2004). Calcium carbonate dissolution in deep sea sediments: Reconciling microelectrode, pore water and benthic flux chamber results. *Geochimica et Cosmochimica Acta*, 68, 47–59. [https://doi.org/10.1016/s0016-7037\(03\)00260-6](https://doi.org/10.1016/s0016-7037(03)00260-6)
- Jiang, Z., Tyrrell, T., Hydes, D. J., Dai, M., & Hartman, S. E. (2014). Variability of alkalinity and the alkalinity-salinity relationship in the tropical and subtropical surface ocean. *Global Biogeochemical Cycles*, 28, 729–742. <https://doi.org/10.1002/2013GB004678>
- Jørgensen, B. B. (1977). Sulphur cycle of a coastal marine sediment (Limfjorden, Denmark). *Limnology and Oceanography*, 22, 814–832. <https://doi.org/10.4319/lm.1977.22.5.0814>
- Jørgensen, B. B. (1982). Mineralization of organic matter in the sea bed—The role of sulphate reduction. *Nature*, 296, 643–645.

- Kanamori, S., & Ikegami, H. (1982). Calcium-alkalinity relationship in the North Pacific. *Journal of Oceanography*, 38, 57–62. <https://doi.org/10.1007/bf02110291>
- Kim, H.-C., Lee, K., & Choi, W. (2006). Contribution of phytoplankton and bacterial cells to the measured alkalinity of seawater. *Limnology and Oceanography*, 51, 331–338. <https://doi.org/10.4319/lo.2006.51.1.0331>
- Kinniburgh, D. G., van Riemsdijk, W. H., Koopal, L. K., Borkovec, M., Benedetti, M. F., & Avena, M. J. (1999). Ion binding to natural organic matter: Competition, heterogeneity, stoichiometry and thermodynamic consistency. *Colloids and Surfaces A: Physicochemical and Engineering Aspects*, 151(1–2), 147–166. [https://doi.org/10.1016/S0927-7757\(98\)00637-2](https://doi.org/10.1016/S0927-7757(98)00637-2)
- Kinsey, D. W. (1978). Alkalinity changes and coral reef calcification. *Limnology and Oceanography*, 23. <https://doi.org/10.4319/lo.1978.23.5.0989>
- Ko, Y. H., Lee, K., Eom, K. H., & Han, I. (2016). Organic alkalinity produced by phytoplankton and its effect on the computation of ocean carbon parameters. *Limnology and Oceanography*, 61, 1462–1471. <https://doi.org/10.1002/lno.10309>
- Koppel, M., & Spiro, K. (1914). Über die Wirkung von Moderatoren (Puffern) bei der Verschiebung des Säure-Basengleichgewichtes in biologischen Flüssigkeiten. *Biochemische Zeitschrift*, 65, 409–439.
- Kroeker, K. J., Kordas, R. C., Crim, R., Hendriks, I. E., Ramajo, L., Singh, F. S., et al. (2013). Impacts of ocean acidification on marine organisms: Quantifying sensitivities and interaction with warming. *Global Change Biology*, 19, 1884–1896. <https://doi.org/10.1111/gcb.12179>
- Krumins, V., Gehlen, M., Arndt, S., Van Cappellen, P., & Regnier, P. (2013). Dissolved inorganic carbon and alkalinity fluxes from coastal marine sediments: Model estimates for different shelf environments and sensitivity to global change. *Biogeosciences*, 10, 371–398. <https://doi.org/10.5194/bg-10-371-2013>
- Kuliński, K., Schneider, B., Hammer, K., Machulik, U., & Schulz-Bull, D. (2014). The influence of dissolved organic matter on the acid–base system of the Baltic Sea. *Journal of Marine Systems*, 132, 106–115. <https://doi.org/10.1016/j.jmarsys.2014.01.011>
- Kwiatkowski, L., & Orr, J. C. (2018). Diverging seasonal extremes for ocean acidification during the twenty-first century. *Nature Climate Change*, 8(2), 141–145. <https://doi.org/10.1038/s41558-017-0054-0>
- Lasaga, A. C. (1998). Kinetic theory in the earth sciences. In *Princeton Series in Geochemistry*. x+811 pp. Princeton, Chichester: Princeton.
- Le Chatelier, H. (1884). Sur un énoncé général des lois des équilibres chimiques. *Comptes Rendus de l'Académie des Sciences*, 99, 786–789.
- Lee, K., Tong, L. T., Millero, F. J., Sabine, C. L., Dickson, A. G., Goyet, C., et al. (2006). Global relationships of total alkalinity with salinity and temperature in surface waters of the world's oceans. *Geophysical Research Letters*, 33, L19605. <https://doi.org/10.1029/2006GL027207>
- Leenheer, J. A., & Croué, J.-P. (2003). Characterizing aquatic dissolved organic matter. *Environmental Science & Technology*, 37(1), 18A–26A. <https://doi.org/10.1021/es032333c>
- Li, M., Peng, C., Wang, M., Xue, W., Zhang, K., Wang, K., et al. (2017). The carbon flux of global rivers: A re-evaluation of amount and spatial patterns. *Ecological Indicators*, 80(September), 40–51. <https://doi.org/10.1016/j.ecolind.2017.04.049>
- Ludwig, W., Suchet, P. A., & Probst, J. L. (1996). River discharges of carbon to the world's oceans: Determining local inputs of alkalinity and of dissolved and particulate organic carbon. *Comptes Rendus de l'Académie des Sciences*, 323, 1007–1014.
- Lukawska-Matuszewska, K. (2016). Contribution of non-carbonate inorganic and organic alkalinity to total measured alkalinity in pore waters in marine sediments (Gulf of Gdansk, S-E Baltic Sea). *Marine Chemistry*, 186, 211–220. <https://doi.org/10.1016/j.marchem.2016.10.002>
- Lukawska-Matuszewska, K., Grzybowski, W., Szweczun, A., & Tarasiewicz, P. (2018). Constituents of organic alkalinity in pore water of marine sediments. *Marine Chemistry*, 200, 22–32. <https://doi.org/10.1016/j.marchem.2018.01.012>
- Luo, Y., Boudreau, B. P., Dickens, G. R., Sluijs, A., & Middelburg, J. J. (2016). An alternative model for CaCO₃ over-shooting during the PETM—Biological carbonate compensation. *Earth and Planetary Science Letters*, 453, 223–233. <https://doi.org/10.1016/j.epsl.2016.08.012>
- Mackenzie, F. T., & Garrels, R. M. (1966). Chemical mass balance between rivers and oceans. *American Journal of Science*, 264, 507–525. <https://doi.org/10.2475/ajs.264.7.507>
- Marsigli, L. F. C. (1725). *Histoire Physique de la Mer, Ouvrage Enrichi de figures dessinees d'apres le Naturel, Seconde partie. De Son Eau, Aus dens de la Compagnie* (pp. 18–43). Amsterdam.
- Martin, J. M., & Meybeck, M. (1979). Elemental mass-balance of material carried by major world rivers. *Marine Chemistry*, 7(3), 173–206. [https://doi.org/10.1016/0304-4203\(79\)90039-2](https://doi.org/10.1016/0304-4203(79)90039-2)
- Meister, P. (2013). Two opposing effects of sulfate reduction on carbonate precipitation in normal marine, hypersaline, and alkaline environments. *Geology*, 41, 499–502. <https://doi.org/10.1130/G34185.1>
- Meybeck, M. (1982). Carbon, nitrogen, and phosphorus transport by world rivers. *American Journal of Science*, 282(4), 401–450. <https://doi.org/10.2475/ajs.282.4.401>
- Meybeck, M. (1987). Global chemical weathering of surficial rocks estimated from river dissolved loads. *American Journal of Science*, 287(5), 401–428. <https://doi.org/10.2475/ajs.287.5.401>
- Michalopoulos, P., & Aller, R. C. (1995). Rapid clay mineral formation in Amazon delta sediments: Reverse weathering and oceanic elemental cycles. *Science*, 270, 614–617. <https://doi.org/10.1126/science.270.5236.614>
- Michalopoulos, P., & Aller, R. C. (2004). Early diagenesis of biogenic silica in the Amazon delta: Alteration, authigenic clay formation, and storage. *Geochimica et Cosmochimica Acta*, 68, 1061–1085. <https://doi.org/10.1016/j.gca.2003.07.018>
- Michalowski, T., & Asuero, A. G. (2012). new approaches in modeling carbonate alkalinity and total alkalinity. *Critical Reviews in Analytical Chemistry*, 42, 220–244. <https://doi.org/10.1080/10408347.2012.660067>
- Middelburg, J. J. (2019). Marine carbon biogeochemistry. In *A primer for Earth System scientists* (126 pp.). Springer Nature Switzerland AG: Springer Verlag.
- Millero, F. J. (2007). The marine inorganic carbon cycle. *Chemical Reviews*, 107, 308–341. <https://doi.org/10.1021/cr0503557>
- Millero, F. J., Lee, K., & Roche, M. (1998). Distribution of alkalinity in the surface waters of the major oceans. *Marine Chemistry*, 60(1–2), 111–130. [https://doi.org/10.1016/S0304-4203\(97\)00084-4](https://doi.org/10.1016/S0304-4203(97)00084-4)
- Milliman, J. D. (1993). Production and accumulation of calcium carbonate in the ocean: Budget of a nonsteady state. *Global Biogeochemical Cycles*, 7, 927–957. <https://doi.org/10.1029/93GB02524>
- Milliman, J. D., & Droxler, A. W. (1996). Neritic and pelagic carbonate sedimentation in the marine environment: Ignorance is not bliss. *Geologische Rundschau*, 85, 496–504. <https://doi.org/10.1007/BF02369004>
- Mitchell, P. H., & Rakestraw, N. W. (1933). Buffer capacity of sea water. *Biological Bulletin Publications - Marine Biological Laboratory Woods Hole*, 65, 435–451. <https://doi.org/10.3390/w5031303>

- Morel, F., McDuff, R. E., & Morgan, J. J. (1976). Theory of Interaction intensities, buffer capacities, and pH stability in aqueous systems, with application to the pH of seawater and a heterogeneous model ocean system. *Marine Chemistry*, 4(1), 1–28. [https://doi.org/10.1016/0304-4203\(76\)90031-1](https://doi.org/10.1016/0304-4203(76)90031-1)
- Morel, F. M. M., & Hering, J. G. (1993). *Principles and applications of aquatic chemistry*. Hoboken, N. J.: John Wiley.
- Morse, J. W., Arvidson, R. S., & Lüttge, A. (2007). Calcium carbonate formation and dissolution. *Chemical Reviews*, 107, 342–381. <https://doi.org/10.1021/cr050358j>
- Morse, J. W., & Mackenzie, F. T. (1990). *Geochemistry of sedimentary carbonates*. Amsterdam: Elsevier.
- Muller, F. L., & Bleie, B. (2008). Estimating the organic acid contribution to coastal seawater alkalinity by potentiometric titrations in a closed cell. *Analytica Chimica Acta*, 619, 183–191. <https://doi.org/10.1016/j.aca.2008.05.018>
- O'Mara, N., & Dunne, J. P. (2019). Hot spots of carbon and alkalinity cycling in the coastal oceans. *Scientific Reports*, 9(1), 4434. <https://doi.org/10.1038/s41598-019-41064-w>
- Omta, A. W., Goodwin, P., & Follows, M. J. (2010). Multiple regimes of air-sea carbon partitioning identified from constant-alkalinity buffer factors. *Global Biogeochemical Cycles*, 24, GB3008. <https://doi.org/10.1029/2009GB003726>
- Orr, J. C., Epitalon, J.-M., Dickson, A. G., & Gattuso, J.-P. (2018). Routine uncertainty propagation for the marine carbon dioxide system. *Marine Chemistry*, 207, 84–107. <https://doi.org/10.1016/j.marchem.2018.10.006>
- Perdue, E. M., Reuter, J. H., & Parrish, R. S. (1984). A statistical model of proton binding by humus. *Geochimica et Cosmochimica Acta*, 48, 1257–1263. [https://doi.org/10.1016/0016-7037\(84\)90060-7](https://doi.org/10.1016/0016-7037(84)90060-7)
- Pytkowicz, R. M. (1967). Carbonate cycle and the buffer mechanism of recent oceans. *Geochimica et Cosmochimica Acta*, 31(73), 63. [https://doi.org/10.1016/0016-7037\(67\)90098-1](https://doi.org/10.1016/0016-7037(67)90098-1)
- Rahman, S., Aller, R. C., & Cochran, J. K. (2016). Cosmogenic ^{32}Si as a tracer of biogenic silica burial and diagenesis: Major deltaic sinks in the silica cycle. *Geophysical Research Letters*, 43, 7124–7132. <https://doi.org/10.1002/2016GL069929>
- Raymond, P. A., & Hamilton, S. K. (2018). Anthropogenic influences on riverine fluxes of dissolved inorganic carbon to the oceans. *Limnology and Oceanography*, 3, 143–155. <https://doi.org/10.1002/lo.120069>
- Reeburgh, W. S., Whalen, S. C., & Alperin, M. J. (1993). The role of methylothrophy in the global methane budget. In J. C. Murrell, & D. P. Kelly (Eds.), *Microbial Growth on C-1 Compounds* (pp. 1–14). Andover, UK: Intercept.
- Renforth, P., & Henderson, G. (2017). Assessing ocean alkalinity for carbon sequestration. *Reviews of Geophysics*, 55, 636–674. <https://doi.org/10.1002/2016RG000533>
- Revelle, R., & Suess, H. E. (1957). Carbon dioxide exchange between atmosphere and ocean and the question of an increase of atmospheric CO_2 during the past decades. *Tellus*, 9, 18–27. <https://doi.org/10.3402/tellusa.v9i1.9075>
- Richier, S., Achterberg, E. P., Humphreys, M. P., Poulton, A. J., Suggett, D. J., Tyrrell, T., & Moore, C. M. (2018). Geographical CO_2 sensitivity of phytoplankton correlates with ocean buffer capacity. *Global Change Biology*, 24, 4438–4452. <https://doi.org/10.1111/gcb.14324>
- Ridgwell, A. (2007). Interpreting transient carbonate compensation depth changes by marine sediment core modeling. *Paleoceanography*, 22, PA4102. <https://doi.org/10.1029/2006PA001372>
- Ridgwell, A., & Zeebe, R. (2005). The role of the global carbonate cycle in the regulation and evolution of the Earth system. *Earth and Planetary Science Letters*, 234, 299–315. <https://doi.org/10.1016/j.epsl.2005.03.006>
- Riebesell, U., Körtzinger, A., & Oschlies, A. (2009). Sensitivities of marine carbon fluxes to ocean change. *Proceedings of the National Academy of Sciences of the United States of America*, 106(49), 20,602–20,609. <https://doi.org/10.1073/pnas.0813291106>
- Sarmiento, J. L., & Gruber, N. (2006). *Ocean biogeochemical dynamics*, 528 pp. Princeton, N. J.: Princeton University Press.
- Schlunegger, S., Rodgers, K. B., Sarmiento, J. L., Frolicher, T. L., Dunne, J. P., Ishii, M., & Slater, R. (2019). Emergence of anthropogenic signals in the ocean carbon cycle. *Nature Climate Change*, 9, 719–725. <https://doi.org/10.1038/s41558-019-0553-2>
- Schulz, K. G., & Riebesell, U. (2013). Diurnal changes in seawater carbonate chemistry speciation at increasing atmospheric carbon dioxide. *Marine Biology*, 160, 1889. <https://doi.org/10.1007/s00227-012-1965-y>
- Seitzinger, S., Harrison, J. A., Bohlke, J. K., Bouwman, A. F., Lowrance, R., Peterson, B., et al. (2006). Denitrification across landscapes and waterscapes: A synthesis. *Ecological Applications*, 16, 2064–2090. [https://doi.org/10.1890/1051-0761\(2006\)016\[2064:DALAWA\]2.0.CO;2](https://doi.org/10.1890/1051-0761(2006)016[2064:DALAWA]2.0.CO;2)
- Sigman, D. M., McCorkle, D. C., & Martin, W. R. (1998). The calcite lysocline as a constraint on glacial/interglacial low-latitude production changes. *Global Biogeochemical Cycles*, 12(3), 409–427. <https://doi.org/10.1029/98GB01184>
- Sillén, L. G. (1967). The ocean as a chemical system. *Science*, 156, 1189–1197. <https://doi.org/10.1126/science.156.3779.1189>
- Slomp, C. P., & Van Cappellen, P. (2004). Nutrient inputs to the coastal ocean through submarine groundwater discharge: Controls and potential impact. *Journal of Hydrology*, 295(1–4), 64–86. <https://doi.org/10.1016/j.jhydrol.2004.02.018>
- Smith, S. V. (2013). *Parsing the oceanic calcium carbonate cycle: A net atmospheric carbon dioxide source, or a sink?*. Waco, TX: L&O e-Books. Association for the Sciences of Limnology and Oceanography (ASLO). <https://doi.org/10.4319/svsmith.2013.978-0-9845591-2-1>
- Smith, S. V., & Mackenzie, F. T. (2016). The role of CaCO_3 reactions in the contemporary oceanic CO_2 cycle. *Aquatic Geochemistry*, 22, 153. <https://doi.org/10.1007/s10498-015-9282-y>
- Smith, W. R., & Missen, R. W. (2003). Sensitivity analysis in ChE education: Part 1. Introduction and application to explicit models. *Chemical Engineering Education*, 37(3), 222–227.
- Soetaert, K., Hofmann, A. F., Middelburg, J. J., Meysman, F. J. R., & Greenwood, J. (2007). The effect of biogeochemical processes on pH. *Marine Chemistry*, 105, 30–51. <https://doi.org/10.1016/j.marchem.2006.12.012>
- Song, S., Wang, Z. A., Gonnea, M. E., Kroeger, K. D., Chu, S. N., Li, D., & Liang, H. (2020). An important biogeochemical link between organic and inorganic carbon cycling: Effects of organic alkalinity on carbonate chemistry in coastal waters influenced by intertidal salt marshes. *Geochimica et Cosmochimica Acta*, 275, 123–139. <https://doi.org/10.1016/j.gca.2020.02.013>
- Staudigel, H., Hart, S. R., Schmincke, H.-U., & Smith, B. M. (1989). Cretaceous ocean crust at DSDP Sites 417 and 418: Carbon uptake from weathering versus loss by magmatic outgassing. *Geochimica et Cosmochimica Acta*, 53, 3091–3094. [https://doi.org/10.1016/0016-7037\(89\)90189-0](https://doi.org/10.1016/0016-7037(89)90189-0)
- Stumm, W., & Morgan, J. W. (1981). *Aquatic chemistry, chemical equilibria and rates in natural waters* (2nd ed.). New York: John Wiley & Sons, Inc.
- Suchet, A., Probst, J. L., & Ludwig, W. (2003). Worldwide distribution of continental rock lithology: Implications for the atmospheric/soil CO_2 uptake by continental weathering and alkalinity river transport to the oceans. *Global Biogeochemical Cycles*, 17, 1038–1051. <https://doi.org/10.1029/2002gb001891>
- Sundquist, E. T., & Plummer, L. N. (1981). Carbon dioxide in the ocean surface layer: Some modelling considerations. In B. Bolin (Ed.), *SCOPE 16 Carbon Cycle Modelling*. John Wiley & Sons. 259–270 pp.
- Sundquist, E. T., Plummer, L. N., & Wigley, T. M. I. (1979). Carbon dioxide in the ocean surface: The homogeneous buffer factor. *Science*, 204, 1203–1205. <https://doi.org/10.1126/science.204.4398.1203>

- Takahashi, T., Olafsson, J., Goddard, J. G., Chipman, D. W., & Sutherland, S. C. (1993). Seasonal variation of CO₂ and nutrients in the high-latitude surface oceans: A comparative study. *Global Biogeochemical Cycles*, 7(4), 843–878. <https://doi.org/10.1029/93GB02263>
- Takahashi, T., Sutherland, S. C., Chipman, D. W., Goddard, J. G., Ho, C., Newberger, T., et al. (2014). Climatological distributions of pH, pCO₂, total CO₂, alkalinity, and CaCO₃ saturation in the global surface ocean, and temporal changes at selected locations. *Marine Chemistry*, 164, 95–125. <https://doi.org/10.1016/j.marchem.2014.06.004>
- Thompson, T. G., & Bonnar, R. U. (1931). The buffer capacity of seawater. *Industrial and Engineering Chemistry, Analytical Edition*, 3, 393–395. <https://doi.org/10.1021/ac50076a029>
- Turchyn, A. V., & DePaolo, D. J. (2019). Seawater chemistry through Phanerozoic Time. *Annual Review of Earth and Planetary Sciences*, 47, 197–224. <https://doi.org/10.1146/annurev-earth-082517-010305>
- Tyrrell, T., & Zeebe, R. E. (2004). History of carbonate ion concentration over the last 100 million years. *Geochimica et Cosmochimica Acta*, 68, 3521–3530. <https://doi.org/10.1016/j.gca.2004.02.018>
- Ulfso, A., Kuliński, K., Anderson, L. G., & Turner, D. R. (2015). Modelling organic alkalinity in the Baltic Sea using a Humic-Pitzer approach. *Marine Chemistry*, 168, 18–26. <https://doi.org/10.1016/j.marchem.2014.10.013>
- Urbansky, E. T., & Schock, M. R. (2000). Understanding, deriving, and computing buffer capacity. *Journal of Chemical Education*, 77, 1640–1644. <https://doi.org/10.1021/ed077p1640>
- Urey, H. C. (1952). On the early chemical history of the earth and the origin of life. *Proceedings of the National Academy of Sciences*, 38, 352–363. <https://doi.org/10.1073/pnas.38.4.351>
- van der Ploeg, R., Boudreau, B. P., Middelburg, J. J., & Sluijs, A. (2019). Cenozoic carbonate burial along continental margins. *Geology*. <https://doi.org/10.1130/G46418.1>
- van Slyke, D. D. (1922). On the measurement of buffer values and on the relationship of buffer value to the dissociation constant of the buffer and the concentration and reaction of the buffer solution. *The Journal of Biological Chemistry*, 52, 525–570.
- Wallmann, K., Aloisi, G., Haeckel, M., Tishchenko, P., Pavlova, G., Greinert, J., et al. (2008). Silicate weathering in anoxic marine sediments. *Geochimica et Cosmochimica Acta*, 72, 2895–2918. <https://doi.org/10.1016/j.gca.2008.03.026>
- Wallmann, K., Pinero, E., Burwicz, E., Haeckel, M., Hensen, C., Dale, A., & Ruepke, L. (2012). The global inventory of methane hydrate in marine sediments: A theoretical approach. *Energies*, 5(12), 2449–2498. <https://doi.org/10.3390/en5072449>
- Weber, W., & Stumm, W. (1963). Mechanism of hydrogen ion buffering in natural waters. *Journal - American Water Works Association*, 55, 1553–1578. <https://doi.org/10.1002/j.1551-8833.1963.tb01178.x>
- Wolf-Gladrow, D. A., Zeebe, R. E., Klaas, C., Kortzinger, A., & Dickson, A. G. (2007). Total alkalinity: The explicit conservative expression and its application to biogeochemical processes. *Marine Chemistry*, 106, 287–300. <https://doi.org/10.1016/j.marchem.2007.01.006>
- Wollast, R. (1994). The relative importance of biomineralization and dissolution of CaCO₃ in the global carbon cycle. In F. Doumenge (Ed.), *N° Special 13 Past and Present Biomineralization Processes. Consideration about the carbonate cycle* (pp. 11–35). Monaco: Institute of Oceanography.
- Yang, B., Byrne, R. H., & Lindemuth, M. (2015). Contributions of organic alkalinity to total alkalinity in coastal waters: A spectrophotometric approach. *Marine Chemistry*, 176, 199–207. <https://doi.org/10.1016/j.marchem.2015.09.008>
- Zeebe, R. E., & Wolf-Gladrow, D. (2001). CO₂ in seawater: Equilibrium, kinetics, isotopes. In *Elsevier Oceanography Series* (360 pp.). Amsterdam: Elsevier Science B.V.
- Zeebe, R. E., Zachos, J. C., & Dickens, G. R. (2009). Carbon dioxide forcing alone insufficient to explain Palaeocene–Eocene Thermal Maximum warming. *Nature Geoscience*, 2, 576–580. <https://doi.org/10.1038/ngeo578>
- Zeebe, R. W., & Tyrrell, T. (2019). History of carbonate ion concentration over the last 100 million years II: Revised calculations and new data. *Geochimica et Cosmochimica Acta*, 257, 373–392. <https://doi.org/10.1016/j.gca.2019.02.041>
- Zhang, S., & Planavsky, N. J. (2019). Revisiting groundwater carbon fluxes to the ocean with implications for the carbon cycle. *Geology*, 48. <https://doi.org/10.1130/G46408.1>
- Zhou, Y. Q., Sawyer, A. H., David, C. H., & Famiglietti, J. S. (2019). Fresh submarine groundwater discharge to the near-global coast. *Geophysical Research Letters*, 46, 5855–5863. <https://doi.org/10.1029/2019GL082749>

References From the Supporting Information

- Li, Y. H., Takahashi, T., & Broecker, W. S. (1969). Degree of saturation of CaCO₃ in the oceans. *Journal of Geophysical Research*, 74, 5507–5525. <https://doi.org/10.1029/jc074i023p05507>
- Ludwig, W., Suchet, P., Munhoven, G., & Probst, J.-L. (1998). Atmospheric CO₂ consumption by continental erosion: Present-day controls and implications for the last glacial maximum. *Global and Planetary Change*, 16, 107–120. [https://doi.org/10.1016/S0921-8181\(98\)00016-2](https://doi.org/10.1016/S0921-8181(98)00016-2)