TROPICAL FOREST

Long-term thermal sensitivity of Earth's tropical forests

Martin J. P. Sullivan^{1,2}*, Simon L. Lewis^{1,3}, Kofi Affum-Baffoe⁴, Carolina Castilho⁵, Flávia Costa⁶, Aida Cuni Sanchez^{7,8}, Corneille E. N. Ewango^{9,10,11}, Wannes Hubau^{1,12,13}, Beatriz Marimon¹⁴, Abel Monteagudo-Mendoza¹⁵, Lan Qie¹⁶, Bonaventure Sonké¹⁷, Rodolfo Vasquez Martinez¹⁵, Timothy R. Baker¹, Roel J. W. Brienen¹, Ted R. Feldpausch¹⁸, David Galbraith¹, Manuel Gloor¹, Yadvinder Malhi¹⁹, Shin-Ichiro Aiba²⁰, Miguel N. Alexiades²¹, Everton C. Almeida²², Edmar Almeida de Oliveira²³, Esteban Álvarez Dávila²⁴, Patricia Alvarez Loayza²⁵, Ana Andrade²⁶, Simone Aparecida Vieira²⁷, Luiz E. O. C. Aragão^{18,28}, Alejandro Araujo-Murakami²⁹, Eric J. M. M. Arets³⁰, Luzmila Arroyo³¹, Peter Ashton³², Gerardo Aymard C.³³, Fabrício B. Baccaro³⁴, Lindsay F. Banin³⁵, Christopher Baraloto³⁶, Plínio Barbosa Camargo³⁷, Jos Barlow³⁸, Jorcely Barroso³⁹, Jean-François Bastin^{40,41}, Sarah A. Batterman^{1,42,43,44}, Hans Beeckman¹², Serge K. Begne^{1,17}, Amy C. Bennett¹, Erika Berenguer^{19,38}, Nicholas Berry⁴⁵, Lilian Blanc⁴⁶, Pascal Boeckx⁴⁷, Jan Bogaert⁴⁸, Damien Bonal⁴⁹, Frans Bongers⁵⁰, Matt Bradford⁵¹, Francis Q. Brearley², Terry Brncic⁵², Foster Brown⁵³, Benoit Burban⁵⁴, José Luís Camargo²⁶, Wendeson Castro⁵⁵†, Carlos Céron⁵⁶, Sabina Cerruto Ribeiro⁵⁷, Victor Chama Moscoso¹⁵, Jerôme Chave⁵⁸, Eric Chezeaux⁵⁹, Connie J. Clark²⁵, Fernanda Coelho de Souza¹, Murray Collins^{60,61}, James A. Comiskey^{62,63}, Fernando Cornejo Valverde⁶⁴, Massiel Corrales Medina⁶⁵, Lola da Costa⁶⁶, Martin Dančák⁶⁷, Greta C. Dargie¹, Stuart Davies⁶⁸, Nallaret Davila Cardozo⁶⁹, Thales de Haulleville^{12,48}, Marcelo Brilhante de Medeiros⁷⁰, Jhon del Aguila Pasquel⁷¹, Géraldine Derroire⁷², Anthony Di Fiore⁷³‡, Jean-Louis Doucet⁷⁴, Aurélie Dourdain⁷², Vincent Droissart⁷⁵, Luisa Fernanda Duque⁷⁶, Romeo Ekoungoulou⁷⁷, Fernando Elias⁷⁸, Terry Erwin⁷⁹, Adriane Esquivel-Muelbert⁸⁰, Sophie Fauset⁸¹, Joice Ferreira⁸², Gerardo Flores Llampazo⁸³, Ernest Foli⁸⁴, Andrew Ford⁵¹, Martin Gilpin¹, Jefferson S. Hall⁸⁵, Keith C. Hamer⁸⁶, Alan C. Hamilton⁸⁷, David J. Harris⁸⁸, Terese B. Hart^{89,90}, Radim Hédl^{91,92}, Bruno Herault^{47,93,94}, Rafael Herrera⁹⁵, Niro Higuchi⁶, Annette Hladik⁹⁶, Eurídice Honorio Coronado⁷¹, Isau Huamantupa-Chuquimaco⁹⁷, Walter Huaraca Huasco⁹⁷, Kathryn J. Jeffery⁹⁸, Eliana Jimenez-Rojas⁹⁹, Michelle Kalamandeen^{1,100,101}, Marie Noël Kamdem Djuikouo^{11,13,17,102}, Elizabeth Kearsley⁴¹, Ricardo Keichi Umetsu¹⁰³, Lip Khoon Kho¹⁰⁴, Timothy Killeen¹⁰⁵, Kanehiro Kitayama¹⁰⁶, Ponte Klitsgaral¹⁰⁷, Alexander Kosh¹⁰⁸, Nicolas Labrière⁵⁸, William Laurange¹⁰⁹, Sugna Laurange¹⁰⁹, Miguel E. Legi¹¹⁰, Aurena Laurange¹⁰⁹, Pagel Region and Region Bente Klitgaard¹⁰⁷, Alexander Koch¹⁰⁸, Nicolas Labrière⁵⁸, William Laurance¹⁰⁹, Susan Laurance¹⁰⁹, Miguel E. Leal¹¹⁰, Aurora Levesley¹, Adriano J. N. Lima⁶, Janvier Lisingo¹¹, Aline P. Lopes²⁸, Gabriela Lopez-Gonzalez¹, Tom Lovejoy¹¹¹, Jon C. Lovett^{1,107}, Richard Lowe¹¹², William E. Magnusson¹¹³, Jagoba Malumbres-Olarte^{114,115}, Ângelo Gilberto Manzatto¹¹⁶, Ben Hur Marimon Jr.¹¹⁷, Andrew R. Marshall^{8,118,119}, Toby Marthews¹²⁰, Simone Matias de Almeida Reis^{14,19}, Colin Maycock¹²¹, Karina Melgaço¹, Casimiro Mendoza¹²², Faizah Metali¹²³, Vianet Mihindou^{124,125}, William Milliken¹⁰⁷, Edward T. A. Mitchard⁶¹, Paulo S. Morandi¹⁴, Hannah L. Mossman², Laszlo Nagy¹²⁶, Henrique Nascimento⁶, David Neill¹²⁷, Reuben Nilus¹²⁸, Percy Núñez Vargas⁹⁵, Walter Palacios¹²⁹, Nadir Pallqui Camacho^{1,95}, Julie Peacock¹, Colin Pendry⁸⁸, Maria Cristina Peñuela Mora¹³⁰, Georgia C. Pickavance¹, Percy Núñez Vargas⁹⁹, Walter Palacios¹²⁹, Nadir Pallqui Camacho^{1,99}, Julie Peacock¹, Colin Pendry⁸⁸, Maria Cristina Peñuela Mora¹³⁰, Georgia C. Pickavance¹, John Pipoly¹³¹, Nigel Pitman¹³², Maureen Playfair¹³³, Lourens Poorter⁵⁰, John R. Poulsen²⁵, Axel Dalberg Poulsen⁸⁸, Richard Preziosi², Adriana Prieto¹³⁴, Richard B. Primack¹¹³⁵, Hirma Ramírez-Angulo¹¹³⁶, Jan Reitsma¹¹³⁷, Maxime Réjou-Méchain¹⁷⁵, Zorayda Restrepo Correa¹¹³⁸, Thaiane Rodrigues de Sousa⁶, Lily Rodriguez Bayona¹¹³⁹, Anand Roopsind¹¹⁴⁰, Agustín Rudas¹¹³⁴, Ervan Rutishauser⁴^{31,41}, Kamariah Abu Salim¹¹²³, Rafael P. Salomão¹^{142,143}, Juliana Schietti⁶, Douglas Sheil¹¹⁴⁴, Richarlly C. Silva⁵^{57,145}, Javier Silva Espejo¹¹⁴⁶, Camila Silva Valeria³⁸, Marcos Silveira⁵⁵⁷, Murielle Simo-Droissart¹⁷, Marcelo Fragomeni Simon⁷⁰, James Singh¹¹⁴⁷, Yahn Carlos Soto Shareva¹⁵, Clement Stahl⁵⁵⁴, Juliana Stropp¹¹⁴⁸, Rahayu Sukri¹¹²³, Terry Sunderland¹^{149,150}, Martin Svátek¹¹⁵, Michael D. Swaine¹¹⁵², Varun Swamy¹¹⁵³, Hermann Taedoumg¹^{154,155}, Joey Talbot¹§, James Taplin¹¹⁵⁶, David Taylor¹¹⁵⁷, Hans ter Steege¹^{158,159}, John Terborgh²⁵, Raquel Thomas¹¹⁴⁰, Sean C. Thomas¹⁶⁰, Armando Torres-Lezama¹⁶¹, Peter Umunay¹^{162,163}, Luis Valenzuela Gamarra¹⁵, Geertje van der Heijden¹⁶⁴, Peter van der Hout¹⁶⁵, Peter van der Meer¹⁶⁶, Mark van Nieuwstadt¹⁶⁷, Hans Verbeeck⁴¹, Ronald Vernimmen¹⁶⁸, Alberto Vicentini⁶, Ima Célia Guimarães Vieira¹⁴⁵, Emilio Vilanova Torre¹⁶⁹, Jason Vleminckx³⁶, Vincent Vos^{170,171}, Ophelia Wang¹⁷², Lee J. T. White^{98,124,173}, Simon Willcock¹⁷⁴, John T. Woods¹⁷⁵, Verginia Wortel¹⁷⁶, Kenneth Young¹⁷⁷, Roderick Zagt¹⁷⁸, Lise Zemagho¹⁷, Pieter A. Zuidema⁵⁰, Joeri A. Zwerts^{133,167}, Oliver L. Phillips¹

The sensitivity of tropical forest carbon to climate is a key uncertainty in predicting global climate change. Although short-term drying and warming are known to affect forests, it is unknown if such effects translate into long-term responses. Here, we analyze 590 permanent plots measured across the tropics to derive the equilibrium climate controls on forest carbon. Maximum temperature is the most important predictor of aboveground biomass (–9.1 megagrams of carbon per hectare per degree Celsius), primarily by reducing woody productivity, and has a greater impact per °C in the hottest forests (>32.2°C). Our results nevertheless reveal greater thermal resilience than observations of short-term variation imply. To realize the long-term climate adaptation potential of tropical forests requires both protecting them and stabilizing Earth's climate.

he response of tropical terrestrial carbon to environmental change is a critical component of global climate models (1). Land-atmosphere feedbacks depend on the balance of positive biomass growth stimulation by CO_2 fertilization (i.e., β) and negative responses to warmer temperatures and any change in precipitation (i.e., γ). Yet the climate response is so poorly constrained that it remains one of the largest uncertainties in Earth system models (2, 3), with the temperature sensitivity of tropical land carbon

Author affiliations are listed at the end of this paper.
*Corresponding author. Email: m.j.sullivan@leeds.ac.uk

stocks alone differing by >100 Pg C $^{\circ}$ C⁻¹ among models (2). Such uncertainty impedes our understanding of the global carbon cycle, limiting our ability to simulate the future of the Earth system under different long-term climate mitigation strategies. A critical long-term control on tropical land-atmosphere feedbacks is the sensitivity to climate of tropical forests (a key component of γ), where about 40% of the world's vegetation carbon resides (4).

The sensitivity to environmental change of tropical biomass carbon stocks, rates of production, and the persistence of fixed carbon can all be estimated by relating their short-term and interannual responses to variation in climate (5–7). These sensitivities are then used to con-

strain longer-term projections of climate responses (2). Such approaches typically find that higher minimum temperatures are strongly associated with slower tree growth and reduced forest carbon stocks, likely owing to increased respiration at higher temperatures (7–9). Tropical forest carbon is also sensitive to precipitation (10), with, for example, increased tree mortality occurring during drought events (11).

Yet the sensitivity of ecosystems to interannual fluctuations may be an unreliable guide to their longer-term responses to climate change. Such responses will also be influenced by physiological acclimation (12), changes in demographic rates (13), and shifts in species composition (14). For example, both respiration




Fig. 1. Spatial variation in tropical forest carbon. (**A**) The RAINFOR (South America), AfriTRON (Africa), T-FORCES (Asia), and Australian plot networks. Filled symbols show 590 multicensus plots used in the main analysis; open symbols show 223 single-census plots used as an independent dataset. Symbol color indicates the region: green, South America; orange, Africa; purple, Asia; and pink, Australia. (**B**) Variation in carbon among continents. Boxplots show raw variation, whereas blue points show estimated mean values (±SE) after accounting for environmental variation. Letters denote statistically significant differences between continents (*P* < 0.05) based on raw data (black) or after accounting for environmental effects (blue in brackets).

and photosynthesis can acclimate under sustained temperature increases (15-17), tropical trees exhibit physiological plasticity (18), and shifts in species composition occur (14) under sustained drought. These processes could mean that tropical forests are less sensitive to climate than estimates derived from interannual variability imply. An alternative, complementary approach to assessing sensitivity to climate is to measure and analyze spatial variation in tropical ecosystems across climate gradients as a space-for-time substitution. Such biome-wide spatial variation in forest carbon stocks, fluxes, and persistence offers a distinctive and largely unexplored window into the potential equilibrium sensitivity of tropical forest vegetation to warming, because it captures real-world vegetation responses that allow for physiological and ecological adaptation (12).

To assess the long-term climate controls on tropical forest growth and carbon stocks, we assembled, measured, and analyzed a pantropical network of 590 permanent, long-term inventory plots (Fig. 1; see figs. S1 and S2 for ability to capture biome climate space). Our analysis combines standardized measurements from across South American, African, Asian, and Australian tropical lowland forests (273, 239, 61, and 17 plots, respectively). For every plot, we calculated aboveground carbon stocks

(19). Then, to better assess the dynamic controls on aboveground carbon stocks, we also computed the rate of carbon gained by the system (aboveground woody carbon production, calculated as tree growth plus newly recruited trees, in Mg C ha⁻¹ year⁻¹) and the carbon residence time in living biomass (calculated as the ratio of living carbon stocks to carbon gains, in years).

We found considerable variation in biomass carbon among continents, with lower stocks per unit area in South America compared with the Paleotropics, even after accounting for environmental variables (Fig. 1). Continents with high carbon stocks had either large carbon gains (Asia) or long carbon residence times (Africa) (Fig. 1). Because of these differences among continents, which are potentially due to differences in evolutionary history (20), we analyzed the environmental drivers of spatial variation in carbon stocks while accounting for biogeographical differences. We fitted linear models with explanatory variables representing hypothesized mechanistic controls of climate on tropical forest carbon (table S1). We also included soil covariates, continent intercepts, and eigenvectors describing spatial relationships among plots to account for other sources of variation (21).

Forest carbon stocks were most strongly related to maximum temperature [Fig. 2; -5.9% per 1° C increase in mean daily maximum tem-

perature in the warmest month with a 95% confidence interval (CI) = -8.6 to -3.1%, which is equivalent to -9.1 Mg C ha⁻¹ oC⁻¹ for a stand with the mean carbon stock in our dataset, 154.6 Mg C ha⁻¹] followed by rainfall (Fig. 2; +2.4% per 100-mm increase in precipitation in the driest quarter with a 95% CI = 0.6 to 4.3%, equivalent to 0.04 Mg C ha⁻¹ mm⁻¹ for a stand with the mean carbon stocks in our dataset), with no statistically significant relationship with minimum temperature, wind speed, or cloud cover (Fig. 2). The effects of maximum temperature and precipitation are also evident in an analysis considering a wider suite of climate variables than those tied to hypothesized mechanisms (fig. S3) and in an additional independent pantropical dataset of 223 single-census plots (for which carbon gains and residence time cannot be assessed, fig. S4).

The negative effect of maximum temperature on above ground carbon stocks mainly reflects reduced carbon gains with increasing temperature (-4.0% per 1°C, 95% CI = -6.2 to -1.8%; Fig. 2), whereas the positive effect of precipitation emerges through longer carbon residence times with increasing precipitation in the driest quarter (3.3% per 100 mm, 95% CI = 0.9 to 5.7%; Fig. 2). Carbon residence time also increased with the proportion of clay in the soil (Fig. 2). The additive effects of precipitation and temperature on carbon stocks were modified by an interaction between them [change in Akaike information criterion (Δ AIC) = 15.4 comparing the full linear model with or without interaction], with temperature effects more negative when precipitation is low (fig. S6). The interaction was through shortening carbon residence time (Δ AIC = 11.9) rather than reducing carbon gains (model without interaction performed better, Δ AIC = 1.4).

An alternative analysis using decision-tree algorithms (22) also showed maximum temperature and precipitation to be important (fig. S7). This decision-tree approach, which can capture complex nonlinear relationships (22), indicated potential nonlinearity in the relationships between carbon stocks and both temperature and precipitation, with the positive effect of increasing dry-season precipitation on residence times strengthening when precipitation was low and the negative effect of maximum temperature intensifying at high temperatures (fig. S7).

We further investigated nonlinearity in the temperature relationship using breakpoint regression (supported over linear regression based on lower AIC, \triangle AIC = 15.0), which revealed that above 32.2 °C (95% CI = 31.7 ° to 32.6°C), the relationship between carbon stocks and maximum temperature became more negative (cooler than breakpoint, -3.8% °C⁻¹, and warmer than breakpoint, -14.7% °C⁻¹; Fig. 3). By partitioning carbon stocks into their production and persistence, we found that this nonlinearity reflects changes to carbon residence time (\triangle AIC = 10.6) rather than gains $(\Delta AIC = 1.7)$. Overall, our results thus indicate two separate climate controls on carbon stocks: a negative linear effect of maximum temperature through reduced carbon gains and a nonlinear negative effect of maximum temperature, ameliorated by high dry-season precipitation, through reduced carbon residence time.

The effect of temperature on carbon residence time only emerges when dry-season precipitation is low; this is consistent with theoretical expectations that negative effects of temperature on tree longevity are exacerbated by moisture limitation, rather than being independent of it and only due to increased respiration costs (23). This could occur through high vapor pressure deficits in hot and dry forests increasing mortality risk by causing hydraulic stress (23, 24) or carbon starvation due to limited photosynthesis as a result of stomatal closure (23). Notably, the temperatureprecipitation interaction we found for aboveground stocks is in the opposite direction to temperature-precipitation interactions reported for soil carbon (25). In soils, moisture limitation suppresses the temperature response of heterotrophic respiration, whereas in trees, moisture limitation increases the mortality risks of high temperatures.

The negative effects of temperature on biomass carbon stocks and gains are primarily due to maximum rather than minimum temperature. This is consistent with high daytime temperatures reducing CO₂ assimilation rates, for example, owing to increased photorespiration or longer duration of stomatal closure (26, 27), whereas if negative temperature effects were to have increased respiration rates, there should be a stronger relationship with minimum (i.e., nighttime) temperature. Critically, minimum temperature is unrelated to aboveground carbon stocks both pantropically and in one continent, South America, where maximum and minimum temperature are largely de-

coupled [correlation coefficient (r) = 0.33; fig. S8]. Although carbon gains are negatively related to minimum temperature (fig. S9), this bivariate relationship is weaker than with maximum temperature and disappears once the effects of other variables are accounted for (Fig. 2). Finally, in Asia, the tropical region that experiences the warmest minimum temperatures of all, both carbon stocks and carbon gains are highest (Fig. 1 and fig. S11).

Overall, our results suggest that tropical forests have considerable potential to acclimate and adapt to the effects of nighttime minimum temperatures but are clearly sensitive to the effects of daytime maximum temperature. This is consistent with ecophysiological observations suggesting that the acclimation potential of respiration (15) is greater than that of photosynthesis (17). The temperature sensitivity revealed by our analysis is also considerably weaker than short-term sensitivities associated with interannual climate variation (7-9). For example, by relating shortterm annual climate anomalies to responses in plots, the effect of a 1°C increase in temperature on carbon gains has been estimated as more than threefold our long-term, pantropical result (28). This stronger, long-term thermal resilience is likely due to a combination of individual acclimation and plasticity (15-17), differences in species' climate responses (29) leading to shifts in community composition due to changing demographic rates (12), and the immigration of species with higher performance at high temperatures (12).

Our pantropical analysis of the sensitivity to climate of aboveground forest carbon stocks, gains, and persistence shows that warming reduces carbon stocks and woody productivity. Using a reference carbon stock map (30) and applying our estimated temperature sensitivity (including nonlinearity) while holding other variables constant leads to an eventual biome-wide reduction of 14.1 Pg C in live biomass (including scaling to estimate carbon in roots) for a 1°C increase in mean daily maximum temperature in the warmest month (95% CI = 6.9 to 20.7 Pg). This compares with a large range of projected sensitivities in coupled climate carbon cycle models that report vegetation carbon (1 to 58 Pg C °C⁻¹), although these models have not been run to equilibrium (see supplementary methods).

Our results suggest that stabilizing global surface temperatures at 2°C above preindustrial levels will cause a potential long-term biome-wide loss of 35.3 Pg C (95% CI = 20.9 to 49.0 Pg, estimates with alternative baseline biomass maps of 24.0 to 28.4 Pg; fig. S12). The greatest long-term reductions in carbon stocks are projected in South America, where baseline temperatures and future warming are both highest (Fig. 4 and fig. S13). This warming would push 71% of the biome beyond




Fig. 2. Correlates of spatial variation in tropical forest carbon. Points show coefficients from model-averaged general linear models. Variables that did not occur in well-supported models are shrinkage-adjusted toward zero. Coefficients are standardized so that they represent change in the response variable for one standard deviation change in the explanatory variable. Error bars show standard errors (thick lines) and 95% confidence intervals (thin lines); error bar color is for illustrative purposes to reflect the category of variable. Soil texture is represented by the percentage clay and soil fertility by cation exchange capacity. The full models explained 44.1, 31.4, and 30.9% of spatial variation in carbon stocks, gains, and residence time, respectively. Coefficients are shown in table S2. Results are robust to using an alternative allometry to estimate tree biomass (fig. S5).




Fig. 3. Temperature effects on tropical forest carbon stocks, carbon gains, and carbon residence time. Black lines show the best pantropical relationships accounting for environmental covariates. The gray line additionally shows the linear pantropical relationship for carbon stocks. Colored lines show bivariate relationships within each continent, as identified in the legend. Statistically significant relationships are shown with solid lines; nonsignificant relationships are shown with dashed lines. The y axis is on a log scale. Symbol point size is proportional to weights used in model

fitting based on plot size and monitoring length; see supplementary materials and methods. For stocks and gains, linear and breakpoint pantropical relationships are all statistically significant (P < 0.001) as are better-sampled continents. For carbon residence time, relationships with temperature are nonsignificant (ns), but there is a statistically significant interaction between maximum temperature and precipitation in the driest quarter (fig. S6). Relationships with other variables are shown in figs. S8 to S10. ***P < 0.001; **P < 0.01; *P < 0.05; ns, $P \ge 0.05$.




Fig. 4. Long-term change in carbon stocks due to temperature effects alone for global surface air temperature warming of 2°C. Maps show the predicted absolute and relative change in tropical forest carbon stocks. Parts of the biome become warmer than observed now in our dataset (fig. S14). See fig. S12 for predictions using alternative carbon reference maps. Predictions are based on temperature alone and do not include precipitation changes (for which future patterns of change are uncertain) or moderation by increased CO₂. (See fig. S15 for analysis incorporating this.)

the thermal threshold—a maximum temperature of 32.2°C—where larger long-term reductions in biomass are expected (fig. S14). Of course, growth stimulation by carbon dioxide (31) will partially or wholly offset the effect of this temperature increase, depending on both the level of atmospheric carbon dioxide that limits warming to 2°C above preindustrial levels and the fertilization effect of this carbon dioxide on tropical trees. Although

 ${\rm CO_2}$ fertilization will reduce temperature-induced carbon losses from biomass across the tropics (table S3), our analysis indicates that ${\rm CO_2}$ fertilization will not completely offset long-term temperature-induced carbon losses within Amazonia (fig. S15), consistent with a recent decadal-scale analysis of inventory data (32).

The long-term climate sensitivities derived from our pantropical field measurements

incorporate ecophysiological and ecological adaptation and so provide an estimate of the long-term, quasi-equilibrium response of tropical vegetation to climate. This thermal adaptation potential may not be fully realized in future responses because (i) the speed of temperature rises may exceed species' adaptive capabilities, (ii) habitat fragmentation may limit species' ability to track changes in the environment, and (iii) other human impacts

such as logging and fire can increase the vulnerability of forest carbon stocks to high temperatures. Although many tropical forests are under severe threat of conversion, our results show that, in the long run, tropical forests that remain intact can continue to store high levels of carbon under high temperatures. Achieving the biome-wide climate resilience potential that we document depends on limiting heating and on large-scale conservation and restoration to protect biodiversity and allow species to move.

REFERENCES AND NOTES

- P. M. Cox, R. A. Betts, C. D. Jones, S. A. Spall, I. J. Totterdell, Nature 408, 184–187 (2000).
- 2. P. M. Cox et al., Nature **494**, 341–344 (2013).
- 3. B. B. B. Booth et al., Environ. Res. Lett. 7, 024002 (2012).
- 4. K.-H. Erb et al., Nature 553, 73-76 (2018).
- W. Wang et al., Proc. Natl. Acad. Sci. U.S.A. 110, 13061–13066 (2013).
- J. Liu et al., Science 358, eaam5690 (2017).
- D. A. Clark, S. C. Piper, C. D. Keeling, D. B. Clark, Proc. Natl. Acad. Sci. U.S.A. 100, 5852–5857 (2003).
- W. R. L. Anderegg et al., Proc. Natl. Acad. Sci. U.S.A. 112, 15591–15596 (2015).
- 9. A. Ballantyne et al., Nat. Clim. Chang. 7, 148-152 (2017).
- 10. J. K. Green et al., Nature 565, 476-479 (2019).
- 11. O. L. Phillips et al., Science **323**, 1344–1347 (2009)
- M. D. Smith, A. K. Knapp, S. L. Collins, *Ecology* 90, 3279–3289 (2009)
- J. H. Brown, T. J. Valone, C. G. Curtin, Proc. Natl. Acad. Sci. U.S.A. 94, 9729–9733 (1997).
- 14. S. Fauset et al., Ecol. Lett. 15, 1120-1129 (2012).
- C. A. Gunderson, K. H. O'Hara, C. M. Campion, A. V. Walker, N. T. Edwards, Glob. Chang. Biol. 16, 2272–2286 (2010).
- N. 1. Edwards, Glob. Chang. Biol. **16**, 2272–2286 (2010). 16. M. Slot et al., Glob. Chang. Biol. **20**, 2915–2926 (2014).
- L. F. Ow, K. L. Griffin, D. Whitehead, A. S. Walcroft, M. H. Turnbull, New Phytol. 178, 123–134 (2008).
- 18. T. F. Domingues *et al.*, *Oecologia* **187**, 933–940 (2018).
- 19. See supplementary materials.
- 20. J. W. F. Slik et al., Proc. Natl. Acad. Sci. U.S.A. 115, 1837-1842 (2018).
- S. Dray, P. Legendre, P. R. Peres-Neto, *Ecol. Modell.* 196, 483–493 (2006).
- 22. L. Breiman, Mach. Learn. 45, 5-32 (2001).
- 23. N. McDowell et al., New Phytol. 219, 851-869 (2018).
- C. G. Fontes et al., Philos. Trans. R. Soc. Lond. B Biol. Sci. 373, 20180209 (2018).
- 25. P. Ciais et al., Nature 437, 529-533 (2005).
- M. E. Dusenge, A. G. Duarte, D. A. Way, New Phytol. 221, 32–49 (2019).
- 27. S. Pau, M. Detto, Y. Kim, C. J. Still, Ecosphere 9, e02311 (2018).
- D. A. Clark, D. B. Clark, S. F. Oberbauer, J. Geophys. Res. Biogeosci. 118, 783–794 (2013).
- 29. W. R. L. Anderegg et al., Nature **561**, 538–541 (2018).
- 30. V. Avitabile et al., Glob. Chang. Biol. 22, 1406-1420 (2016).
- 31. S. Piao et al., Glob. Chang. Biol. 19, 2117–2132 (2013).
- 32. W. Hubau et al., Nature 579, 80-87 (2020).
- M. J. P. Sullivan, Data for "Long-term thermal sensitivity of Earth's tropical forests," ForestPlots.NET (2020); doi: 10.5521/forestplots.net/2020 2.

ACKNOWLEDGMENTS

This paper is a product of the RAINFOR, AfriTRON and T-FORCES networks, as well as ForestPlots.net data management for supporting international science synergies. These initiatives have been supported by numerous people and grants since their inception, but we are particularly indebted to hundreds of institutions, field assistants, botanists, and local communities for help in establishing and maintaining the plots. We thank W. Lopez Oviedo for assistance with access to datasets. For additional assistance, see the extended acknowledgments in the supplementary materials. We thank J. Lloyd and C. Quesada for discussions and three anonymous reviewers for helpful comments and suggestions. Funding: Our plot monitoring networks have been supported by multiple grants from a large number of funding bodies: European Research Council: Natural Environment Research Council; European Union's Fifth, Sixth, and Seventh Framework Programme; Royal Society, Leverhulme Trust; Gordon and Betty Moore Foundation; David and Lucile Packard Foundation; State of São Paulo Research Foundation (FAPESP); National Council for

Science and Technology Development of Brazil (CNPq); Agence Nationale de la Recherche; Conservation International; Missouri Botanical Garden: Smithsonian Institution: Wildlife Conservation Society; National Geographic Society; Centre for International Forestry; and Gabon's National Park Agency. Data from RAINFOR, AfriTRON, and T-FORCES are stored and curated by ForestPlots. net, a cyber-infrastructure initiative developed at the University of Leeds that unites permanent plot records and their contributing scientists from the world's tropical forests. Many of the bodies listed above funded the development of ForestPlots.net and curation of most data analyzed here. For details of additional support and grant references, see the extended acknowledgments in the supplementary materials. Author contributions: M.J.P.S., O.L.P., and S.L.L. conceived, designed, and wrote the study. O.L.P. S.L.L., and Y.M. conceived the RAINFOR, AfriTRON, and T-FORCES forest census network programs. L.A., A.A.-M., T.R.B., R.J.W.B., S.K.B., K.A.-B., F.C., C.C., E.A.D., A.C.S., C.E.N.E., T.R.F., W.H., S.L.L., A.M.M., B.S.M., O.L.P., L.Q., B.S., T.S., R.V., and L.J.T.W. coordinated data collection with all authors and ForestPlots.net. M.J.P.S. analyzed the data. Full details of author contributions for data collection, management, and analysis tools are listed in the supplementary materials. All authors commented on or approved the manuscript. Competing interests: The authors declare no competing financial interests. Data and materials availability: Plot-level input data and R scripts are available as a data package on ForestPlots.net (33).

¹School of Geography, University of Leeds, Leeds, UK. ²Department of Natural Sciences, Manchester Metropolitan University, Manchester, UK. ³Department of Geography, University College London, London, UK. 4Mensuration Unit, Forestry Commission of Ghana, Kumasi, Ghana. 5Embrapa Roraima, Brazilian Agricultural Research Corporation (EMBRAPA), Brasília, Brazil. ⁶Instituto Nacional de Pesquisas da Amazônia (INPA), Manaus, Brazil. ⁷Department of Ecosystem Science and Sustainability, Colorado State University, Fort Collins, CO, USA. 8Department of Environment and Geography, University of York, York, UK. ⁹DR Congo Programme, Wildlife Conservation Society, Kisangani, Democratic Republic of Congo. 10 Centre de Formation et de Recherche en Conservation Forestiere (CEFRECOF), Epulu, Democratic Republic of Congo. 11 Faculté de Gestion de Ressources Naturelles Renouvelables, Université de Kisangani, Kisangani, Democratic Republic of Congo. 12 Service of Wood Biology, Royal Museum for Central Africa, Tervuren, Belgium. ¹³Department of Environment, Laboratory of Wood Technology (Woodlab), Ghent University, Ghent, Belgium. ¹⁴UNEMAT - Universidade do Estado de Mato Grosso, Nova Xavantina-MT, Brazil. ¹⁵ Jardín Botánico de Missouri, Oxapampa, Peru. ¹⁶School of Life Sciences, University of Lincoln, Lincoln, UK. ¹⁷Plant Systematics and Ecology Laboratory, Higher Teachers' Training College, University of Yaoundé I, Yaoundé, Cameroon. ¹⁸Geography, College of Life and Environmental Sciences, University of Exeter, Exeter, UK. 19 Environmental Change Institute, School of Geography and the Environment, University of Oxford, Oxford, UK. ²⁰Faculty of Environmental Earth Science, Hokkaido University, Sapporo, Japan. ²¹School of Anthropology and Conservation, University of Kent, Canterbury, UK. ²²Instituto de Biodiversidade e Florestas, Universidade Federal do Oeste do Pará, Santarém - PA, Brazil. ²³Universidade do Estado de Mato Grosso, Cáceres MT, Brazil. ²⁴Escuela de Ciencias Agrícolas, Pecuarias y del Medio Ambiente, National Open University and Distance, Bogotá, Colombia. ²⁵Nicholas School of the Environment, Duke University, Durham, NC, USA. ²⁶Projeto Dinâmica Biológica de Fragmentos Florestais, Instituto Nacional de Pesquisas da Amazônia, Manaus, Brazil. 27 Universidade Estadual de Campinas, Campinas, SP, Brazil. 28 National Institute for Space Research (INPE), São José dos Campos, SP. Brazil. ²⁹Museo de Historia Natural Noel Kempff Mercado, Universidad Autónoma Gabriel René Moreno, Santa Cruz, Bolivia. 30 Wageningen Environmental Research, Wageningen, Netherlands. 31 Dirección de la Carrera de Biología, Universidad Autónoma Gabriel René Moreno, Santa Cruz, Bolivia. 32Department of Organismic and Evolutionary Biology, Harvard University, Cambridge, MA, USA ³³Programa de Ciencias del Agro y el Mar, Herbario Universitario, Guanare, Venezuela. ³⁴Departamento de Biologia, Universidade Federal do Amazonas, Manaus, Brazil. UK Centre of Ecology and Hydrology, Penicuik, UK. ³⁶International Center for Tropical Botany, Department of Biological Sciences, Florida International University, Miami,

FL, USA. ³⁷Centro de Energia Nuclear na Agricultura, Universidade de São Paulo, São Paulo, SP, Brazil. 38Lancaster Environment Centre, Lancaster University, Lancaster, UK. ³⁹Centro Multidisciplinar, Universidade Federal do Acre, Cruzeiro do Sul, AC, Brazil. ⁴⁰Institure of Integrative Biology, ETH Zurich, Zurich, Switzerland. 41Department of Environment, Computational and Applied Vegetation Ecology (CAVELab), Ghent University, Ghent, Belgium. 42Priestley International Centre for Climate, University of Leeds, Leeds, UK. ⁴³Smithsonian Tropical Research Institute, Panama, Panama. ⁴⁴Cary Institute of Ecosystem Studies, Millbrook, NY, USA. ⁴⁵The Landscapes and Livelihoods Group, Edinburgh, UK. 46UR Forest and Societies, CIRAD Montpellier, France. 47 Isotope Bioscience Laboratory (ISOFYS), Ghent University, Ghent, Belgium. ⁴⁸Gembloux Agro-Bio Tech, University of Liège, Liège, Belgium. ⁴⁹UMR Silva, INRAE, Nancy, France. ⁵⁰Forest Ecology and Forest Management Group, Wageningen University, Wageningen, Netherlands. 51CSIRO Land and Water, Australia. 52Congo Programme, Wildlife Conservation Society, Brazzavile, Republic of Congo. 53Woods Hole Research Center, Falmouth, MA, USA. ⁵⁴INRAE, UMR EcoFoG, CNRS, CIRAD, AgroParisTech, Université des Antilles, Université de Guyane, 97310 Kourou, French Guiana. ⁵⁵Programa de Pós-Graduação Ecologia e Manejo de Recursos Naturais, Universidade Federal do Acre, Rio Branco, AC, Brazil ⁵⁶Herbario Alfredo Paredes, Universidad Central del Ecuador. Ouito. Ecuador. 57Centro de Ciências Biológicas e da Natureza, Universidade Federal do Acre, Rio Branco, AC Brazil. ⁵⁸Laboratoire Évolution et Diversité Biologique, UMR 5174 (CNRS/IRD/UPS), CNRS, Toulouse, France. 59Rougier-Gabon, Libreville, Gabon. 60 Grantham Research Institute on Climate Change and the Environment, London, UK. 61School of Geosciences, University of Edinburgh, Edinburgh, UK. ⁶²Inventory and Monitoring Program, National Park Service, Fredericksburg, VA, USA. ⁶³Smithsonian Institution, Washington, DC, USA. ⁶⁴Proyecto Castaña, Made de Dios, Peru. 65 Universidad Nacional de San Agustín de Arequipa, Areguipa, Peru. 66 Instituto de Geociências, Faculdade de Meteorologia, Universidade Federal do Para, Belém, PA, Brazil. ⁶⁷Faculty of Science, Department of Ecology and Environmental Sciences, Palacký University Olomouc, Olomouc, Czech Republic. ⁶⁸Center for Tropical Forest Science, Smithsonian Tropical Research Institute, Panama, Panama. 69 Facultad de Ciencias Biológicas, Universidad Nacional de la Amazonía Peruana, Iquitos, Peru. 70 Embrapa Genetic Resources and Biotechnology, Brazilian Agricultural Research Corporation (EMBRAPA), Brasília, Brazil. 71 Instituto de Investigaciones de la Amazonía Peruana, Iquitos, Peru. ⁷²Cirad, UMR EcoFoG (AgroParisTech, CNRS, INRAE, Université des Antilles, Université de Guyane), Kourou, French Guiana. ⁷³Department of Anthropology, The University of Texas at Austin, Austin, TX, USA. ⁷⁴Forest Resources Management, Gembloux Agro-Bio Tech, University of Liège, Liège, Belgium. ⁷⁵AMAP, Universite de Montpellier, IRD, CNRS, CIRAD, INRAE, Montpellier, France. ⁷⁶Socioecosistemas y Cambio Climatico, Fundacion con Vida, Medellín, Colombia. ⁷⁷School of Forestry, Beijing Forestry University, Beijing, China. ⁷⁸Institute of Biological Sciences, Universidade Federal do Pará, Belém, PA, Brazil. 79 National Museum of Natural History, Smithsonian Institution, Washington, DC, USA. 80 School of Geography, Earth and Environmental Sciences, University of Birmingham, Birmingham, UK, 81School of Geography, Earth and Environmental Sciences, University of Plymouth, Plymouth, UK. 82 Embrapa Amazônia Oriental, Brazilian Agricultural Research Corporation (EMBRAPA), Brasília, Brazil. ⁸³Universidad Nacional Jorge Basadre de Grohmann (UNJBG), Tacna, Peru. ⁸⁴Forestry Research Institute of Ghana (FORIG), Kumasi, Ghana. ⁸⁵Smithsonian Institution Forest Global Earth Observatory (ForestGEO), Smithsonian Tropical Research Institute, Washington, DC, USA, ⁸⁶School of Biology, University of Leeds, Leeds, UK. ⁸⁷128 Busbridge Lane, Godalming, Surrey, UK. ⁸⁸Royal Botanic Garden Edinburgh, Edinburgh, UK. ⁸⁹Lukuru Wildlife Research Foundation, Kinshasa, Democratic Republic of Congo. ⁹⁰Division of Vertebrate Zoology, Yale Peabody Museum of Natural History, New Haven, CT, USA. 91Institute of Botany, Czech Academy of Sciences, Brno, Czech Republic. ⁹²Department of Botany, Palacký University in Olomouc, Olomouc, Czech Republic. 93CIRAD, UPR Forêts et Sociétés, Yamoussoukro, Côte d'Ivoire. 94 Institut National Polytechnique Félix Houphouët-Boigny, INP-HB,

Yamoussoukro, Côte d'Ivoire. 95 Instituto Venezolano de Investigaciones Científicas (IVIC), Caracas, Venezuela. ⁹⁶Département Hommes, Natures, Sociétés, Muséum National d'Histoire Naturel, Paris, France. ⁹⁷Universidad Nacional de San Antonio Abad del Cusco, Cusco, Peru. ⁸Biological and Environmental Sciences, University of Stirling, Stirling, UK. ⁹Instituto IMANI, Universidad Nacional de Colombia, Leticia, Colombia. ¹⁰⁰Living with Lakes Centre, Laurentian University, Sudbury, Canada. ¹⁰¹Department of Plant Sciences, University of Cambridge, Cambridge, UK. 102 Faculty of Science, Department of Botany and Plant Physiology, University of Buea, Buea, Cameroon. 103 PELD, Universidade do Estado de Mato Grosso, Nova Xavantina-MT, Brazil. ¹⁰⁴Tropical Peat Research Institute, Malaysian Palm Oil Board, Selangor, Malaysia. ¹⁰⁵Agteca, Santa Cruz, Bolivia. ¹⁰⁶Graduate School of Agriculture, Kvoto University, Kvoto, Japan. 107 Royal Botanic Gardens Kew, Richmond, London, UK. ¹⁰⁸Department of Earth Sciences, University of Hong Kong, Pok Ful Lam, Hong Kong Special Administrative Region, China. ¹⁰⁹Centre for Tropical Environmental and Sustainability Science (TESS) and College of Marine and Environmental Sciences, James Cook University, Douglas, OLD, Australia. 110 Uganda Programme, Wildlife Conservation Society, Kampala, Uganda. ¹¹¹Environmental Science and Policy, George Mason University, Fairfax, VA, USA. ¹¹²Botany Department, University of Ibadan, Ibadan, Nigeria. 113 Coordenação da Biodiversidade, Instituto Nacional de Pesquisas da Amazônia (INPA), Mauaus, Brazil. 114cE3c -Centre for Ecology, Evolution and Environmental Changes / Azorean Biodiversity Group, Universidade dos Açores, Angra do Heroísmo, Azores, Portugal. ¹¹⁵LIBRe – Laboratory for Integrative Biodiversity Research, Finnish Museum of Natural History, University of Helsinki, Helsinki, Finland. 116 Laboratório de Biogeoquímica Ambiental Wolfgang C. Pfeiffer, Universidade Federal de Rondônia, Porto Velho - RO, Brazil. 117 Faculdade de Ciências Agrárias, Biológicas e Sociais Aplicadas, Universidad do Estado de Mato Grosso, Nova Xavantina-MT, Brazil. ¹¹⁸Tropical Forests and People Research Centre, University of the Sunshine Coast, Sippy Downs, QLD, Australia. 119 Flamingo Land Ltd., North Yorkshire, UK. 120 UK Centre for Ecology and Hydrology, Wallingford, UK. ¹²¹School of International Tropical Forestry, Universiti Malaysia Sabah, Kota Kinabalu, Malaysia. ¹²²Escuela de Ciencias Forestales, Unidad Académica del Trópico, Universidad Mayor de San Simón, Sacta, Bolivia. 123 Faculty of Science, Universiti Brunei Darussalam, Brunei. ¹²⁴Agence Nationale des Parcs Nationaux, Libreville, Gabon.

125 Ministère de la Forêt, de la Mer, de l'Environnement, Chargé du Plan Climat, Libreville, Gabon. 126 Institute of Biology, University of Campinas, Campinas, SP, Brazil.

127 Facultad de Ingeniería Ambiental, Universidad Estatal Amazónica, Puyo, Pastaza, Ecuador. 128 Forest Research Centre, Sabah Forestry Department, Sepilok, Malaysia. ¹²⁹Carrera de Ingeniería Forestal, Universidad Tecnica del Norte, Ibarra, Ecuador. ¹³⁰Grupo de Ecosistemas Tropicales y Cambio Global, Universidad Regional Amazónica Ikiam, Tena, Ecuador. 131 Public Communications and Outreach Group, Parks and Recreation Division, Oakland Park, FL, USA. ¹³²Keller Science Action Center, Field Museum, Chicago, IL, USA. ¹³³Centre for Agricultural Research in Suriname (CELOS), Paramaribo, Suriname. ¹³⁴Instituto de Ciencias Naturales, Universidad Nacional de Colombia, Leticia, Colombia. 135 Department of Biology, Boston University, Boston, MA, USA, 136 Institute of Research for Forestry Development (INDEFOR), Universidad de los Andes, Mérida, Venezuela. ¹³⁷Bureau Waardenburg, Culemborg, Netherlands. ¹³⁸Socioecosistemas y Cambio Climatico, Fundacion Con Vida, Medellín, Colombia. 139 Centro de Conservacion, Investigacion y Manejo de Areas Naturales, CIMA Cordillera Azul, Lima, Peru, 140 lwokrama International Centre for Rainforest Conservation and Development, Georgetown. Guyana. ¹⁴¹Carboforexpert, Geneva, Switzerland. ¹⁴²Universidade Federal Rural da Amazônia/CAPES, Belém, PA, Brazil. 143 Museu Paraense Emílio Goeldi, Belém, PA, Brazil. 144 Faculty of Environmental Sciences and Natural Brazii. Facuity of Eliminimental Sciences and Natural Resource Management, Norwegian University of Life Sciences, As, Norway. ¹⁴⁵Instituto Federal do Acre, Rio Branco, AC, Brazii. ¹⁴⁶Universidad de San Antonio Abad del Cusco, Cusco, Peru. ¹⁴⁷Guyana Forestry Commission, Georgetown, Guyana. ¹⁴⁸Departamento de Biogeografía y Cambio Global, Museo Nacional de Ciencias Naturales, Consejo Superior de Investigaciones Científicas (MNCN-CSIC), Madrid, Spain. 149 Sustainable Landscapes and Food Systems, Center for International Forestry Research, Bogor, Indonesia. ¹⁵⁰Faculty of Forestry, University of British Columbia, Vancouver, Canada. ¹⁵¹Department of Forest Botany, Dendrology and Geobiocoenology, Mendel University in Brno, Brno, Czech Republic. 152 Department of Plant and Soil Science, School of Biological Sciences, University of Aberdeen, Aberdeen, UK. ¹⁵³Institute for Conservation Research, San Diego Zoo, San Diego, CA. USA. ¹⁵⁴Department of Plant Biology, Faculty of Sciences, University of Yaounde 1, Yaoundé, Cameroon. 155 Bioversity International, Yaoundé, Cameroon. 156UK Research and

Innovation, Innovate UK, London, UK. 157 Department of Geography, National University of Singapore, Singapore. ¹⁵⁸Naturalis Biodiversity Center, Leiden, Netherlands. ¹⁵⁹Systems Ecology, Vrije Universiteit Amsterdam, Amsterdam, Netherlands. ¹⁶⁰Faculty of Forestry, University of Toronto, Toronto, Canada. ¹⁶¹Universidad de los Andes, Merida, Colombia. ¹⁶²Wildlife Conservation Society, New York, NY, USA. ¹⁶³Yale School of Forestry and Environmental Studies, Yale University, New Haven, CT, USA. 164School of Geography, University of Notingham, Nottingham, UK. 16 der Hout Forestry Consulting, Rotterdam, Netherlands. ¹⁶⁶Van Hall Larenstein University of Applied Sciences, Velp, Netherlands. ¹⁶⁷Utrecht University, Utrecht, Netherlands. ¹⁶⁸Data for Sustainability, Axel, Netherlands. ¹⁶⁹School of Environmental and Forest Sciences, University of Washington, Seattle, OR, USA, 170 Centro de Investigación v Promoción del Campesinado, La Paz, Bolivia. ¹⁷¹Universidad Autónoma del Beni José Ballivián, Riberalta, Bolivia, ²School of Earth Sciences and Environmental Sustainability, Northern Arizona University, Flagstaff, AZ, USA. 173 Institut de Recherche en Ecologie Tropicale, Libreville, Gabon. 174School of Natural Sciences, University of Bangor, Bangor, UK. ¹⁷⁵University of Liberia, Monrovia, Liberia. ¹⁷⁶Forest Management, Centre for Agricultural Research in Suriname (CELOS), Paramaribo, Suriname. ¹⁷⁷Department of Geography and The Environment, University of Texas at Austin, Austin, TX, USA. ¹⁷⁸Tropenbos International, Wageningen, Netherlands.

†Present address: Amazonia Green Landscape Protection and Governance Programme, SOS Amazônia, 61 Pará St., Rio Branco, AC 69905-082, Brazil. ‡Present address: College of Biological and Environmental Sciences, Universidad San Francisco de Quito, Cumbayá, Ecuador. §Present address: Institute for Transport Studies, University of Leeds, Leeds, UK.

SUPPLEMENTARY MATERIALS

science.sciencemag.org/content/368/6493/869/suppl/DC1 Extended Acknowledgments Materials and Methods Figs. S1 to S15 Tables S1 to S3 References (34–87)

10 May 2019; accepted 5 March 2020 10.1126/science.aaw7578



Long-term thermal sensitivity of Earth's tropical forests

Martin J. P. Sullivan, Simon L. Lewis, Kofi Affurn-Baffoe, Carolina Castilho, Flávia Costa, Aida Cuni Sanchez, Corneille E. N. Ewango, Wannes Hubau, Beatriz Marimon, Abel Monteagudo-Mendoza, Lan Qie, Bonaventure Sonké, Rodolfo Vasquez Martinez, Timothy R. Baker, Roel J. W. Brienen, Ted R. Feldpausch, David Galbraith, Manuel Gloor, Yadvinder Malhi, Shin-Ichiro Aiba, Miguel N. Alexiades, Everton C. Almeida, Edmar Almeida de Oliveira, Esteban Álvarez Dávila, Patricia Alvarez Loayza, Ana Andrade, Simone Aparecida Vieira, Luiz E. O. C. Aragão, Alejandro Araujo-Murakami, Eric J. M. M. Arets, Luzmila Arroyo, Peter Ashton, Gerardo Aymard C., Fabricio B. Baccaro, Lindsay F. Banin, Christopher Baraloto, Plinio Barbosa Camargo, Jos Barlow, Jorcely Barroso, Jean-François Bastin, Sarah A. Baterman, Hans Beeckman, Serge K. Begne, Amy C. Bennett, Erika Berenguer, Nicholas Berry, Lilian Blanc, Pascal Boeckx, Jan Bogaert, Damien Bonal, Frans Bongers, Matt Bradford, Francis Q. Brearley, Terry Brncic, Foster Brown, Benoit Burban, José Luís Camargo, Wendeson Castro, Carlos Céron, Sabina Cerruto Ribeiro, Víctor Chama Moscoso, Jerôme Chave, Eric Chezeaux, Connie J. Clark, Fernanda Coelho de Souza, Murray Collins, James A. Comiskey, Fernando Cornejo Valverde, Massiel Corrales Medina, Lola da Costa, Martin Dancák, Greta C. Dargie, Stuart Davies, Nallaret Davia Cardozo, Thales de Haulleville, Marcelo Brilhante de Medeiros, Jhon del Aguila Pasquel, Géraldine Derroire, Anthony Di Fiore, Jean-Louis Doucet, Aurélie Dourdain, Vincent Droissart, Luisa Fernanda Duque, Romeo Ekoungoulou, Fernando Elias, Terry Erwin, Adriane Esquivel-Muelbert, Sophie Fauset, Joice Ferreira, Gerardo Flores Llampazo, Ernest Foli, Andrew Ford, Martin Glipin, Jefferson S. Hall, Keith C. Hamer, Alan C. Hamilton, David J. Harris, Terese B. Hart, Radim Hedl, Bruno Herault, Rafael Herrera, Niro Higuchi, Annette Hladik, Euridice Honorio Coronado, Isau Huamantupa-Chuquimaco, Walter Huaraca Huaeso, Kathryn J. Jeffery, Eliana Jimenez-Rojas, Michelle Kalamandeen, Marie Noël

Science **368** (6493), 869-874. DOI: 10.1126/science.aaw7578

Use of this article is subject to the Terms of Service

Thermal sensitivity of tropical trees

A key uncertainty in climate change models is the thermal sensitivity of tropical forests and how this value might influence carbon fluxes. Sullivan et al. measured carbon stocks and fluxes in permanent forest plots distributed globally. This synthesis of plot networks across climatic and biogeographic gradients shows that forest thermal sensitivity is dominated by high daytime temperatures. This extreme condition depresses growth rates and shortens the time that carbon resides in the ecosystem by killing trees under hot, dry conditions. The effect of temperature is worse above 32°C, and a greater magnitude of climate change thus risks greater loss of tropical forest carbon stocks. Nevertheless, forest carbon stocks are likely to remain higher under moderate climate change if they are protected from direct impacts such as clearance, logging, or fires.

Science, this issue p. 869

ARTICLE TOOLS http://science.sciencemag.org/content/368/6493/869

SUPPLEMENTARY http://science.sciencemag.org/content/suppl/2020/05/20/368.6493.869.DC1

RELATED http://science.sciencemag.org/content/sci/368/6493/813.full

REFERENCES This article cites 87 articles, 11 of which you can access for free

http://science.sciencemag.org/content/368/6493/869#BIBL

PERMISSIONS http://www.sciencemag.org/help/reprints-and-permissions

Use of this article is subject to the Terms of Service