



Prevalence, risk factors and genetic traits of *Salmonella* Infantis in Dutch broiler flocks

Lapo Mughini-Gras^{a,b,*}, Angela H.A.M. van Hoek^a, Tryntsje Cuperus^a, Cecile Dam-Deisz^a, Wendy van Overbeek^a, Maaïke van den Beld^a, Ben Wit^c, Michel Rapallini^d, Bart Wullings^d, Eelco Franz^a, Joke van der Giessen^{a,e}, Cindy Dierikx^a, Marieke Opsteegh^a

^a Centre for Infectious Disease Control (CIb), National Institute for Public Health and the Environment (RIVM), Bilthoven, the Netherlands

^b Institute for Risk Assessment Sciences (IRAS), Utrecht University, Utrecht, the Netherlands

^c Netherlands Food and Consumer Product Safety Authority (NVWA), Utrecht, the Netherlands

^d Wageningen Food Safety Research (WFSR), Wageningen, the Netherlands

^e Wageningen Bioveterinary Research (WBVR), Lelystad, the Netherlands

ARTICLE INFO

Keywords:
Salmonella
 Infantis
 Epidemiology
 Genomics
 Broilers

ABSTRACT

Salmonella Infantis is a poultry-adapted *Salmonella enterica* serovar that is increasingly reported in broilers and is also regularly identified among human salmonellosis cases. An emerging *S. Infantis* mega-plasmid (pESI), carrying fitness, virulence and antimicrobial resistance genes, is also increasingly found. We investigated the prevalence, genetic characteristics and risk factors for (pESI-carrying) *S. Infantis* in broilers. Faecal samples from 379 broiler flocks (in 198 farms with ≥ 3000 birds) in the Netherlands were tested. A questionnaire about farm characteristics was also administered. Sampling was performed in July 2018-May 2019, three weeks before slaughter. Fourteen flocks (in 10 farms) were *S. Infantis*-positive, resulting in a 3.7 % flock-level and 5.1 % farm-level prevalence. Based on multi-locus sequence typing (MLST), all isolates belonged to sequence type 32. All but one isolate carried a pESI-like mega-plasmid. Core-genome MLST showed considerable heterogeneity among the isolates, even within the same farm, with a few small clusters detected. The typical pESI-borne multi-resistance pattern to aminoglycosides, sulphonamide and tetracycline (93 %), as well as trimethoprim (71 %), was found. Additionally, resistance to (fluoro)quinolones based on *gyrA* gene mutations was detected. *S. Infantis* was found more often in flocks using salinomycin as coccidiostat, where flock thinning was applied or litter quality was poor, whereas employing external cleaning companies, wheat in feed, and vaccination against infectious bronchitis, were protective. Suggestive evidence for vertical transmission from hatcheries was found. A heterogeneous (pESI-carrying) *S. Infantis* population has established itself in Dutch broiler flocks, calling for further monitoring of its spread and a comprehensive appraisal of control options.

1. Introduction

In recent years, the ranking of the most common *Salmonella enterica* subspecies *enterica* serovars isolated from broilers has changed in the Netherlands and in other industrialized countries. Specifically, serovar Paratyphi B variant Java was the dominant serovar in broilers until about the year 2010, when the serovar Infantis began to increase sharply (EFSA/ECDC, 2019). Since 2014, *S. Infantis* has been the most commonly isolated serovar in broilers in the Netherlands, accounting for 78 % of all *Salmonella* isolates from broilers reported in 2018

(Vlaanderen et al., 2019). Also Europe-wide, *S. Infantis* is currently the most common serovar in broilers (46–57 %) and broiler meat (37–51 %), and its prevalence has increased in breeding flocks as well (EFSA/ECDC, 2018, 2019). Moreover, *S. Infantis* ranked fourth among the serovars isolated from human salmonellosis cases in the Netherlands and in Europe in the past few years, after *S. Enteritidis*, *S. Typhimurium* and its monophasic variant (Vlaanderen et al., 2019).

S. Infantis is also increasingly associated with antimicrobial resistance. In 2014, a multi-drug resistant *S. Infantis* strain was identified in Israel carrying a mega-plasmid of approximately 300 Kb named pESI

* Corresponding author at: Centre for Infectious Disease Control (CIb), National Institute for Public Health and the Environment (RIVM), Antonie van Leeuwenhoeklaan 9, 3721MA Bilthoven, the Netherlands.

E-mail address: lapo.mughini.gras@rivm.nl (L. Mughini-Gras).

<https://doi.org/10.1016/j.vetmic.2021.109120>

Received 17 February 2021; Accepted 13 May 2021

Available online 15 May 2021

0378-1135/© 2021 The Author(s). Published by Elsevier B.V. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

(plasmid of emerging *S. Infantis*) (Aviv et al., 2014); genes associated with antibiotic resistance, virulence and fitness were located on this plasmid. *S. Infantis* isolates carrying the pESI were shown to have increased ability to form biofilms and attach to host cells, leading to raised pathogenicity in mouse models (Aviv et al., 2016). After the first detection, pESI-carrying *S. Infantis* isolates have been reported worldwide, particularly in Europe and in the United States of America (USA) (Alba et al., 2020; Franco et al., 2015; Hindermann et al., 2017; Tate et al., 2017). Extended spectrum β -lactamase (ESBL) genes have also been found on the pESI of some *S. Infantis* isolates from Europe and the USA. Moreover, a *S. Infantis* clone was found in Italy that contained an *mcr-1* gene on a second plasmid, which causes resistance to the last-resort antibiotic colistin (Carfora et al., 2018). In a mouse model, it has also been shown that pESI can be transferred from *S. Infantis* to gut bacteria or to other *Salmonella* serovars through horizontal gene transfer during co-infection (Aviv et al., 2016), which entails that multi-resistance and increased virulence can be spread to other pathogens as well.

To provide baseline epidemiological data on (pESI-carrying) *S. Infantis*, this study investigated the prevalence, antibiotic resistance profiles and genetic characteristics of (pESI-carrying) *S. Infantis* in broiler flocks in the Netherlands, as well as the risk factors associated with its occurrence in those flocks.

2. Materials and methods

2.1. Sample collection and analysis

Fecal samples from 198 Dutch broiler farms with at least 3000 birds included in the annual surveillance system for zoonotic pathogens in farm animals in the Netherlands were collected for *Salmonella* testing. In parallel with sample collection, a comprehensive questionnaire about farm characteristics (e.g. farm infrastructure, husbandry practices, health management, etc.) was completed by the farmer. Sampling was performed between July 2018 and May 2019 within three weeks before slaughter. If available, two poultry houses (i.e. flocks) per farm were sampled. This was possible on 181 farms, whereas on 17 farms, only one poultry house was present, resulting in 379 flocks sampled. Per flock, three fecal samples (each consisting of 12 fresh droppings) were taken, i.e. at the front, central, and hindmost parts of the poultry house itself, and individually frozen with an equal amount of glycerol pending further investigation.

Subsequently, the three samples from each flock were pooled together by mixing 10 g of feces/glycerol material from each individual sample. This pooled sample of 30 g of feces/glycerol per flock was then suspended into 270 ml of Buffered Peptone Water (Biotrading) and incubated overnight at 37 °C. Afterwards, 100 μ l of the suspension was transferred to Modified Semi-solid Rappaport-Vassiliadis (Difco) plates with novobiocin (Oxoid). After incubation for 24–48 hours at 41.5 °C, *Salmonella*-suspected colonies were re-incubated on Brilliance *Salmonella* Agar (BSA; Oxoid) plates for 18–22 hours at 37 °C. A maximum of three colonies per BSA plate were then grown on blood agar plates (Oxoid), after which they were serotyped using the xMAP *Salmonella* serotyping assay (Luminex). For isolates with negative or inconclusive reactions with O and H antisera, MALDI-TOF was used for confirmation of genus *Salmonella*. Minimum Inhibitory Concentrations (MIC) were determined by the dilution method using the EUVSEC panel for *Salmonella* and *E. coli* (Sensititre). This panel contains the following antibiotics: ampicillin (AMP), azithromycin (AZI), cefotaxime (CTX), ceftazidime (CAZ), chloramphenicol (CHL), ciprofloxacin (CIP), colistin (COL), gentamicin (GEN), meropenem (MER), nalidixic acid (NAL), sulphamethoxazole (SMX), tetracycline (TET), tigecycline (TIG), trimethoprim (TMP). The epidemiological cut-off values (ECOFFs) published by EUCAST (http://www.eucast.org/mic_distributions_and_coffs/) were applied.

2.2. Sequencing and bioinformatics

S. Infantis isolates were grown overnight in 9 ml Brain Heart Infusion tubes (Oxoid) at 37 °C. Cell pellets, obtained from 1.8 ml of the culture, were resuspended and stored in 450 μ l DNA/RNA Shield (Zymo) at 4 °C. The DNA was extracted and sequenced by BaseClear (Leiden, the Netherlands). Paired-end 2 \times 150bp short-reads were generated using a Nextera XT library preparation (Illumina) and were sequenced on a NovaSeq 6000 system (Illumina). The obtained reads were first trimmed and filtered with ERNE-filter (Del Fabbro et al., 2013) and *de novo* assembled with SPAdes 3.10.0 (Bankevich et al., 2012).

The sequence type (ST) was determined from the assembled genomes with *in silico* multi-locus sequence typing (MLST) using the traditional seven house-keeping gene scheme (Achtman et al., 2012). The *S. Infantis* isolates were also analyzed with the Enterobase *Salmonella* core genome MLST scheme (Alikhan et al., 2018) using Ridom SeqSphere⁺ 6.0.2 (Ridom; Münster, Germany). The assembled genomes were screened for antimicrobial resistance genes (acquired as well as point mutations) based on the entries in the ResFinder database (Zankari et al., 2012) and for the presence of 233 *Salmonella* virulence factors (Kuijpers et al., 2019). To investigate whether the *S. Infantis* isolates harbored a pESI-like plasmid, the coding DNA sequences (CDS) of 283 genes were extracted from the pESI reference plasmid of *S. Infantis* strain 119944 (accession number: CP047882) and implemented as a scheme in SeqSphere⁺. This included, for example, determinants specific for IncI1 plasmids and several toxin-antitoxin systems, K88 and other fimbrial factors, genes from the mercury resistance operon (*merTPCBDE*), but also the antimicrobial resistance genes *aadA1*, *dfrA14*, *sul1* and *tet(A)*. These CDS also allowed to perform an *in silico* IncI1 plasmid MLST (pMLST) analysis (Carattoli et al., 2014). Nucleotide BLAST was used to confirm and investigate the pESI screening results of the broiler *S. Infantis* isolates in more detail. All the genomic sequences are available at the European Nucleotide Archive (ENA) within the study accession number PRJEB42821.

2.3. Risk factor analysis

Risk factors for the occurrence of *S. Infantis* in the flocks were assessed using logistic regression analysis in which the binary outcome variable was the presence or absence of *S. Infantis* in the flock. In total, 138 independent variables were obtained from the questionnaire and tested for association with the outcome variable. Flock age at sampling was always controlled in the analyses. A cluster-robust sandwich variance estimator was used to account for clustering of flocks at the farm level. Associations were expressed as odds ratios (OR) with 95 % confidence intervals (95 % CI). A complete-record analysis was performed. All models showed overall statistical significance (likelihood-ratio test, $p < 0.05$) and goodness-of-fit (Hosmer–Lemeshow test, $p > 0.05$). Pairwise correlations between independent variables were also assessed. Analyses were performed using Stata v. 16 (StataCorp, College Station, TX, USA). Overall, a $p < 0.05$ was considered statistically significant.

3. Results

3.1. Prevalence and resistance phenotype

There were 35 *Salmonella*-positive flocks (corresponding to 23 farms), resulting in a flock-level prevalence of 9.2 % (95 %CI 6.1–13.8 %) and a farm-level prevalence of 11.6 % (95 %CI 7.5–16.9 %) for *Salmonella* spp. Serotyping of these *Salmonella* isolates at the farm-level revealed the presence of *S. Paratyphi* B variant Java (6.1 %, 95 %CI 3.5–10.3), *S. Infantis* (5.1 %, 95 %CI 2.4–9.1 %) and *S. Agona* (0.5 %, 95 %CI 0.1–2.8 %). This corresponded to a flock-level prevalence of respectively 5.3 % (95 %CI 3.4–8.0 %), 3.7 % (95 %CI 1.9–7.0 %) and 0.3 % (95 %CI 0.0–1.5 %). For *S. Infantis* in particular, 14 flocks of ten farms were positive for this serovar.

The antimicrobial resistance phenotype of these 14 *S. Infantis* isolates revealed that one isolate was fully susceptible while the other 13 isolates were resistant to either four or five antimicrobials (Table 1). The most common phenotypic profile encountered among the 13 resistant isolates was CIP-NAL-SUL-TET-TMP (77 %). The other phenotype was CIP-NAL-SUL-TET (23 %).

3.2. Genetic traits of *S. Infantis* isolates

In silico MLST showed that all 14 *S. Infantis* isolates belonged to the same sequence type (ST), i.e. ST32. The cgMLST analysis of these isolates, including a pESI-positive reference (*S. Infantis* strain 119944; accession number: CP047881) showed considerable variation among the isolates included in this study. Even isolates from different flocks of the same farm (indicated with A and B in the similarly colored sphere of Fig. 1) were never 100 % identical, although they clustered together. Overall, three clusters and four isolates not belonging to any of these were identified (Fig. 1). The largest cluster (Cluster 1) contained six isolates originating from four different farms.

In silico screening of the assembled genomes showed a high prevalence of five antibiotic resistance genes. They encoded resistance against four different classes of antibiotics; aminoglycosides (*aadA1* and *aph(3')-Ic*), sulphonamide (*sul1*), tetracycline (*tet(A)*) and trimethoprim (*drfA14*). An indication for (fluoro)quinolone resistance was also observed among the *S. Infantis* isolates due to mutations at common positions in *gyrA*, which results in amino acid substitutions in *gyrA* (Table 1).

In silico screening for the presence of 233 *Salmonella* virulence factors revealed that all isolates included in this study harbored 157 of these genes (Supplementary Table S1). Most of these virulence gene sequences were identical (92.7 %, 144/157) to the virulence determinants in the control *S. Infantis* strain 119944, but 13 of them showed genetic differences in one or more isolates (Table 2). Five genes contained a single nucleotide polymorphism (SNP), which did not result in an amino acid substitution because they were synonymous SNPs. The other eight virulence genes contained non-synonymous SNPs. In the case of the outer membrane protein encoded by *sinI*, this SNP resulted in a premature stop codon (PMSC) that could attenuate virulence for the 11 isolates in which this was found. Only in the isolates from farms D and J this SNP was not found. As shown by the cgMLST-based MST (Fig. 1), the isolates from these farms were the most dissimilar when compared to the other isolates (Table 1).

In silico screening of the assembled genomes for the presence of a pESI-like mega-plasmid showed that only one of the 14 isolates did not contain this plasmid, i.e. 367SI-1 (farm J). In the other isolates, over 95 % of the 283 genes of the pESI reference genome were found

(Supplementary Table S2). However, three isolates (from farms D and H) appeared to miss a class 2 integron that includes the trimethoprim resistance gene *drfA14* (Supplementary Table S2). This corresponds to the phenotype, as well as the screening results for antibiotic resistance genes (Table 1). Because of the use of short-read Illumina sequencing and due to working with *de novo* assemblies, some results regarding common IS family transposases were ambiguous. The *in silico* IncI1 pMLST analysis did not reveal the presence of *repI1*, which prevented determination of a ST. However, the other genes of the IncI1 pMLST scheme were identified and showed for all except one isolate the same allelic profile, i.e. *ardA-2*, *trbA-21*, *sogS-9*, and *pilL-3*. The isolate from farm E (130SI-1) differed in the pMLST results for the *trbA* gene: *trbA-46*.

3.3. Risk factors

Given the limited number of outcome events (i.e. 14 *S. Infantis* positive flocks), only univariable associations (adjusted for broiler age at sampling and clustering of flocks at the farm level) were assessed. Three factors were significantly associated with increased occurrence of *S. Infantis* in broiler flocks (Table 3): use of the coccidiostat drug salinomycin (OR 4.4), thinning of birds once (OR 2.2) or twice (OR 5.9) before sampling vs. no thinning at all, and litter quality being mostly 'wet and sticky' (OR 5.2) rather than 'dry and fresh'. Three factors were significantly associated with decreased occurrence of *S. Infantis* (Table 3): cleaning and disinfection operations performed by both farm personnel and external cleaning companies (OR 0.2) vs. farm personnel only, providing wheat in the feed (OR 0.2), and vaccinating birds against infectious bronchitis (OR 0.2).

Of these factors, only three were significantly and negatively associated with each other. These were 'cleaning and disinfection' and 'wheat feed provided' (phi-coefficient -0.14, $p = 0.006$) and 'cleaning and disinfection' and 'coccidiostat salinomycin used' (phi-coefficient -0.15, $p = 0.004$).

4. Discussion

Salmonella *Infantis* is one of the most common serovars causing human salmonellosis in Europe and is strongly associated with poultry (Alba et al., 2020). In this study, we determined the prevalence, risk factors and molecular characteristics of *S. Infantis*, including pESI-carrying *S. Infantis*, in broiler farms in the Netherlands.

A *S. Infantis* prevalence of 5.1 % at farm-level and of 3.7 % at flock-level was found. Moreover, a pESI-like mega-plasmid was present in 92.9 % of the obtained *S. Infantis* isolates. As the overall *Salmonella* prevalence was only twice as much as that of *S. Infantis*, it was confirmed that *S. Infantis* indeed is a dominant serovar in broilers. This

Table 1

Phenotypic resistance profile and *in silico* screening results for antibiotic resistance genes of the fourteen *S. Infantis* isolates from broiler flocks.

Isolate	Farm	Flock	Phenotypic profile [§]	<i>aadA1</i> [†]	<i>aph(3')-Ic</i> [†]	<i>drfA14</i> [†]	<i>sul1</i> [†]	<i>tet(A)</i> [†]	GyrA position 83*	GyrA position 87*
15SI-1	A	A	CIP-NAL-SUL-TET-TMP	+	-	+	+	+	wt	G
16SI-1	A	B	CIP-NAL-SUL-TET-TMP	+	-	+	+	+	wt	G
38SI-1	B	A	CIP-NAL-SUL-TET-TMP	+	+	+	+	+	wt	G
39SI-1	B	B	CIP-NAL-SUL-TET-TMP	+	+	+	+	+	wt	G
46SI-1	C	A	CIP-NAL-SUL-TET-TMP	+	+	+	+	+	wt	G
86SI-1	D	A	CIP-NAL-SUL-TET	+	-	-	+	+	Y	wt
87SI-1	D	B	CIP-NAL-SUL-TET	+	-	-	+	+	Y	wt
130SI-1	E	A	CIP-NAL-SUL-TET-TMP	+	+	+	+	+	wt	G
144SI-1	F	A	CIP-NAL-SUL-TET-TMP	+	+	+	+	+	wt	G
234SI-1	G	A	CIP-NAL-SUL-TET-TMP	+	+	+	+	+	wt	G
235SI-1	G	B	CIP-NAL-SUL-TET-TMP	+	+	+	+	+	wt	G
275SI-1	H	A	CIP-NAL-SUL-TET	+	-	-	+	+	wt	G
302SI-1	I	A	CIP-NAL-SUL-TET-TMP	+	+	+	+	+	wt	G
367SI-1	J	A	Fully susceptible	-	-	-	-	-	wt	wt

Note: [§] CIP, ciprofloxacin; NAL, nalidixic acid; SMX, sulfamethoxazole; TET, tetracycline; TMP, trimethoprim. [†]*aadA1*: aminoglycoside resistance gene, *aph(3')-Ic*: aminoglycoside resistance gene, *drfA14*: trimethoprim resistance gene, *sul1*: sulfonamide resistance gene, *tet(A)*: tetracycline resistance gene. *wt = wildtype (GyrA position 83 is a serine (S) while position 87 is aspartic acid (D), Y = tyrosine, G = glycine).




Fig. 1. Minimum Spanning Tree (MST) of the cgMLST analysis of fourteen *S. Infantis* isolates (ST32) from broilers. The MST is based on 2963 genes of the total of 3002 present in the official cgMLST scheme. Each sphere represents an isolate and the number on the connecting lines shows the number of genes different between the ST32 isolates. Clusters are indicated with a grey background color, with a cluster definition set to a maximum distance of seven genes. The different colors of the spheres display the ten positive farms, while the A and B in the spheres indicate from which flock of a farm the *S. Infantis* was isolated.

Table 2
Variants of virulence determinants in comparison to *S. Infantis* strain 119944.

Gene	Function	SNP ^S	AA ^S substitution	Isolate (Farm)
<i>fimF</i>	Type I fimbriae adaptor protein	T62C	V21A	367SI-1 (J)
<i>hilD</i>	Invasion regulatory protein	G468A	–	275SI-1 (H)
<i>invA</i>	Cell invasion protein	A1218G	–	367SI-1 (J)
<i>invE</i>	Cell Invasion protein	C456T	–	302SI-1 (I)
<i>ratB</i>	Outer membrane protein	T3898C	Y1300H	86SI-1 (D), 87SI-1 (D)
<i>sifB</i>	Secreted effector protein	G766A	V256I	6SI-1 (D), 87SI-1 (D)
<i>sinI</i>	Outer membrane protein	T589G	E197Stop	15SI-1 (A), 16SI-1 (A), 38SI-1 (B), 39SI-1 (B), 46SI-1 (C), 130SI-1 (E), 144SI-1 (F), 234SI-1 (G), 235SI-1 (G), 275SI-1 (H), 302SI-1 (I)
<i>strP</i>	E3 ubiquitin-protein ligase	C178A	A60I	86SI-1 (D), 87SI-1 (D)
<i>sscB</i>	Secretion system chaperone	T238A	F80I	15SI-1 (A), 16SI-1 (A), 38SI-1 (B), 39SI-1 (B), 46SI-1 (C), 130SI-1 (E), 144SI-1 (F), 234SI-1 (G), 235SI-1 (G), 275SI-1 (H), 302SI-1 (I)
<i>ssrA</i>	Sensor kinase	C1894T	H632Y	130SI-1 (E)
<i>stdB</i>	Outer membrane usher protein	C438T	–	275SI-1 (H)
<i>stfA</i>	Fimbrial subunit	G192A	M64I	86SI-1 (D), 87SI-1 (D)
<i>sthD</i>	Fimbrial subunit	C420T	–	275SI-1 (H)

Note: ^S SNP = single nucleotide polymorphism; AA = amino acid.

also concurs with findings from poultry meat at retail (Vlaanderen et al., 2019) and reflects the current situation in several other European countries (Alba et al., 2020; EFSA/ECDC, 2018, 2019; Pate et al., 2019). Overall, these findings confirm that *S. Infantis* has successfully established itself in the Dutch broiler sector.

The rapid spread of *S. Infantis* in broilers has been related to the

occurrence of the pESI mega-plasmid carrying genes associated with fitness, virulence and antimicrobial resistance (Aviv et al., 2016, 2014). In this study, pESI was found in all but one isolate. Yet, despite the ubiquity of pESI and the presence of only one ST among these isolates (i. e. ST32), the *S. Infantis* population was heterogeneous in terms of molecular relatedness at the cgMLST level, with some variation even between isolates from the same farm. This is consistent with the conclusions of a recent molecular analysis of *S. Infantis* isolates from all over Europe (Alba et al., 2020). Indeed, that study found the European *S. Infantis* population to be heterogeneous, with different genetic clusters defined at core genome level. However, the same study reported that pESI-like variants were present in 64% of the isolates and were more genetically homogeneous and capable of infecting different clonal lineages in most of the countries. The same was observed here, as the identified genes of the IncI1 pMLST scheme revealed that all but one isolate had the same allelic profile. Therefore, pESI appears to be able to colonize different clonal lines of *S. Infantis* and to spread widely in poultry populations.

Of the three clusters of *S. Infantis* isolates found here, the largest included six isolates from four different farms. From the questionnaires completed by these farms, it appeared that they received chicks from the same two hatcheries. Although these hatcheries also provided chicks to flocks on other farms that tested negative, the observation is suggestive of possible vertical transmission. Together with Enteritidis, Typhimurium, Hadar and Virchow, *Infantis* is one of the target serovars included in the current control programs for *Salmonella* in breeding flocks of *Gallus gallus* (Commission Regulation (EU) No 200/2010), but not for broiler flocks (Commission Regulation (EU) No 200/2012) in the European Union.

The public health implications of *S. Infantis* in broiler flocks, particularly when the isolates carry pESI, are unclear. The proportion of *S. Infantis* infections among human salmonellosis patients in the Netherlands has stabilized in recent years and these infections do not appear to be associated with increased invasiveness in humans (Mughini-Gras et al., 2020). Moreover, in a mouse experiment, *S. Infantis* was found to be less virulent than *S. Typhimurium* (Aviv et al., 2019). However, *S. Infantis* without pESI was used in that experiment. In a different mouse experiment, it appeared that infection with *S. Infantis* causes a more severe inflammation when carrying pESI (Aviv et al., 2014). The consequences of infections with *S. Infantis* carrying pESI in humans, including the severity of infection and the implication of antibiotic resistance, have not yet been investigated.

It has been proven that pESI frequently harbors resistance genes

Table 3
Factors significantly associated with increased or decreased occurrence of *S. Infantis* in broiler flocks.

Risk factor	<i>S. Infantis</i> positive flocks	<i>S. Infantis</i> negative flocks	OR ¹	95 % CI ¹		p
Cleaning and disinfection Performed by farm personnel only	7 (50 %)	78 (22 %)	Reference			
Performed by farm personnel and professional cleaners	7 (50 %)	285 (78 %)	0.233	0.057	0.955	0.043
Wheat feed provided						
No	6 (43 %)	52 (14 %)	Reference			
Yes	8 (57 %)	313 (86 %)	0.235	0.055	0.999	0.049
Coccidiostat salinomycin used						
No	4 (29 %)	205 (56 %)	Reference			
Yes	10 (71 %)	160 (44 %)	4.372	1.101	17.363	0.036
Chickens vaccinated against infectious bronchitis						
No	2 (14 %)	11 (3%)	Reference			
Yes	12 (86 %)	352 (97 %)	0.170	0.034	0.855	0.031
Thinning of flock before sampling						
No	7 (50 %)	268 (75 %)	Reference			
Yes, once	4 (29 %)	78 (22 %)	2.182	0.382	12.451	0.380
Yes, twice	3 (21 %)	13 (3%)	5.883	1.025	33.760	0.047
Litter quality						
Mostly dry and fresh	10 (77 %)	334 (94 %)	Reference			
Mostly wet and sticky	3 (23 %)	21 (6%)	5.183	1.167	23.023	0.031

Note: ¹Estimates are adjusted for broilers' age at sampling and clustering of flocks at the holding level. OR = Odds Ratio. 95 % CI = 95 % Confidence Interval. Totals do not add up because of missing values.

against at least three antimicrobial classes (sulphonamides, tetracyclines and trimethoprim), with variable carriage of the aminoglycosides resistance gene *aadA1* and resistance to heavy metals and several toxin/anti-toxin systems (Alba et al., 2020; Aviv et al., 2014; Franco et al., 2015). The typical pESI-associated genes *aadA1*, *dfrA14*, *sul1*, and *tet(A)* were found in the isolates included in the present study, with resistance gene rates being particularly high (92.9 %) for aminoglycosides, sulphonamides and tetracycline, and somewhat lower (71.4 %) for trimethoprim. The presence of nearly all of these genes can be linked to the antimicrobial resistance phenotypes of the *S. Infantis* isolates (Table 1). Unfortunately, *aadA1* encodes for resistance to aminoglycoside antibiotics not included in the EUVSEC panel, i.e. streptomycin and spectinomycin. This is also the case for *aph(3')-Ic* (not part of pESI) which causes resistance to the aminoglycosides; kanamycin, neomycin and paromycin. Otherwise, a 100 % match between antimicrobial phenotype and genotype was found. The SUL-TET-TMP profile can be linked to the presence of *sul1*, *tet(A)*, and *dfrA14*, respectively. Moreover, in all isolates with pESI and a CIP-NAL phenotype, resistance to these (fluoro)quinolones was the result of mutations in the chromosomal

gyrA gene. The phenotypic and genotypic resistance profiles were consistent with the previously described pESI-carrying *S. Infantis* isolates from several countries (Alba et al., 2020; Franco et al., 2015; Hindermann et al., 2017). Resistance genes to macrolides, which have been described in 18 % of the isolates in an European study (Alba et al., 2020), were not found in the isolates investigated here. Interestingly, ESBL genes were not found either, whereas they have been reported in *S. Infantis* isolates from human patients in the Netherlands (Carfora et al., 2018). Moreover, the *mcr-1* gene causing resistance to colistin, which was reported previously (Carfora et al., 2018), was also not encountered in this study.

An increased risk for *S. Infantis* colonization was found for flocks in which salinomycin was used, as well as flocks where thinning was applied or where litter quality was poor, whereas the use of professional cleaners (i.e. employing external cleaning companies vs. own personnel only), providing wheat in the feed, and vaccinating against infectious bronchitis, had protective effects. Salinomycin is an ionophoric coccidiostat that is widely used as a supplement in poultry feed. Although it has been reported that salinomycin may reduce *Salmonella* prevalence in broilers (Bolder et al., 1999), other authors found no reduction in *Salmonella* prevalence or load when using this coccidiostat (Scalzo et al., 2004). On the contrary, they observed an increase of one log unit on *Salmonella* colony forming units between control and salinomycin-treated chickens. In addition, experimentally infected chickens (with *S. Typhimurium*) showed higher *Salmonella* isolation rates when treated with salinomycin vs. control group during the 3rd week post-dosing (Ford et al., 1981). Moreover, the latter study reported that salinomycin caused the decline of salmonellae to be more gradual, although at the end of the experiment the treatment and control groups were comparable. It is therefore difficult to explain why salinomycin treatment would be a risk factor for *S. Infantis* colonization, but it might be related to the overall farm management quality rather than the effect of the drug itself. Indeed, the fact that poor litter quality was also a risk factor and that employing professional cleaners was protective (as also found previously in Belgian broiler flocks as a risk factor for the presence of *Salmonella* spp. (Namata et al., 2009)) and negatively associated with salinomycin treatment itself, seems to point to (suboptimal) farm management practices as the main driver of *S. Infantis* colonization in broilers. The hypothesis is therefore that the use of coccidiostat drugs act as a proxy for lower quality farm management, such as lower biosecurity, as the need of using coccidiostat drugs implies the existence of that issue (i.e. coccidial infection risk) that cannot be controlled otherwise (e.g. with biosecurity) in the farm. However, it is also true that almost all (95 %) of the farms in this study used a coccidiostat (salinomycin or other drugs), so it cannot be excluded that the issue concerns specifically this component and that the use of it, alike other antimicrobials, might play a role (e.g. selective advantage) as to favor the establishment of (pESI-carrying) *S. Infantis* in the flocks.

Also flock thinning is by its nature a well-known 'breach of biosecurity', as well as a stressor for the birds, and a clear increase in *S. Infantis* colonization risk was observed as more thinning cycles were applied. The risks concern mainly the passive transfer of microorganisms, including *Salmonella* among others, from previously visited farms to the remaining flock during catching via, e.g., vehicles, crates, crews (boots, clothes, etc.) (FAO/WHO, 2009). Less straightforward is the interpretation of the protective effects associated with the use of wheat in the feed and vaccination against infectious bronchitis, which might, once again, mirror the overall farm management standards, as well as have genuine effects due to, e.g., improved health or welfare of the broilers resulting from the applied dietary and immunization schemes.

In conclusion, this study shows that a heterogeneous (pESI-carrying) *S. Infantis* population has successfully established itself in broiler flocks in the Netherlands, with a prevalence of around 5% and some indications of potential vertical transmission. The isolates analyzed here were characterized by a high rate of resistance genes and phenotypical resistance to the typical pattern of multi-drug resistance to

aminoglycosides, (fluoro)quinolones, sulphonamides, tetracycline, and trimethoprim. Although the incidence of human *S. Infantis* infections in the Netherlands appears to be stable (Vlaanderen et al., 2019), the virulence and antimicrobial resistance genes harbored by pESI and the increasing trends observed in poultry worldwide call for continuous monitoring of the spread of *S. Infantis* and its large virulence plasmid, including a comprehensive appraisal of the options for control. These options should consider in particular biosecurity, as most of the risk factors identified seemed to be related, directly or indirectly, to this topic and less to other farm characteristics.

Declaration of Competing Interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Acknowledgements

This study was supported by the Dutch Ministry of Agriculture, Nature and Food Quality (grant number 9.2.24.B). The authors are thankful to Anjo Verbruggen and Sjoerd Kuiling for their contribution to the laboratory work.

Appendix A. Supplementary data

Supplementary material related to this article can be found, in the online version, at doi:<https://doi.org/10.1016/j.vetmic.2021.109120>.

References

- Achtman, M., Wain, J., Weill, F.X., Nair, S., Zhou, Z., Sangal, V., Krauland, M.G., Hale, J. L., Harbottle, H., Uesbeck, A., Dougan, G., Harrison, L.H., Brisse, S., 2012. Multilocus sequence typing as a replacement for serotyping in *Salmonella enterica*. *PLoS Pathog.* 8, e1002776.
- Alba, P., Leekitcharoenphon, P., Carfora, V., Amoroso, R., Cordaro, G., Di Matteo, P., Ianzano, A., Iurescia, M., Diaconu, E.L., Study Group, E.-E.-A.N, Pedersen, S.K., Guerra, B., Hendriksen, R.S., Franco, A., Battisti, A., 2020. Molecular epidemiology of *Salmonella infantis* in Europe: insights into the success of the bacterial host and its parasitic pESI-like megaplasmid. *Microb. Genom.* 6.
- Alikhan, N.F., Zhou, Z., Sergeant, M.J., Achtman, M., 2018. A genomic overview of the population structure of *Salmonella*. *PLoS Genet.* 14.
- Aviv, G., Tsyba, K., Steck, N., Salmon-Divon, M., Cornelius, A., Rahav, G., Grassl, G.A., Gal-Mor, O., 2014. A unique megaplasmid contributes to stress tolerance and pathogenicity of an emergent *Salmonella enterica* serovar *Infantis* strain. *Environ. Microbiol.* 16, 977–994.
- Aviv, G., Rahav, G., Gal-Mor, O., 2016. Horizontal transfer of the *Salmonella enterica* serovar *infantis* resistance and virulence plasmid pESI to the gut microbiota of warm-blooded hosts. *mBio* 7.
- Aviv, G., Cornelius, A., Davidovich, M., Cohen, H., Suwandi, A., Galeev, A., Steck, N., Azriel, S., Rokney, A., Valinsky, L., Rahav, G., Grassl, G.A., Gal-Mor, O., 2019. Differences in the expression of SPI-1 genes pathogenicity and epidemiology between the emerging *Salmonella enterica* serovar *Infantis* and the model *Salmonella enterica* serovar *Typhimurium*. *J. Infect. Dis.* 220, 1071–1081.
- Bankevich, A., Nurk, S., Antipov, D., Gurevich, A.A., Dvorkin, M., Kulikov, A.S., Lesin, V. M., Nikolenko, S.I., Pham, S., Prjibelski, A.D., Pyshkin, A.V., Sirotkin, A.V., Vyahhi, N., Tesler, G., Alekseyev, M.A., Pevzner, P.A., 2012. SPAdes: a new genome assembly algorithm and its applications to single-cell sequencing. *J. Comput. Biol.* 19, 455–477.
- Bolder, N.M., Wagenaar, J.A., Putirulan, F.F., Veldman, K.T., Sommer, M., 1999. The effect of flavophospholipol (Flavomycin) and salinomycin sodium (SacoX) on the excretion of *Clostridium perfringens*, *Salmonella enteritidis*, and *Campylobacter jejuni* in broilers after experimental infection. *Poult. Sci.* 78, 1681–1689.
- Carattoli, A., Zankari, E., García-Fernández, A., Larsen, M.V., Lund, O., Villa, L., Aarestrup, F.M., Hasman, H., 2014. *In silico* detection and typing of plasmids using plasmidfinder and plasmid multilocus sequence typing. *Antimicrob. Agents Chemother.* 58, 3895–3903.
- Carfora, V., Alba, P., Leekitcharoenphon, P., Ballaro, D., Cordaro, G., Di Matteo, P., Donati, V., Ianzano, A., Iurescia, M., Stravino, F., Tagliaferri, T., Battisti, A., Franco, A., 2018. Colistin resistance mediated by mcr-1 in ESBL-Producing, multidrug resistant *Salmonella infantis* in broiler chicken industry, Italy (2016–2017). *Front. Microbiol.* 9, 1880.
- Del Fabbro, C., Scalabrini, S., Morgante, M., Giorgi, F.M., 2013. An extensive evaluation of read trimming effects on illumina NGS data analysis. *PLoS One* 8.
- EFSA/ECDC, 2018. The European Union summary report on trends and sources of zoonoses, zoonotic agents and food-borne outbreaks in 2017. *EFSA J.* 16, 262.
- EFSA/ECDC, 2019. The European Union one health 2018 zoonoses report. *EFSA J.* 17, e05926.
- FAO/WHO, 2009. *Salmonella and Campylobacter in Chicken Meat in Microbiological Risk Assessment Series 19*. FAO/WHO, Rome, Italy.
- Ford, A.M., Fagerberg, D.J., Quarles, C.L., George, B.A., McKinley, G.A., 1981. Influence of salinomycin on incidence, shedding, and antimicrobial resistance of *Salmonella typhimurium* in experimentally infected broiler chicks. *Poult. Sci.* 60, 2441–2453.
- Franco, A., Leekitcharoenphon, P., Feltrin, F., Alba, P., Cordaro, G., Iurescia, M., Tolli, R., D'Incau, M., Staffolani, M., Di Giannatale, E., Hendriksen, R.S., Battisti, A., 2015. Emergence of a clonal lineage of multidrug-resistant ESBL-Producing *Salmonella infantis* transmitted from broilers and broiler meat to humans in Italy between 2011 and 2014. *PLoS One* 10, e0144802.
- Hindermann, D., Gopinath, G., Chase, H., Negrete, F., Althaus, D., Zurfluh, K., Tall, B.D., Stephan, R., Nuesch-Inderbinen, M., 2017. *Salmonella enterica* serovar *infantis* from food and human infections, Switzerland, 2010–2015: poultry-related multidrug resistant clones and an emerging ESBL producing clonal lineage. *Front. Microbiol.* 8, 1322.
- Kuijpers, A.F.A., Marinovic, A.A.B., Wijnands, L.M., Delfgou-Van Asch, E.H.M., Van Hoek, A.H.A.M., Franz, E., Pielat, A., 2019. Phenotypic prediction: linking in vitro virulence to the genomics of 59 *Salmonella enterica* strains. *Front. Microbiol.* 10.
- Mughini-Gras, L., Pijnacker, R., Duijster, J., Heck, M., Wit, B., Veldman, K., Franz, E., 2020. Changing epidemiology of invasive non-typhoid *Salmonella* infection: a nationwide population-based registry study. *Clin. Microbiol. Infect.* 26, 941 e949–941.e914.
- Namata, H., Welby, S., Aerts, M., Faes, C., Abrahantes, J.C., Imberechts, H., Vermeersch, K., Hooyberghs, J., Méroc, E., Mintiens, K., 2009. Identification of risk factors for the prevalence and persistence of *Salmonella* in Belgian broiler chicken flocks. *Prev. Vet. Med.* 90, 211–222.
- Pate, M., Mićunović, J., Golob, M., Vestby, L.K., Očepek, M., 2019. *Salmonella infantis* in broiler flocks in Slovenia: the prevalence of multidrug resistant strains with high genetic homogeneity and low biofilm-forming ability. *Biomed Res. Int.* 2019, 4981463.
- Scalzo, S., Corkill, J.E., Shanks, D.J., Rowan, T.G., Delaval, J., Fleetwood, A., Murphy, M., Hart, C.A., 2004. Phenotypic and genotypic changes in *Salmonella enterica* subsp. *Enterica* serotype *typhimurium* during passage in intestines of broiler chickens fed on diets that included ionophore anticoccidial supplements. *J. Clin. Microbiol.* 42, 3399–3405.
- Tate, H., Folster, J.P., Hsu, C.H., Chen, J., Hoffmann, M., Li, C., Morales, C., Tyson, G.H., Mukherjee, S., Brown, A.C., Green, A., Wilson, W., Dessai, U., Abbott, J., Joseph, L., Haro, J., Ayers, S., McDermott, P.F., Zhao, S., 2017. Comparative analysis of extended-spectrum-beta-Lactamase CTX-M-65-Producing *Salmonella enterica* serovar *infantis* isolates from humans, food animals, and retail chickens in the United States. *Antimicrob. Agents Chemother.* 61.
- Vlaanderen, F., Uiterwijk, M., Cuperus, T., Keur, I., De Rosa, M., Rozendaal, H., Koene, M.G.J., Schreurs, H., Nijssse, R., Nielen, M., Friesema, I., Van Pelt, W., Franz, E., Hogerwerf, L., Opsteegh, M., Maassen, C.B., 2019. *Staat van Zoonosen 2018*. RIVM ed. (Bilthoven, RIVM).
- Zankari, E., Hasman, H., Cosentino, S., Vestergaard, M., Rasmussen, S., Lund, O., Aarestrup, F.M., Larsen, M.V., 2012. Identification of acquired antimicrobial resistance genes. *J. Antimicrob. Chemother.* 67, 2640–2644.