

1 ***BrphyB* is critical for rapid recovery to darkness in mature *Brassica rapa***  
2 **leaves**

3 Andrej A. Arsovski<sup>1</sup>, Joseph E. Zemke<sup>1</sup>, Morgan Hamm<sup>1</sup>, Lauren Houston<sup>1</sup> Andrés  
4 Romanowski<sup>2</sup>, Karen J. Halliday<sup>2</sup>, Nathalie Nesi<sup>3</sup> and Jennifer L. Nemhauser<sup>1</sup>

5 <sup>1</sup>Dept of Biology, University of Washington, Seattle, WA 98102

6 <sup>2</sup>Institute for Molecular Plant Sciences, School of Biological Sciences, University of  
7 Edinburgh, Edinburgh EH9 3BF, United Kingdom

8 <sup>3</sup>IGEPP, INRA, Agrocampus Ouest, Université de Rennes, Université Bretagne-  
9 Loire, 35650 Le Rheu, France.

10

11 **HIGHLIGHT**

12 *BrphyB* plays a central role in recovery from darkness and return to carbon fixation  
13 by regulating photosynthesis and light response genes with those targeted to the  
14 chloroplast especially affected.

15

16 **ABSTRACT**

17 Crop biomass and yield are tightly linked to how the light signaling network translates  
18 information about the environment into allocation of resources, including  
19 photosynthates. Once activated, the phytochrome (phy) class of photoreceptors  
20 signal and re-deploy carbon resources to alter growth, plant architecture, and  
21 reproductive timing. *Brassica rapa* has been used as a crop model to test for  
22 conservation of the phytochrome–carbon network. *B. rapa phyB* mutants have  
23 significantly decreased or absent CO<sub>2</sub>-stimulated growth responses in seedlings, and  
24 adult plants have reduced chlorophyll levels, photosynthetic rate, stomatal index, and  
25 seed yield. Here, we examine the transcriptomic response of adult wild-type and  
26 *BrphyB* leaves to darkening and recovery in light. Three days of darkness was  
27 sufficient to elicit a response in wild type leaves suggesting a shift from carbon  
28 fixation and nutrient acquisition to active redistribution of cellular resources. Upon a  
29 return to light, wild-type leaves appeared to transcriptionally return to a pre-darkness  
30 state restoring a focus on nutrient acquisition. Overall, *BrphyB* mutant plants have a  
31 similar response with key differences in genes involved in photosynthesis and light  
32 response which deviate from the wild type transcriptional dynamics. Genes targeted  
33 to the chloroplast are especially affected. Adult *BrphyB* mutant plants had fewer,

34 larger chloroplasts, further linking phytochromes, chloroplast development,  
35 photosynthetic deficiencies and optimal resource allocation.

36

37 **KEY WORDS:** Brassicaceae, chloroplast development, gene regulation, light  
38 response, photosynthesis, phytochrome B.

39

## 40 **INTRODUCTION**

41 Light plays at least two distinct roles in shaping plant form and productivity. First,  
42 light is essential for photosynthesis, which allows plants to convert the energy held in  
43 photons into the high potential energy found in the chemical bonds of sugars.  
44 Second, light provides information on how a plant can optimize its architecture to  
45 maximize photosynthetic potential in a given environment. How these two light  
46 systems are coordinated remains largely unknown, especially in mature leaves.

47 Limited light supply by an established canopy triggers a rapid shade-avoidance  
48 response that is characterized by increased elongation growth rate of stems and  
49 petioles, decreased leaf surface area and thickness, and delayed leaf yellowing  
50 (Casal, 2012; Franklin and Whitelam, 2005). On the other hand, partial plant shading  
51 or darkening will induce a range of responses between acclimation to leaf  
52 senescence (Weaver and Amasino, 2001; Brouwer *et al.*, 2012). These processes  
53 directly reduce the impact of shade or dark while additional responses such as  
54 acclimation of the photosynthetic apparatus rather help to fine tune the use of  
55 resources under shade/dark.

56 Plants use an array of photoreceptors to capture and transduce the light signal in  
57 diverse responses known collectively as photomorphogenesis. Photoreceptors'  
58 absorbance properties span most of the visible light spectrum, from the  
59 phytochromes that absorb in the red (R)/far-red (FR) to the cytochromes and  
60 phototropins that absorb in the blue/near-ultraviolet to the UV-receptors. Among  
61 these, the phytochromes (phys) are among the best characterized (Chen *et al.*,  
62 2004). Upon illumination, phys undergo conformational changes from an inactive (Pr)  
63 to an active (Pfr) form (Fraser *et al.*, 2016), which is subsequently translocated into  
64 the nucleus and participates in transcriptional regulation (Chen *et al.*, 2005; Castillon  
65 *et al.*, 2007). Five *PHY* genes have been described in the *A.thaliana* genome  
66 (*PHYA-PHYE*) with partial overlapping functions (reviewed in Chen *et al.*, 2004).

67 Phytochrome-dependent light signaling that initiates photomorphogenesis has been  
68 extensively studied using the seedling model (reviewed in Arsovski *et al.*, 2012). In  
69 addition, it is clear from work in *A.thaliana* that phytochromes control chloroplast  
70 gene expression, as well as nuclear-encoded factors involved in chloroplast  
71 development (Oh and Montgomery, 2014; Nevarez *et al.*, 2017). Recent studies in  
72 *A.thaliana* and *Brassica rapa* showed that adult *phyB* mutants have reduced  
73 chlorophyll levels, photosynthetic rate, and stomatal index. Work by a number of  
74 groups has connected PhyB to biomass accumulation, carbon resource  
75 management, seed yield and changes in metabolism across the plant life cycle  
76 (Yang *et al.*, 2016; Krahmer *et al.*, 2018; Arsovski *et al.*, 2018; Wies *et al.*, 2019).

77 To date, most of our knowledge about the roles of phytochromes in the dark-to-light  
78 transition primarily came from experiments focused on the de-etiolation process of  
79 seedlings (Li *et al.*, 2011). This has left a gap in our understanding about the role of  
80 phytochromes in light-activated transcription of genes in mature leaves. This is  
81 important because several light-regulatory mechanisms essential for photosynthetic  
82 efficiency and adaptation occur only in mature leaves. For example, Chory *et al.*  
83 demonstrated that the primary role of phytochrome in greening *A. thaliana* plants is  
84 in modulating the degree rather than the initiation of chloroplast development (Chory  
85 *et al.*, 1989).

86

87 In this study, we investigated the effects of *phyB* on gene expression upon dark-to-  
88 light transition in the mature leaf of *B. rapa* by comparing the transcriptomic  
89 responses between wild-type and a *phyB* mutant. *B. rapa* is closely related to  
90 *A.thaliana* (Wang *et al.*, 2011) but its leaves are significantly larger. Larger leaves  
91 cause more self-shading, and, in combination with the longer life of *B. rapa*  
92 compared to *A.thaliana*, there is more total demand for resources. As the *B. rapa*  
93 genome contains only one *PhyB* ortholog and no likely ortholog for the closely  
94 related *AtPhyD*, we took advantage of the *BrphyB3* mutant allele described  
95 previously (Arsovski *et al.*, 2018). Wild-type and *BrphyB* leaves exhibited significant  
96 overlaps in their transcriptomic response to dark and recovery; however, gene  
97 ontology analyses pointed out important misregulations in *BrphyB* mutant for genes  
98 involved in nitrogen metabolism, light harvesting and photosynthesis. Altogether  
99 these results support a role for PhyB in chloroplast development and resource

100 allocation, and have implications for increasing the resource-efficiency of Brassica  
101 crops.

102

## 103 **MATERIALS AND METHODS**

104

### 105 **Growth conditions of *B. rapa* adult plants**

106

107 The *B. rapa* wild-type R-o-18 and *BrphyB* mutants were originally from the John  
108 Innes Center's RevGenUK resource. The *BrphyB*-3 previously described in Arsovski  
109 *et al.*, 2018 was used for RNAseq experiments. *BrphyB*-1 was also previously  
110 described in Arsovski *et al.*, 2018. Seeds were planted directly into our standard soil  
111 mix of 1:1 Sunshine Mix #4 (SunGro Horticulture):vermiculite. Plants were grown in  
112 2.6 liter square pots (McConkey Grower Products; Sumner, WA, USA) and bottom-  
113 watered daily in long day conditions (16 h light, 8 h dark,  $\sim 115 \mu\text{mol}\cdot\text{m}^{-2}\cdot\text{s}^{-1}$  light  
114 intensity) in a Percival E-30B growth chamber (<https://www.percival-scientific.com/>)  
115 set to 20°C. Experiments were conducted at 3 weeks and the plants were then  
116 moved to growth room until seed harvest.

117

### 118 **Leaf sample preparation**

119

120 Three weeks after sowing, two developmentally matched leaves from each wild-type  
121 and *BrphyB*-3 plant were tagged that corresponded to the first and second true  
122 leaves. Samples were collected at ZT 5 using a standard hole punch (28 mm<sup>2</sup>  
123 circular area of leaf blade tissue) with symmetrical harvest (a second hole punch on  
124 the other side of the other side of the mid-vein of the same leaf) for chlorophyll  
125 assay, chloroplast measurements and transcriptome analysis. Tissue from 3  
126 individual plants was combined to make one biological replicate. At 3 weeks of age  
127 the "Pre" sample was collected from the first leaf while the second one was covered  
128 with tinfoil. 24 hours later the "24hr" sample was harvested from the uncovered leaf,  
129 the same leaf that provided the "Pre" sample. Then, 48 hours later the tinfoil was  
130 removed from the covered leaf and the "dark" samples were similarly collected.  
131 Finally, 24 hours later the "recovery" samples were collected from this same leaf.  
132 Samples were immediately flash frozen in liquid nitrogen (Fig.1). In total, three  
133 biological replicates were collected in similar fashion.

134

### 135 **Chlorophyll measurement**

136

137 For chlorophyll measurement, ethanol extractions were done as in (Yang *et al.*,  
138 2016). Determinations were run by measuring optical density at 645 nm and 665 nm  
139 using an Epoch Microplate Spectrophotometer ([www.biotek.com](http://www.biotek.com)). Values were  
140 obtained using the following formulas:  $\text{Chl } a = 5.21A_{665} - 2.07A_{645}$ ;  
141  $\text{Chl } b = 9.29A_{645} - 2.74A_{665}$ , for Chlorophyll A and B, respectively. Three individual  
142 biological replicates were used for this assay.

143

### 144 **RNAseq**

145

146 Leaf tissue was disrupted with Zirconia/Silica beads for 1 minute in a  
147 MiniBeadbeater-96 (BioSpec Products, Inc.) while frozen. After adding 500  $\mu\text{L}$  of  
148 Lysis/Binding Buffer to each sample and vortexing until homogeneous, samples  
149 were run on the MiniBeadbeater-96 for an additional minute. Following tissue  
150 disruption samples were centrifuged at 16,000 x g for 10 minutes at 20 °C. For each  
151 sample, a 50  $\mu\text{L}$  aliquot of the supernatant was added to 50  $\mu\text{L}$  of NEB RNA binding  
152 buffer and mRNA isolated as per the NEBNext® Poly(A) mRNA Magnetic Isolation  
153 Module manual.

154

155 RNA-seq libraries were prepared by using the Full Transcript mode YourSeq Dual  
156 (FT & 3'-DGE) RNAseq Library Kit (Amaryllis Nucleics). A Bioanalyzer 2100 (Agilent,  
157 High Sensitivity DNA Kit) was used for library quality control, to determine average  
158 library size, and together with concentration data from a Qubit 2.0 Fluorometer (Life  
159 Technologies, dsDNA High Sensitivity Assay Kit) to determine individual library  
160 molarity and pooled library molarity. Pooled libraries were sequenced on a NextSeq  
161 500 (Illumina, High Output v2 75 cycle kit) to yield single-read 80 bp reads.

162

163 FASTQ processing was performed by Amaryllis Nucleics (Oakland, CA). Sequence  
164 files were preprocessed in two steps. A Python library (`clipper.py`,  
165 <https://github.com/mfcovington/clipper>) was used to trim off the first 8 nucleotides of  
166 each read to remove potential mismatches to the reference sequence caused by

167 annealing of a random hexamer required for library synthesis. Trimmomatic v0.36  
168 (<http://www.usadellab.org/cms/?page=trimmomatic>) was used to remove adapter  
169 sequences and trim or filter reads based on quality. The parameters used for  
170 Trimmomatic were ILLUMINACLIP:TruSeq3-PE-2.fa:2:30:10 LEADING:3  
171 TRAILING:3 SLIDINGWINDOW:4:15 MINLEN:50.

172 Preprocessed reads were mapped to the *Brassica rapa* v2.5 genomic reference  
173 sequence  
174 ([http://brassicadb.org/brad/datasets/pub/Genomes/Brassica\\_rapa/V2.0/V2.5/Chr/Bra](http://brassicadb.org/brad/datasets/pub/Genomes/Brassica_rapa/V2.0/V2.5/Chr/Bra)  
175 [paV2.5\\_Chchr.fa.gz](http://brassicadb.org/brad/datasets/pub/Genomes/Brassica_rapa/V2.0/V2.5/Chr/Bra)) using bowtie2. Read counts for each gene in the gene annotation  
176 file  
177 ([http://brassicadb.org/brad/datasets/pub/Genomes/Brassica\\_rapa/V2.0/V2.5/Chr/Bra](http://brassicadb.org/brad/datasets/pub/Genomes/Brassica_rapa/V2.0/V2.5/Chr/Bra)  
178 [paV2.5\\_Chchr.gene.gff.gz](http://brassicadb.org/brad/datasets/pub/Genomes/Brassica_rapa/V2.0/V2.5/Chr/Bra)) were calculated using htseq-count (with the -s yes  
179 parameter to enforce strand-specific alignment) from the HTSeq Python library  
180 (<https://academic.oup.com/bioinformatics/article/31/2/166/2366196>;  
181 <http://htseq.readthedocs.io/en/master/index.html>).

182

183 The package edgeR (Robinson *et al.*, 2010) was used to process the expression  
184 matrix and identify differentially expressed genes between treatments and  
185 genotypes. For the main analysis, the generalized linear model functionality of this  
186 package, based on a negative binomial distribution model for gene expression was  
187 used to identify differentially expressed genes. Genes were considered significantly  
188 differentially expressed based on having a fold change greater than 2-fold up or  
189 down between conditions, and a q-value (adjusted p-value by Benjamini-Hochberg  
190 procedure - (Benjamini and Hochberg, 1995) less than 0.01.

191

192 Contrasts between the Pre and 24hr timepoints were used to identify genes that  
193 could be exhibiting differential expression caused by wound response from the tissue  
194 sampling rather than response to darkness and recovery. Genes identified in these  
195 wound control contrasts were tagged but not excluded from the rest of the analysis.  
196 Dispersion was estimated independently for the wound control contrasts based on  
197 the Pre and 24hr timepoints only. This was done because using the dispersion  
198 estimates from the main analysis, including the Dark and Recovery timepoints  
199 resulted in biased p-value distributions due to the significant change in expression of

200 most genes in the dark timepoint as compared to the other timepoints (68% of genes  
201 between wild type-Dark and wild type-Pre).

202

203 Complete data can be accessed from the Gene Expression Omnibus (GEO) under  
204 the entry GSE135955.

205

## 206 **Venn diagrams and gene ontology (GO) analysis to prioritize the DEG**

207

208 To find differences between the wild-type and *phyB* response to darkness and re-  
209 illumination we looked at sets of genes that were significantly changed in these  
210 contrasts (darkened vs. pre or recovery vs. darkened) in one genotype but not the  
211 other. For each up or down Venn-diagram used in the GO analysis, all genes which  
212 either the wild-type or *BrphyB* mutant were greater (or less for down regulated Venn)  
213 than the log fold change cutoff of 2 were considered. These genes were then put into  
214 4 categories: wild-type significant only, *BrphyB* significant only, both wild-type and  
215 *BrphyB* significant, and neither genotype significant.

216

217 To obtain a comprehensive list of all *B. rapa* presumptive orthologs in *A. thaliana* we  
218 used each *B. rapa* protein as a query to perform local homology searches. Briefly, for  
219 each protein in the *B. rapa* v2.5 proteome, the best *A. thaliana* hit was retrieved by  
220 sequence similarity search using a local installation of the BlastP algorithm (Protein-  
221 Protein BLAST 2.7.1+) of the NCBI tool BLAST  
222 (<http://blast.ncbi.nlm.nih.gov/Blast.cgi>) against the *A. thaliana* ARAPORT11  
223 proteome, with default parameters and the outfmt parameter set to 7 (to obtain a  
224 tabular output with comment lines). This resulted in a collection of the best *A.*  
225 *thaliana* hits for each *B. rapa* protein. This output was further processed to limit each  
226 ortholog to the best hit, using a custom BASH script (available upon request). GO  
227 term enrichment analysis was performed against the well annotated *A.thaliana*  
228 genome using the PANTHER classification System (v.14.1 available at  
229 <http://pantherdb.org/>; (Mi *et al.*, 2019).

230

231

## 232 **Law 2018 comparison**

233

234 For the data in Table S3, we created a list by matching the *A. thaliana* gene IDs from  
235 the (Law *et al.*, 2018) analysis up to the *B. rapa* genes they were found to be the  
236 nearest homolog of (see GO analysis section above). We considered the genes that  
237 were differentially expressed after 3 days (D3) of darkness applied on an  
238 individualized leaf (IDL) or a whole plant (DP). *A. thaliana* genes that did not match  
239 to *B. rapa* genes in our set were dropped from the comparison, *A. thaliana* genes with  
240 multiple brassica homologs were listed multiple times in this list. The elements of this  
241 list were then broken down into a Venn diagram based on whether they were  
242 considered significantly differentially expressed up or down in our darkened vs. pre  
243 contrast and Law's IDL\_D3 and DP\_D3 contrasts. Each category list was then  
244 reduced to only contain unique *A. thaliana* IDs. These counts are displayed in Table  
245 S3.

246

#### 247 **RNA extraction and quantitative real-time PCR (qRT-PCR) analysis**

248 Expression analysis was performed using 4 biological replicate samples collected  
249 identically as described for RNA sequencing. Each sample was immediately frozen  
250 in liquid nitrogen and stored at  $-80^{\circ}\text{C}$  until processing. Frozen tissue was ground in  
251 liquid nitrogen and total RNA was extracted using the GE Illustra RNA kit (GE Life  
252 Sciences), and 2  $\mu\text{g}$  of eluted RNA was used for cDNA synthesis employing iScript  
253 (Bio-Rad). Samples were analyzed using SYBR Green Supermix (Bio-Rad) reactions  
254 run in a C100 Thermal Cycler (Bio-Rad) fitted with a CFX96 Real-Time Detection  
255 System (Bio-Rad). Relative expression levels were calculated using the formula  
256  $(E_{\text{target}})^{-C_{\text{Ptarget}}}/(E_{\text{ref}})^{-C_{\text{Pref}}}$  (Pfaffl, 2001) and normalized to the *B. rapa* *PP2A*  
257 (Brara.F00691) reference gene. qPCR primer sequences are as follows: *BrPP2A*  
258 (forward 5'-TCGGTGGTAACGCCCCCGAT-3'; reverse 5'-  
259 CGACTCTCGTGGTTCCCTCGC-3'); *BrMGT6* (Brara.E02300) (forward 5'-  
260 CAGCATCCGCCACCGCAAGA-3'; reverse 5'-GCCTTCGCAACAACCGCAGC-3');  
261 *BrRLK4* (Brara.I00004) (forward 5'- TCCGCCGTCGCGATCTCTCT-3'; reverse 5'-  
262 CCCGCTCCAAACGCTTGTCCA-3'); *BrPIFI* (forward 5'-  
263 GCCACCACTCTTGACCCCC-3'; reverse 5'- CCGCGTTGGAGGAAGACCG-3').

264

#### 265 **Code**

266 The R code used to generate all the analysis results is provided in supplement X and  
267 can also be found on [github.com/nemhauser-lab/brassica\\_rna\\_seq](https://github.com/nemhauser-lab/brassica_rna_seq).



268

## 269 **Methods, Motif Enrichment**

270 A set of binding motifs for 619 *A.thaliana* transcription factors was downloaded from  
271 plantTFDB (<http://planttfdb.cbi.pku.edu.cn/>), (Jin *et al.*, 2014, 2015, 2017). The RSAT  
272 matrix clustering tool ([http://rsat.eead.csic.es/plants/matrix-clustering\\_form.cgi](http://rsat.eead.csic.es/plants/matrix-clustering_form.cgi))  
273 (Castro-Mondragon *et al.*, 2017) was used with default parameters to group the  
274 motifs into 56 clusters based on similarity of aligned position weight matrices. Genes  
275 that were significantly differentially expressed from conditions pre to dark, and from  
276 dark to recovery were divided into groups depending on 3 factors: genotype,  
277 direction of regulation (up or down), and whether they are annotated with the GO  
278 term “chloroplast” (GO:0009507). The package “motifmatchr” (Schep, 2019) was  
279 used to count the number of genes in each set with promoter sequences with  
280 matches to each of the 619 motifs. Promoters sequences were defined as the 1000  
281 base pairs immediately upstream of the gene start position as defined by the  
282 genome annotation file. The fisher exact test was then used to determine if there  
283 was significant enrichment for each motif between genes only significantly regulated  
284 in WT and genes only significantly regulated in *BrphyB* in these sets. The raw p-  
285 values from these tests were adjusted using the by Benjamini-Hochberg procedure  
286 (Benjamini and Hochberg, 1995). Adjusted p-values of <0.01 were considered  
287 significant.

288

289 All motifs found significant from these tests closely matched to three canonical motifs  
290 described in the literature: the G-box motif CACGTG, the Evening Element  
291 AGATATTTT, and the Telo-box motif AAACCCTAA. The proportions of promoters  
292 with one or more exact matches to three canonical motifs were found for the gene  
293 sets described above.

294

## 295 **Chloroplast Measurement**

296 Tissue was immediately cleared after collection and fixed using ClearSee solution as  
297 described in (Kurihara *et al.*, 2015). Images were taken using a Leica TCS SP5 II  
298 laser scanning confocal microscope (<https://www.leica-microsystems.com>).  
299 Chloroplast number, area, and density were determined using ImageJ software.

300

301

## 302 RESULTS

303

### 304 **Mature *BrPhyB* mutant leaves have significant transcript reductions of** 305 **chloroplast targeted genes.**

306

307 Loss of phyB leads to significant reductions in both chlorophyll levels and rates of  
308 photosynthesis in three-week-old *B. rapa* plants (Arsovski *et al.*, 2018). To further  
309 understand the link between phyB, chloroplast development, and photosynthesis we  
310 examined the transcriptomic response of mature leaves that were subjected to three  
311 days of darkness before being reintroduced into the light. As part of this experiment  
312 we first compared the transcriptome of three-week-old *B.rapa* wild type and *BrphyB*  
313 leaves. Genes were considered differentially expressed if the fold change between  
314 timepoints or genotypes was greater than 2, and the significance (adjusted p-value)  
315 was less than 0.01. 114 genes were significantly upregulated in *BrphyB* leaves  
316 compared to wild type. Unsurprisingly, these include *B.rapa* orthologs to *A.thaliana*  
317 *LONG HYPOCOTYL IN FAR-RED(HFR1)*, *PHYTOCHROME INTERACTING*  
318 *FACTOR 3-LIKE 1 (PIL1)*, *PHYTOCHROME-INTERACTING FACTOR 6 (PIF6)*, and  
319 *INDOLE-3-ACETIC ACID INDUCIBLE 29(IAA29)*. 79 genes were significantly  
320 downregulated in *BrphyB* leaves compared to the wild type. Gene Ontology (GO)  
321 analysis of cellular location annotations showed a strong enrichment for the  
322 chloroplast envelope, stroma, and photosystem II. These include *B.rapa* orthologs  
323 for *A.thaliana* *PHOTOSYSTEM II SUBUNIT P*, *PHOTOSYSTEM II BY*, *LIGHT-*  
324 *HARVESTING CHLOROPHYLL B-BINDING PROTEIN 3*, and *CHLOROPHYLL A/B*  
325 *BINDING PROTEIN 1* (Table S1).

326

### 327 **Darkening of individual leaves for three days initiates resource reallocation**

328

329 Samples were taken from the first or second true leaf of three-week-old wild-type  
330 and *BrphyB-3* plants (hereafter termed “pre”). As a wounding control, a second  
331 sample was taken from the “pre” leaves 24 hours later (hereafter termed “24hr”).  
332 Leaves that were developmentally-matched with those selected for the “pre”  
333 treatment were covered with foil. After three days, the foil was removed and the  
334 “darkened” sample was collected immediately. The “recovery” sample was collected  
335 from the same leaf 24 hours after this timepoint to capture the earliest stages of

336 recovery (Fig. 1A). Matching samples were collected from each leaf to assay  
337 chlorophyll levels. At three weeks old, *BrphyB* mutants are visibly paler compared to  
338 same aged wild-type plants, and have significantly reduced chlorophyll levels (Fig.  
339 1B and Arsovski *et al.*, 2018). Three days of dark resulted in a 23% reduction of  
340 chlorophyll levels in wild type leaves while levels remained low in the mutant. This is  
341 consistent with similar experiments performed on individually darkened *A.thaliana*  
342 leaves where total chlorophyll levels and protein decline was observed after two  
343 days of darkness (Weaver and Amasino, 2001). The 24 hours of light exposure for  
344 the recovery samples was not sufficient to restore chlorophyll levels in either wild-  
345 type or *BrphyB-3* leaves (Fig. 1B).

346

347 Extended darkness of leaves acts as a signal to initiate the organized breakdown  
348 and remobilization of valuable resources to growing vegetative and reproductive  
349 tissues (Himmelblau and Amasino, 2001; Buchanan-Wollaston *et al.*, 2003; Lim *et al.*,  
350 2007). We performed RNAseq analysis on the pre, 24hr, dark and recovery samples  
351 to assess the specific response to darkness and return to light of mature leaves in *B.*  
352 *rapa*. We began our analysis with the response to darkness, as previous studies in  
353 *A. thaliana* had already shown that dark stress is accompanied by dramatic  
354 transcriptional changes, as well as depletion of chlorophyll and large-scale  
355 degradation of proteins (Guo *et al.*, 2004; Keech *et al.*, 2007; Law *et al.*, 2018). The  
356 expression of 6852 *B.rapa* genes was significantly altered in leaves after three days  
357 in darkness. Gene Ontology (GO) analysis of predicted *A. thaliana* orthologs showed  
358 a pattern consistent with the overall expectations of metabolic reprogramming seen  
359 in other species. The 3110 genes up-regulated in response to dark were mainly  
360 involved with autophagy, catabolism, leaf senescence and vesicle fusion (Fig. 2A;  
361 Table S2). Conversely, down-regulated genes were mainly involved in  
362 photosynthesis, biosynthetic processes and plastid translation. Together, these data  
363 suggest a shift from carbon fixation and nutrient acquisition to active redistribution of  
364 cellular resources (Fig. 2B, Table S2).

365

366 A recent experiment in *A.thaliana* found that the effect of darkening individual leaves  
367 was substantially similar to the effect of darkening whole plants, albeit with distinct  
368 timing for peak differences in gene expression between the two treatments Law *et*  
369 *al.*, 2018). When we compared our transcriptomic response to this dataset, we found

370 a substantial overlap. The highest similarity between the *B.rapa* dark response was  
371 to *A.thaliana* individually darkened leaves for 3 days (IDL\_3D). Of the genes  
372 significantly up or down regulated in *A. thaliana* individually darkened leaves after 3  
373 days, 46.7% had *B.rapa* homologs also significantly up or down regulated (in the  
374 same direction) (Table S3). Shared *A. thaliana* genes upregulated in response to  
375 dark were significantly enriched for GO terms such as autophagy, catabolic process,  
376 and leaf senescence (Table S4). The darkening response in *B.rapa* was more  
377 similar to that of individually darkened leaves than whole darkened plants in  
378 *A.thaliana*. Of the genes found to be up/down regulated in individual leaves (IDL\_D3)  
379 but not whole darkened plants (DP\_D3), 35.3% had *B.rapa* homologs with significant  
380 change in the same direction, compared to 25.1% in whole darkened plant unique  
381 genes (Table S3,4). In *A. thaliana*, 167 senescence-associated genes were shown to  
382 change expression in response to darkening (Parlitz *et al.*, 2011). The *B.rapa*  
383 orthologs of 103 of these genes do not show significant changes in response to dark.  
384 Of the remaining 57 senescence-associated genes, 42 show a reversible pattern of  
385 upregulation in the dark and downregulation upon re-illumination. 15 are 'non-  
386 reversible', upregulated in the dark without significant changes upon a return to light  
387 (Table S5).

388

389 Many of the genes that were regulated by returning the leaves to light were similar to  
390 those already identified as light-responsive from experiments in seedlings. In *A.*  
391 *thaliana*, expression of up to one-fourth of the whole genome is altered in seedlings  
392 grown for 4 days in red light compared to those grown in the dark (Shi *et al.*, 2018).  
393 These changes are largely mediated by a small group of transcription factor families  
394 which include the PHYTOCHROME INTERACTING FACTORS (PIFs). Nearly 60%  
395 of PIF-dependent, red light induced genes in *A. thaliana* seedlings have Gene  
396 Ontology (GO) annotations indicating functions related to photosynthesis and  
397 chloroplast (Leivar *et al.*, 2012). In *B.rapa* wild type leaves 3756 genes were  
398 upregulated in the recovery condition when compared to the dark timepoint. The  
399 most significantly enriched GO terms were response to light stimulus,  
400 photosynthesis, translation and metabolism, suggesting a return to a pre-darkness  
401 transcriptional state (Figure 3A, Table S2). The 3299 genes downregulated in  
402 recovery compared to dark were mainly involved in catabolism, vesicle fusion and

403 transport, and protein degradation further supporting a shift from resource  
404 remobilization towards nutrient acquisition (Figure 3B, Table S2).

405

#### 406 ***BrphyB* is critical for full recovery response**

407

408 RNAseq analysis of *BrphyB* individually darkened leaves revealed an essentially  
409 similar response to what was observed in wild-type plants. 7,994 genes were  
410 differentially expressed in wild-type and *BrphyB-3* leaves after 3 days of dark  
411 treatment compared to the pre samples. The vast majority (81.8%) of this response  
412 was shared between the genotypes (Fig. 2A, B). However, analysis of GO terms  
413 significantly enriched in the uniquely wild-type or *BrphyB* differentially regulated gene  
414 sets illustrated phyB-dependent responses to darkness. Unique wild-type enriched  
415 terms were largely related to cellular responses to organic and inorganic  
416 compounds, drugs and stress (Table S6). A closer look at these phyB-controlled  
417 groups revealed orthologs to *A. thaliana* genes REVEILLE2, REVEILLE8, and  
418 CIRCADIAN CLOCK ASSOCIATED1—all key transcriptional regulators of circadian  
419 rhythm, auxin and stress response in *A. thaliana* and known to act downstream of  
420 PhyB (Fig. 2A,C) (Alabadí *et al.*, 2001; Zhang *et al.*, 2007; Rawat *et al.*, 2011;  
421 Farinas and Mas, 2011; Jiang *et al.*, 2016). *BrphyB* unique up-regulated genes were  
422 enriched in genes associated with response to light, including *B.rapa* orthologs of  
423 PHYTOCHROME INTERACTING FACTOR4, PHYTOCHROME INTERACTING  
424 FACTOR5, PHYTOCHROME KINASE SUBSTRATE1 and CRYPTOCHROME 1  
425 (Table S6, Fig. 2C).

426

427 PhyB-repressed genes are enriched for categories such as response to light  
428 stimulus and cellular biosynthetic process, while those activated by phyB are  
429 enriched for categories related to protein synthesis such as cysteine metabolic  
430 process, translational elongation, and peptide biosynthesis (Fig.2B). These include  
431 *B. rapa* orthologs to three *A. thaliana* glutamate-ammonia ligases (GLUTAMINE  
432 SYNTHASE 1;2, 1;3 and 1;4) with roles in nitrogen remobilization and seed yield  
433 (Guan *et al.*, 2015), stress response and pollen viability (Ji *et al.*, 2019). In *B. rapa*,  
434 *BrphyB* mutants have up to a 90% decrease in seed yield (Arsovski *et al.*, 2018);  
435 however, we did not observe a difference in area or weight of seeds compared to  
436 wild type (data not shown). This would suggest that plants are re-calibrating the

437 amount of resources available, and maintaining quality by partitioning them into a  
438 smaller number of seeds.

439

440 While there is substantial overlap between the response of wild type and *BrphyB*  
441 mutants to re-illumination (80.2% of the 7,765 genes are common to both  
442 genotypes), there are also several key differences. To validate the RNAseq results,  
443 we selected three genes whose expression in wild type was significantly  
444 downregulated during darkening compared to pre followed by a significant  
445 upregulation in recovery compared to darkening. qPCR of these genes was done in  
446 wild-type, *BrphyB-3*, and an additional mutant allele *BrphyB-1*. Brara.E02555 is an  
447 ortholog of the *A.thaliana* At3g15840 gene. In *A.thaliana* POST-ILLUMINATION  
448 CHLOROPHYLL FLUORESCENCE INCREASE (PIFI) is a nuclear-encoded  
449 chloroplast protein essential for NDH-mediated non-photochemical reduction of the  
450 plastoquinone pool in chlororespiratory electron transport (Wang and Portis, 2007).  
451 The *A.thaliana* orthologs RECEPTOR-LIKE PROTEIN KINASE 4 (RLK4) and  
452 MAGNESIUM TRANSPORTER 6 (MGT6) are a Ser/Thr receptor-like protein kinase  
453 expressed in the root and a magnesium transporter required for growth in  
454 Magnesium limited conditions, respectively. The qPCR expression closely resembled  
455 the RNAseq results for wild-type and *BrphyB-3* and *BrphyB-1* expression matched  
456 that of *BrphyB-3* for all three genes. *BrPIFI* expression decreases dramatically in  
457 response to darkening in both wild-type and *BrphyB-1* mutant leaves. However,  
458 while *BrPIFI* expression increases in response to the leaf's return to light in wild-type,  
459 it remains low in the mutant. Similarly, *BrRLK4* and *BrMGT6* expression increases in  
460 recovery in wild-type leaves. However, in *BrphyB-3* leaves expression of both  
461 decreases 24 hours after the cover is removed from the leaf (Fig. S1) (Coello *et al.*,  
462 1999; Wang and Portis, 2007).

463

464 *BrphyB* is required for rapid return to the full photosynthetically-active transcriptional  
465 program. There are 404 genes up-regulated in recovery of only wild-type leaves.  
466 These genes are mainly involved in light harvesting in Photosystem I,  
467 photosynthesis, cellular carbohydrate catabolism and generation of precursor  
468 metabolites and energy and are not upregulated in *BrphyB* leaves 24 hours after  
469 return to light (Fig. 3A, Table S6). We previously showed that the expression *B.rapa*  
470 GOLDEN2-LIKE 1 (*BrGLK1*) increases 70% in response to high CO<sup>2</sup> in wild type

471 seedlings but decreases in *BrphyB* mutants (Arsovski *et al.*, 2018). In *A.thaliana*  
472 GLK1 is one of a pair of partially redundant transcription factors that affect the  
473 expression of nuclear photosynthetic genes involved in chloroplast development  
474 (Waters *et al.*, 2008, 2009; Kobayashi *et al.*, 2013). Here the *B.rapa* ortholog of  
475 *A.thaliana* GOLDEN2-LIKE 2 is significantly upregulated upon return to light in wild  
476 type but not *BrphyB* mutant leaves (Table S7). Closer examination of chloroplast  
477 localized genes whose expression significantly changes in response to darkening  
478 and recovery revealed a stark contrast in responsiveness between the two  
479 genotypes. In response to darkening, 1861 chloroplast related genes were  
480 significantly downregulated in either genotype. Of these, 94.7 % (1763 genes) were  
481 unique to wild type leaves and were not significantly downregulated in the mutant.  
482 Upon a return to light 1904 chloroplast localized genes were significantly upregulated  
483 with 85.8% common to both genotypes. However, 131 localized genes were  
484 upregulated only in wild type leaves while 140 were unique to the *BrphyB* mutant.  
485 GO cellular component analysis identified 125 genes with predicted chloroplast-  
486 localization that are up-regulated in wild-type but not *BrphyB* leaves during recovery.  
487 *A.thaliana* Photosystem II genes such as LIGHT-HARVESTING CHLOROPHYLL B-  
488 BINDING PROTEIN 3 (BraA10000990), LIGHT HARVESTING COMPLEX  
489 PHOTOSYSTEM II SUBUNIT 6 (BraA09006187), LIGHT-HARVESTING  
490 CHLOROPHYLL-PROTEIN COMPLEX II SUBUNIT B1( BraA05001183) and three  
491 orthologs to *A.thaliana* CHLOROPHYLL A/B BINDING PROTEIN 1 (BraA04002510,  
492 BraA07001020, BraA08002753) are upregulated on return to light only in wild type  
493 leaves (Figure 3C, Table S7) .

494

495

496 Downregulated genes unique to wild type are enriched in annotations associated  
497 with nuclear transport, ribosome biogenesis, and rRNA processing, while terms  
498 including phototropism and regulation of primary metabolism, cellular biosynthesis  
499 and nitrogen compound metabolism are enriched in the *BrphyB* unique  
500 downregulated genes (Fig. 3B). This overall pattern suggests that *BrphyB* may be  
501 required for effective monitoring and switching between carbon- and nitrogen-  
502 demanding processes, and that this role may be essential for maximal reallocation of  
503 resources to developing seeds.

504

505

506 ***BrphyB* leaves have regulatory motif differences in chloroplast related genes**  
507 **and fewer, larger chloroplasts**

508

509 In the pre condition, the 79 genes significantly downregulated in *BrphyB* leaves  
510 compared to the wild-type were enriched for localization to the chloroplast envelope,  
511 stroma, and photosystem II (Table S1). The recovery condition created a sensitized  
512 environment to detect the more immediate impacts of *BrphyB* on establishing or  
513 maintaining the photosynthetic machinery. To investigate whether there were  
514 regulatory differences in recovery between chloroplast and non-chloroplast genes in  
515 wild-type and *BrphyB* leaves we examined the promoters (1Kb upstream for TSS) of  
516 up and downregulated genes in recovery compared to dark. Genes with the GO  
517 term 0009507: chloroplast were designated as 'chloroplast' and those without it 'non-  
518 chloroplast'.

519

520 The frequency of three major motifs appeared to change in response to dark and in  
521 recovery and between genotypes (Fig.S2A). The G-box element (CACGTC) is a  
522 focal point of light-regulated gene expression. In vitro gel-shift, random DNA-binding  
523 selection, and chromatin immunoprecipitation (ChIP) assays in *A.thaliana* show that  
524 four PIFs (PIF1, PIF3, PIF4, and PIF5) bound to either a G-box (CACGTG) and/or an  
525 E box (CANNTG) (Martínez-García *et al.*, 2000; Huq and Quail, 2002; Huq *et al.*,  
526 2004; Hornitschek *et al.*, 2012). PIFs can also interact with other transcription factors  
527 at the G-box, and these interactions modulate the PIF DNA-binding activity. PIF3  
528 and PIF4 interact with BRASSINAZOLE-RESISTANT 1 (BZR1) and bind to the same  
529 G-box DNA sequence element to regulate genes involved in the light and  
530 brassinosteroid pathways (Oh *et al.*, 2012; Zhang *et al.*, 2013). PIF1 and PIF3 also  
531 interact with the light-regulated activator ELONGATED HYPOCOTYL (HY5) at the  
532 G-box where it can both promote PIF1/3 binding and compete for binding sites  
533 (Chen *et al.*, 2013; Toledo-Ortiz *et al.*, 2014). When *B.rapa* leaves were returned to  
534 light, 37% of chloroplast genes significantly upregulated in wild type but not *BrphyB*  
535 leaves have a G-box in their promoter region compared to only 15% of non-  
536 chloroplast genes. This is not the case with chloroplast genes upregulated only in  
537 *BrphyB*. For these genes, there was essentially no difference in the number of genes



538 a G-box whether or not they were annotated as chloroplast-associated (chloroplast  
539 genes: 13%, non-chloroplast genes: 16%) (Fig.S2B).

540

541 Differences were also present in the frequencies of Evening Element (AAAATATCT)  
542 and Telobox motif (AAACCCTAA) in chloroplast-annotated genes between wild-type  
543 and *BrphyB* leaves in recovery as well. The Evening Element (EE) motif is central to  
544 circadian clock function and environmental and endogenous signal coordination in  
545 *A.thaliana*. Key regulators of the circadian clock CIRCADIAN CLOCK  
546 ASSOCIATED1 (CCA1), LATE ELONGATED HYPOCOTYL (LHY) and REVEILLE 8  
547 (RVE8) bind and regulate genes with EEs in their promoters (Harmer and Kay, 2005;  
548 Hsu *et al.*, 2013). Among chloroplast-annotated genes, the EE motif was present in  
549 promoter regions of 37% of genes which were significantly downregulated in *BrphyB*  
550 but not wild type leaves, while only 7% of those downregulated in wild type but not  
551 *BrphyB* had the same motif (Fig.S2C).

552

553 Short interstitial telomere motifs (telo boxes) are short sequences identical to plant  
554 telomere repeat units. In *A.thaliana* and *O.satīva* genomes telo boxes are associated  
555 with genes involved in the biogenesis of the translational apparatus (Gaspin *et al.*,  
556 2010). Telo box motifs were enriched in the promoters of genes significantly  
557 downregulated in wild type but not *BrphyB*, and genes significantly upregulated in  
558 *BrphyB* but not wild type in recovery, 15% compared to 20%, respectively. Whereas,  
559 of genes that were upregulated in wild type only, and genes downregulated in  
560 *BrphyB* only in recovery, 5% and 7% respectively had Telo-box motifs in their  
561 promoters. The differences between genes with and without the chloroplast GO  
562 annotation was less noticeable for this motif than the other two. Together these  
563 results point to a significant difference in the cis-regulatory landscape of *BrphyB*  
564 leaves (Fig. S2D).

565

566 The chloroplast carries out many functions beyond photosynthetic carbon fixation  
567 that are essential for metabolic homeostasis, including fatty acid synthesis and  
568 fixation of nitrogen and sulfur (Lopez-Juez and Pyke, 2005). Mutants with reduced  
569 phy function have significantly lower chlorophyll levels in *A.thaliana* and *B.rapa*  
570 (Ghassemian *et al.*, 2006; Strasser *et al.*, 2010; Hu *et al.*, 2013; Arsovski *et al.*,  
571 2018). It has been suggested that phyA is primarily responsible for chloroplast

572 maturation during de etiolation in *A.thaliana*, although there are some reports that  
573 phyB might also be involved (McCormac and Terry, 2002; Xu *et al.*, 2019).

574

575 Our results, in combination with our earlier findings that *BrphyB* mutants had  
576 reduced chlorophyll levels and photosynthetic rates, led us to hypothesize that  
577 BrphyB might be required for normal chloroplast development. We found that  
578 chloroplast density was significantly decreased in the mature leaves of the *BrphyB*  
579 mutants. Wild-type leaves had an average of 466 chloroplasts per 0.5mm<sup>2</sup> compared  
580 to 326 and 253 in *BrphyB-1* and *BrphyB-3*, respectively (Fig. 4A, B ANOVA and  
581 Tukey HSD multiple comparison test). Chloroplast area however was significantly  
582 larger in *BrphyB-3* and *BrphyB-1* compared to wild type, 33.6 and 41.4 to 31.3 um<sup>2</sup>  
583 respectively (Fig. 4C ANOVA and Tukey HSD multiple comparison test). In  
584 *A.thaliana*, an investigation into photosynthetic, biochemical, and anatomical traits of  
585 accumulation and replication of chloroplasts (*arc*) mutants found that fewer, enlarged  
586 chloroplasts were less efficient at photosynthesis than more, smaller chloroplasts.  
587 Photosynthetic rate and photosynthetic nitrogen use efficiency were significantly  
588 lower in the mutants than their wild-types likely due to decreases in mesophyll  
589 conductance and chloroplast CO<sub>2</sub> concentration (Xiong *et al.*, 2017). These  
590 functional differences could explain the reduced ability of *BrphyB* leaves to rapidly  
591 switch metabolic functions when exposed to darkness and again with the return to  
592 light.

593

## 594 **DISCUSSION**

595

596 Human-driven climate change, and the associated changes in temperature,  
597 atmospheric CO<sub>2</sub>, and precipitation, are an urgent challenge for plant life on Earth.  
598 Crop yield and global food security will depend on how individual crop species  
599 respond to new and potentially more variable conditions. *B.rapa* is a laboratory crop  
600 model that has been successfully used to study the plant response to environmental  
601 change. phyB is emerging as a key regulator of carbon response and supply,  
602 metabolism and biomass production (Yang *et al.*, 2016; Arsovski *et al.*, 2018). In  
603 addition to a diminished high CO<sub>2</sub> response we previously showed that *B.rapa phyB*  
604 mutants have reduced chlorophyll levels and photosynthetic rate (Arsovski *et al.*,  
605 2018). In this work, we elicited a dark response in individual leaves and examined

606 the transcriptomic response of wild-type and *phyB* leaves as they are darkened and  
607 were subsequently returned to light.

608

609 Wild-type *B.rapa* leaves darkened for three days have a significant upregulation of  
610 genes involved in autophagy, catabolism and leaf senescence while large groups of  
611 genes functioning in photosynthesis, metabolism and translation (Fig. 2A). This  
612 senescence response and redistribution of cell resources is typical of the dark  
613 response observed in various plant systems (Guo *et al.*, 2004; Brouwer *et al.*, 2012;  
614 Song *et al.*, 2014; Law *et al.*, 2018; Sobieszczuk-Nowicka *et al.*, 2018). Upon return  
615 to light upregulated and downregulated functional groups are essentially reversed.  
616 Genes acting mainly in response to light stimulus, photosynthesis, translation and  
617 metabolism were upregulated in leaves, while those with roles in catabolism, vesicle  
618 fusion and transport, and protein degradation were downregulated 24 hours following  
619 return to light (Fig. 3B). In many plants, leaf senescence is reversible in a limited  
620 time span after senescence initiation, leading to ‘regreening’ of the leaves. For  
621 example, a return to light following a 2 day dark period initiates a reconstitution of  
622 photosynthetic capability in *A. thaliana* (Parlitz *et al.*, 2011).

623

624 While there is a significant overlap in the transcriptomic response to dark and  
625 recovery between wild-type and *BrphyB* mutant leaves, there are important  
626 categories of genes that may explain some of phenotypes associated with the  
627 *BrphyB* mutant (Fig. 2,3). After three days of dark the orthologs of *A.thaliana*  
628 PHYTOCHROME INTERACTING FACTORS 4 and 5 are only upregulated in the  
629 *BrphyB* mutant suggesting a misregulation of the light response, as well as likely  
630 impacts on hormone homeostasis. In *A. thaliana* PIFs/EIN3/HY5-regulated genes in  
631 the dark were estimated to account for half of the light-directed transcriptome  
632 changes (Shi *et al.*, 2018). In recovery, *BrphyB* mutant leaves lack the increased  
633 transcription of genes involved in light harvesting and photosynthesis, including 125  
634 chloroplast-targeted genes, and the mutants uniquely down-regulate genes involved  
635 in phototropism suggesting possible “crossed-wires” from mismatches between  
636 different photoreceptor responses (Fig. 3A). Together these finding may connect the  
637 observed reduction of chlorophyll and photosynthetic rate (Arsovski *et al.*, 2018) and  
638 fewer larger chloroplasts in *BrphyB* leaves compared to wild type.

639

640 Nitrogen is a critical resource that governs plant growth. The availability of nitrogen  
641 to the roots plays a particularly significant role in constraining plant growth and crop  
642 yield worldwide (Epstein and Bloom, 2005; Hirel *et al.*, 2011; Alvarez *et al.*, 2012).  
643 Nitrogen deficiency is one of the endogenous and environmental factors that  
644 regulates the onset of leaf senescence (Gregory, 1937; Mei and Thimann, 1984;  
645 Masclaux-Daubresse *et al.*, 2007; Koeslin-Findeklee *et al.*, 2015). In *B.rapa*, nitrogen  
646 availability limits growth increase in high CO<sub>2</sub> (Arsovski *et al.*, 2018). In dark-induced  
647 leaf senescence nitrogen from senescing leaves is mobilized and transported to still  
648 growing vegetative tissues. Wild type *B.rapa* plants darkened for 3 days show a  
649 significant upregulation of genes involved in senescence, catabolism, and vesicle  
650 transport while downregulation of genes involved in protein synthesis indicating and  
651 active export of nitrogen resources (Supplementary File 1, 2). While *BrphyB* mutants  
652 largely share this response, there is evidence that this resource allocation is altered.  
653 After 3 days of dark phyB unique downregulated genes are significantly enriched for  
654 GO terms related to protein synthesis suggesting either a delayed or prolonged  
655 response compared to wild type. In recovery, *BrphyB* uniquely upregulated genes  
656 are significantly enriched for translation and peptide biosynthesis, while wild type  
657 unique genes are photosynthesis-related. *B.napus*, a close relative of *B.rapa* has  
658 poor nitrogen use efficiency (Masclaux-Daubresse *et al.*, 2007; Xu *et al.*, 2012). Only  
659 50–60% of the applied nitrogen is recovered in the plants and at the time of harvest  
660 a relatively low 80% of the total plant nitrogen is localized in the seeds (Schjoerring  
661 *et al.*, 1995; Jensen *et al.*, 1997; Malagoli *et al.*, 2005; Rathke *et al.*, 2006). Here  
662 *BrphyB* mutants had a misregulation of genes orthologous to *A.thaliana*  
663 GLUTAMINE SYNTHASE 1;2, 1;3 and 1;4 that play a role in seed yield and size  
664 (Fig. 3C). While seeds at harvest were not significantly different from wild type in size  
665 and weight, seed yield is dramatically reduced in the mutant (Arsovski *et al.* 2018). A  
666 more detailed understanding of the phyB-regulated network holds the promise of  
667 improved plant growth models and identification of new targets for engineering more  
668 resource-efficient crops.

669

## 670 **Supplemental Figures**

671

672 Figure S1: qPCR validation.

673

674 Figure S2: Transcription factor motifs enriched in promoter regions.

675

676 Table S1: Gene expression comparison between *BrphyB* and wild type leaves in the  
677 Pre condition.

678

679 Table S2: Gene Ontology comparison of differentially expressed genes between Pre  
680 (P), Dark (D), and Recovery (R) in wild type and *BrphyB* leaves.

681

682 Table S3: A comparison of *A.thaliana* gene differentially expressed in whole  
683 darkened plants or individually darkened leaves in response to 3 days of darkness  
684 from Law et al.,2018 and *A.thaliana* orthologs of *B.rapa* genes differentially  
685 expressed in Dark vs. Pre.

686

687 Table S4: Gene Ontology analysis of *A.thaliana* genes differentially expressed in  
688 individually darkened leaves (IDL), or whole darkened plants (DP) from Law et al.  
689 2018 and *A.thaliana* orthologs of *B.rapa* genes differentially expressed in Dark vs.  
690 Pre. Sheets show *A.thaliana* IDs, gene model names, MapMan bins and descriptions  
691 of common up or down regulated genes followed by GO annotations of those genes.

692

693 Table S5: A comparison of genes differentially expressed in dark and re-illumination  
694 from Parlitz et al. 2018 and *A.thaliana* orthologs of *B.rapa* genes in Dark vs. Pre and  
695 Recovery vs. Dark.

696

697 Table S6: Gene Ontology enrichment for DEGs common and unique to wild type and  
698 *BrphyB* in pre (P), dark (D), and recovery (R) conditions.

699

700 Table S7: Differentially expressed genes in pre, dark and recovery. Significance  
701 column denotes whether the significance is common to both wild type and *BrphyB* or  
702 unique to either.

703

## 704 **Acknowledgements**

705 We thank Prof. Mark Stitt and Dr. Virginie Mengin for sharing their insights and  
706 expertise, as well as the other members of the PHYTOCAL consortium, Nemhauser  
707 and Imaizumi labs for feedback and discussion. We also thank the undergraduate  
708 researchers from Dr Arsovski's Spring 2018 Special Field Topics class who carried  
709 out protocol optimization and preliminary measurements of phenotypes presented  
710 here: Ericka Budinich, Jonas Hill, Sean Hoeger, Katrin Hosseini, Nikhil Kaza, Winnie  
711 Kwong, Kellen Larsen, Andrew Lui, Rohan Menon, Claudia Moroney, Anita Nguyen,  
712 Arthur Sargent, Emma Stevens, and Nanami Tsumura. This work was supported by  
713 the National Science Foundation participation in the ERA-CAPS program (IOS-  
714 1539834).

715

716 **Figure 1: RNAseq experimental set-up (A).** At 3 weeks of age the “pre” sample  
717 was collected from the first of the two developmentally matched leaves. Symmetrical  
718 samples from the same leaf was collected for chlorophyll measurement. The second  
719 matched leaf was covered with tinfoil at this time. 24 hours later the “24” sample was  
720 harvested from the uncovered leaf, the same leaf that provided the “pre” sample. 48  
721 hours later the tinfoil was removed from the covered leaf and the “dark” samples  
722 were similarly collected. 24 hours later the “recovery” samples were collected from  
723 this same leaf. Samples were immediately frozen in liquid nitrogen. Three biological  
724 replicated were similarly collected. Total chlorophyll in Pre, dark, and recovery  
725 samples, error bars are SE.

726

727 **Figure 2: Genes differentially expressed in dark.** A Gene Ontology (GO) analysis  
728 of genes uniquely differentially expressed in wild type and *BrphyB* mutant leaves  
729 following 72 hours of dark. A) Upregulated genes. B) Downregulated genes. C)  
730 Expression values of 3 biological replicates in exemplar genes in Pre(P), Dark (D)  
731 and Recovery (R).

732

733 **Figure 3: Genes differentially expressed on return to light.** A Gene Ontology  
734 (GO) analysis of genes uniquely differentially expressed in wild type and *BrphyB*  
735 mutant leaves 24 hours after return to light. A) Upregulated genes. B)  
736 Downregulated genes. C) Expression values of 3 biological replicates in exemplar  
737 genes in Pre(P), Dark (D) and Recovery (R).

738

739 **Figure 4 : BrphyB mutant plants have fewer and larger chloroplasts.** A.  
740 Fluorescent images of chloroplasts in 3 week old *B.rapa* leaves. B. Chloroplast  
741 density in same leaves as A. Chloroplast area of individual chloroplasts in same  
742 plants as A. Lower case letters in B and C indicate significant difference (ANOVA  
743 and Tukey HSD multiple comparison test;  $p < 0.001$ )

744

## References

- Alabadí D, Oyama T, Yanovsky MJ, Harmon FG, Más P, Kay SA.** 2001. Reciprocal regulation between TOC1 and LHY/CCA1 within the Arabidopsis circadian clock. *Science (New York, N.Y.)* **293**, 880–883.
- Alvarez JM, Vidal EA, Gutiérrez RA.** 2012. Integration of local and systemic signaling pathways for plant N responses. *Current Opinion in Plant Biology* **15**, 185–191.
- Arsovski AA, Galstyan A, Guseman JM, Nemhauser JL.** 2012. Photomorphogenesis. *The Arabidopsis Book / American Society of Plant Biologists* **10**.
- Arsovski AA, Zemke JE, Haagen BD, Kim S-H, Nemhauser JL.** 2018. Phytochrome B regulates resource allocation in *Brassica rapa*. *Journal of Experimental Botany* **69**, 2837–2846.
- Benjamini Y, Hochberg Y.** 1995. Controlling the False Discovery Rate: A Practical and Powerful Approach to Multiple Testing. *Journal of the Royal Statistical Society. Series B (Methodological)* **57**, 289–300.
- Brouwer B, Ziolkowska A, Bagard M, Keech O, Gardeström P.** 2012. The impact of light intensity on shade-induced leaf senescence: Light-dependent induction of leaf senescence. *Plant, Cell & Environment* **35**, 1084–1098.
- Buchanan-Wollaston V, Earl S, Harrison E, Mathas E, Navabpour S, Page T, Pink D.** 2003. The molecular analysis of leaf senescence--a genomics approach. *Plant Biotechnology Journal* **1**, 3–22.
- Casal JJ.** 2012. Shade avoidance. *The Arabidopsis Book* **10**, e0157.
- Castillon A, Shen H, Huq E.** 2007. Phytochrome Interacting Factors: central players in phytochrome-mediated light signaling networks. *Trends in Plant Science* **12**, 514–521.
- Castro-Mondragon JA, Jaeger S, Thieffry D, Thomas-Chollier M, van Helden J.** 2017. RSAT matrix-clustering: dynamic exploration and redundancy reduction of transcription factor binding motif collections. *Nucleic Acids Research* **45**, e119–e119.
- Chen M, Chory J, Fankhauser C.** 2004. Light signal transduction in higher plants. *Annual Review of Genetics* **38**, 87–117.
- Chen M, Tao Y, Lim J, Shaw A, Chory J.** 2005. Regulation of phytochrome B nuclear localization through light-dependent unmasking of nuclear-localization signals. *Current biology: CB* **15**, 637–642.
- Chen D, Xu G, Tang W, Jing Y, Ji Q, Fei Z, Lin R.** 2013. Antagonistic basic helix-loop-helix/bZIP transcription factors form transcriptional modules that integrate light and reactive oxygen species signaling in Arabidopsis. *The Plant Cell* **25**, 1657–1673.

**Chory J, Peto CA, Ashbaugh M, Saganich R, Pratt L, Ausubel F.** 1989. Different Roles for Phytochrome in Etiolated and Green Plants Deduced from Characterization of *Arabidopsis thaliana* Mutants. *The Plant Cell* **1**, 867–880.

**Coello P, Sassen A, Haywood V, Davis KR, Walker JC.** 1999. Biochemical characterization and expression of RLK4, a receptor-like kinase from *Arabidopsis thaliana*. *Plant Science* **142**, 83–91.

**Epstein E, Bloom AAJ.** 2005. *Mineral Nutrition Of Plants: Principles And Perspectives*. Sinauer Associates, Incorporated.

**Farinas B, Mas P.** 2011. Functional implication of the MYB transcription factor RVE8/LCL5 in the circadian control of histone acetylation. *The Plant Journal: For Cell and Molecular Biology* **66**, 318–329.

**Franklin KA, Whitelam GC.** 2005. Phytochromes and shade-avoidance responses in plants. *Annals of Botany* **96**, 169–175.

**Fraser DP, Hayes S, Franklin KA.** 2016. Photoreceptor crosstalk in shade avoidance. *Current Opinion in Plant Biology* **33**, 1–7.

**Gaspin C, Rami J-F, Lescure B.** 2010. Distribution of short interstitial telomere motifs in two plant genomes: putative origin and function. *BMC Plant Biology* **10**, 283.

**Ghassemian M, Lutes J, Tepperman JM, Chang H-S, Zhu T, Wang X, Quail PH, Lange BM.** 2006. Integrative analysis of transcript and metabolite profiling data sets to evaluate the regulation of biochemical pathways during photomorphogenesis. *Archives of Biochemistry and Biophysics* **448**, 45–59.

**Gregory FG.** 1937. Mineral Nutrition of Plants. *Annual Review of Biochemistry* **6**, 557–578.

**Guan M, Møller IS, Schjoerring JK.** 2015. Two cytosolic glutamine synthetase isoforms play specific roles for seed germination and seed yield structure in *Arabidopsis*. *Journal of Experimental Botany* **66**, 203–212.

**Guo Y, Cai Z, Gan S.** 2004. Transcriptome of *Arabidopsis* leaf senescence. *Plant, Cell and Environment* **27**, 521–549.

**Harmer SL, Kay SA.** 2005. Positive and Negative Factors Confer Phase-Specific Circadian Regulation of Transcription in *Arabidopsis*. *The Plant Cell* **17**, 1926–1940.

**Himelblau E, Amasino RM.** 2001. Nutrients mobilized from leaves of *Arabidopsis thaliana* during leaf senescence. *Journal of Plant Physiology* **158**, 1317–1323.

**Hirel B, Tétu T, Lea PJ, Dubois F.** 2011. Improving Nitrogen Use Efficiency in Crops for Sustainable Agriculture. *Sustainability* **3**, 1452–1485.

**Hornitschek P, Kohnen MV, Lorrain S, et al.** 2012. Phytochrome interacting factors 4 and 5 control seedling growth in changing light conditions by directly



controlling auxin signaling. *The Plant Journal: For Cell and Molecular Biology* **71**, 699–711.

**Hsu PY, Devisetty UK, Harmer SL.** 2013. Accurate timekeeping is controlled by a cycling activator in *Arabidopsis* (J Chory, Ed.). *eLife* **2**, e00473.

**Hu W, Franklin KA, Sharrock RA, Jones MA, Harmer SL, Lagarias JC.** 2013. Unanticipated regulatory roles for *Arabidopsis* phytochromes revealed by null mutant analysis. *Proceedings of the National Academy of Sciences of the United States of America* **110**, 1542–1547.

**Huq E, Al-Sady B, Hudson M, Kim C, Apel K, Quail PH.** 2004. Phytochrome-interacting factor 1 is a critical bHLH regulator of chlorophyll biosynthesis. *Science (New York, N.Y.)* **305**, 1937–1941.

**Huq E, Quail PH.** 2002. PIF4, a phytochrome-interacting bHLH factor, functions as a negative regulator of phytochrome B signaling in *Arabidopsis*. *The EMBO journal* **21**, 2441–2450.

**Jensen LS, Christensen L, Mueller T, Nielsen NE.** 1997. Turnover of residual <sup>15</sup>N-labelled fertilizer N in soil following harvest of oilseed rape (*t Brassica napus* L.). *Plant and Soil* **190**, 193–202.

**Ji Y, Li Q, Liu G, Selvaraj G, Zheng Z, Zou J, Wei Y.** 2019. Roles of Cytosolic Glutamine Synthetases in *Arabidopsis* Development and Stress Responses. *Plant and Cell Physiology* **60**, 657–671.

**Jiang Z, Xu G, Jing Y, Tang W, Lin R.** 2016. Phytochrome B and REVEILLE1/2-mediated signalling controls seed dormancy and germination in *Arabidopsis*. *Nature Communications* **7**, 12377.

**Jin J, He K, Tang X, Li Z, Lv L, Zhao Y, Luo J, Gao G.** 2015. An *Arabidopsis* Transcriptional Regulatory Map Reveals Distinct Functional and Evolutionary Features of Novel Transcription Factors. *Molecular Biology and Evolution* **32**, 1767–1773.

**Jin J, Tian F, Yang D-C, Meng Y-Q, Kong L, Luo J, Gao G.** 2017. PlantTFDB 4.0: toward a central hub for transcription factors and regulatory interactions in plants. *Nucleic Acids Research* **45**, D1040–D1045.

**Jin J, Zhang H, Kong L, Gao G, Luo J.** 2014. PlantTFDB 3.0: a portal for the functional and evolutionary study of plant transcription factors. *Nucleic Acids Research* **42**, D1182–D1187.

**Keech O, Pesquet E, Ahad A, Askne A, Nordvall D, Vodnala SM, Tuominen H, Hurry V, Dizengremel P, Gardeström P.** 2007. The different fates of mitochondria and chloroplasts during dark-induced senescence in *Arabidopsis* leaves. *Plant, Cell & Environment* **30**, 1523–1534.

**Kobayashi K, Sasaki D, Noguchi K, et al.** 2013. Photosynthesis of Root Chloroplasts Developed in *Arabidopsis* Lines Overexpressing GOLDEN2-LIKE Transcription Factors. *Plant and Cell Physiology* **54**, 1365–1377.

**Koeslin-Findeklee F, Rizi VS, Becker MA, Parra-Londono S, Arif M, Balazadeh S, Mueller-Roeber B, Kunze R, Horst WJ.** 2015. Transcriptomic analysis of nitrogen starvation- and cultivar-specific leaf senescence in winter oilseed rape (*Brassica napus* L.). *Plant Science: An International Journal of Experimental Plant Biology* **233**, 174–185.

**Krahmer J, Ganpudi A, Abbas A, Romanowski A, Halliday KJ.** 2018. Phytochrome, Carbon Sensing, Metabolism, and Plant Growth Plasticity. *Plant Physiology* **176**, 1039–1048.

**Kurihara D, Mizuta Y, Sato Y, Higashiyama T.** 2015. ClearSee: a rapid optical clearing reagent for whole-plant fluorescence imaging. *Development (Cambridge, England)* **142**, 4168–4179.

**Law SR, Chrobok D, Juvany M, et al.** 2018. Darkened leaves use different metabolic strategies for senescence and survival. *Plant Physiology*, pp.00062.2018.

**Leivar P, Tepperman JM, Cohn MM, Monte E, Al-Sady B, Erickson E, Quail PH.** 2012. Dynamic Antagonism between Phytochromes and PIF Family Basic Helix-Loop-Helix Factors Induces Selective Reciprocal Responses to Light and Shade in a Rapidly Responsive Transcriptional Network in *Arabidopsis*. *The Plant Cell* **24**, 1398–1419.

**Li J, Li G, Wang H, Wang Deng X.** 2011. Phytochrome signaling mechanisms. *The Arabidopsis Book* **9**, e0148.

**Lim PO, Kim HJ, Gil Nam H.** 2007. Leaf Senescence. *Annual Review of Plant Biology* **58**, 115–136.

**Lopez-Juez E, Pyke KA.** 2005. Plastids unleashed: their development and their integration in plant development. *The International Journal of Developmental Biology* **49**, 557–577.

**Malagoli P, Laine P, Rossato L, Ourry A.** 2005. Dynamics of nitrogen uptake and mobilization in field-grown winter oilseed rape (*Brassica napus*) from stem extension to harvest: I. Global N flows between vegetative and reproductive tissues in relation to leaf fall and their residual N. *Annals of Botany* **95**, 853–861.

**Martínez-García JF, Huq E, Quail PH.** 2000. Direct targeting of light signals to a promoter element-bound transcription factor. *Science (New York, N.Y.)* **288**, 859–863.

**Masclaux-Daubresse C, Purdy S, Lemaitre T, Pourtau N, Taconnat L, Renou J-P, Winkler A.** 2007. Genetic variation suggests interaction between cold acclimation and metabolic regulation of leaf senescence. *Plant Physiology* **143**, 434–446.

**McCormac AC, Terry MJ.** 2002. Light-signalling pathways leading to the coordinated expression of HEMA1 and Lhcb during chloroplast development in *Arabidopsis thaliana*. *The Plant Journal: For Cell and Molecular Biology* **32**, 549–559.

- Mei H-S, Thimann KV.** 1984. The relation between nitrogen deficiency and leaf senescence. *Physiologia Plantarum* **62**, 157–161.
- Mi H, Muruganujan A, Huang X, Ebert D, Mills C, Guo X, Thomas PD.** 2019. Protocol Update for large-scale genome and gene function analysis with the PANTHER classification system (v.14.0). *Nature Protocols* **14**, 703.
- Nevarez PA, Qiu Y, Inoue H, Yoo CY, Benfey PN, Schnell DJ, Chen M.** 2017. Mechanism of Dual Targeting of the Phytochrome Signaling Component HEMERA/pTAC12 to Plastids and the Nucleus. *Plant Physiology* **173**, 1953–1966.
- Oh S, Montgomery BL.** 2014. Phytochrome-dependent coordinate control of distinct aspects of nuclear and plastid gene expression during anterograde signaling and photomorphogenesis. *Frontiers in Plant Science* **5**, 171.
- Oh E, Zhu J-Y, Wang Z-Y.** 2012. Interaction between BZR1 and PIF4 integrates brassinosteroid and environmental responses. *Nature Cell Biology*.
- Parlitz S, Kunze R, Mueller-Roeber B, Balazadeh S.** 2011. Regulation of photosynthesis and transcription factor expression by leaf shading and re-illumination in *Arabidopsis thaliana* leaves. *Journal of Plant Physiology* **168**, 1311–1319.
- Pfaffl MW.** 2001. A new mathematical model for relative quantification in real-time RT–PCR. *Nucleic Acids Research* **29**, e45.
- Rathke G-W, Behrens T, Diepenbrock W.** 2006. Integrated nitrogen management strategies to improve seed yield, oil content and nitrogen efficiency of winter oilseed rape (*Brassica napus* L.): A review. *Agriculture, Ecosystems & Environment* **117**, 80–108.
- Rawat R, Takahashi N, Hsu PY, Jones MA, Schwartz J, Salemi MR, Phinney BS, Harmer SL.** 2011. REVEILLE8 and PSEUDO-RESPONSE REGULATOR5 form a negative feedback loop within the *Arabidopsis* circadian clock. *PLoS genetics* **7**, e1001350.
- Robinson MD, McCarthy DJ, Smyth GK.** 2010. edgeR: a Bioconductor package for differential expression analysis of digital gene expression data. *Bioinformatics (Oxford, England)* **26**, 139–140.
- Schep A.** 2019. *motifmatchr: Fast Motif Matching in R*.
- Schjoerring JK, Bock JGH, Gammelvind L, Jensen CR, Mogensen VO.** 1995. Nitrogen incorporation and remobilization in different shoot components of field-grown winter oilseed rape (*Brassica napus* L.) as affected by rate of nitrogen application and irrigation. *Plant and Soil* **177**, 255–264.
- Shi H, Lyu M, Luo Y, Liu S, Li Y, He H, Wei N, Deng XW, Zhong S.** 2018. Genome-wide regulation of light-controlled seedling morphogenesis by three families of transcription factors. *Proceedings of the National Academy of Sciences* **115**, 6482–6487.

- Sobieszczuk-Nowicka E, Wrzesiński T, Bagniewska-Zadworna A, Kubala S, Rucińska-Sobkowiak R, Polcyn W, Misztal L, Mattoo AK.** 2018. Physio-Genetic Dissection of Dark-Induced Leaf Senescence and Timing Its Reversal in Barley. *Plant Physiology* **178**, 654–671.
- Song Y, Yang C, Gao S, Zhang W, Li L, Kuai B.** 2014. Age-Triggered and Dark-Induced Leaf Senescence Require the bHLH Transcription Factors PIF3, 4, and 5. *Molecular Plant* **7**, 1776–1787.
- Strasser B, Sánchez-Lamas M, Yanovsky MJ, Casal JJ, Cerdán PD.** 2010. *Arabidopsis thaliana* life without phytochromes. *Proceedings of the National Academy of Sciences of the United States of America* **107**, 4776–4781.
- Toledo-Ortiz G, Johansson H, Lee KP, Bou-Torrent J, Stewart K, Steel G, Rodríguez-Concepción M, Halliday KJ.** 2014. The HY5-PIF regulatory module coordinates light and temperature control of photosynthetic gene transcription. *PLoS genetics* **10**, e1004416.
- Wang J-G, Chen C-H, Chien C-T, Hsieh H-L.** 2011. FAR-RED INSENSITIVE 219 modulates CONSTITUTIVE PHOTOMORPHOGENIC 1 activity via physical interaction to regulate hypocotyl elongation in *Arabidopsis*. *Plant Physiology*.
- Wang D, Portis AR.** 2007. A novel nucleus-encoded chloroplast protein, PIFI, is involved in NAD(P)H dehydrogenase complex-mediated chlororespiratory electron transport in *Arabidopsis*. *Plant Physiology* **144**, 1742–1752.
- Waters MT, Moylan EC, Langdale JA.** 2008. GLK transcription factors regulate chloroplast development in a cell-autonomous manner. *The Plant Journal: For Cell and Molecular Biology* **56**, 432–444.
- Waters MT, Wang P, Korkaric M, Capper RG, Saunders NJ, Langdale JA.** 2009. GLK Transcription Factors Coordinate Expression of the Photosynthetic Apparatus in *Arabidopsis*. *The Plant Cell* **21**, 1109–1128.
- Weaver LM, Amasino RM.** 2001. Senescence Is Induced in Individually Darkened *Arabidopsis* Leaves, but Inhibited in Whole Darkened Plants. *Plant Physiology* **127**, 876–886.
- Wies G, Mantese AI, Casal JJ, Maddonni GÁ.** 2019. Phytochrome B enhances plant growth, biomass and grain yield in field-grown maize. *Annals of Botany* **123**, 1079–1088.
- Xiong D, Huang J, Peng S, Li Y.** 2017. A few enlarged chloroplasts are less efficient in photosynthesis than a large population of small chloroplasts in *Arabidopsis thaliana*. *Scientific Reports* **7**, 5782.
- Xu MY, Dong Y, Zhang QX, Zhang L, Luo YZ, Sun J, Fan YL, Wang L.** 2012. Identification of miRNAs and their targets from *Brassica napus* by high-throughput sequencing and degradome analysis. *BMC genomics* **13**, 421.


**Xu D, Marino G, Klingl A, Enderle B, Monte E, Kurth J, Hiltbrunner A, Leister D, Kleine T.** 2019. Extrachloroplastic PP7L Functions in Chloroplast Development and Abiotic Stress Tolerance1[OPEN]. *Plant Physiology* **180**, 323–341.

**Yang D, Seaton DD, Krahmer J, Halliday KJ.** 2016. Photoreceptor effects on plant biomass, resource allocation, and metabolic state. *Proceedings of the National Academy of Sciences* **113**, 7667–7672.

**Zhang X, Chen Y, Wang ZY, Chen Z, Gu H, Qu LJ.** 2007. Constitutive expression of CIR1 (RVE2) affects several circadian-regulated processes and seed germination in Arabidopsis. *The Plant journal* □: for cell and molecular biology **51**, 512–525.

**Zhang Y, Mayba O, Pfeiffer A, Shi H, Tepperman JM, Speed TP, Quail PH.** 2013. A Quartet of PIF bHLH Factors Provides a Transcriptionally Centered Signaling Hub That Regulates Seedling Morphogenesis through Differential Expression-Patterning of Shared Target Genes in Arabidopsis. *PLOS Genetics* **9**, e1003244.


**A**



**Figure 1: RNAseq experimental set-up (A).** At 3 weeks of age the “pre” sample was collected from the first of the two developmentally matched leaves. Symmetrical samples from the same leaf was collected for chlorophyll measurement. The second matched leaf was covered with tinfoil at this time. 24 hours later the “24” sample was harvested from the uncovered leaf, the same leaf that provided the “pre” sample. 48 hours later the tinfoil was removed from the covered leaf and the “dark” samples were similarly collected. 24 hours later the “recovery” samples were collected from this same leaf. Samples were immediately frozen in liquid nitrogen. Three biological replicated were similarly collected. Total chlorophyll in Pre, dark, and recovery samples, error bars are SE.


**A**

**Genes upregulated in D vs. P**




**B**

**Genes downregulated in D vs. P**




**C**



**Figure 2: Genes differentially expressed in dark.** A Gene Ontology (GO) analysis of genes uniquely differentially expressed in wild type and *BrphyB* mutant leaves following 72 hours of dark. A) Upregulated genes. B) Downregulated genes. C) Expression values of 3 biological replicates in exemplar genes in Pre(P), Dark (D) and Recovery (R).


**A**

**Genes upregulated in R vs. D**




**B**

**Genes downregulated in R vs. D**




**C**




**Figure 3: Genes differentially expressed on return to light.** A Gene Ontology (GO) analysis of genes uniquely differentially expressed in wild type and BrphyB mutant leaves 24 hours after return to light. A) Upregulated genes. B) Downregulated genes. C) Expression values of 3 biological replicates in exemplar genes in Pre (P), Dark (D) and Recovery (R).



A



B



C




Figure 4 : *BrphyB* mutant plants have fewer and larger chloroplasts. A. Fluorescent images of chloroplasts in 3 week old *B. rapa* leaves. B. Chloroplast density in same leaves as A. Chloroplast area of individual chloroplasts in same plants as A. Lower case letters in B and C indicate significant difference (ANOVA and Tukey HSD multiple comparison test; p<0.001)