

1 **High performance machine learning models can fully automate labeling of camera trap images** 2 **for ecological analyses**

3
4 Robin Whytock^{1,2*}†, Jędrzej Świeżewski^{3†}, Joeri A. Zwerts⁴, Tadeusz Bara-Słupski⁴, Aurélie Flore
5 Koumba Pambo², Marek Rogala³, Laila Bahaa-el-din⁵, Kelly Boekee^{6,7}, Stephanie Brittain^{8,9}, Anabelle
6 W. Cardoso¹⁰, Philipp Henschel^{11,12}, David Lehmann^{2,1}, Brice Momboua², Cisquet Kiebou Opepa¹³,
7 Christopher Orbell^{11,1}, Ross T. Pitman¹¹, Hugh S. Robinson^{11,14}, Katharine A. Abernethy^{1,12}

8 † These authors contributed equally to the manuscript.

9 ***Corresponding author:** Robin C. Whytock, robbie.whytock1@stir.ac.uk

10
11 ¹ Faculty of Natural Sciences, University of Stirling, FK9 4LA, UK

12 ² Agence Nationale des Parcs Nationaux, Libreville, Gabon

13 ³ Appsilon AI for Good, Warsaw, Poland

14 ⁴ Utrecht University, Heidelberglaan 8, 3584 CS Utrecht, The Netherlands

15 ⁵ School of Life Sciences, University of KwaZulu-Natal, South Africa

16 ⁶ Program for the Sustainable Management of Natural Resources, South West Region, Buea, Cameroon

17 ⁷ Center for Tropical Forest Science, Smithsonian Tropical Research Institute, Balboa, Ancon,
18 Republic of Panama

19 ⁸ The Interdisciplinary Centre for Conservation Science, Department of Zoology, University of Oxford,
20 11a Mansfield Road, Oxford, OX1 3SZ

21 ⁹ The Institute of Zoology, Zoological Society of London, Outer Circle, London NW1 4RY

22 ¹⁰ Yale University, New Haven, CT 06520, USA

23 ¹¹ Panthera, 8 West 40th Street, 18th Floor, New York, NY 10018, USA

24 ¹² Institut de Recherche en Ecologie Tropicale, CENAREST, BP 842 Libreville, Gabon

25 ¹³ Wildlife Conservation Society, Republic of the Congo

26 ¹⁴ Wildlife Biology Program, W.A. Franke College of Forestry and Conservation, University of
27 Montana, Natural Sciences Room 205, Missoula, Montana, 59812, USA

29 **ORCID IDs**

30
31 **Robin C. Whytock:** 0000-0002-0127-6071

32 **Jędrzej Świeżewski:** not available

33 **Joeri A. Zwerts:** 0000-0003-3841-6389

34 **Tadeusz Bara-Słupski:** not available

35 **Aurélie Flore Koumba Pambo:** not available

36 **Marek Rogala:** 0000-0002-9949-4551

37 **Laila Bahaa-el-din:** not available

38 **Kelly Boekee:** 0000-0001-8131-5204

39 **Stephanie Brittain:** 0000-0002-7865-0391

40 **Anabelle W. Cardoso:** 0000-0002-4327-7259

41 **Philipp Henschel:** not available

42 **David Lehmann:** 0000-0002-4529-8117

43 **Brice Momboua:** not available

44 **Cisquet Kiebou Opepa:** not available

45 **Christopher Orbell:** not available

46 **Ross. T. Pitman:** 0000-0002-3574-0063

47 **Hugh S. Robinson:** 0000-0002-4060-3143

48 **Katharine A. Abernethy:** 0000-0002-0393-9342

49

50 **Keywords:** artificial intelligence, birds, biodiversity, Central Africa, mammals

51 **Author contributions:** R.C.W., J.S., M.R., A.F.K.P., P.H., C.O., R.P., H.R., K.A. T.B. designed
52 research; R.C.W., J.S., M.R. performed research; R.C.W., J.S. analyzed data; R.C.W., J.A.W., L.B.,
53 K.B., A.C., D.L., B.M., C.K.O., C.O. collected data; R.W., J.S., J.A.Z., T.B., A.F.K.P., M.R., L.B.,
54 K.B., S.B., A.C., P.H., D.L., C.O., H.R., K.A. wrote the paper.

55 **This PDF file includes:**

56 Main text

57 Figures 1 to 7

58 Tables 1 and 2

59 Supplementary Tables S1 to S4

60 Supplementary Figures S1 to S9

61 **Data and code availability statement:** All data and code will be publicly archived in the University of
62 Stirling's publicly accessible data repository and given a unique DOI on acceptance. Code for the
63 machine learning model is available for review online at
64 https://github.com/Appsilon/gabon_wildlife_training. Code for the offline application to run the model
65 is available at <https://github.com/Appsilon/wildlife-explorer>. R code for the ecological analyses are
66 available for review online at https://github.com/rcwhytock/Whytock_and_Swiezewski_et_al_2020/.

66 Abstract

67 1. Ecological data are increasingly collected over vast geographic areas using arrays of digital sensors.
 68 Camera trap arrays have become the ‘gold standard’ method for surveying many terrestrial mammals
 69 and birds, but these arrays often generate millions of images that are challenging to process. This
 70 causes significant latency between data collection and subsequent inference, which can impede
 71 conservation at a time of ecological crisis. Machine learning algorithms have been developed to
 72 improve camera trap data processing speeds, but these models are not considered accurate enough for
 73 fully automated labeling of images.

74 2. Here, we present a new approach to building and testing a high performance machine learning model
 75 for fully automated labeling of camera trap images. As a case-study, the model classifies 26 Central
 76 African forest mammal and bird species (or groups). The model was trained on a relatively small
 77 dataset (*c.*300,000 images) but generalizes to fully independent data and outperforms humans in several
 78 respects (e.g. detecting ‘invisible’ animals). We show how the model’s precision and accuracy can be
 79 evaluated in an ecological modeling context by comparing species richness, activity patterns ($n = 4$
 80 species tested) and occupancy ($n = 4$ species tested) derived from machine learning labels with the
 81 same estimates derived from expert labels.

82 3. Results show that fully automated labels can be equivalent to expert labels when calculating species
 83 richness, activity patterns ($n = 4$ species tested) and estimating occupancy ($n = 3$ of 4 species tested) in
 84 completely out-of-sample test data ($n = 227$ camera stations, $n = 23868$ images). Simple thresholding
 85 (discarding uncertain labels) improved the model’s performance when calculating activity patterns and
 86 estimating occupancy, but did not improve estimates of species richness.

87 4. We provide the user-community with a multi-platform, multi-language user interface for running the
 88 model offline, and conclude that high performance machine learning models can fully automate
 89 labeling of camera trap data.

90 **Introduction**

91 The urgent need to understand how ecosystems are responding to rapid environmental change has
 92 driven a ‘big data’ revolution in ecology and conservation (Farley, Dawson, Goring, & Williams,
 93 2018). High resolution ecological data are now streamed in real-time from satellites, Global Positioning
 94 System tags, bioacoustic detectors, cameras and other sensor arrays. The data generated offer consider-
 95 able opportunities to ecologists, but challenges such as data processing, data storage and data sharing
 96 cause latency between data gathering and ecological inference (i.e. creating derived ecological metrics,
 97 testing ecological hypotheses and quantifying ecological change), sometimes in the order of years or
 98 more. Overcoming these challenges could open the gateway to ecological ‘forecasting’, where direc-
 99 tional changes in ecological processes are detected in real time and near-term responses are predicted
 100 effectively using an iterative data gathering, model updating and model prediction approach (Dietze et
 101 al., 2018).

102
 103 Digital camera traps or wildlife ‘trail cams’ have revolutionized wildlife monitoring and are now the
 104 ‘gold standard’ for monitoring many medium to large terrestrial mammals (Glover-Kapfer, Soto
 105 Navarro, & Wearn, 2019). Animals and their behavior are identified in images either by manual label-
 106 ing, using citizen science platforms (Swanson et al., 2015) or, more recently, by using machine learning
 107 models (Norouzzadeh et al., 2018; Tabak et al., 2019; Willi et al., 2019). Machine learning models can
 108 at minimum separate true animal detections from non-detections (Wei, Luo, Ran, & Li, 2020) or in the
 109 most advanced examples identify species, count individuals and describe behavior (Norouzzadeh et al.,
 110 2018). These recent advances in machine learning have increased the speed at which camera trap data
 111 are analyzed but, in all cases we are aware of, the outputs (e.g. species labels) are not used to make eco-
 112 logical inference directly. Instead, machine learning models are typically used as a ‘first pass’ to iden-
 113 tify and group images belonging to individual species for full or partial manual validation at a later
 114 stage, or to cross-validate labels from citizen science platforms (Willi et al., 2019). This can substan-

tially reduce manual labeling effort but many hundreds or thousands of photos might still need to be labeled manually. Thus, although machine learning models are reducing manual data processing times, ecologists are not yet comfortable using the outputs (e.g. species labels) as part of a completely automated workflow. This is despite the development of advanced machine learning models that classify species in camera trap images with accuracy that matches or exceeds humans (Norouzzadeh et al., 2018; Tabak et al., 2019).

One significant challenge limiting the application of machine learning models to camera trap data is that models rarely generalize well to completely out-of-sample data (i.e. data from new, spatially and temporally independent studies), particularly when used to classify animals to species level (Beery, Van Horn, & Perona, 2018). Models can quickly learn the features of specific camera ‘stations’ (the spatial replicate in camera trap studies) such as the general background instead of learning features of the animal itself. This problem is further amplified by the fact that rare species in the training data might only ever appear at a limited number of camera stations, so training and validation data are rarely independent. Various approaches can be used to reduce these biases, such as carefully ensuring that training and validation data are independent (e.g. by using data from multiple studies), and by using data augmentation such as adding noise to training data in the form of image transformations. Until the problem of generalization can be overcome, machine learning models for classifying camera trap images will remain an important tool for reducing manual labeling effort, but they will not achieve their full potential for creating fully automated pipelines for data analysis.

Machine learning models also have the potential to be deployed inside camera trap hardware in the field at the ‘edge’ (i.e. on micro-computers installed inside hardware that collects data), with summarized results (e.g. species labels) transmitted in real-time via a Global System for Mobile Communications networks or via satellite (Glover-Kapfer et al., 2019). In geographically remote areas or time-sensitive situations (e.g. law enforcement) this would greatly reduce the latency between data capture and

interpretation, and reduce the expense and effort required to collect data in remote regions by removing the need to transfer data-heavy images across wireless networks. However, before ‘smart’ cameras become a reality, it is essential that users understand how uncertainty in machine learning model predictions might impact derived ecological metrics and analyses, which are often sensitive to biases (e.g. false positives in occupancy models). To achieve this, there is a need to develop workflows that test the performance of machine learning models in an ecological modeling context that goes beyond simple measures of precision and accuracy.

Ideally, if machine learning models had 100% precision and accuracy (e.g. for species identification), camera trap data could be collected, labeled automatically using the model and the results used to directly calculate ecological metrics or as variables in ecological models. However, the reality is that machine learning models are imperfect. It is therefore uncertain what levels of precision and accuracy are needed to meet the requirements of ecological analyses. This is particularly the case for the spatial and temporal analyses of animal distributions in camera trap data, which require specialized ecological models (e.g. occupancy models) that account for imperfect detection (MacKenzie et al., 2002).

In this paper, we describe the approach used to build a new high-performance machine learning model that identifies species in camera trap images (26 species/groups of Central African forest mammals and birds) that generalizes to spatially independent data. To evaluate how well the machine learning model labeling precision and accuracy performs in an ecological modeling context, we (1) evaluate how uncertainties in the precision and accuracy of machine learning labels affect ecological inference (derived metrics of species richness, activity patterns and occupancy) compared to the same metrics calculated using expert, manually generated labels, and (2) propose a workflow to ‘ground truth’ the performance of machine learning models for camera trap data in an ecological modeling context. We discuss the implications of these results for making fully automated ecological inference from camera trap data using the outputs of machine learning models. We also provide the user community with an easily installed,

open-source graphical user interface that needs no understanding of machine learning to run the model offline on both camera trap images and videos.

Methods

Data preparation

As a case study, the model was developed for classifying terrestrial forest mammals and birds in Central Africa (see Table S1 for further details on species and groups), where camera traps are now frequently deployed over large spatial scales to survey secretive birds and mammals in remote and inaccessible landscapes (Bahaa-el-din & Cusack, 2018; Bessone et al., 2020; O'Brien et al., 2020). Training data were obtained from multiple countries and sources (c.1.6 million images; reduced to $n = 347120$ images after data processing; Table 1). Each source used different camera trap models (Reconyx, Bushnell, Cuddeback, Panthera Cams) and images were diverse in resolution, quality (e.g. sharpness, illumination) and color. Individual studies also used different field protocols for camera deployment but all were focused on detecting terrestrial forest mammals, with cameras installed on trees approximately 30 - 40 cm above ground level. The exception to this was data from (Cardoso et al., 2020) who installed cameras at a height of approximately 1 m for the primary purpose of detecting forest elephants *Loxodonta cyclotis*. Camera trap configuration was set to be highly sensitive in some cases and images were often captured in a series of rapid, short bursts (e.g. taking 10 images consecutively). This resulted in long sequences of very similar images, for example showing an animal walking in front of the camera (Figure S1).

Table 1. Sources of training data used to train the machine learning model for classifying species in camera trap images, sorted by number of images provided. The final subset of data used to train the model was $n = 347120$ images (see later).

Source	Country	Reference	n images
Anabelle Cardoso	Gabon	(Cardoso et al., 2020)	102418
Kelly Boekee	Cameroon	-	123954
Cisquet Kiebou Opepa	Republic of Congo	-	60393
Joeri Zwerts	Cameroon	-	36027
Laila Bahaa-el-Din	Gabon	(Bahaa-el-din et al., 2013)	16558
Stephanie Brittain	Cameroon	-	7770

It was important to account for image sequences when selecting a validation set during the model training phase, since there was a risk of highly similar images being present in both the training and validation sets. To address this issue, the training and validation split was performed based on image meta-data (timing of images and image source) to identify unique ‘events’ and camera locations that were not replicated across the training and validation split (Norouzzadeh et al., 2018). This solution posed a challenge for maintaining class balances in the training and validation sets, but it reduced the risk non-independent training and validation sets. A total of 27 classes were used to train the model, which were mostly mammals or mammal groups ($n = 21$), birds ($n = 4$), humans ($n = 1$) and ‘blank’ images (i.e. no mammal, bird or human). Details of taxonomy and justification for species groups are in Table S1.

Issues identified in the training data

Our ‘real-life’ training data had not been pre-processed or professionally curated for the purposes of training machine learning models and naturally contained errors that arise from hardware faults, human error and different approaches to manual species labeling by experts. We identified three primary sources of error. The first was over-exposed images (a hardware fault) where the image foreground was ‘flooded’ by the flash (usually at night), making the image appear mostly white. Animals in these

images were sometimes partially visible and could be classified by a skilled human observer, despite the loss of color information, texture and other detail. However, over-exposed images presented a challenge for the machine learning model because white dominated the image regardless of the species.

The second main source of error was caused by under-exposed images. This error was revealed after inspecting model outputs during the training phase, and showed that highly under-exposed images appeared almost entirely or entirely black to a human observer, but the machine learning model was capable of using information in the image to detect and correctly classify the species (Figure 1).




Figure 1. (a) Raw image from the dataset, labeled by experts as "blank", but classified by the machine learning model with high certainty as a red duiker. (b) The same image as in (a), but manually brightened by narrowing the displayed color spectrum, reveals a red duiker is present and the model was correct.

The final source of error in the training data was mis-labeled images (e.g. confusing similar species, such as chimpanzee *Pan troglodytes* and gorilla *Gorilla gorilla*) and using different approaches to labeling, for example one data source combined all primates into 'monkey', whereas other data sources separated apes from other primates.

We used an iterative approach to address these issues that consisted of model training, validation, error

correction (correcting mis-labeled images in the training data) and model updating. In particular, we carefully inspected images that appeared to be incorrectly labeled by the model, but which were labeled with high confidence. This approach revealed hidden problems in the data, such as the presence of animals in under-exposed images that would have otherwise led us to underestimate the model's performance.

Machine learning model

We chose the established ResNet50 architecture to build the model (He, Zhang, Ren, & Sun, 2016). Transfer learning was used to speed up training and we used weights pre-trained on the ImageNet dataset. We identified species using the entire image frame without using bounding boxes and used basic augmentation (horizontal flips, rotations, zoom, lighting and contrast adjustments, and warps) during training, but not during model validation. We used one-cycle policy training (Smith, 2018) and trained using progressive resizing in two stages. Details on the training scheme and implementation can be found in our GitHub repository (https://github.com/Appsilon/gabon_wildlife_training). It is worth noting that most of the training approaches and many of the mechanisms we used to enhance training were taken directly or almost directly from the fast.ai Python library (<https://github.com/fastai>), exemplifying how exceptionally robust the library is. We trained the models on various virtual machines equipped with GPU processing units, run on Google Cloud Platform with resources granted by a Google Cloud Education grant.

Out-of-sample test data

One of the major limitations to model performance for camera trap images is the ability to generalize predictions to new, independent camera stations, i.e. unique locations with different backgrounds not seen during model training (Beery et al., 2018). Since our objective was to create a model that could generalize well to new study sites, we tested the final model's performance using a new out-of-sample

dataset that was completely spatially and temporally independent from the data used to train the model. These out-of-sample data consisted of images from 227 camera stations surveyed between 16 January 2018 and 4 October 2019 in central and southern Gabon in closed canopy forest. Cameras also differed from the models used in the training data (Panthera Cams V4 and V5), but field protocols were similar and cameras were placed approximately 30 cm above the ground on a tree at a distance of c. 3 – 5 m perpendicular to the center of animal trails. Single-frame images were captured using medium sensitivity settings, and images were separated by a minimum of 1 s. The aim of the study was to survey the small-to-large mammal community, with a particular focus on great apes (*Pan troglodytes*, *Gorilla gorilla*), forest elephants *Loxodonta cyclotis*, leopard *Panthera pardus* and golden cat *Caracal aurata*. These data ($n = 23868$ images, median 75, range 1 - 545 images per station) were manually labeled by an expert (co-author CO).

Summary of model's general performance

To allow general comparison of our model's performance with other similar models in the literature (Norouzzadeh et al., 2018; Tabak et al., 2019; Willi et al., 2019) we calculated top-one and top-five accuracies using the out-of-sample data. Top-one accuracy is the percent of expert labels that match the top-ranking label generated by the machine learning model. Top-five accuracy calculates the percent of expert labels that match any of the top five ranking machine learning generated labels. Top-one accuracy for the overall machine learning model was 77.63% and top-five accuracy was 94.24% (Table S2; Figures S2 & S3). After aggregating labels of similar species that were frequently mis-classified by the model into a reduced set of 11 classes, top-one and top-five accuracies increased to 79.92% and 95.99%, respectively (Figure S4). The model can classify around 4000 images (c.0.5 MB in size) per hour using an Intel® Core™ i7-8665U CPU @ 1.90GHz × 8 and the model can operate 24/7 if necessary. For comparison, based on our experience, manual labeling can be done at speeds ranging from 125 to 500 images per hour depending on the quality of the images and if images are captured in

275 sequences (which can be faster to label manually).

276

277 We also compared the precision and recall for each species from our optimal model (see later, Table 2)
 278 with precision and recall for the same species reported for the model used by the WildlifeInsights web-
 279 platform (www.wildlifeinsights.org). This global project uses a deep convolutional neural network
 280 trained using Google's Tensorflow framework and a training dataset of 8.7M images, comprising 614
 281 species.

282

283 ***Comparing derived ecological metrics using machine learning labels and expert labels***

284 We calculated three common ecological metrics for the out-of-sample data (raw species richness at in-
 285 dividual camera stations, activity patterns for four focal species, and occupancy for four focal species)
 286 separately using the manually generated, expert labels and the machine learning generated labels.

287 Species richness (the number of species in a discrete unit of space and time) can be used to quantify
 288 temporal and spatial changes in biodiversity. Although other measures of species diversity exist, we
 289 chose this simple metric because it is widely used in the ecology literature despite its limitations. Activ-
 290 ity patterns describe the diel activity patterns of focal species (M. Rowcliffe, 2019) and are typically
 291 calculated to understand fundamental life history traits and behavior such as temporal niche partition-
 292 ing. Occupancy models are hierarchical models commonly fitted to camera trap data because they can
 293 account for imperfect detection (which rarely equals 1) to estimate the conditional probability that a
 294 site is 'occupied' by a species given it was not detected (MacKenzie et al., 2002). Covariates such as
 295 measures of vegetation cover can be included in both the detection and occupancy component models.
 296 These models are relatively complex, and small changes in detection histories (presence or absence of a
 297 species during a discrete time interval), false positives or false negatives can dramatically affect results
 298 (Royle & Link, 2006). We therefore predicted that occupancy estimates obtained using machine learn-
 299 ing generated labels would compare poorly with estimates using expert, manually generated labels.

300

301 The four focal species used for calculating activity patterns and occupancy were African golden cat,
302 chimpanzee, leopard and African forest elephant. These species were chosen because they were the
303 focus of the camera trap survey that generated the out-of-sample test data and because they are
304 conservation priority species in Central Africa. We also initially included western lowland gorilla but
305 we had too few unique captures of this species (only seven of 227 stations having > 5 captures) to fit
306 either activity pattern models or occupancy models.

307

308 ***Thresholding and overall model performance***

309 All three metrics derived from machine learning labels were re-calculated using a threshold approach,
310 where labels were excluded if the model's predicted confidence was below a given threshold. The
311 thresholds tested ranged from 0 (no threshold) to 90%, increasing in 10% intervals. For each of the
312 three ecological metrics, we then re-calculated results using the machine learning labels and compared
313 these with results from the expert labeled dataset using various statistical measures (see later). We also
314 calculated the effect of removing data on sample size, top-one balanced accuracy and top-five accuracy
315 for the overall model, and on four standard measures of model precision and accuracy (precision, re-
316 call, F1 score, and balanced accuracy for each species using the `confusionMatrix` function in the
317 `caret` R package (Kuhn, 2020).

318

319 Estimated species richness from machine learning generated labels and expert labels was compared
320 using linear regression fitted by least squares. Species richness from expert labels was used as the
321 predictor variable and species richness from machine learning labels was used as the response. For each
322 threshold, we evaluated how well species richness from machine learning labels correlated with expert
323 labels by calculating the slope coefficient and variance explained (R^2).

324

Diel activity patterns were calculated for all four focal species using the `fitact` function (200 bootstrap replicates from the model) using the `activity` R package (J. M. Rowcliffe, Kays, Kranstauber, Carbone, & Jansen, 2014; M. Rowcliffe, 2019). For each species and threshold combination, we tested if there was a significant difference in diel activity (proportion of 24 h day active) estimated by machine learning labels and expert labels using the `compareAct` function, expecting no difference using an alpha level of 0.05.

Single season, single species occupancy models were fitted using the `occu` function from the `unmarked` R package (Fiske & Chandler, 2011). Detection histories were collapsed to five-day occasion lengths as a compromise between achieving model stability and ensuring an adequate number of replicates for each site. In the detection component model, we included Elevation (m), Date (first day of the five day occasion length) and Date^2 (to allow for non-linear, seasonal changes in detection) as covariates. In the occupancy component model, Elevation (m), Distance to the Nearest River (m), Distance to the Nearest Road (m) and mean distance to the Nearest Village (m) were included as continuous predictors without interactions. All covariates were mean-centered and scaled by 1 SD to prevent convergence issues. We did not perform model selection and predicted occupancy for the 227 camera stations using the full model. We then compared occupancy predictions ($n = 227$ camera stations) for no threshold (i.e. using all data), and the nine thresholds using linear regression fitted by least squares as described previously for the species richness comparison.

Results

Effect of thresholding on overall model performance

Regardless of the threshold used, top-five accuracy for the overall model predictions on the out-of-sample data were consistently close to or above 95% (Figure 2). To achieve a top-one balanced accuracy of 90% or more for the overall model, a threshold of $\geq 70\%$ confidence was required and >

25% of the data were discarded (Figure 2). With a threshold of 70% confidence (i.e. excluding labeled images below 70% confidence), top-one balanced accuracies for 16 of the 27 classes were > 90% and a further five were > 75% (Table 2). Top-one balanced accuracies for the remaining seven classes ranged from 50% to 70% (Table 2). All other measures of accuracy and precision at all thresholds are in Table S3 and Figure 3 shows the confusion matrix for the out-of-sample data after excluding labels below 70% confidence (see Figure S5 for the confusion matrix of aggregated labels after thresholding).




Figure 2. Relationship between threshold level to accept top label, % of data discarded and overall top-five and top-one balanced accuracy (+/- 95% CI) for predictions on out-of-sample test data.

Table 2. Precision, recall, accuracy, F1 score and prevalence (%) for the 27 classes (Table S1) in the out-of-sample test data after removing labels with a predicted confidence < 70%. Species are sorted from lowest to highest balanced accuracy. For comparison, the precision and recall for the model used by the wildlifeinsights.org web platform are given in brackets. Orange indicates our model performed worse than the WildlifeInsights model for a given species, and purple indicates our model performed better. Note that this comparison should be interpreted with caution. Ideally, we would run the WildlifeInsights model on our out-of-sample test data, but data sharing restrictions prevented this. Where our species or groups could not be compared with an equivalent class on WildlifeInsights this is indicated as no equivalent class (NE). If precision and recall cannot be estimated because of

369 insufficient training and validation data this is indicated as ‘needs more data’ (NMD).

Species	Precision %	Recall %	F1	Prevalence	Balanced Accuracy
Civet_African_Palm	NMD (<i>NMD</i>)	NMD (<i>NMD</i>)	NA	NA	NA
Gorilla	NMD (<i>NMD</i>)	NMD (<i>NMD</i>)	NA	0.4	50
Rail_Nkulengu	0.0 (<i>47.2</i>)	0.0 (<i>48.6</i>)	NA	NA	50
^a Guineafowl_Crested	100 (<i>99.8</i>)	5.3 (<i>91.2</i>)	10	0.1	52.6
Mandrillus	83.9 (<i>96.1</i>)	29 (<i>72.3</i>)	43.1	1.8	64.5
Blank	98.1 (<i>98.3</i>)	40.3 (<i>78.7</i>)	57.1	3.6	70.1
Buffalo_African	97.5 (<i>91.1</i>)	55.7 (<i>73.6</i>)	70.9	1.2	77.8
Bird	11.2 (<i>NE</i>)	60.0 (<i>NE</i>)	18.9	0.1	79.7
Chevrotain_Water	100 (<i>NMD</i>)	67.4 (<i>NMD</i>)	80.6	0.2	83.7
Guineafowl_Black	70.6 (<i>79.6</i>)	72.7 (<i>79.5</i>)	71.6	0.2	86.3
Cat_Golden	96.0 (<i>NMD</i>)	78.0 (<i>NMD</i>)	86.1	1	89
Pangolin	94.1 (<i>NMD</i>)	80.0 (<i>NMD</i>)	86.5	0.1	90
Duiker_Yellow_Backed	97.5 (<i>88.8</i>)	83.8 (<i>72.3</i>)	90.2	2.9	91.9
Human	78.4 (<i>84.8</i>)	87.4 (<i>75.2</i>)	82.6	4	93.2
Chimpanzee	83.5 (<i>87</i>)	88.4 (<i>71.4</i>)	85.9	2.2	94
Monkey	70.7 (<i>NE</i>)	92.0 (<i>NE</i>)	80	2.9	95.4
Mongoose	83.5 (<i>NMD</i>)	91.0 (<i>NMD</i>)	87.1	0.4	95.5
Rat_Giant	68.2 (<i>76</i>)	93.8 (<i>75.8</i>)	78.9	0.1	96.9
^b Duiker_Red	95.9 (<i>95.6</i>)	96.5 (<i>79.6</i>)	96.2	30.8	97.3
Duiker_Blue	90.04 (<i>98.2</i>)	97.0 (<i>65.7</i>)	93.6	17.6	97.4
Hog_Red_River	97.0 (<i>82.7</i>)	95.7 (<i>84.7</i>)	96.3	6.5	97.7
Squirrel	85.9 (<i>98.6</i>)	95.8 (<i>67.6</i>)	90.6	0.9	97.8
Leopard_African	92.8 (<i>85.2</i>)	96.0 (<i>61.4</i>)	94.4	2.2	97.9
Elephant_African	91.9 (<i>94.4</i>)	98.4 (<i>84.2</i>)	95.1	19.3	98.2
Porcupine_Brush_Tailed	93.9 (<i>89.4</i>)	98.9 (<i>42.1</i>)	96.3	0.5	99.4
Genet	95.3 (<i>89.2</i>)	99.3 (<i>65.6</i>)	97.2	0.8	99.6
Mongoose_Black_Footed	92.9 (<i>NMD</i>)	100 (<i>NMD</i>)	96.3	0.1	100

Figure 3. Confusion matrix (% correct labels for each species/group) showing model performance on out of sample test data after excluding labels below a confidence threshold of 70% (each row is normalized independently). Figure S6 shows the confusion matrix with absolute numbers.

378

Species richness

Species richness estimated by machine learning labels and expert labels was strongly correlated at all thresholds used (Figure 4). There was a general tendency for species richness to be underestimated by machine learning as the threshold increased, and the slope of the relationship was close to 1 with no threshold.




Figure 4. Relationship between species richness at each camera station ($n = 227$) predicted by the machine learning model (y-axis) and species richness predicted from expert labels (x-axis) for no threshold and the nine thresholds used after predicting on the out-of-sample test data. The dotted line shows where a 1:1 relationship would fit the data.

Activity patterns

Above a threshold of 70% there was no significant difference between diel activity patterns estimated by machine learning labels and expert labels for all four focal species in the out-of-sample test data (Figure 5; Table S4).




Figure 5. Estimated activity patterns for the four focal species in the out-of-sample test data using machine learning labels (orange; $n = 18078$ observations after excluding labels below 70% confidence) and expert labels (blue; $n = 23868$ observations).

Occupancy models

As expected, occupancy estimates made using machine learning labels were sometimes inconsistent with those made using expert labels, and thresholding had a dramatic impact on inference in some cases (Figure 6). For golden cat and leopard, which are predicted with high accuracy and precision by our machine learning model, occupancy estimates from machine learning labels and expert labels were highly correlated at all thresholds (Figure S8). African elephant occupancy estimates using machine learning labels improved dramatically as the threshold increased, but chimpanzee occupancy estimates from machine learning labels were consistently uncorrelated with those estimated using expert labels (Figure 6).




Figure 6. Relationship between estimated occupancy probability for $n = 227$ camera stations (points) from machine learning (ML) labels (y-axis) and expert labels (x-axis) for the four focal species after discarding labels below a 90% threshold of predicted confidence. Plots for all thresholds tested are shown in Figure S8.

Discussion

Machine learning models have the potential to fully automate labeling of camera trap images without the need for manual validation. This would allow ecologists to rapidly process data and use the outputs (e.g. species labels) directly in ecological analyses, but it has been uncertain how this can be achieved.

In particular, models published to date do not evaluate their predictive performance in an ecological modeling context (Beery et al., 2018; Norouzzadeh et al., 2018; Tabak et al., 2019; Willi et al., 2019). Here, we compared ecological metrics calculated on an out-of-sample test dataset using machine learning labels with the same metrics calculated using expert, manually generated labels. Using our new, high performance species classification model that generalizes to out-of-sample data, we show machine learning labels can be used in a fully automated workflow that removes the need for manual validation prior to conducting ecological analyses.

We used an established architecture for the machine learning model. However, other more recent architectures could yield further increases in performance. The ResNeXt (Xie, Girshick, Dollar, Tu, & He, 2017), the ResNeSt (Zhang et al., 2020) and the EfficientNet (Tan & Le, 2020) families of network architectures are particularly worth exploring in this context. Another avenue of possible further improvement is to use an approach based on a sequence of models. One natural step is to first detect a bounding box for an animal with a localisation model (Beery et al., 2019) and later classify only the content found in that box. Independently, another step can be introduced where a model is trained to first identify an aggregated species class (comprised of species that share similar characteristics; e.g. see Figure S4), and later dedicated models are trained to identify the individual species within these aggregated classes.

We used a relatively small training set (c.300,000 images here vs 3.2 million in (Norouzzadeh et al., 2018) and 8.7M used by (Ahumada et al., 2020)) and a large number of individual classes, yet our model achieved high precision and accuracy even when tested on completely out-of-sample data, which is considered a significant challenge for the field (Beery et al., 2018, 2019). We believe this encouraging result can be explained both by the machine learning approaches used (e.g. the fast.ai framework and image augmentation), and because forest camera traps in the tropics are often deployed in very similar settings, with animals captured at a predictable distance from the camera (usually on a path)

with a general background of green and brown vegetation. This is in contrast to camera trap images from more open habitats, where animals are detected across a wide range of distances and backgrounds (Beery et al., 2018). On the other hand, informational richness in the background of photos taken in forest settings poses a significant challenge to machine learning models as well as human experts, as illustrated in Figure 7.



Figure 7. An image correctly classified as nkulengu rail by our machine learning model but marked as blank by an expert. The bird is visible slightly right of center. The dark beak is pointing left and most of the body is hidden behind branches and leaves. A section of its characteristic red legs is visible between the leaves. The model used features from the beak and head region to identify the bird (see Figure S9).

Thresholding improved the overall performance of the model and its performance for individual species. In our tests we ‘discarded’ labels with low confidence but these data could equally be

classified manually if sample sizes were small. It is important to note, however, that this additional effort to manually label low confidence images would not have improved inference in our example ecological analyses, with the exception of chimpanzee occupancy estimates. Chimpanzee images had the lowest measure of precision among the four focal species, which suggests that true detection events were probably missed frequently, resulting in false negatives (Figure S2). Species that were classified with the highest precision and accuracy were either relatively unique in their shape, color and pattern (e.g. African leopard, the ‘Genet’ group) or were well represented in the training data. We recommend that users of our model in Central Africa use a threshold of 70% to accept labels and have created an offline, multi-platform software tool that can label large batches of images or videos, and display simple maps of species presence/absence and species richness (available at <https://github.com/Appsilon/wildlife-explorer>). The software also outputs the labels in a format that can be used for calculating activity patterns or for use in occupancy models. We do not fully automate these analyses at present (in part because of logistical constraints and delays caused by the COVID19 pandemic), but we anticipate these features will be integrated into future releases.

If machine learning models can fully automate labeling of camera trap images, the first question likely to be posed by most ecologists is ‘Should we?’. Camera trap images contain a wealth of information beyond species identity that would be missed using our model such as behavior, demography, individual phenotype and body condition. A trained model is also limited to detecting and classifying the species in the training dataset, and by definition cannot detect new species. Some machine learning models can already classify behavior (Norouzzadeh et al., 2018) and other future models will achieve this and much more. In our opinion fully automated labels can and should be used in ecological analyses, but only after validation (and re-validation) from an ecological perspective, and to answer clearly defined questions. Each use-case will also differ in the benefits that can be gained from fully automated analysis. A conservation manager with tens of thousands of images collected on a rolling basis might

accept a trade-off between increased speed of data analysis and having to discard images with uncertain labels, but a scientist testing hypotheses for peer-reviewed publication might prefer to view all of the images manually. We recommend that in all cases models should be validated on a continual basis using sub-sampled data to detect potentially new or hidden biases. Model accuracy could change if field protocols or environmental conditions change in unexpected ways (e.g. heavy snowfall in temperate zones). However, during model evaluation we found that expert labels in the training and validation data were also never themselves ‘perfect’, and perhaps high performance machine learning models offer a more consistent means of analyzing camera trap data than manual labeling because biases are predictable and can be quantified explicitly.

Camera traps are commonly used worldwide by conservation practitioners whose normal scope of work might not allow sufficient time for the handling, processing, and analyzing of large quantities of digital data. The authors personally know of several large camera trap databases that have not been analyzed years after data collection ended, often because of a lack of resources or technical expertise. New web-based platforms for ecological data are seeking to address this problem by allowing users to upload data to the cloud where it is stored and analyzed using machine learning models (Aide et al., 2013; Ahumada et al., 2020), but a lack of fast internet access can be a barrier to using such platforms and our offline application can fill this important gap. The next generation of camera traps will also have embedded machine learning models following the current rise in edge-computing technology. Together, edge and cloud computing will open the door to national and international real-time ecological forecasting at unprecedented spatial and temporal scales. We anticipate that the model, software and validation workflow presented here could revolutionize how camera trap data are processed and analyzed, and conclude that high performance machine learning models can be used for fully automated labeling of camera trap data for ecological analyses.

Acknowledgments

RW was funded by the EU 11eme FED ECOFAC6 program grant to the National Parks Agency of Gabon. Appsilon Data Science funded the machine learning model and software development costs. Cloud computing costs were funded by a Google Cloud Education Grant awarded to KA. Camera trap data from co-authors KB and CKO were kindly made available by the Tropical Ecology Assessment and Monitoring Network (now <https://wildlifeinsights.org>).

References

- Ahumada, J. A., Fegraus, E., Birch, T., Flores, N., Kays, R., O'Brien, T. G., ... Dancer, A. (2020). Wildlife Insights: A Platform to Maximize the Potential of Camera Trap and Other Passive Sensor Wildlife Data for the Planet. *Environmental Conservation*, 47(1), 1–6.
doi:10.1017/S0376892919000298
- Aide, T. M., Corrada-Bravo, C., Campos-Cerqueira, M., Milan, C., Vega, G., & Alvarez, R. (2013). Real-time bioacoustics monitoring and automated species identification. *PeerJ*, 1, e103.
doi:10.7717/peerj.103
- Bahaa-el-din, L., & Cusack, J. J. (2018). Camera trapping in Africa: Paving the way for ease of use and consistency. *African Journal of Ecology*, 56(4), 690–693. doi:10.1111/aje.12581
- Bahaa-el-din, L., Henschel, P., Aba'a, R., Abernethy, K., Bohm, T., Bout, N., ... Hunter, L. (2013). Notes on the distribution and status of small carnivores in Gabon, 48, 11.
- Beery, S., Morris, D., Yang, S., Simon, M., Norouzzadeh, A., & Joshi, N. (2019). Efficient Pipeline for Automating Species ID in new Camera Trap Projects. *Biodiversity Information Science and Standards*, 3, e37222. doi:10.3897/biss.3.37222
- Beery, S., Van Horn, G., & Perona, P. (2018). Recognition in Terra Incognita. In *Proceedings of the European Conference on Computer Vision (ECCV)* (pp. 456–473). Retrieved from https://openaccess.thecvf.com/content_ECCV_2018/html/Beery_Recognition_in_Terra_ECCV_2018_paper.html
- Bessone, M., Kühl, H. S., Hohmann, G., Herbinger, I., N'Goran, K. P., Asanzi, P., ... Fruth, B. (2020). Drawn out of the shadows: Surveying secretive forest species with camera trap distance

- sampling. *Journal of Applied Ecology*, 57(5), 963–974. doi:10.1111/1365-2664.13602
- Cardoso, A. W., Malhi, Y., Oliveras, I., Lehmann, D., Ndong, J. E., Dimoto, E., ... Abernethy, K. (2020). The Role of Forest Elephants in Shaping Tropical Forest–Savanna Coexistence. *Ecosystems*, 23(3), 602–616. doi:10.1007/s10021-019-00424-3
- Dietze, M. C., Fox, A., Beck-Johnson, L. M., Betancourt, J. L., Hooten, M. B., Jarnevich, C. S., ... White, E. P. (2018). Iterative near-term ecological forecasting: Needs, opportunities, and challenges. *Proceedings of the National Academy of Sciences*, 115(7), 1424–1432. doi:10.1073/pnas.1710231115
- Farley, S. S., Dawson, A., Goring, S. J., & Williams, J. W. (2018). Situating Ecology as a Big-Data Science: Current Advances, Challenges, and Solutions. *BioScience*, 68(8), 563–576. doi:10.1093/biosci/biy068
- Fiske, I., & Chandler, R. (2011). unmarked: An R Package for Fitting Hierarchical Models of Wildlife Occurrence and Abundance. *Journal of Statistical Software*, 43(10), 1–23.
- Glover-Kapfer, P., Soto Navarro, C. A., & Wearn, O. R. (2019). Camera-trapping version 3.0: current constraints and future priorities for development. *Remote Sensing in Ecology and Conservation*, 5(3), 209–223. doi:10.1002/rse2.106
- He, K., Zhang, X., Ren, S., & Sun, J. (2016). Identity Mappings in Deep Residual Networks. In B. Leibe, J. Matas, N. Sebe, & M. Welling (Eds.), *Computer Vision – ECCV 2016* (pp. 630–645). Cham: Springer International Publishing. doi:10.1007/978-3-319-46493-0_38
- Kuhn, M. (2020). caret: Classification and Regression Training. *R Package Version 6.0-86*. Retrieved from <https://CRAN.R-project.org/package=caret>
- MacKenzie, D. I., Nichols, J. D., Lachman, G. B., Droege, S., Royle, J. A., & Langtimm, C. A. (2002). Estimating Site Occupancy Rates When Detection Probabilities Are Less Than One. *Ecology*, 83(8), 2248–2255. doi:10.1890/0012-9658(2002)083[2248:ESORWD]2.0.CO;2
- Norouzzadeh, M. S., Nguyen, A., Kosmala, M., Swanson, A., Palmer, M. S., Packer, C., & Clune, J. (2018). Automatically identifying, counting, and describing wild animals in camera-trap images with deep learning. *Proceedings of the National Academy of Sciences*, 115(25), E5716–E5725. doi:10.1073/pnas.1719367115
- O’Brien, T. G., Ahumada, J., Akampurila, E., Beaudrot, L., Boekee, K., Brncic, T., ... Strindberg, S.

- (2020). Camera trapping reveals trends in forest duiker populations in African National Parks. *Remote Sensing in Ecology and Conservation*, 6(2), 168–180. doi:10.1002/rse2.132
- Rowcliffe, J. M., Kays, R., Kranstauber, B., Carbone, C., & Jansen, P. A. (2014). Quantifying levels of animal activity using camera trap data. *Methods in Ecology and Evolution*, 5(11), 1170–1179. doi:10.1111/2041-210X.12278
- Rowcliffe, M. (2019). activity: Animal Activity Statistics. *R Package v 1.3*. Retrieved from <https://CRAN.R-project.org/package=activity>
- Royle, J. A., & Link, W. A. (2006). Generalized Site Occupancy Models Allowing for False Positive and False Negative Errors. *Ecology*, 87(4), 835–841. doi:10.1890/0012-9658(2006)87[835:GSOMAF]2.0.CO;2
- Smith, L. N. (2018). A disciplined approach to neural network hyper-parameters: Part 1 -- learning rate, batch size, momentum, and weight decay. *ArXiv:1803.09820 [Cs, Stat]*. Retrieved from <http://arxiv.org/abs/1803.09820>
- Swanson, A., Kosmala, M., Lintott, C., Simpson, R., Smith, A., & Packer, C. (2015). Snapshot Serengeti, high-frequency annotated camera trap images of 40 mammalian species in an African savanna. *Scientific Data*, 2(1), 150026. doi:10.1038/sdata.2015.26
- Tabak, M. A., Norouzzadeh, M. S., Wolfson, D. W., Sweeney, S. J., Vercauteren, K. C., Snow, N. P., ... Miller, R. S. (2019). Machine learning to classify animal species in camera trap images: Applications in ecology. *Methods in Ecology and Evolution*, 10(4), 585–590. doi:10.1111/2041-210X.13120
- Tan, M., & Le, Q. V. (2020). EfficientNet: Rethinking Model Scaling for Convolutional Neural Networks. *ArXiv:1905.11946 [Cs, Stat]*. Retrieved from <http://arxiv.org/abs/1905.11946>
- Wei, W., Luo, G., Ran, J., & Li, J. (2020). Zilong: A tool to identify empty images in camera-trap data. *Ecological Informatics*, 55, 101021. doi:10.1016/j.ecoinf.2019.101021
- Willi, M., Pitman, R. T., Cardoso, A. W., Locke, C., Swanson, A., Boyer, A., ... Fortson, L. (2019). Identifying animal species in camera trap images using deep learning and citizen science. *Methods in Ecology and Evolution*, 10(1), 80–91. doi:10.1111/2041-210X.13099
- Xie, S., Girshick, R., Dollar, P., Tu, Z., & He, K. (2017). Aggregated Residual Transformations for Deep Neural Networks (pp. 1492–1500). Presented at the Proceedings of the IEEE Conference

on Computer Vision and Pattern Recognition. Retrieved from

https://openaccess.thecvf.com/content_cvpr_2017/html/Xie_Aggregated_Residual_Transformations_CVPR_2017_paper.html

Zhang, H., Wu, C., Zhang, Z., Zhu, Y., Zhang, Z., Lin, H., ... Smola, A. (2020). ResNeSt: Split-

Attention Networks. *ArXiv:2004.08955 [Cs]*. Retrieved from <http://arxiv.org/abs/2004.08955>

Supplementary Information for:

High performance machine learning models can fully automate labeling of camera trap images for ecological analyses

Robin Whytock^{1,2*†}, Jędrzej Świeżewski^{3†}, Joeri A. Zwerts⁴, Tadeusz Bara-Słupski⁴, Aurélie Flore Koumba Pambo², Marek Rogala³, Laila Bahaa-el-din⁵, Kelly Boeke^{6,7}, Stephanie Brittain^{8,9}, Anabelle W. Cardoso¹⁰, Philipp Henschel^{11,12}, David Lehmann^{2,1}, Brice Momboua², Cisquet Hector Ror Kiebou Opepa¹³, Christopher Orbell^{11,1}, Ross T. Pitman¹¹, Hugh S. Robinson^{11,14}, Katharine A. Abernethy^{1,12}

† These authors contributed equally to the manuscript.

***Corresponding author:** Robin C. Whytock, robbie.whytock1@stir.ac.uk

Paste manuscript title here

¹ Faculty of Natural Sciences, University of Stirling, FK9 4LA, UK

² Agence Nationale des Parcs Nationaux, Libreville, Gabon

³ Appsilon AI for Good, Warsaw, Poland

⁴ Utrecht University, Heidelberglaan 8, 3584 CS Utrecht, The Netherlands

⁵ School of Life Sciences, University of KwaZulu-Natal, South Africa

⁶ Program for the Sustainable Management of Natural Resources, South West Region, Buea, Cameroon

⁷ Center for Tropical Forest Science, Smithsonian Tropical Research Institute, Balboa, Ancon, Republic of Panama

⁸ The Interdisciplinary Centre for Conservation Science, Department of Zoology, University of Oxford, 11a Mansfield Road, Oxford, OX1 3SZ

⁹ The Institute of Zoology, Zoological Society of London, Outer Circle, London NW1 4RY

¹⁰ Yale University, New Haven, CT 06520, USA

¹¹ Panthera, 8 West 40th Street, 18th Floor, New York, NY 10018, USA

¹² Institut de Recherche en Ecologie Tropicale, CENAREST, BP 842 Libreville, Gabon

¹³ Wildlife Conservation Society, Republic of the Congo

¹⁴ Wildlife Biology Program, W.A. Franke College of Forestry and Conservation, University of Montana, Natural Sciences Room 205, Missoula, Montana, 59812, USA

This PDF file includes:

Tables S1 to S4

Figures S1 to S9

552 **Table S1.** Species taxonomy, label descriptions and justification for species/class groups

Species class	Scientific name	Justification
Civet_African_Palm	<i>Nandinia binotata</i>	-
Gorilla	<i>Gorilla gorilla gorilla</i>	-
Rail_Nkulengu	<i>Himantornis haematopus</i>	-
Guineafowl_Crested	<i>Guttera pucherani</i>	-
Mandrillus	<i>Mandrillus sphinx</i>	-
Blank		No animal or human
Buffalo_African	<i>Cyncerus cafer nanus</i>	-
Bird		Any other bird
Chevrotain_Water	<i>Hymenoschus aquaticus</i>	-
Guineafowl_Black	<i>Agelastes niger</i>	-
Cat_Golden	<i>Caracal aurata</i>	-
Pangolin		Identifies any pangolin but trained mainly on <i>Smutsia gigantea</i>
Duiker_Yellow_Backed	<i>Cephalophus silvicultor</i>	-
Human	<i>Homo sapiens</i>	-
Chimpanzee	<i>Pan troglodytes</i>	-
Monkey		Any guenon, colobus or mangabey
Mongoose		Marsh mongoose <i>Atilax paludinosus</i> or long-nosed mongoose <i>Herpestes naso</i>
Rat_Giant	<i>Cricetomys emini</i>	-
Duiker_Red	<i>Cephalophus</i> sp.	Any of the red <i>Cephalophus</i> sp. duikers
Duiker_Blue	<i>Philantomba monticola</i>	-
Hog_Red_River	<i>Potamochoerus porcus</i>	-
Squirrel		Any squirrel but most training data are <i>Protoxerus stangeri</i>
Leopard_African	<i>Panthera pardus</i>	-
Elephant_African	<i>Loxodonta cyclotis</i>	-
Porcupine_Brush_Tailed	<i>Atherurus africanus</i>	-
Genet	<i>Genetta</i> sp.	Most training data are <i>Genetta servalina</i>
Mongoose_Black_Footed	<i>Bdeogale nigripes</i>	-

553

554

Table S2. Measures of precision, accuracy and prevalence (%) for the 27 species/groups (see Table S1 for further details on species groups) in the out-of-sample test data.

Species class	Precision	Recall	F1	Prevalence	Balanced Accuracy
Bird	6.4	35.6	10.9	0.3	67
Blank	96.3	31.2	47.1	13.1	65.5
Buffalo_African	90.6	43.3	58.6	1.6	71.6
Cat_Golden	86.5	68.1	76.2	1.1	84
Chevrotain_Water	96.2	37.3	53.8	0.6	68.7
Chimpanzee	65.3	74.5	69.6	2.4	86.8
Civet_African_Palm	9.1	100	16.7	< 0.1	100
Duiker_Blue	73.1	91.3	81.2	14.9	92.7
Duiker_Red	87.5	91.8	89.6	26	93.6
Duiker_Yellow_Backed	88.7	71.2	79	2.9	85.5
Elephant_African	83	95	88.6	15.9	95.6
Genet	87.3	93.8	90.4	0.7	96.8
Gorilla	50	15.7	23.9	0.8	57.8
Guineafowl_Black	22.1	60.4	32.3	0.2	80
Guineafowl_Crested	100	16.1	27.8	0.1	58.1
Hog_Red_River	89.9	89.6	89.8	5.9	94.5
Human	51.5	79.9	62.6	3.6	88.6
Leopard_African	87	85.9	86.4	2	92.8
Mandrillus	73.5	26.1	38.6	2.7	62.9
Mongoose	48.2	80.4	60.3	0.4	90
Mongoose_Black_Footed	72.5	64.4	68.2	0.2	82.2
Monkey	59.9	81	68.9	2.9	89.7
Pangolin	76.7	62.2	68.7	0.2	81.1
Porcupine_Brush_Tailed	86.6	83.3	84.9	0.6	91.6
Rail_Nkulengu	0	0	0	0	50
Rat_Giant	39	88.5	54.1	0.1	94.2
Squirrel	59.9	78.2	67.8	1	88.8

559 **Table S3.** Precision, recall, F1 score and prevalence (%) for the 27 species/groups (see Table S1 for
560 further details on species groups) in the out-of-sample test data at all thresholds used (10 – 90% confi-
561 dence).

Species class	Threshold	Precision	Recall	F1	Prevalence	Balanced Accuracy
Bird	10	0.064	0.356	0.109	0.003	0.670
Blank	10	0.963	0.312	0.471	0.131	0.655
Buffalo_African	10	0.906	0.433	0.586	0.016	0.716
Cat_Golden	10	0.865	0.681	0.762	0.011	0.840
Chevrotain_Water	10	0.962	0.373	0.538	0.006	0.687
Chimpanzee	10	0.653	0.745	0.696	0.024	0.868
Civet_African_Palm	10	0.091	1.000	0.167	0.000	1.000
Duiker_Blue	10	0.731	0.913	0.812	0.149	0.927
Duiker_Red	10	0.875	0.918	0.896	0.260	0.936
Duiker_Yellow_Backed	10	0.887	0.712	0.790	0.029	0.855
Elephant_African	10	0.830	0.950	0.886	0.159	0.956
Genet	10	0.873	0.938	0.904	0.007	0.968
Gorilla	10	0.500	0.157	0.239	0.008	0.578
Guineafowl_Black	10	0.221	0.604	0.323	0.002	0.800
Guineafowl_Crested	10	1.000	0.161	0.278	0.001	0.581
Hog_Red_River	10	0.899	0.896	0.898	0.059	0.945
Human	10	0.515	0.799	0.626	0.036	0.886
Leopard_African	10	0.870	0.859	0.864	0.020	0.928
Mandrillus	10	0.735	0.261	0.386	0.027	0.629
Mongoose	10	0.482	0.804	0.603	0.004	0.900
Mongoose_Black_Footed	10	0.725	0.644	0.682	0.002	0.822
Monkey	10	0.599	0.810	0.689	0.029	0.897
Pangolin	10	0.767	0.622	0.687	0.002	0.811
Porcupine_Brush_Tailed	10	0.866	0.833	0.849	0.006	0.916
Rail_Nkulengu	10	0.000	0.000	NA	0.000	0.500
Rat_Giant	10	0.390	0.885	0.541	0.001	0.942
Squirrel	10	0.599	0.782	0.678	0.010	0.888
Bird	20	0.063	0.352	0.107	0.003	0.668
Blank	20	0.966	0.316	0.476	0.128	0.657
Buffalo_African	20	0.906	0.433	0.586	0.016	0.716
Cat_Golden	20	0.865	0.688	0.767	0.011	0.844
Chevrotain_Water	20	0.961	0.380	0.544	0.005	0.690
Chimpanzee	20	0.659	0.745	0.699	0.024	0.868
Civet_African_Palm	20	0.111	1.000	0.200	0.000	1.000
Duiker_Blue	20	0.734	0.915	0.814	0.149	0.928
Duiker_Red	20	0.877	0.919	0.898	0.261	0.937
Duiker_Yellow_Backed	20	0.888	0.714	0.792	0.029	0.856
Elephant_African	20	0.830	0.951	0.886	0.160	0.957
Genet	20	0.893	0.938	0.915	0.007	0.969
Gorilla	20	0.482	0.148	0.227	0.008	0.574
Guineafowl_Black	20	0.225	0.604	0.328	0.002	0.800
Guineafowl_Crested	20	1.000	0.161	0.278	0.001	0.581
Hog_Red_River	20	0.900	0.896	0.898	0.059	0.945
Human	20	0.524	0.800	0.633	0.036	0.886
Leopard_African	20	0.872	0.861	0.866	0.020	0.929
Mandrillus	20	0.735	0.263	0.388	0.027	0.630
Mongoose	20	0.516	0.804	0.628	0.004	0.900

Species class	Threshold	Precision	Recall	F1	Prevalence	Balanced Accuracy
Mongoose_Black_Footed	20	0.763	0.674	0.716	0.002	0.837
Monkey	20	0.601	0.814	0.691	0.029	0.899
Pangolin	20	0.821	0.622	0.708	0.002	0.811
Porcupine_Brush_Tailed	20	0.866	0.853	0.859	0.005	0.926
Rail_Nkulengu	20	0.000	0.000	NA	0.000	0.500
Rat_Giant	20	0.386	0.880	0.537	0.001	0.939
Squirrel	20	0.608	0.782	0.684	0.010	0.888
Bird	30	0.065	0.377	0.111	0.003	0.681
Blank	30	0.968	0.329	0.491	0.112	0.664
Buffalo_African	30	0.911	0.446	0.599	0.016	0.722
Cat_Golden	30	0.885	0.708	0.787	0.011	0.853
Chevrotain_Water	30	0.980	0.403	0.571	0.005	0.702
Chimpanzee	30	0.683	0.762	0.720	0.024	0.876
Civet_African_Palm	30	0.200	1.000	0.333	0.000	1.000
Duiker_Blue	30	0.754	0.921	0.829	0.152	0.934
Duiker_Red	30	0.888	0.922	0.905	0.268	0.940
Duiker_Yellow_Backed	30	0.909	0.726	0.807	0.029	0.862
Elephant_African	30	0.840	0.955	0.894	0.164	0.960
Genet	30	0.904	0.950	0.926	0.007	0.974
Gorilla	30	0.519	0.153	0.237	0.008	0.576
Guineafowl_Black	30	0.283	0.604	0.386	0.002	0.800
Guineafowl_Crested	30	1.000	0.161	0.278	0.001	0.581
Hog_Red_River	30	0.911	0.902	0.907	0.061	0.948
Human	30	0.551	0.802	0.653	0.037	0.888
Leopard_African	30	0.887	0.872	0.879	0.020	0.935
Mandrillus	30	0.764	0.266	0.395	0.026	0.632
Mongoose	30	0.599	0.828	0.695	0.004	0.913
Mongoose_Black_Footed	30	0.763	0.829	0.795	0.002	0.914
Monkey	30	0.612	0.831	0.705	0.029	0.908
Pangolin	30	0.815	0.667	0.733	0.001	0.833
Porcupine_Brush_Tailed	30	0.858	0.858	0.858	0.005	0.929
Rail_Nkulengu	30	0.000	0.000	NA	0.000	0.500
Rat_Giant	30	0.449	0.917	0.603	0.001	0.958
Squirrel	30	0.645	0.801	0.715	0.010	0.898
Bird	40	0.078	0.423	0.132	0.002	0.706
Blank	40	0.976	0.352	0.518	0.090	0.676
Buffalo_African	40	0.929	0.473	0.627	0.015	0.736
Cat_Golden	40	0.905	0.722	0.803	0.011	0.860
Chevrotain_Water	40	0.977	0.452	0.618	0.004	0.726
Chimpanzee	40	0.725	0.788	0.755	0.024	0.890
Civet_African_Palm	40	0.200	1.000	0.333	0.000	1.000
Duiker_Blue	40	0.791	0.934	0.857	0.157	0.944
Duiker_Red	40	0.904	0.930	0.917	0.279	0.946
Duiker_Yellow_Backed	40	0.924	0.751	0.829	0.030	0.875
Elephant_African	40	0.860	0.962	0.908	0.170	0.965
Genet	40	0.921	0.950	0.935	0.007	0.975
Gorilla	40	0.528	0.119	0.195	0.007	0.559
Guineafowl_Black	40	0.375	0.600	0.462	0.002	0.799
Guineafowl_Crested	40	1.000	0.161	0.278	0.001	0.581
Hog_Red_River	40	0.930	0.911	0.920	0.063	0.953
Human	40	0.593	0.811	0.685	0.038	0.895
Leopard_African	40	0.897	0.903	0.900	0.020	0.951
Mandrillus	40	0.795	0.288	0.422	0.025	0.643
Mongoose	40	0.704	0.835	0.764	0.004	0.917

Species class	Threshold	Precision	Recall	F1	Prevalence	Balanced Accuracy
Mongoose_Black_Footed	40	0.800	0.903	0.848	0.001	0.951
Monkey	40	0.632	0.856	0.727	0.029	0.920
Pangolin	40	0.880	0.733	0.800	0.001	0.867
Porcupine_Brush_Tailed	40	0.888	0.904	0.896	0.005	0.951
Rail_Nkulengu	40	0.000	0.000	NA	0.000	0.500
Rat_Giant	40	0.537	0.917	0.677	0.001	0.958
Squirrel	40	0.715	0.843	0.774	0.010	0.920
Bird	50	0.084	0.450	0.142	0.002	0.720
Blank	50	0.981	0.378	0.546	0.072	0.689
Buffalo_African	50	0.938	0.503	0.655	0.014	0.751
Cat_Golden	50	0.923	0.741	0.822	0.011	0.870
Chevrotain_Water	50	0.974	0.500	0.661	0.004	0.750
Chimpanzee	50	0.753	0.824	0.787	0.024	0.909
Civet_African_Palm	50	0.500	1.000	0.667	0.000	1.000
Duiker_Blue	50	0.825	0.945	0.881	0.162	0.953
Duiker_Red	50	0.920	0.939	0.930	0.288	0.953
Duiker_Yellow_Backed	50	0.942	0.773	0.849	0.029	0.886
Elephant_African	50	0.879	0.968	0.921	0.177	0.969
Genet	50	0.932	0.962	0.947	0.008	0.981
Gorilla	50	0.583	0.107	0.181	0.006	0.553
Guineafowl_Black	50	0.492	0.638	0.556	0.002	0.818
Guineafowl_Crested	50	1.000	0.138	0.242	0.001	0.569
Hog_Red_River	50	0.946	0.922	0.934	0.064	0.959
Human	50	0.646	0.827	0.725	0.039	0.904
Leopard_African	50	0.902	0.921	0.912	0.021	0.959
Mandrillus	50	0.816	0.292	0.430	0.023	0.645
Mongoose	50	0.775	0.868	0.819	0.004	0.934
Mongoose_Black_Footed	50	0.903	0.933	0.918	0.001	0.967
Monkey	50	0.654	0.879	0.750	0.029	0.932
Pangolin	50	0.913	0.724	0.808	0.001	0.862
Porcupine_Brush_Tailed	50	0.902	0.953	0.927	0.005	0.976
Rail_Nkulengu	50	0.000	0.000	NA	0.000	0.500
Rat_Giant	50	0.625	0.909	0.741	0.001	0.954
Squirrel	50	0.758	0.888	0.818	0.010	0.943
Bird	60	0.103	0.552	0.174	0.002	0.772
Blank	60	0.985	0.399	0.568	0.052	0.699
Buffalo_African	60	0.957	0.545	0.694	0.013	0.772
Cat_Golden	60	0.935	0.768	0.844	0.011	0.884
Chevrotain_Water	60	0.970	0.582	0.727	0.003	0.791
Chimpanzee	60	0.809	0.848	0.828	0.023	0.921
Civet_African_Palm	60	0.500	1.000	0.667	0.000	1.000
Duiker_Blue	60	0.870	0.960	0.912	0.169	0.965
Duiker_Red	60	0.943	0.952	0.948	0.298	0.964
Duiker_Yellow_Backed	60	0.962	0.802	0.875	0.029	0.901
Elephant_African	60	0.897	0.975	0.934	0.186	0.975
Genet	60	0.943	0.980	0.961	0.008	0.990
Gorilla	60	0.583	0.065	0.118	0.006	0.533
Guineafowl_Black	60	0.614	0.711	0.659	0.002	0.855
Guineafowl_Crested	60	1.000	0.087	0.160	0.001	0.543
Hog_Red_River	60	0.962	0.942	0.952	0.065	0.970
Human	60	0.714	0.850	0.777	0.039	0.918
Leopard_African	60	0.916	0.945	0.930	0.022	0.971
Mandrillus	60	0.809	0.276	0.411	0.021	0.637
Mongoose	60	0.794	0.895	0.842	0.004	0.947

Species class	Threshold	Precision	Recall	F1	Prevalence	Balanced Accuracy
Mongoose_Black_Footed	60	0.900	1.000	0.947	0.001	1.000
Monkey	60	0.680	0.898	0.774	0.029	0.942
Pangolin	60	0.905	0.731	0.809	0.001	0.865
Porcupine_Brush_Tailed	60	0.923	0.970	0.946	0.005	0.985
Rail_Nkulengu	60	0.000	0.000	NA	0.000	0.500
Rat_Giant	60	0.633	0.950	0.760	0.001	0.975
Squirrel	60	0.810	0.938	0.869	0.009	0.968
Bird	70	0.112	0.600	0.189	0.001	0.797
Blank	70	0.981	0.403	0.571	0.036	0.701
Buffalo_African	70	0.975	0.557	0.709	0.012	0.778
Cat_Golden	70	0.960	0.780	0.861	0.010	0.890
Chevrotain_Water	70	1.000	0.674	0.806	0.002	0.837
Chimpanzee	70	0.835	0.884	0.859	0.022	0.940
Civet_African_Palm	70	NA	NA	NA	0.000	NA
Duiker_Blue	70	0.904	0.970	0.936	0.176	0.974
Duiker_Red	70	0.959	0.965	0.962	0.308	0.973
Duiker_Yellow_Backed	70	0.975	0.838	0.902	0.029	0.919
Elephant_African	70	0.919	0.984	0.951	0.193	0.982
Genet	70	0.953	0.993	0.972	0.008	0.996
Gorilla	70	0.000	0.000	NA	0.004	0.500
Guineafowl_Black	70	0.706	0.727	0.716	0.002	0.863
Guineafowl_Crested	70	1.000	0.053	0.100	0.001	0.526
Hog_Red_River	70	0.970	0.957	0.963	0.065	0.977
Human	70	0.784	0.874	0.826	0.040	0.932
Leopard_African	70	0.928	0.960	0.944	0.022	0.979
Mandrillus	70	0.839	0.290	0.431	0.018	0.645
Mongoose	70	0.835	0.910	0.871	0.004	0.955
Mongoose_Black_Footed	70	0.929	1.000	0.963	0.001	1.000
Monkey	70	0.707	0.920	0.800	0.029	0.954
Pangolin	70	0.941	0.800	0.865	0.001	0.900
Porcupine_Brush_Tailed	70	0.939	0.989	0.963	0.005	0.994
Rail_Nkulengu	70	0.000	0.000	NA	0.000	0.500
Rat_Giant	70	0.682	0.938	0.789	0.001	0.969
Squirrel	70	0.859	0.958	0.906	0.009	0.978
Bird	80	0.151	0.786	0.253	0.001	0.891
Blank	80	0.986	0.363	0.530	0.023	0.681
Buffalo_African	80	1.000	0.596	0.747	0.010	0.798
Cat_Golden	80	0.962	0.839	0.896	0.009	0.919
Chevrotain_Water	80	1.000	0.750	0.857	0.002	0.875
Chimpanzee	80	0.875	0.919	0.897	0.019	0.958
Civet_African_Palm	80	NA	NA	NA	0.000	NA
Duiker_Blue	80	0.932	0.981	0.956	0.183	0.982
Duiker_Red	80	0.973	0.975	0.974	0.315	0.981
Duiker_Yellow_Backed	80	0.988	0.885	0.934	0.028	0.942
Elephant_African	80	0.940	0.991	0.965	0.203	0.987
Genet	80	0.949	1.000	0.974	0.008	1.000
Gorilla	80	0.000	0.000	NA	0.003	0.500
Guineafowl_Black	80	0.769	0.690	0.727	0.002	0.845
Guineafowl_Crested	80	1.000	0.067	0.125	0.001	0.533
Hog_Red_River	80	0.980	0.971	0.975	0.065	0.985
Human	80	0.853	0.892	0.872	0.040	0.943
Leopard_African	80	0.952	0.974	0.963	0.023	0.987
Mandrillus	80	0.888	0.305	0.454	0.014	0.652
Mongoose	80	0.829	0.944	0.883	0.004	0.972

Species class	Threshold	Precision	Recall	F1	Prevalence	Balanced Accuracy
Mongoose_Black_Footed	80	1.000	1.000	1.000	0.001	1.000
Monkey	80	0.756	0.928	0.833	0.029	0.959
Pangolin	80	1.000	0.824	0.903	0.001	0.912
Porcupine_Brush_Tailed	80	0.935	0.989	0.961	0.005	0.994
Rail_Nkulengu	80	NA	NA	NA	0.000	NA
Rat_Giant	80	0.737	0.933	0.824	0.001	0.967
Squirrel	80	0.879	0.979	0.926	0.008	0.989
Bird	90	0.220	0.900	0.353	0.001	0.949
Blank	90	1.000	0.320	0.484	0.011	0.660
Buffalo_African	90	1.000	0.647	0.785	0.008	0.823
Cat_Golden	90	0.980	0.897	0.937	0.007	0.949
Chevrotain_Water	90	1.000	0.833	0.909	0.001	0.917
Chimpanzee	90	0.914	0.922	0.918	0.016	0.960
Civet_African_Palm	90	NA	NA	NA	0.000	NA
Duiker_Blue	90	0.957	0.990	0.973	0.196	0.989
Duiker_Red	90	0.984	0.984	0.984	0.317	0.988
Duiker_Yellow_Backed	90	0.994	0.912	0.952	0.026	0.956
Elephant_African	90	0.961	0.994	0.977	0.218	0.991
Genet	90	0.957	1.000	0.978	0.007	1.000
Gorilla	90	NA	0.000	NA	0.002	0.500
Guineafowl_Black	90	0.864	0.826	0.844	0.002	0.913
Guineafowl_Crested	90	NA	0.000	NA	0.001	0.500
Hog_Red_River	90	0.986	0.982	0.984	0.063	0.990
Human	90	0.918	0.923	0.920	0.040	0.960
Leopard_African	90	0.973	0.979	0.976	0.025	0.989
Mandrillus	90	0.900	0.298	0.448	0.010	0.649
Mongoose	90	0.855	0.967	0.908	0.004	0.983
Mongoose_Black_Footed	90	1.000	1.000	1.000	0.001	1.000
Monkey	90	0.811	0.952	0.876	0.028	0.973
Pangolin	90	1.000	0.813	0.897	0.001	0.906
Porcupine_Brush_Tailed	90	0.934	1.000	0.966	0.005	1.000
Rail_Nkulengu	90	NA	NA	NA	0.000	NA
Rat_Giant	90	0.846	0.917	0.880	0.001	0.958
Squirrel	90	0.938	0.981	0.959	0.007	0.990

562

563


Table S4. Difference in proportion of day (24 h) active for each species and threshold combination showing standard error (SE), Wald test statistic (W) and p value (p).

Species	Threshold	Difference	SE	W	p
Elephant_African	0.00	0.08	0.03	5.97	0.01
Leopard_African	0.00	0.16	0.08	3.99	0.05
Cat_Golden	0.00	0.06	0.09	0.42	0.52
Chimpanzee	0.00	0.05	0.03	2.92	0.09
Elephant_African	0.10	0.08	0.04	5.47	0.02
Leopard_African	0.10	0.16	0.08	4.26	0.04
Cat_Golden	0.10	0.06	0.09	0.45	0.50
Chimpanzee	0.10	0.05	0.03	2.71	0.10
Elephant_African	0.20	0.08	0.03	6.04	0.01
Leopard_African	0.20	0.16	0.08	4.15	0.04
Cat_Golden	0.20	0.06	0.09	0.38	0.54
Chimpanzee	0.20	0.04	0.03	2.33	0.13
Elephant_African	0.30	0.08	0.03	6.44	0.01
Leopard_African	0.30	0.16	0.08	4.53	0.03
Cat_Golden	0.30	0.04	0.09	0.20	0.66
Chimpanzee	0.30	0.04	0.03	2.07	0.15
Elephant_African	0.40	0.07	0.03	4.15	0.04
Leopard_African	0.40	0.16	0.08	4.08	0.04
Cat_Golden	0.40	0.08	0.10	0.67	0.41
Chimpanzee	0.40	0.04	0.03	1.91	0.17
Elephant_African	0.50	0.05	0.03	2.64	0.10
Leopard_African	0.50	0.17	0.08	4.48	0.03
Cat_Golden	0.50	0.09	0.09	0.96	0.33
Chimpanzee	0.50	0.04	0.03	1.77	0.18
Elephant_African	0.60	0.04	0.03	1.64	0.20
Leopard_African	0.60	0.15	0.08	3.53	0.06
Cat_Golden	0.60	0.06	0.10	0.44	0.50
Chimpanzee	0.60	0.04	0.03	1.52	0.22
Elephant_African	0.70	0.04	0.03	1.21	0.27
Leopard_African	0.70	0.15	0.08	3.08	0.08
Cat_Golden	0.70	0.10	0.10	1.00	0.32
Chimpanzee	0.70	0.03	0.03	1.36	0.24
Elephant_African	0.80	0.03	0.03	0.69	0.41
Leopard_African	0.80	0.16	0.08	3.86	0.05
Cat_Golden	0.80	0.15	0.10	2.14	0.14
Chimpanzee	0.80	0.03	0.03	0.93	0.33
Elephant_African	0.90	0.02	0.03	0.57	0.45
Leopard_African	0.90	0.16	0.08	4.51	0.03
Cat_Golden	0.90	0.16	0.10	2.70	0.10
Chimpanzee	0.90	0.03	0.03	1.09	0.30



Figure S1. Three example photos taken from a burst of 10 images, showing a porcupine *Atherurus africanus* walking in front of the camera.

572



573


574 **Figure S2.** Confusion matrix showing model performance on out of sample test data (each row is
575 normalized independently). Figure S7 shows the confusion matrix with absolute numbers.

576

577

578


579



580

581 **Figure S3.** Histograms showing the frequency distribution (normalized density) of label confidence
582 from the machine learning model for the 27 classes in the out-of-sample test data.

583




584

585 **Figure S4.** Confusion matrix showing model performance for an aggregated set of 11 classes.

586

587



588

589 **Figure S5.** Confusion matrix showing model performance for an aggregated set of 11 classes after
590 removing labels with a predicted confidence < 70%

591


		Confusion matrix																															
Actual	Bird	12	0	2	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	0	5	0	0							
	Guineafowl_Crested	16	1	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	0						
	Guineafowl_Black	9	0	24	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0						
	Rail_Nkulengu	1	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0						
	Blank	56	0	3	2	265	1	0	0	46	66	0	4	3	0	0	21	3	4	0	4	0	11	0	51	116	2						
	Leopard_African	1	0	0	0	0	387	3	0	8	3	0	0	0	0	0	0	0	0	0	0	0	0	0	1	0	0						
	Cat_Golden	0	0	0	1	0	28	145	0	0	2	0	0	0	0	0	0	0	1	0	0	0	2	0	1	6	0						
	Chevrotain_Water	0	0	2	0	0	0	0	29	0	0	0	0	0	0	0	0	1	0	0	2	0	0	0	9	0	0						
	Elephant_African	0	0	0	0	0	0	0	0	3438	45	0	0	0	1	1	1	0	0	0	0	0	3	0	3	0	1						
	Human	1	0	0	0	0	0	0	0	82	631	0	1	4	0	0	0	0	0	0	0	0	1	0	0	2	0						
	Pangolin	0	0	0	0	0	1	0	0	0	2	16	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0						
	Monkey	0	0	0	0	0	0	0	0	3	0	0	483	2	0	3	2	0	0	0	0	12	3	0	7	10	0						
	Chimpanzee	0	0	1	0	0	0	0	0	29	4	0	4	344	2	1	0	0	0	0	0	1	0	0	1	2	0						
	Gorilla	0	0	0	0	0	0	0	15	0	0	3	47	0	13	0	0	0	0	0	0	0	0	0	1	0							
	Mandrillus	0	0	0	0	0	0	2	0	2	0	0	182	8	0	94	1	0	0	1	0	1	2	0	6	25	0						
	Squirrel	2	0	0	0	1	0	0	0	0	1	0	0	0	0	0	159	0	0	0	0	0	0	0	0	3	0						
	Porcupine_Brush_Tailed	0	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0	92	0	0	0	0	0	0	0	0	0						
	Rat_Giant	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	15	0	0	0	0	0	0	0	0						
	Mongoose_Black_Footed	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	26	0	0	0	0	0	0						
	Genet	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	0	141	0	0	0	0	0						
	Mongoose	2	0	0	0	0	0	0	0	1	0	0	0	2	0	0	0	1	0	1	0	71	0	0	0	0	0						
	Hog_Red_River	0	0	0	0	0	0	0	15	12	0	0	1	0	0	0	0	0	0	0	0	0	1124	0	22	1	0						
	Buffalo_African	0	0	0	0	2	0	0	0	30	4	0	0	0	0	0	0	0	0	0	0	0	10	118	36	6	6						
	Duiker_Red	0	0	0	0	1	0	0	0	27	15	1	0	0	0	0	0	0	1	0	0	0	3	1	5364	145	2						
	Duiker_Blue	7	0	2	0	0	0	1	0	11	5	0	5	0	0	0	1	0	1	0	0	0	0	0	0	61	3080	0					
	Duiker_Yellow_Back	0	0	0	0	0	0	0	0	32	13	0	0	1	0	0	0	0	0	0	0	0	0	0	2	27	9	436					
		Predicted																															
		Bird	Guineafowl_Crested	Guineafowl_Black	Rail_Nkulengu	Blank	Leopard_African	Cat_Golden	Chevrotain_Water	Elephant_African	Human	Pangolin	Monkey	Chimpanzee	Gorilla	Mandrillus	Squirrel	Porcupine_Brush_Tailed	Rat_Giant	Mongoose_Black_Footed	Genet	Mongoose	Hog_Red_River	Buffalo_African	Duiker_Red	Duiker_Blue	Duiker_Yellow_Back						

592

593 **Figure S6.** Confusion matrix showing model performance on out of sample test data after excluding
594 labels below a confidence threshold of 70% (with absolute numbers). Figure 3 shows the confusion
595 matrix with each row normalized independently.

		Confusion matrix																											
Actual	Bird	26	0	5	1	4	1	0	0	2	9	0	1	0	1	0	2	0	0	0	0	0	0	1	0	11	9	0	
	Guineafowl_Crested	24	5	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	0	
	Guineafowl_Black	17	0	32	0	1	0	0	0	2	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	0	
	Rail_Nkulengu	1	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	0	0	
	Blank	233	0	75	14	974	4	0	0	193	292	3	17	31	7	8	99	3	17	0	1	7	45	30	2	326	722	22	
	Leopard_African	6	0	3	0	2	401	10	0	26	7	0	4	2	0	0	0	0	0	0	0	0	0	0	0	3	2	1	
	Cat_Golden	1	0	0	2	0	40	179	0	2	10	0	2	2	0	0	0	0	4	1	1	2	1	3	0	4	9	0	
	Chevrotrain_Water	6	0	4	0	0	3	0	50	3	1	2	0	1	0	0	0	5	1	1	0	8	3	5	0	37	2	2	
	Elephant_African	3	0	5	0	7	3	1	0	3610	115	1	0	6	2	2	9	0	2	0	0	0	1	4	2	18	6	5	
	Human	5	0	3	0	0	0	0	0	130	679	0	1	12	2	2	1	0	0	0	0	0	0	1	0	6	6	2	
	Pangolin	0	0	0	0	1	1	0	0	2	5	23	0	0	0	0	0	0	0	0	0	3	0	1	0	1	0	0	
	Monkey	6	0	2	1	0	0	4	0	8	1	0	562	18	0	17	8	1	1	1	1	0	17	7	0	18	21	0	
	Chimpanzee	11	0	2	0	1	0	2	0	71	11	0	12	427	11	6	1	0	0	1	1	0	2	4	0	4	5	1	
	Gorilla	0	0	0	1	1	0	0	0	29	2	0	9	84	29	21	0	0	0	0	0	0	1	4	0	2	2	0	
	Mandrillus	3	0	0	2	0	0	5	0	13	0	0	310	37	1	166	4	1	0	4	2	0	3	10	0	19	55	0	
	Squirrel	15	0	1	0	2	0	0	0	1	4	0	3	0	0	0	194	2	2	0	0	0	2	2	0	2	18	0	
	Porcupine_Brush_Tailed	5	0	0	0	4	0	0	0	1	1	0	0	0	0	0	2	110	2	0	0	2	4	0	0	0	1	0	
	Rat_Giant	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	1	23	0	0	0	0	0	0	1	0	0	
	Civet_African_Palm	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	
	Mongoose_Black_Footed	0	0	0	0	0	0	0	0	3	1	0	2	1	0	0	0	0	1	2	29	0	1	4	0	1	0	0	
	Genet	0	0	0	0	0	6	0	0	2	1	0	0	0	0	0	0	0	0	0	0	151	0	0	0	1	0	0	
	Mongoose	6	0	0	0	0	0	1	0	2	1	0	0	2	0	0	0	2	0	0	4	0	82	1	0	0	1	0	
	Hog_Red_River	1	0	1	1	0	0	0	0	33	29	0	2	4	0	0	0	0	1	0	0	0	0	1258	2	61	11	0	
	Buffalo_African	3	0	0	0	2	0	0	1	53	14	0	2	1	3	0	0	0	0	0	0	0	0	29	164	76	14	17	
	Duiker_Red	11	0	2	1	6	0	0	0	80	68	1	2	15	0	2	0	0	2	0	1	0	4	11	3	5700	287	11	
	Duiker_Blue	22	0	9	0	4	2	5	1	24	39	0	7	9	1	0	4	0	3	0	0	0	4	16	0	157	3237	2	
	Duiker_Yellow_Back	1	0	1	0	2	0	0	0	58	27	0	2	2	1	2	0	1	0	0	0	0	0	8	8	64	22	493	
		Bird	Guineafowl_Crested	Guineafowl_Black	Rail_Nkulengu	Blank	Leopard_African	Cat_Golden	Chevrotrain_Water	Elephant_African	Human	Pangolin	Monkey	Chimpanzee	Gorilla	Mandrillus	Squirrel	Porcupine_Brush_Tailed	Rat_Giant	Civet_African_Palm	Mongoose_Black_Footed	Genet	Mongoose	Hog_Red_River	Buffalo_African	Duiker_Red	Duiker_Blue	Duiker_Yellow_Back	
		Predicted																											

Figure S7. Confusion matrix showing model performance on out of sample test data (absolute numbers). Figure S2 shows the confusion matrix with each row normalized independently.



ABOVE: Figure S8. Relationship between estimated occupancy probability for $n = 227$ camera stations (points) from machine learning (ML) labels (y-axis) and expert labels (x-axis) for the four focal species at each threshold (row) from 0 to 90%, in 10% intervals.




Figure S9. The image from Figure 9 with an added layer illustrating the most important regions of the image for the model when identifying the nkulengu rail. The brightest spot (yellow) near the center of the image encompasses a part of the bird's beak and head, which apparently were crucial during identification. We used the Grad-CAM (1) technique to create this image.

SI References

1. R. R. Selvaraju, *et al.*, Grad-CAM: Visual Explanations From Deep Networks via Gradient-Based Localization in (2017), pp. 618–626.