

A Research Pathway Model for evaluating the implementation of practice-based research: The case of self-management health innovations

Wilke van Beest ^{1,2,3,*}, Wouter P. C. Boon ², Daan Andriessen¹, Harald Pol³, Gerrita van der Veen³ and Ellen H. M. Moors²

¹Research Group Research Competence, HU University of Applied Science Utrecht, Padualaan 99, Utrecht CH 3584, The Netherlands, ²Innovation Studies, Copernicus Institute of Sustainable Development, Faculty of Geosciences, Utrecht University, Princetonlaan 8a, Utrecht CB 3584, The Netherlands and ³Research Group Marketing & Customer Experience, HU University of Applied Science Utrecht, Heidelberglaan 15, Utrecht CS 3584, The Netherlands

*Corresponding author. Email: Wilke.vanbeest@hu.nl

Abstract

This study explores the evaluation of research pathways of self-management health innovations from discovery to implementation in the context of practice-based research. The aim is to understand how a new process model for evaluating practice-based research provides insights into the implementation success of innovations. Data were collected from nine research projects in the Netherlands. Through document analysis and semi-structured interviews, we analysed how the projects start, evolve, and contribute to the healthcare practice. Building on previous research evaluation approaches to monitor knowledge utilization, we developed a Research Pathway Model. The model's process character enables us to include and evaluate the incremental work required throughout the lifespan of an innovation project and it helps to foreground that innovation continues during implementation in real-life settings. We found that in each research project, pathways are followed that include activities to explore a new solution, deliver a prototype and contribute to theory. Only three projects explored the solution in real life and included activities to create the necessary changes for the solutions to be adopted. These three projects were associated with successful implementation. The exploration of the solution in a real-life environment in which users test a prototype in their own context seems to be a necessary research activity for the successful implementation of self-management health innovations.

Key words: implementation; pathway model; self-management health innovation; practice-based research; process model; normal impact

1. Introduction

For dealing with complex societal problems, such as the ageing population and rising healthcare costs, much is expected from self-management health innovations (Lorig and Holman 2003). Self-management concerns a series of activities performed by patients to manage their symptoms, support treatment, and alleviate physical and psychosocial consequences outside the clinical setting and

integral to their daily lives (Barlow et al. 2002). Innovations to support self-management range from digital tools, for example monitoring body indicators to health education programmes, and potentially improve health outcomes and reduce costs (Jonkman et al. 2016). However, there are barriers to self-management health innovations, which are associated with their highly various character (Kennedy et al. 2014). The highly contextualized setting makes the generation of evidence on efficacy and safety difficult, which is

needed for healthcare professionals to prescribe and insurance companies to reimburse these products (Sivertsen and Meijer 2020). Moreover, such setting requires these products to be constantly adapted during implementation (Rosenberg 1994; Candy and Edmonds 2010; Bornmann 2013), calling for studying the entire innovation process.

Many self-management health innovations are studied and developed in the so-called practice-based research (PBR) projects at Universities of Applied Sciences (UASs). We define PBR as a co-creation process in which the implementation of results into practice is realized by integrating research results throughout the entire process in the form of activities, interventions, and interactions with stakeholders (McCull-Kennedy et al. 2012). The practice-based way of doing research and the research pathways involved with PBR is understudied phenomena though (Coombs and Meijer 2021).

Both developing and implementing self-management health innovations and the nature of PBR projects underline that the steps in the innovation are heavily interconnected. Thus, we evaluate research pathways using previous conceptual approaches to knowledge utilization by employing a process-based approach (Langley and Tsoukas 2012; De Jong et al. 2014). Employing a process perspective enables us to include and evaluate the incremental work of researchers and stakeholders required throughout the lifespan of an innovation project and it helps to foreground that innovation continues during implementation in real-life settings (e.g. Janssen 2016), emphasizing the ‘daily’ practical organizational level of interaction (Sivertsen and Meijer 2020). To address the gap in the literature of ways to evaluate research pathways involved with PBR, our study focuses on the following question: *What are the requirements for a process model to evaluate the implementation success of self-management health innovations developed in PBR projects?*

2. Theoretical framework

This study focuses on the pathway to successful implementation of self-management health innovation in the context of PBR projects. The aim of such PBR projects is to develop a prototype of a self-management device, training protocol, etc. and related knowledge on safety, usability, and so on. As discussed above, we apply a process perspective to evaluate successful implementation. Section 2.1 elaborates on this. Section 2.2 discusses examples of existing research impact models that use a process-based perspective. As implementation is an integral part of PBR, this requires an approach to evaluate the impact step by step throughout the entire research process. Therefore, Section 2.3 proposes a dedicated process model to describe research pathways of self-management health innovation in PBR projects.

2.1 Successful implementation of self-management health innovations

Several studies about the implementation of self-management health innovations have led to the identification of success factors for innovation, such as the ‘involvement of end-users, partners, or stakeholders’ (Van Beest et al. 2020). The evaluation of self-management health innovations should be extended and tailored to those originating from PBR. PBR projects have specific characteristics, such as the involvement of and high degree of interaction between many different stakeholders in the project and the frequent touching base

with practice, i.e. the specific context in which the project is intended for.

Self-management health innovations aim to equip individual’s ability into their daily lives to manage the symptoms, treatment, and physical and psychosocial consequences, as well as the lifestyle changes inherent in living with a chronic condition (Barlow et al. 2002). As such, these innovations focus on achieving ‘normal impact’, i.e. influencing daily activities and how well they are organized, not about individual cases of particularly interesting or impressive impact (Sivertsen and Meijer 2020).

Moreover, what is regarded as successful implementation varies from one context to another (Janssen, Stoopendaal and Putters 2015). The value assigned to an innovation is the result of interactive processes in which actors are involved and which are context-dependent. Not only does this mean that valuation of PBR projects varies across contexts but also that innovation activities do not stop at diffusion and continue throughout implementation (Janssen 2016).

So, PBR projects should be valued by taking into account particular contexts as well as using a process perspective describing the activities that lead to implementation success. Studying the implementation of innovations as a result of research projects is not a new issue. Over the last decades, scholars have analysed the social, cultural, environmental, and economic returns on research investment (Bornmann 2013). The following section presents an overview of existing research impact models that follow a process approach.

2.2 Impact of research

Within the scientific and policy community, there is a growing concern about assessing the impact of research. This has resulted in emphasizing methods to evaluate the process and impact of research. An emerging stream of literature has focused on interaction between researchers and stakeholders, taking slightly different perspectives. Examples are multidimensional models focused on the social impact of research, like the Payback model (Donovan and Hanney 2011) and Public Value Mapping (Bozeman and Sarewitz 2011). Other models aim to uncover the contribution of interactions, such as the Social Impact Assessment Methods through Productive Interactions (SIAMPI, Spaapen and van Drooge 2011) and Contribution Mapping (Kok and Schuit 2012). A third group is the actor- and process-oriented frameworks focused on diverse outcomes and interactions, such as Participatory Impact Pathway Analysis (PIPA, Douthwaite et al. 2007) and Socio-Economic Analysis of the Impacts of Public Agricultural Research (ASIRPA, Joly et al. 2015). This section discusses these models and their relevance for assessing PBR.

The Payback Framework was created by Buxton and Hanney (1996) to assess the outcomes of health research structures. It consists of a logic model of the research processes, and various categories of research paybacks and anticipated impacts. The Payback Framework recognizes and explicitly makes room for a plurality of types of impact, and has been applied in a range of different contexts both within and beyond healthcare (Donovan and Hanney 2011). The framework is relevant for PBR because PBR involves different contexts, types of impacts, and ways of doing research. At the same time, it is not possible to tie the categories of benefits to specific stages of the model, as the Payback Framework is mainly focused on the outcomes of research. A focus on the entire process is important, though, because implementation of PBR into follow-up research and

daily organizational practice is realized during the whole research process. Bozeman and Sarewitz (2011) propose Public Value Mapping (PVM) as an approach to assess what they call public values of science. PVM is, just like PBR, based on the idea that science outcomes should serve and advance public values. Public values include outcomes like environmental quality, environmental sustainability, healthcare, and meeting basic needs, e.g. housing and food. PVM does not *a priori* define the different dimensions of effects (the set of public values), ensuring that PVM can take a context-specific approach. At the same time, PVM makes it difficult to compare projects.

Two approaches that are more focused on the contribution of interactions during the process are SIAMPI (Spaapen and van Drooge 2011) and Contribution Mapping (Kok and Schuit 2012). Both approaches focus on interactions during the process which are better traceable and provide more insightful information regarding implementation. Central to the SIAMPI framework is the concept of productive interactions: the mechanisms through which research activities lead to a socially relevant application. The SIAMPI framework introduces the notion of impact pathways in which interactions occur during or after a research project, or even before it has started. Using process indicators does help anticipating societal impacts that may not yet have materialized at the moment of evaluating (De Jong et al. 2014). SIAMPI is an inspiration for capturing impact throughout the entire PBR process. Contribution Mapping is a three-phase process. Each phase is associated with its own research activities, goals, and interactions between different stakeholders. Contribution Mapping emphasizes that different activities and different types of interactions may be important during different stages of a research project. Both frameworks (SIAMPI and Contribution Mapping) are focused on efforts instead of results. For PBR, it is relevant to use a framework that emphasizes that each phase requires different interactions through activities, without reducing the complexity of interactions. The types of impact described in these two frameworks are primarily about knowledge transactions in interactions whereby knowledge could be implicitly contained in products or services. In PBR, impact is just as often explicitly achieved through interventions during the research with the aim of initializing change in practice. Recognizing the different ways in which PBR generates impact is essential to evaluate the impact of this type of research.

PIPA (Douthwaite et al. 2007) is an actor- and process-oriented framework focused on diverse outcomes and interactions. For PBR, this integration is helpful because PBR is a co-creation process between actors with the aim to directly contribute to practice. However, PIPA does not make explicit the different contexts, types of impacts, and ways of doing research. Essential for PBR is the way in which practical real-life problems are analysed and how research contributes to practice by means of generating actionable knowledge and developing practical solutions, methods, and instruments. ASIRPA (Joly et al. 2015) pays attention to the process of transformation of knowledge actionable by incorporating it into new products, processes, or governing things. ASIRPA is multidimensional; based on the involvement of networks of actors, at different stages and playing a variety of roles, and over a non-linear impact pathway. The method is useful for PBR where it offers assessment of the societal impact from design and scaling-up to the level of the organization. The ASIRPA framework contains all factors in the research process to achieve impact and is therefore comprehensive. At the same time, the way in which impact is incrementally realized during

the whole process is not part of the model. In PBR, the impact is achieved through steps from the early start through the implementation. Mapping the entire pathway provides insights into the different ways in which PBR generates impact which is essential to evaluate the impact of this type of research. It is therefore important for PBR to obtain an overview of the entire research path, whereby during the project it is made explicit which output can be attributed to a certain impact area such as further research or practice. Table 1 gives an overview of the above-mentioned models and shows which elements for a model to analyse the pathway of the implementation of PBR resulting in self-management health innovations can be used.

The existing models share a process-based approach, but the models are not sufficient for PBR because of the missing elements as mentioned in Table 1. Coombs and Meijer (2021) emphasize that there is a need for a new process-based evaluation approach for PBR, which makes the inclusion for stakeholders in the process explicit. In Section 2.3, we describe the contours of such a process model building on the elements as mentioned in Table 1.

2.3 A process model to study innovation processes in PBR

There is a need for a process model that makes explicit what type of impact can be attributed to what research step to achieve a better understanding of the pathway to a successful implementation through PBR. We propose a Research Pathway Model as a step in this direction.

A Dutch version of the *Research Pathway Model (RPM)*, PRO-model, was developed in an earlier research project (Van Beest, Balje and Andriessen 2017) for research in different domains to discuss the readiness of research projects for real-life practice before, during and after the project. The early version of the RPM uses the Technology Readiness Levels (TRL) as a structuring device (Mankins 2009). TRL is meant to assess the readiness of a particular technology from the discovery phase to the implementation in the market. TRL indicates the degree of a technology in nine levels, whereby TRL 1 stands for technology at the start of development and TRL 9 captures a technology that is technically and commercially ready for upscaling (Mankins 2009). However, TRL could not be used one-on-one because the linearity of the model does not fit into the implementation process and TRL is only focused on technological innovations, whereas the self-management health innovations could also be non-technological. The main difference between TRL and the RPM is that TRL is a linear model with nine consecutive levels, while the RPM is a process model placed in a matrix (Figure 1) without imposing a predefined order of activities. The model was discussed, updated, and translated into English in two co-design sessions together with a group of Dutch researchers from the research groups Co-design, Physiotherapy, Education and Research impact at an UAS who have been working with the Dutch version of the RPM and in a third session with a group of experts on PBR. The model consists of two axes (research contexts and research activities) and nine research goals (Figure 1).

We discern three research contexts: *theoretical context*, *conceptual context*, and *real-life practice context*. The contexts are based on Dalsgaard and Dinder (2014) who distinguish 'theory', 'bridging concepts' as one form of intermediary knowledge bridging between theory and practice and 'real-life practice', and Hevner (2007) who demarcated the 'knowledge base' (theoretical context), the design cycle which iterates between research processes and the core

Table 1. Overview of models to evaluate research projects

Model	Description	Helpful elements for PBR	Missing elements for PBR
ASIRPA (Joly et al. 2015)	Comprehensive approach for assessing the socio-economic impact of public-sector research organizations through theory-based case studies, selected to characterize the diversity of the broader impacts, and standardized to allow the scaling-up of the analysis of impact to the level of the organization.	<ul style="list-style-type: none"> - Actor- and process-oriented framework focused on diverse outcomes and interactions. - Pays attention to the process of transformation of knowledge actionable by incorporating it into new products, processes or governing things. - Multidimensional; based on the involvement of networks, at different stages, playing a variety of roles, over a non-linear impact pathway. 	The way how impact step by step is realized during the whole process is not part of the model.
Contribution Mapping (Kok and Schuit 2012)	A three-phase process. Each phase is associated with its own research activities, goals and interactions between different stakeholders. The phases are separated by soft, blurry borders as activities of one phase may continue into the next.	Focus on interactions during the process which are traceable and provide insightful information regarding the process of implementation.	Primarily about knowledge transactions in interactions.
Payback Framework (Donovan and Hanney 2011)	Facilitate data collection and cross-case analysis by providing a common structure and consists of two elements: <ol style="list-style-type: none"> 1. A logical structure which describes the various stages of outputs, outcomes and dissemination. 2. An element describing the outputs, outcomes and impact. 	<ul style="list-style-type: none"> - Recognizes a plurality types of impact, explicitly makes room for these and has been applied in a range of different contexts both within and beyond the health service. - It is possible to do a cross-case analysis by providing a common structure. 	<ul style="list-style-type: none"> - It is not possible to tie the categories of benefits to specific stages of the model. - It is focused on the outcomes (goals) of research.
PIPA (Douthwaite et al. 2007)	Evolving tool that offers a deeper understanding of the results that projects might attain with specific partners to help set priorities and support funding proposals.	<ul style="list-style-type: none"> - Actor- and process-oriented framework focused on diverse outcomes and interactions. - Possible to use before, during and after the project. 	The model does not make explicit the different contexts, types of impacts, and ways of doing research.
Public Value Mapping (Bozeman and Sarewitz, 2011)	Conceptual tool for developing systematic understanding of the multiple determinants of social outcomes and the role of science as part of institutional arrangements and networks. The focus is on social impacts rather than scientific and economic impacts.	<ul style="list-style-type: none"> - Case-based approach. - Focusing on assessing the impacts of a given research endeavour on public values. - Based on the idea that science outcomes should serve and advance public values. 	Difficult to compare cases.
Social Impact Assessment SIAMPI model (Spaapen and van Drooge 2011)	Central to the framework is the concept of productive interactions. Involves two central tasks: <ol style="list-style-type: none"> 1. to enlighten the mechanisms by which social impact occurs; 2. to develop methods to assess social impact. 	<ul style="list-style-type: none"> - Focus on interactions during the process which are traceable and provide insightful information regarding the process of implementation. - Can occur during or after a research project, or even before it has started. 	Primarily about knowledge transactions in interactions. In PBR, impact is just as often achieved through the development of products, the personal development of the involved people and through interventions during the research with the aim of initialize change in practice.

activities of building and evaluating the design artefacts (conceptual context), and environment (real-life practice context). The RPM extends these concepts from design studies and makes them applicable to PBR. In the theoretical context, the research is focused on creating, exploring, and delivering a better understanding of

problems and related propositions for solutions, which are advanced and not verified in practice yet. The conceptual context makes the translation from presupposed solutions to a more specific prototype that is created, explored, and made. In this context, researchers, project partners, and other stakeholders bring in experiential

Research contexts Research activities	Theoretical context	Conceptual context	Real-life practice context
Create	A.1 Create theoretical understanding	B.1 Create a concept	C.1 Create understanding of real-life practice
Explore	A.2 Explore theory or a concept in a controlled situation	B.2 Explore a concept	C.2 Explore a solution in real-life practice
Deliver	A.3 Deliver theory or knowledge	B.3 Deliver a concept	C.3 Deliver change in real-life practice

Figure 1. Research Pathway Model.

knowledge to translate theory to prototype, by themselves, together with the consortium and with or without the end-users, but always in a protected niche as a brainstorm room, a pilot environment, or in the context of a pilot organization. The real-life practice context concerns the context in which the prototype is tested in the world of professional practice and/or the living environment of end-users.

In addition to the focus areas, the model discerned three overarching research activities: *Create*, *Explore*, and *Deliver*, which have been explored and validated during previous application in PBR innovation processes (see also above for how this validation has been done). *Create* refers to research goals with the aim to create more understanding of the problem and to create an idea to solve the problem. These goals can be realized by research activities throughout the whole research pathway, whereby it is possible that a research goal returns during the iterative process. Examples are performing a literature review (create theoretical understanding), co-designing a prototype (create a concept), observing end-users in their own context (create understanding of real-life practice).

Explore refers to research goals with the aim of exploring an idea, concept, construct, or solution by research activities. Examples are laboratory research and tests in a controlled environment (explore theory or a concept in a controlled situation), evaluate a prototype in a pilot (explore a concept), and testing a prototype in the environment of the end-user (explore a solution in real-life practice).

Deliver refers to goals related to the delivery of the insights during the research pathway. Examples are: publishing research papers or grey literature (Deliver theory or knowledge), deliver a prototype to the market (deliver a concept), and working on an implementation strategy in an organization (deliver change in real-life practice).

Based on the characterization of pathways in e.g. SIAMPI and ASIRPA, we define a research pathway as the order or sequence in which a project goes through the nine research goals, where it is possible to have goals iterate several times. The matrix does not prescribe an ideal sequence of goals. The RPM can be used as a process model mapping activities, patterns, and the linkage between them. By using the RPM, impact analysts are able to visualize goals and the underlying activities. As such, the activities that need to be done to implement an innovation in practice become visible for the researchers themselves and communicable to other stakeholders. By using such a process model, researchers are able to *ex ante* design their research pathways to plan their actual research activities, to monitor them during the project, and to evaluate their pathway after concluding the research project. The matrix makes it possible to map the activities that contribute to the implementation of the

innovation project whereby the contributions to the real-life practice, conceptual, and theoretical context can be made explicit.

Inspired by and based on the impact models presented in Section 2.2, we offer RPM as a research evaluation framework suited for PBR. Following a process approach, we are able to analyse how the activities that contribute to the implementation of research in practice give meaning to a self-management health innovation and how this work is made manageable. Thus, we apply the RPM to nine illustrative self-management health research projects.

3. Methods

To explore how a process model for innovation provides insights into the implementation success of self-management health innovations developed in research projects, we used a qualitative multiple case-study approach (Stake 1995). Empirical data in this study came from nine research projects on the development of self-management health innovations, with consortia that consist of practice-based researchers, healthcare professionals, and/or healthcare entrepreneurs. The research design is exploratory due to the newness of the *Research Pathway Model*.

3.1 Case selection

For this multiple case study, we selected nine illustrative self-management health research projects from the HU University of Applied Sciences Utrecht with consortia consisting of researchers, healthcare professionals, and entrepreneurs (see Appendix 1). We focused on only one UAS because we are studying implementation success in a particular context. Cases were selected from the project databases of websites of the subsidy providers ZonMw and the Taskforce for Applied Research SIA and the website of the HU University of Applied Sciences Utrecht. The criteria for including a case were based on criteria in the self-management definition by Jonkman et al. (2016, pp. 36):

'self-management innovations aim to equip patients with skills to actively participate and take responsibility in the management of their chronic condition in order to function optimally through at least two of the following aspects: (1) providing knowledge about the condition and/or treatment, (2) active stimulation of symptom monitoring, (3) enhancing problem solving skills (self-treatment, resource utilisation, stress/symptom management), (4) enhancing physical activity, (5) enhancing dietary intake, (6)

enhancing smoking cessation, (7) enhancing medication adherence.’

The project description was assessed by one researcher, and iteratively validated during a three-hour group discussion with the other authors. This resulted in the selection of nine projects. They were funded between 2010 and 2020, with a lead time of 2–5 years, and are briefly described in [Appendix 1](#).

3.2 Implementation success

We define implementation success as the actual use or integration of an innovation within a specific setting in an organization ([Rabin et al. 2008](#)). Implementation success in this study meets at least one of the following criteria: 1, the innovation is used in a specific programme of at least one organization by the end-users themselves, 2, the innovation is integrated in daily organizational routines of end-users in a specific setting in or related to an organization, 3, the innovation is sold to organizations through a (healthcare) entrepreneur or publisher, and 4, the innovation is made available in an organization whereby the actual use is monitored by the organization as well.

[Table 2](#) presents the nine projects and shows which projects reach a successful implementation at the moment of data collection (February–May 2019) related to the criteria of [Rabin et al. \(2008\)](#). Three projects (Projects 1, 2, and 3) achieved a successful implementation of a self-management health innovation. In Project 1, the intervention tool is made freely available and can be downloaded. The intervention is integrated with a different working method with in a participating partner and a non-partner organization. The intervention has also been implemented in an education programme of another university. The main researcher provides training to interested organizations on the use of the intervention. The researchers obtained a grant for a follow-up study with the aim of digitalizing the tool. In Project 2, the mobile phone app is implemented by a business partner. The researchers have applied for and received

grants for an extension of the app so that it can help to self-manage other disorders. In Project 3, the tool is made available via a publisher. Participating speech therapists and speech therapists in their network will implement the tool in their treatments.

In five of the projects (4, 5, 7, 8, and 9) that have not reached a successful implementation yet, one or more project members are still active in the implementation process of the innovation. This means that they are still engaged in innovation activities (see [Appendix 2](#)), and that the implementation success of these projects is thus not guaranteed (yet). The development of the tool was discontinued in Project 6 with no further work being planned.

3.3 Data collection

To explore the contributions of self-management health innovation projects to the implementation success of the developed innovations, we used a qualitative multiple case-study approach, including document analysis and semi-structured interviews ([Stake 1995](#)). For the document analysis, we manually analysed documents from the project website or the public website of the UAS to gain insights into the published activities of the projects. To acquire information about additional information of activities and researchers’ motives, we participated in activities during the project and in interactions with stakeholders. We also conducted semi-structured interviews with an interview schedule ([Appendix 4](#)) with the most involved researchers (project leaders or PhD students) of the nine projects, i.e. healthcare entrepreneurs (n=3), healthcare professionals (n=4), and researchers with a different expertise (co-design and information systems) from the main researcher (n=2). In addition, we conducted evaluation interviews after the project with healthcare professionals of one of the projects (n=6). Interviewees were asked to describe how the research pathway had evolved. Interviewees were encouraged to be specific about the research processes and to provide examples. To increase the reliability of the data, the interviews were audio-recorded and transcribed verbatim. Respondents

Table 2. Implementation success in the nine cases

Project	Type of innovation	Implementation stage during data collection	Implementation success
1	Intervention tool	The innovation is used in a specific programme of at least one organization by the end-users themselves.	Yes
2	App	The innovation is made available in an organization whereby the actual use is monitored by the organization as well.	Yes
3	Talking tool	The innovation is sold to organizations through a (healthcare) entrepreneur or publisher.	Yes
4	Technology for stimulating self-management	Multiple prototypes are available for research in follow-up projects. There is market demand for one of the prototypes. The researchers are looking for an entrepreneur who wants to market the product.	No
5	App	There is a prototype and the researchers are working on a grant application. One of the healthcare entrepreneurs is collaborating with the healthcare entrepreneur of project 2 to implement the app in their system.	No
6	Talking touchscreen	There is a prototype.	No
7	Tool to monitor conditions	There is a prototype.	No
8	Digital platform	A final prototype is delivered and the business partner is project leader of this project and will decide about the further development.	No
9	mHealth intervention	There is a prototype of the product. The researchers are looking for an entrepreneur who wants to market the product.	No

gave approval to record and use the data, and checked the manuscript for inaccuracies.

3.4 Data analysis

Data analysis was done in two steps: 1, a detailed qualitative within-case analysis to analyse each research pathway, and 2, a cross-case analysis to compare the different pathways and to discover patterns (Stake 1995). The documents and transcripts were used to iteratively analyse the research pathways of the projects.

The first step of the analysis consisted of coding the documents and interviews through a coding scheme (see Appendix 2) which we developed based on the RPM in Section 2.3. The documents and interview transcripts were read and the content was highlighted using Atlas-ti software version 8.4.15. Coding was done by one researcher. The second step of the analysis was clustering all relevant data in the form of quotes from both interviews and documents. For the within-case analysis, we analysed the presence of the selected process features, and structured them per project in separate tables (Appendix 3). The transcripts of the interviews with researchers were compared to the interviews with the other involved interviewees of the same case to complete the pathway. Process features could only be analysed if the project explicitly undertook a research activity, or named an event or interaction related to the predefined research goals. We then conducted a systematic cross-case analysis and clustered all relevant data in the form of quotes both from interviews and documents in a data matrix (Miles and Huberman 1994). This clustering was structured by the codes of the RPM. To be able to gain insights into the pathways of the projects, we needed to combine the data of the codes. Our analysis was recursive, constantly moving from the specific cases, to the more general, with the aim of identifying commonalities and patterns across the variety of cases.

4. Results

The aim of this article is to understand what requirements are needed for a process model to evaluate the implementation success of self-management health innovations developed in PBR projects. We applied the Research Pathway Model on nine cases. Section 4.1 describes the nine cases using the RPM and explicates how the model can illustrate the steps taken. Section 4.2 then explores how the RPM can be used to demarcate activities and requirements that lead to implementation success. Section 4.3 then emphasizes the sequencing of these activities and puts these activities into a process perspective.

4.1 Using the Research Pathway Model

The *Research Pathway Model* (Section 2.3) was used for evaluating research pathways in PBR projects which aimed at developing and implementing self-management health innovations.

Figure 2 illustrates the research pathway of one of the nine cases (Project 1).

A number of rehabilitation centres in the Netherlands had questions about the long-term effectiveness of the existing self-management chronic pain programmes. These treatment programmes focus on patients with chronic musculoskeletal pain associated with personal and social dysfunction. The interventions are aimed at changing the behaviour of patients so that they can better deal with pain complaints in their daily lives. To address the

articulated problem, the general aim of the project was to develop and bring together knowledge and skills in the field of self-management and chronic pain rehabilitation.

In *step 1* (C.1, Figure 2) of the project, the researchers conducted a retrospective survey of the effects of pain rehabilitation, the degree of self-management, and the percentage of relapse within 6 months. The survey took place at the rehabilitation centre involved, amongst patients who had completed a chronic pain rehabilitation programme indicating that this research step takes place in the real-life practice context. According to the research protocol, these patients completed a short questionnaire 3, 6, and 12 months after the start of rehabilitation. After these research activities, the investigators had an understanding of the effects of the existing programme. In *step 2* (B.1, Figure 2), professionals and patients with chronic pain were asked about the underlying reasons that could possibly cause the relapse and about methods that could be useful to prevent such a relapse. Through co-design, in the conceptual context, a prototype for an intervention was developed to help people with chronic pain to retain the results of their rehabilitation. This is an intervention tool in which values and goals can be formulated and important moments or experiences can be saved during rehabilitation and afterwards, as a reminder and support in the event of a relapse. In a parallel process (Numbered as *step 3*; A.1, Figure 2), a systematic literature study, in a theoretical context, was conducted covering the effects of a self-management intervention on self-efficacy, daily activities, and limitations in patients with chronic pain complaints. In *step 4*, prototypes developed by students were presented at the rehabilitation centre. The exploration was not just about the prototype but, by hidden design and in line with step 2, about the principle of the presented interventions in a conceptual context. In *step 5*, the prototype was tested by eight patients in a feasibility study (C.2, Figure 2) in a real-life context. They rated the usability of the tool on average with a 7.8 and motivated them to actually carry out their goals in the treatment programme. The practitioners of the rehabilitation centres have also been asked about their experiences. They were positive about the possibilities of the tool. Based on the experiences of users, the tool was then adapted and made available for all rehabilitation centres. The pathway was followed by *step 6* (B.3, Figure 2) in which a game was developed and delivered (in a conceptual context) that could support people with chronic pain to make their lives with chronic pain clear to their relatives. The research project was followed up by a PhD project to capture all studies in academic publications. The results were shared during a symposium and in a few newsletters (*step 7*, A.3, Figure 2) in the theoretical context. The pathway ended with *step 8* in which the intervention is made freely available for downloading (C.3, Figure 2). The intervention is integrated in the real-life practice context within a participating partner and a non-participating organization. The principal investigator provides training to new interested organizations on the use of the intervention. The intervention has also been implemented in a teaching programme of another university.

A description of the research path of all cases can be found in Appendix 3. To illustrate the use of the RPM, Figure 3 shows activities with examples from each case to cover all nine research goals.

By structuring activities into the RPM, a research pathway can be drawn made for each case. Up to this point, examples from the cases were used to provide insights into the working of the model. The next section discusses the activities and requirements to reach implementation success by using the RPM.




Figure 2. Steps taken in Project 1 following the Research Pathway Model. In Project 1, the following steps were taken in this sequence: *step 1* C.1 Create understanding of real-life practice, *step 2* B.1 Create a concept, *step 3* A.1 Create theoretical understanding, *step 4* B.2 Explore a concept, *step 5* C.2 Explore a solution in real-life practice, *step 6* B.3 Deliver a concept, *step 7* A.3 Deliver theory or knowledge and *step 8* C.3 Deliver change in real-life practice. A.2 ‘Explore theory or a concept in a controlled environment’ is no part of this project.

	Theoretical context	Conceptual context	Real-life practice context
Create	Create theoretical understanding Literature review or a Delphi study: <i>“We started with a literature review on self-management behaviour, followed by a Delphi study with international experts on the topic.”</i> (Project 9)	Create a concept Co-design session or interviews with the aim to find a solution: <i>“We evaluated the co-design sessions with the [target group]. (...) It became clear that it had to be a touchscreen.”</i> (Project 6)	Create understanding of real-life practice Observations or interviews in a real-life setting with the aim to understand the context: <i>“Because the system needs to be integrated into the homes and lives of elderly, we visited them at home and asked them about hobbies, daytime activities and time schedules.”</i> (Project 8)
Explore	Explore theory or a concept in a controlled situation Laboratory research or taking measurements in a controlled area such as a hospital or physiotherapy practice: <i>“Exercise therapists and physical therapists test with 50 participants (tested twice, with seven days between the two measurements) in an exercise laboratory”</i> (Project 7)	Explore a concept Testing a low fidelity prototype in a pilot or test environment with different stakeholders: <i>“We tested the prototype in an iterative process with speech therapists and parents and incorporated the recommendations in each round into a new prototype.”</i> (Project 3)	Explore a solution in real-life practice Testing a high fidelity prototype in real life: <i>“People have downloaded the app that can advise them to go to a physiotherapist. But of course you don’t want to do that too early, nor too late. We will discuss what the indicators are based on the use of the app and what is possible to integrate into the app.”</i> (Project 2)
Deliver	Deliver theory Sharing information via a scientific or professional journal, information folders or congresses: <i>“We have organised a number of conferences for the professional field and others interested to share knowledge and have prototypes experienced. It makes people think about the possibilities of technology.”</i> (Project 5)	Deliver a concept Transferring the prototype to the next project, a developer or a company: <i>“The goal was to make a prototype. But I see that it is in demand through guest lectures, presentations and workshops that I have given and now I have a follow-up grant to develop it further.”</i> (Project 4)	Deliver change in real-life practice Implementing and/or evaluating the solution in real life: <i>“It is currently used in three pain rehabilitation centres, in the current version. People see it as an added value for current practice. If people are interested, I come by, give a training and then they get that workbook.”</i> (Project 1)

Figure 3. Examples out of the nine projects for each step in the RPM.

4.2 Explaining implementation success by using the Research Pathway Model

We discuss the model by focusing on the three research activities (Create, Explore, Deliver) to explore which insights about project success can be derived from using the process model.

4.2.1 Create

'Create' refers to research goals with the aim of deepening the understanding of the problem (in theory or in real-life practice) and to develop ideas to solve the problem. The research goal of creating a concept and/or understanding the problem was the start in all nine cases. Surprisingly, only two projects included a study with the goal to understand the real-life situation of the end-users before they started creating the idea. More often the projects involved end-users creating a prototype or product in co-design sessions at a later stage. But in the projects doing research activities to understand the real-life situation of end-users, researchers themselves have developed the idea for a prototype or product without the use of input from the end-users: *'The idea of finding the solution in a tool, was our own. The speech therapists have made the problems clearer. And they were very recognizable to me, because I had actually experienced the same problems as a speech therapist'* (project 3). Although the idea in this quote did not arrive from practice, the researcher had experience in practice. All projects created a concept with researchers from different disciplines and/or together with end-users and/or stakeholders. The involvement of stakeholders is seen in the research goals 'create a concept' (n = 8) and 'create understanding of real-life practice' (n = 7) and not in the goal 'create theoretical understanding', which confirms that practical relevance in research done in the 'theoretical context' is limited. Creating an idea or understanding a problem is not just a goal in the beginning of the process. Throughout the whole process, activities with the goal to create a concept or understand the problem occur: *'the researchers collaborated with business partners to keep the tool online after the project had finished, but this partner ended the cooperation. Then we found each other (...) and started a collaboration and have integrated the online tool as a module in our own tool'* (Project 2). This fragment shows that understanding the problem and creating ideas could be an activity in the implementation phase. With this the quote also shows that there are different levels of creating an idea or understanding a problem. Sometimes it can be about understanding a fundamental question and sometimes about adapting an existing idea or practical solution.

4.2.2 Explore

'Explore' refers to research goals that cover research activities in which concepts, constructs, or solutions are explored. All projects included research activities to test or develop their prototype together with the target group (explore a concept). In this step, researchers mostly test a low fidelity prototype. One of the interviewees from a healthcare organization was critical about testing with the target group in early stages of the project and suggested more testing in a controlled environment without end-users: *'sometimes a product must be more developed before you introduce it. If you have consulted or tried out the concept with someone and the basic idea could work well, then you should continue with a product that is actually good and not try it again with some adjustments. Overall it is disappointing for clients that a product is still not available for them'* (Project 4). The quote shows that exploring the

concept could lead to frustrations among the end-users involved, because the prototype did not work well and/or because the prototype is only available for a short period. Four of the projects included a research activity with the goal to explore theory or a concept in a controlled environment. In three projects, researchers explored the solution in a real-life environment in which the users tried the last version of a prototype in their own context. The projects that succeeded in implementing their project benefited from the exploration with the target group and the enthusiasm for the final prototype.

4.2.3 Deliver

'Deliver' refers to goals related to the articulation and diffusion of the insights during the research pathway. All cases delivered theory and a prototype. Three out of nine cases delivered real-life changes by implementing a product. In these three cases, the product is also scaled up or it is possible for healthcare organizations or individuals to buy or download the intervention. Four projects did involve a business partner, aiming for the business partner to play a role in developing the innovation or putting the innovation into production or onto the market. Some of these business partners dropped out during the project because the further development of a prototype is more expensive than the expected reward. One of the business partners who decided not to develop the tool yet stated (Project 5): *'the target group is too small if you want to start selling an app. (...) It would be better if we could add it to the existing service and make those functionalities not only available for the target group, but also for others.'* Some projects did not aim to deliver a market-ready product: *'the goal was to make a prototype'* (Project 4). In this case, a plan or idea for developing the prototype to a product was missing. Other researchers integrated the implementation process in their study, for example, by participating in a competition in order to collect money and help for further development and implementation: *'We estimate that the app will be successfully implemented. Yet there are many challenges, such as the growth of similar applications on the market. Some of them already have agreements with hospitals. On the other hand, we have noticed as a result of the challenge we have won, there is a lot of attention for our project and there are parties who contact us'* (researcher project 9). Two projects (both having succeeded in implementation) changed their first idea of making a digital product to creating a non-technical product to increase the chance of success. Time and a limited budget are the main reasons to change plans. The other project that succeeded in implementation was a follow-up project. All researchers interviewed indicated that they could use more help with the delivery of their prototype or product to real-life contexts.

The described activities = that contribute to implementation success of self-management health innovations developed in PBR should be perceived through a process lens, which is done in the next section.

4.3 Patterns in research pathways

Table 3 presents the pathways the projects followed using the RPM. A research pathway contains all research goals (and underlying activities) in all conceivable sequences. Table 3 shows the nine projects on the vertical axis and the sequences of research goals (in numbers) on the horizontal axis. The table does not provide insights in the underlying activities, but reports at the level of the research goals.

Table 3. Research pathways of the nine cases following the Research Pathway Model (Figure 1)

Pathway projects	1	2	3	4	5	6	7	8	9	10	11	Implementation success
Project 1	C.1	B.1	A.1	B.2	C.2	B.3	A.3	C.3				Yes
Project 2	B.1	A.1	B.2	C.2	C.2	A.2	B.2	C.2	B.3	A.3	C.3	Yes
Project 3	B.1	B.1	C.1	B.1	B.2	B.2	B.2	C.2	A.3	B.3	C.3	Yes
Project 4	B.1	C.1	A.1	A.3	B.1	B.2	A.3	B.3				No
Project 5	B.1	B.2	B.2	C.1	A.1	B.2	A.1	B.1	A.3	B.3		No
Project 6	C.1	C.1	A.1	B.2	B.3	A.3						No
Project 7	B.1	A.1	A.2	B.1	B.2	B.2	A.3	B.3				No
Project 8	C.1	B.1	A.2	B.1	B.2	B.1	B.3	A.3				No
Project 9	A.1	B.1	B.1	A.1	C.1	A.2	B.2	A.3	B.3			No

‘A’ stands for the *theoretical context*, ‘B’ for the *conceptual context* and ‘C’ for the *real life practice context*. ‘1’ stands for activities with the goal to Create, ‘2’ stands for Explore and ‘3’ stands for Deliver (Table 1).

Although each research pathway to reach innovation success differs, Table 3 shows some similarities in these nine pathways: each of them achieved the research goals to create a concept (B.1), explore a concept (B.2), to deliver a concept (B.3) and to Deliver theory or knowledge (A.3). The research goal of creating a research idea and understanding the problem (A.1, B.1, C.1) was the start in all nine cases.

The fact that all projects created a concept (B.1) means that in these nine cases a prototype or product is created by researchers from different disciplines and/or together with end-users and/or stakeholders. All nine projects included the exploration of their prototype in a pilot or test environment with potential users associated with one of the consortium partners (B.2). All projects have delivered theory (A.3), which means that they shared their knowledge and results via, for example publications in national and international papers, newsletters, project websites and/or congress presentations. All cases completed activities with the goal to deliver a concept (B.3). In Projects 1, 2 and 3, the prototype was adopted by either a publisher, a business partner or directly by the healthcare organization. The other projects produced either a prototype which could be adopted by developers and business partners, or a product which is ready for the market, but not yet implemented. Some researchers indicated that delivering the prototype was the goal of the project. Another pattern is that most research activities do have an explorative character and/or are part of the *conceptual context* where researchers involved end-users.

What stands out in Table 3 is that three of the nine cases delivered changes in practice by implementing a product (C.3). Interestingly, in these three projects researchers explored the solution in a real-life environment in which the users tried the last version of a prototype in their own context through a feasibility study. In these three cases, the product is also scaled up or it is possible for other healthcare organizations or individuals to buy or download the intervention. The activities underlying *deliver change in real-life practice* are related to working on an implementation strategy in an organization with the goal to use or integrate the innovation in the setting or to introduce a product to the market. Throughout their research pathways, the three projects that actually successfully implemented the innovation were working on activities with the aim to *explore the solution in real-life practice* and to *deliver change in real-life practice*.

By studying the different research goals and their underlying activities from early discovery all the way to implementation, the

focus was not only on knowledge transfer (*deliver a theory*) but also on the activities to realize actual use in practice (*deliver change in real-life practice*).

5. Conclusions and discussion

The insights provided in this article form a first attempt to understand how a process model provides insights into the evaluation of implementation success of health self-management innovations developed in PBR projects and what the requirements are for such a process model. The process perspective enables us to include and evaluate the incremental work required during the whole innovation project. To evaluate research pathways which focus on PBR and ‘normal impact’, a process-based approach is missing (Sivertsen and Meijer 2020). By performing a multiple case-study approach, we were able to answer the following question: *What are the requirements for a process model to evaluate the implementation success of self-management health innovations developed in PBR projects?*

Through a process-based approach, we were able to recognize three requirements for a process model to evaluate the implementation success of self-management health innovations developed in PBR projects. The first requirement is the specification of research goals. By mapping a process consisting of specific steps in the pathway that influences successful implementation of a self-management health innovation, we were able to recognize the incremental work of researchers and stakeholders.

The second requirement is the specification of the research contexts and research activities to structure the research goals. By making the research contexts explicit, we were able to visualize the focus on theoretical contributions, contributions to further development (concept) or contributions to one or more specific organizations (real-life practice). The focus on specific contributions is helpful to evaluate impact, especially in the case of PBR projects as they have the ambition to go beyond theoretical contributions and have practical use. The research activities *create*, *explore*, and *deliver* provide insights into the pathway to implementation success of self-management health innovations developed in PBR projects. Further research could possibly lead to refinements of these three research activities.

A third requirement is the possibility of the creation of sequences of research activities and the comparison of these pathways vis-à-vis the reported implementation success. We found that each project used a different sequence of research activities, resulting in different pathways. What these pathways have in common is that they include activities to explore a new solution, deliver a prototype construct, and contribute to theory. Another insight is that most

research activities did have the goal to explore whether an innovation could possibly work and/or is part of the 'conceptual context'. In these activities, researchers involved end-users. However, only three projects that include the exploration of the solution in a real-life environment in which the users tried a prototype in their own situation, succeeded in implementing their product. Also activities with the goal to deliver real-life change, such as preparing health practitioners and organizations for using the solution through training and integration in the existing workflow, are prerequisites for success.

Several limitations of our analysis could be mentioned. First, process features could only be analysed if the available documents or interviewee explicitly described a research activity or factor. Second, as we evaluated the process in a retrospective way for each project, future research should focus on the perspective of the different stakeholders before or during the process. Third, for all projects, we interviewed the main researcher and another stakeholder, but we are aware that we missed the perspectives of other involved stakeholders. This might result in a biased picture of the projects. However, we do think that this bias may be limited because we always included multiple perspectives, interviewing at least two stakeholders per project. In all cases both perspectives did not reveal real differences about the research pathway. In addition, some researchers, business partners and healthcare professionals were also involved in other cases and were able to compare cases and add information about other cases.

Subsidy providers and researchers may benefit from the success stories and challenges of these nine cases and use the RPM as a toolbox to prepare, monitor and evaluate their research projects together with involved stakeholders and to discuss their research pathways to successful implement innovations. The role of the researcher in the implementation of innovations is limited and ended at a certain moment. Other stakeholders are part of an ongoing process of developing the innovation further in their own or other specific contexts. An implemented innovation, therefore, is the starting point of a process in which practice continuously develops the innovation (Landry, Amara and Lamari 2002). The Research Pathway Model aims to understand such continuous implementation process of PBR.

Acknowledgements

The authors would like to thank Lisette Munneke PhD, Manon Bloemen PhD, Marlies Welbie PhD, Jasper Huitink, Niek Pronk, and the research group of Research Competence for their contribution to the improvements and the design of the Research Pathway Model.

Conflict of interest statement. None declared.

Availability of data and materials

The dataset analysed and data extraction tools used during the current study are available from the corresponding author on reasonable request.

Authors' contributions

W.v.B., E.M., W.B., and D.A. conceptualized the study. W.v.B. interviewed the respondents and analysed the data. All authors (W.v.B., E.M., D.A.,

W.B., G.v.d.V., and H.P.) contributed to the final version and read and approved the final manuscript.

Ethics approval and consent to participate

The interviewees all agreed with their participation.

References

- Barlow, J., Wright, C., Sheasby, J., Turner, A., and Hainsworth, J. (2002) 'Self-Management Approaches for People with Chronic Conditions: A Review', *Patient Education and Counselling*, 48: 177–87.
- Bornmann, L. (2013) 'What is Societal Impact of Research and How Can It Be Assessed? A Literature Survey', *Journal of the American Society for Information Science and Technology*, 64: 217–33.
- Bozeman, B., and Sarewitz, D. (2011) 'Public Value Mapping and Science Policy Evaluation', *Minerva*, 49: 1–23.
- Buxton, M., and Hanney, S. (1996) 'How Can Payback from Health Services Research Be Assessed?', *Journal of Health Services Research & Policy*, 1: 35–43.
- Candy, L., and Edmonds, E. A. (2010) 'The Role of the Artefact and Frameworks for Practice-Based Research' in Biggs M., Larsson H. (ed.) *The Routledge Companion to Research in the Arts*, pp. 120–37. Routledge: New York.
- Coombs, S. K., and Meijer, I. (2021) 'Towards Evaluating the Research Impact Made by Universities of Applied Sciences', *Science and Public Policy*, 48: 226–234.
- Dalsgaard, P., and Dinder, C. (2014). 'Between Theory and Practice: Bridging Concepts in HCI Research', In Proceedings of the 32Nd Annual ACM Conference on Human Factors in Computing Systems (CHI '14), 1635–1644. 10.1145/2556288.2557342
- De Jong, S., Barker, K., Cox, D., Sveinsdottir, T., and Van den Besselaar, P. (2014) 'Understanding Societal Impact through Productive Interactions: ICT Research as a Case', *Research Evaluation*, 23: 89–102.
- Donovan, C., and Hanney, S. (2011) 'The "Payback Framework" Explained', *Research Evaluation*, 20: 181–3.
- Douthwaite, B., Alvarez, S., Cook, S., Davies, R., George, P., Howell, J., Mackay, R., and Rubiano, J. (2007) 'Participatory Impact Pathway Analysis: A Practical Application of Program Theory in Research-for-Development', *Canadian Journal of Program Evaluation*, 22: 127–59.
- Hevner, A. R. (2007) 'A Three Cycle View of Design Science Research', *Scandinavian Journal of Information Systems*, 19: 87–92.
- Janssen, M. (2016) 'Situating Novelty: A Study on Healthcare Innovation and Its Governance', Ph.D. diss., Erasmus University, Rotterdam.
- Janssen, M., Stoopendaal, A. M. V., and Putters, K. (2015) 'Situating Novelty: Introducing a Process Perspective on the Study of Innovation', *Research Policy*, 44: 1974–84.
- Joly, P.-B., Gaunand, A., Colinet, L., Larédo, P., Lemarié, S., and Matt, M. (2015) 'ASIRPA: A Comprehensive Theory-Based Approach to Assessing the Societal Impacts of a Research Organization', *Research Evaluation*, 24: 440–543.
- Jonkman, N., Schuurmans, M., Jaarsma, T., Shortridge-Baggett, L., Hoes, A., and Trappenburg, J. (2016) 'Self-Management Interventions: Proposal and Validation of a New Operational Definition', *Journal of Clinical Epidemiology*, 80: 34–42.
- Kennedy, A., Rogers, A., Bowen, R., Lee, V., Blakeman, R., Gardner, C., Morris, R., Protheroe, J., and Chew-Graham, C. (2014) 'Implementing, Embedding and Integrating Self-Management Support Tools for People with Long-Term Conditions in Primary Care Nursing: A Qualitative Study', *International Journal of Nursing Studies*, 51: 1103–13.
- Kok, M. O., and Schuit, A. J. (2012) 'Contribution Mapping: A Method for Mapping the Contribution of Research to Enhance Its Impact', *Health Research Policy and Systems*, 10: 21.

- Landry, R., Amara, N., and Lamari, M. (2002) 'Does Social Capital Determine Innovation? To What Extent?', *Technological Forecasting and Social Change*, 69: 681–701.
- Langley, A., and Tsoukas, H. (2012) 'Introducing "Perspectives on Process Organization Studies"', in Hernes T., Maitlis S. (ed.). *Process, Sensemaking and Organizing*, Oxford: Oxford University Press.
- Lorig, K. R., and Holman, H. (2003) 'Self-Management Education: History, Definition, Outcomes, and Mechanisms', *Annals of Behavioral Medicine*, 26: 1–7.
- Mankins, J. C. (2009) 'Technology Readiness Assessments: A Retrospective', *Acta Astronautica*, 65: 1216–23.
- McColl-Kennedy, J., Vargo, S., Dagger, T., Sweeney, J., and Van Kasteren, Y. (2012) 'Healthcare Customer Value Cocreation Practice Styles', *Journal of Service Research*, 15: 370–89.
- Miles, M. B., and Huberman, A. M. (1994) *Qualitative Data Analysis: An Expanded Sourcebook*. Thousand Oaks, CA: Sage.
- Rabin, B. A., Brownson, R. C., Haire-Joshu, D., Kreuter, M. W., and Weaver, N. L. (2008) 'A Glossary for Dissemination and Implementation Research in Health', *Journal of Public Health Management Practice*, 14: 117–23.
- Rosenberg, N. (1994). *Exploring the Black Box: Technology, Economics, and History*. Cambridge University Press: Cambridge.
- Sivertsen, G., and Meijer, I. (2020) 'Normal versus Extraordinary Societal Impact: How to Understand, Evaluate, and Improve Research Activities in Their Relation to Society?', *Research Evaluation*, 29: 66–70
- Spaapen, J., and van Drooge, L. (2011) 'Productive Interactions as a Tool for Social Impact Assessment of Research', *Research Evaluation*, 20: 211–8.
- Stake, R. (1995). *The Art of Case Study Research*, pp. 49–68. Thousand Oaks, CA: Sage.
- Van Beest, W., Balje, J., and Andriessen, D. (2017) 'Hoe Meet je de Praktische Relevantie?', *TH&MA-Tijdschrift Voor Hoger Onderwijs & Management*, 4: 52–7.
- Van Beest, W., Boon, W. P. C., Andriessen, D., Moors, E., Van der Veen, G., and Pol, H. (2020) 'Successful Implementation of Self-Management Health Innovations', *Journal of Public Health (Berl.)*. 10.1007/s10389-020-01330-y

Appendix 1: Case selection

Table A.1. Case selection

Project	Aim of the project	Subsidy provider	Time period	Type of innovation	Co-investigators	Characteristics interviews main researchers	Characteristics subsequent interviews
1	To develop and bring together knowledge and skills in the field of self-management and chronic pain rehabilitation.	SIA Raak Publiek	s–2020	Intervention toolbox	Research group physiotherapy UAS Research group co-design UAS Research group communications UAS Academic hospital Research university Three rehabilitation centres Two associations	Main researcher (PhD) from the research group of physiotherapy	Designer from the co-design research group
2	To improve an app with a self-management programme for people with knee and hip osteoarthritis.	ZonMW	2018–9	App	Research group physiotherapy UAS Other UAS Two companies	Main researcher (PhD) from the research group of physiotherapy	Commercial director of a physiotherapy company
3	To create consensus about the concept of communication ability and the development of usable tools to chart communicative ability.	FNO	2016–8	Talking tool	Research group speech therapy UAS Research group co-design UAS Six associations	Main researcher (PhD) from the research group of speech therapy	Speech therapist
4	To co-design and co-evaluate technological interventions for persons with a mild intellectual disability or autism.	SIA RAAK Publiek	2016–8	Technology for stimulating self-management	Research group co-design UAS Research group social work other UAS Five healthcare organizations Research institute Research university Three companies Network organization	Main researcher from the co-design research group	Healthcare professional
5	To optimize an app for the monitoring of babies with a need for physiotherapy, developed in a previous project, with appropriate implementation strategies and business modelling.	SIA RAAK MKB	2016–8	App	Research group physiotherapy UAS Research group ICT UAS Research group co-design UAS Research group process of information systems Dutch Association of child physiotherapist Academic hospital, physiotherapy for children Research university Work group of child physiotherapy in the centres of the Netherlands	Main researcher from the physiotherapy research group	Physiotherapist Researcher from the research group process of information systems
6	To develop a talking touchscreen, which allows limited health-conscious patients to independently manage the Patient Specific Complaints List (PSK) to fill in.	SIA RAAK MKB	2010–2	Talking touch screen	Research group physiotherapy UAS Research group co-design UAS Research university Research institute Research institute for healthcare Work group of	Main researcher (PhD) from the research group of physiotherapy	Design researcher of the research institute

(continued)

Table A.1. Continued

Project	Aim of the project	Subsidy provider	Time period	Type of innovation	Co-investigators	Characteristics interviews main researchers	Characteristics subsequent interviews
7	To develop a tool for physiotherapists to get and keep patients active after a stroke.	SIA RAAK Publiek	2017–9	Tool to monitor conditions	physiotherapy in disadvantage areas Healthcare insurance Four associations Research group physiotherapy UAS Research group co-design UAS Research group communications UAS Academic hospital Research university Rehabilitation centre Two companies Two associations A group of physiotherapists	Professor from the research group of physiotherapy	Valorization interviews with six
8	physiotherapists To develop an innovative approach to social networking and e-learning focused on polypharmacy management, to support informal and formal caregivers of seniors with dementia. Seniors will be supported in their medication adherence using sensor technology and smart pillboxes. The tool is a digital platform that will bring together informal caregivers, medical professionals, pharmacists and the seniors with dementia themselves.	ZonMW/AAL	2017–9	Digital platform	Dutch partners: Research group technology for health innovations Research group pharmacy Company Institute for medicine Four other participating countries	Main researcher (PhD) from the research group of pharmacy	Project member from the Institute for medicine
9	To develop a mHealth intervention for COPD patients.	ZonMW	2014–9	mHealth intervention	Research group of Chronic Illness UAS Chair of Nursing science at Utrecht University The department of Revalidation, Nursing Science and Sport at the University Medical Centre of Utrecht Several healthcare institutions	Main researcher (PhD) from the research group of Chronic Illness UAS	Designer

Appendix 2. Coding scheme

Table A.2. Constructs of the Research Pathway Model

Main topic/code	Subtopic
Create theoretical understanding	To create understanding by study of the cause of problems To create understanding by studying existing literature
Create a concept	To develop new concepts To develop new theories to explain problems To develop new concept solutions for problems
Create real-life understanding	Through observations To determine why a solution is problematic To prove that a solution could work in principle
Explore theory or a concept or in a controlled situation	Through laboratory trials By taking measurements via a standardized test
Explore a concept	By performing a sample study By conducting a pilot
Explore a solution in real-life practice	By testing a solution in the environment of the end-user By observing a solution in the environment of the end-user
Deliver theory	By publishing the results
Deliver a concept	By delivering a prototype to a business partner By delivering a prototype to one or more specific context(s)
Deliver a solution in real-life practice	By supporting a training to help the implementation process By supporting the adaption of the product, knowledge or procedures By supporting research needed for the upscaling of the project

Appendix 3. Pathways of cases

Case 1

Research Pathway Model case 1

Research contexts/ Research activities	Theoretical context	Conceptual context	Real-life practice context
Create	A.1 Create theoretical understanding	B.1 Create a concept	C.1 Create real-life understanding
Explore	A.2 Explore theory or a concept in a controlled situation	B.2 Explore a concept	C.2 Explore a solution in real-life practice
Deliver	A.3 Deliver theory or knowledge	B.3 Deliver a concept	C.3 Deliver change in real-life practice real-life

Research pathway and activity description Research Pathway Model case 1

Research goal	Example
Step 1: C.1 Create real-life understanding	The researchers conducted a retrospective survey of the effects of pain rehabilitation, degree of self-management and the percentage of relapse in 6 months. This survey takes place at the rehabilitation centres involved among all patients who have completed a chronic pain rehabilitation programme. According to the protocol, these patients complete a short questionnaire after 3 and 12 months after the start of rehabilitation. For this research it is requested to also participate in a measurement after 6 months. The Dutch Pain Rehabilitation Data Set is used for this, supplemented by the Pain Self-Efficacy Questionnaire (PSEQ) and the Illness Perception Questionnaire (short version).
Step 2: B.1 Create a concept	A design-oriented study was carried out using co-design. Co-design aims to achieve a product through shared vision, social learning and mutual understanding between all major stakeholders, patients and professionals, taking into account the different perspectives and expectations. In this type of qualitative study, professionals and patients with chronic pain were asked about the underlying reasons that could possibly cause the relapse. Questions were also asked about methods that could be useful to prevent such a relapse. Through co-design, a prototype for an intervention has been developed to help people with chronic pain to retain the results of their rehabilitation. This is a workbook in which values and goals can be formulated and important moments or experiences can be saved during rehabilitation and afterwards, as a reminder and support in the event of a relapse.
Step 3: A.1 Create theoretical understanding	In a parallel process, a systematic literature study has been conducted into the effects of a self-management intervention on self-efficacy, daily activities and limitations in patients with chronic pain complaints. From this study, it can be concluded that a customized intervention is probably more effective than a generic approach that is the same for all patients.
Step 4: B.2 Explore a concept	With 60 students, the prototypes developed by students were presented at the rehabilitation centre, the exploration was not just about the prototype but, by hidden design, about the principle of the presented interventions. Based on the study, the intervention tool was developed in co-design.
Step 5: C.2 Explore a solution in real-life practice	The intervention tool was tested by 8 patients in a feasibility study. They rated the usability of the workbook on average with a 7.8 and motivated people to actually carry out their goals: 'The workbook was a guideline for me during rehabilitation'. The practitioners of the rehabilitation centres have also been asked about their experiences. They were positive about the possibilities of the intervention tool, in particular because a number of components fit well with their treatment. Based on the experiences of patients and professionals, the intervention tool has been adapted and available for all rehabilitation centres.
Step 6: B.3 Deliver a concept	In the context of the project, a game has also been developed that can support people with chronic pain to make their lives with chronic pain clear to their relatives.
Step 7: A.3 Deliver theory or knowledge	The research project was followed up by a PhD project. All studies are or will be published in national and international papers and collected in a manuscript. The results were shared during a symposium and in a couple of newsletters.
Step 8: C.3 Deliver change in real-life practice real-life	The intervention is made freely available and can be downloaded. The intervention is integrated with a different working method within a participating partner organization and within a non-partner organization. The intervention has also been implemented in an education programme of another university. The main researcher provides a training to interested organizations on the use of the intervention. The researchers applied for a grant for a follow-up study and were also awarded this.

Case 2

Research Pathway Model case 2

Research contexts/ Research activities	Theoretical context	Conceptual context	Real-life practice context
Create	A.1 Create theoretical understanding	B.1 Create a concept	C.1 Create real-life understanding
Explore	A.2 Explore theory or a concept in a controlled situation	B.2 Explore a concept	C.2 Explore a solution in real-life practice
Deliver	A.3 Deliver theory or knowledge	B.3 Deliver a concept	C.3 Deliver change in real-life practice

Research pathway and activity description Research Pathway Model case 2

Research goal	Example
Step 1: B.1 Create a concept	The previous project was about a self-management programme for people with knee and hip osteoarthritis. So that was actually a programme that was independent of the physical therapist. Based on that research, this project started because the researchers discovered that the self-management app was suitable for a certain group, but a lot of people discontinued the programme and need guidance from a physiotherapist. That led to this new application for the e-Exercise project.
Step 2: A.1 Create theoretical understanding	The researchers conducted a systematic review in blended behaviour change interventions in chronic somatic disorders.
Step 3: B.2 Explore a concept	Interviews, a focus group and discussions with a stakeholder-committee were conducted to explore the needs, values and requirements with respect to our to-be-developed blended intervention. The first version of e-Exercise was tested in a pre- and post-test pilot study. Feasibility outcomes, including recruitment rates within each practice, website usage (assignments completed and website visits), health-related outcomes (physical activity, physical functioning pain and fatigue) and user satisfaction, were measured. In addition, therapists and patients from the pilot study were interviewed to investigate users' experiences.
Step 4: C.2 Explore a solution in real-life practice	A protocol of a prospective, single-blinded, multi-centre cluster randomized controlled trial in the (cost)-effectiveness (3 and 12 months) of the app compared to usual physiotherapy with 200 patients. All measurements were performed at baseline, 3 and 12 months after inclusion. Retrospective cost questionnaires were sent at 3, 6, 9 and 12 months and used for the cost-effectiveness and cost-utility analysis.
Step 5: C.2 Explore a solution in real-life practice	The short- and long-term effectiveness of the app were compared to usual physiotherapy. The group who used the app received (in accordance to the study protocol) on average 5 face-to face sessions, the usual physiotherapy group received on average 12 sessions. Within group analyses showed for both groups a significant improvement in physical functioning. After 3 months, the group who used the app reported an increase in physical activity; however, no objectively measured physical activity differences were found. With respect to the secondary outcomes, after 12 months sedentary behaviour significantly increased in the e-Exercise group compared to usual physiotherapy.
Step 6: A.2 Explore theory or a concept in a controlled environment	The short- and long-term effectiveness of the app were compared to usual physiotherapy in patients. The group who used the app received (in accordance to the study protocol) on average 5 face-to face sessions, the usual physiotherapy group received on average 12 sessions. Within group analyses showed for both groups a significant improvement in physical functioning. After 3 months, the group who used the app reported an increase in physical activity; however, no objectively measured physical activity differences were found. With respect to the secondary outcomes, after 12 months sedentary behaviour significantly increased in the app users group compared to usual physiotherapy.
Step 7: B.2 Explore a concept	A mixed-methods study which aimed to explore which patient-, intervention- and environment-related factors are determinants of adherence to the online component of the app. The study contains a randomized controlled trial and interviews.
Step 8: C.2 Explore a solution in real life practice	A mixed-methods study which aimed to identify the determinants that promote or hinder physiotherapists in the use of a blended intervention. Prior to the study, many therapists were interested in blended physiotherapy. Of the 123 physiotherapists allocated the app, 54 recruited one or more eligible patients, 10 physiotherapists used the app after the study period. Determinants related to intervention usage were appropriateness, added value, time, workload, professional autonomy, environmental factors and financial consequences. Therapists recommended to improve the ability to tailor the app to patients' individual needs. Before implementation in physiotherapy practice, the researchers need to integrate more flexibility into the online programme and provide education about how to integrate an online programme within physiotherapy to obtain maximal benefit from both delivery modes.
Step 9: B.3 Deliver a concept	The researchers found a business partner who adopted the app, build the app again in their own system, implemented it in their physiotherapy company and make it available for other physiotherapists.
Step 10: A.3 Deliver theory or knowledge	The research project was part of a PhD project. All studies are published in national and international papers and collected in a manuscript. Besides, there is a website about the app and information on websites of the partners, the UAS and the subsidy provider.
Step 11: C.3 Deliver change in real-life practice real-life	The app is implemented by a business partner. The researchers have applied for and receive grants for an extension of the app to other disorders.

Case 3

Research Pathway Model case 3

Research contexts/ Research activities	Theoretical context	Conceptual context	Real-life practice context
Create	A.1 Create theoretical understanding	B.1 Create a concept	C.1 Create real-life understanding
Explore	A.2 Explore theory or a concept in a controlled situation	B.2 Explore a concept	C.2 Explore a solution in real-life practice
Deliver	A.3 Deliver theory or knowledge	B.3 Deliver a concept	C.3 Deliver change in real-life practice

Research pathway and activity description Research Pathway Model case 3

Research goal	Example
Step 1: B.1 Create a concept	Together with a panel consisting of parents of children with TOS and professionals from different disciplines, a definition of the concept of ‘communication ability’ has been developed.
Step 2: B.1 Create a concept	To develop a user-friendly tool for SLTs to facilitate the dialogue between SLT and parents on the needs of a child with language disorder, a prototype of the tool was developed during co-creation meetings with speech therapists.
Step 3: C.1 Create real-life understanding	A co-design workshops with SLTs ($n = 8$), resulting in: <ul style="list-style-type: none"> insight in target group (parents of children with LD)
Step 4: B.1 Create a concept	A co-design workshops with SLTs ($n = 8$), resulting in: <ul style="list-style-type: none"> design guidelines for development tool usability and product requirements for tool
Step 5: B.2 Explore a concept	Researchers changed roles from facilitators to designers. <ul style="list-style-type: none"> Prototyping with items on communicative participation from Delphi study. Evaluation prototypes with design guidelines, usability and product requirements, resulting in one prototype to elaborate on. The prototype from stage 2 was rated by SLTs ($n = 22$) on functionality, user friendliness, attractiveness and safety.
Step 5: B.2 Explore a concept	Usability testing <ul style="list-style-type: none"> SLT’s ($n = 4$) and parents ($n = 11$) tested the prototype in two iterative rounds and provided feedback that was used to adjust the prototype. The prototype from stage 3 was rated by SLTs ($n = 42$) on functionality, user-friendliness, attractiveness and safety.
Step 6: B.2 Explore a concept	The prototype was further improved in a usability study with parents and speech therapists.
Step 7: C.2 Explore a solution in real-life practice	To assess which changes scores reported by parents on a Likert-scale are clinically relevant a prospective cohort study was conducted for goal setting and evaluation with parents. The goal was evaluated with a Likert scale at the start of the treatment, after one week and after the treatment period. <p>The study sample divided in two groups: children that improved and children that did not.</p>
Step 8: A.3 Deliver theory or knowledge	The research project was followed up by a PhD project. All studies are or will be published in national and international papers and collected in a manuscript. The results were shared during a symposium and in a couple of newsletters.
Step 9: B.3 Deliver a concept	The conversation tool will be available from a publisher from December 2019. With the tool, the speech & language therapist can give the perspective of the patient a central place in the treatment and respond optimally to the request for help. The project thus ties in with healthcare development to involve patients in shaping treatment (shared decision making) and with the new definition of health: ‘Conducting one’s own management in the light of physical, social and emotional challenges of life’.
Step 10: C.3 Deliver change in real-life practice real-life	The tool is available via the publisher. Participating speech therapists and speech therapists in their network will implement the tool in their treatments.

Case 4

Research Pathway Model case 4

Research contexts/ Research activities	Theoretical context	Conceptual context	Real-life practice context
Create	A.1 Create theoretical understanding	B.1 Create a concept	C.1 Create real-life understanding
Explore	A.2 Explore theory or a concept in a controlled situation	B.2 Explore a concept	C.2 Explore a solution in real-life practice
Deliver	A.3 Deliver theory or knowledge	B.3 Deliver a concept	C.3 Deliver change in real-life practice

Research pathway and activity description Research Pathway Model case 4

Research goal	Example
Step 1: B.1 Create a concept	In this project, the consortium have collaborated on co-designing and co-evaluating technological interventions for persons with a mild intellectual disability or autism. Co-design and co-evaluation were central concepts during the project. Together with healthcare organizations and patients, the tools were figured out.
Step 2: C.1 Create real-life understanding	The project started with two projects to observe and co-design together with clients tools that might be worked for the target group.
Step 3: A.1 Create theoretical understanding	Literature research was conducted by students for their thesis. The outcomes were used in the project.
Step 4: A.3 Deliver theory or knowledge	The thesis of the main researcher was used as basis for the three substantive topics in the project: stress management, planning and communication.
Step 5: B.1 Create a concept	Based on the topics and wishes of the participating partners, a selection of tools for the co-evaluation was made.
Step 6: B.2 Explore a concept	The selected products were tested via co-design and co-evaluation by students in student projects in different rounds in the three participating healthcare organizations.
Step 7: A.3 Deliver theory or knowledge	Multiple student publications, student thesis and an online toolkit were published.
Step 8: B.3 Deliver a concept	Multiple prototypes were delivered and research projects were continued.

Case 5

Research Pathway Model case 5

Research contexts/ Research activities	Theoretical context	Conceptual context	Real-life practice context
Create	A.1 Create theoretical understanding	B.1 Create a concept	C.1 Create real-life understanding
Explore	A.2 Explore theory or a concept in a controlled situation	B.2 Explore a concept	C.2 Explore a solution in real-life practice
Deliver	A.3 Deliver theory or knowledge	B.3 Deliver a concept	C.3 Deliver change in real-life practice

Research pathway and activity description Research Pathway Model case 5

Research goal	Example
Step 1: B.1 Create a concept	Software was designed that makes safe receiving and storing of video images made by parents possible and attractive to use for child physiotherapy practices. One part of the study focuses on hard technology, adjustments for availability, scale, security and management; The other part focuses on soft technology, so adjustments to improve the user interface, ease of use and attractiveness of the application.
Step 2: B.2 Explore a concept	Context analysis by qualitative research, using a focus group meeting consisting of paediatric physical therapists from the research consortium.
Step 3: B.2 Explore a concept	Context analysis by qualitative research using semi-structured interviews with parents from a purposeful sample.
Step 4: C.1 Create real-life understanding	A student focused in her thesis on the parent instruction by participative research with parents in the pediatric physical therapist clinic.
Step 5: A.1 Create theoretical understanding	To develop a business and service model for primary care practices, researchers conducted examples of other business and service models at the level of other stakeholders such as health insurers, the trade association and possible service providers. The researchers chose for this approach because they did not find research examples of this specific theme.
Step 6: B.2 Explore a concept	To develop a business and service model for primary care practice, data collection takes place at the level of physiotherapy practices and at the level of other stakeholders such as health insurers, the trade association and possible service providers.
Step 7: A.1 Create theoretical understanding	A Community of Practice was developed with the aim to follow up on the network that has been built up in both projects. A literature review was conducted into suitable models and best practices.
Step 8: B.1 Create a concept	The paediatric physical therapist, stakeholders involved, the research group and experts from the Master's degree programme created a work plan for the formation of a Community of Practice-oriented research for infants.
Step 9: A.3 Deliver theory or knowledge	A scientific publication and a student thesis were published, the researchers present the results on a congress and in the workgroup, all publications are available via a public project website. The researchers organized a festival to launch their prototype and share the results.
Step 10: B.3 Deliver a concept	During the project, a valorization group was active with the goal to find a business model to implement and scale up the tool. Implementation and evaluation of the final implementation plan within the professional field was not a goal of the project. The Community of Practice gives the opportunity to work on valorization of the tool.

Case 6

Research Pathway Model case 6

Research contexts/ Research activities	Theoretical context	Conceptual context	Real-life practice context
Create	A.1 Create theoretical understanding	B.1 Create a concept	C.1 Create real-life understanding
Explore	A.2 Explore theory or a concept in a controlled situation	B.2 Explore a concept	C.2 Explore a solution in real-life practice
Deliver	A.3 Deliver theory or knowledge	B.3 Deliver a concept	C.3 Deliver change in real-life practice real-life

Research pathway and activity description Research Pathway Model case 6

Research goal	Example
Step 1: C.1 Create real-life understanding	Physiotherapists of multiple physiotherapy practices in the same region, who knows each other from a workshop collaborated to help low-literate patients from deprived areas. The professor of the research group was in contact with this group of physiotherapists. The main researcher started as a master student with talking with this group to understand the problem.
Step 2: C.1 Create real-life understanding	Together with the physiotherapists, the researchers discussed the existing questionnaires and decided to test the existing questionnaire at the target group. In a focus group, the researcher asked the respondents about their experiences with the existing questionnaire and their option towards the questionnaire.
Step 4: A.1 Create theoretical understanding	A literature research showed that talking touchscreens has good results for the target group of low-literate patients in a school. The researchers create the idea of a talking touchscreen as an alternative for the regular questionnaire.
Step 5: B.2 Explore a concept	The research institute had made a mock-up version of the tool together with students. It was already clear then that it was a touchscreen. Because after such a session, the mock-up was evaluated. After the evaluation, prototypes were made and a new evaluation round started, all in co-design together with the target group in the school. After that evaluation round, the researchers tested the prototype at 25 physiotherapy practices.
Step 6: B.3 Deliver a concept	None of the project partners wants to develop the prototype further. One of the students who had developed the tool technically, start a kind of start-up. He was involved in a start-up programme for students. But with him it didn't go well psychologically, so he didn't know how to oversee it all and stopped with developing the tool. The researchers won the RAAK awards with their prototype, the third prize. One member of the jury, who was also a CEO, was interested in the tool and helped with finding a business partner, but without success.
Step 7: A.3 Deliver theory or knowledge	The research project was part of a PhD project. All studies are published in national and international papers and collected in a manuscript. The knowledge institute still uses the project as an example project for co-designing with this target group.

Case 7

Research Pathway Model case 7

Research contexts/ Research activities	Theoretical context	Conceptual context	Real-life practice context
Create	A.1 Create theoretical understanding	B.1 Create a concept	C.1 Create real-life understanding
Explore	A.2 Explore theory or a concept in a controlled situation	B.2 Explore a concept	C.2 Explore a solution in real-life practice
Deliver	A.3 Deliver theory or knowledge	B.3 Deliver a concept	C.3 Deliver change in real-life practice

Research pathway and activity description Research Pathway Model case 7

Step 1: B.1 Create a concept	This project starts with the previous project, a project in which the researchers observed that people often become less active at home after a stroke. The solution for this problem is a FIT-stroke intervention in which people are trained to have sufficient capacities to be able to move, but in the previous project the researchers observed that they do not use their capabilities. The researchers start with an idea to focus on physical fitness, towards behavioural change and more exercise.
Step 2: A.1 Create theoretical understanding	For the development of the intervention, the researchers use existing knowledge about physical activity and the hindering and facilitating factors for physical activity. The researchers found in literature a positive effect on giving feedback on a performance. Based on literature they decided to use accelerometers to provide that feedback. In literature, they do not find an existing system that can give feedback on performance. The technology for reliably and validly measuring physical activity is available and there is insight into obstructing and facilitating factors for physical activity. However, there is no proven effective intervention for learning and maintaining a physically active lifestyle for patients after stroke.
Step 3: A.2 Explore theory or a concept in a controlled situation	To determine the training intensity and to be able to measure the outcome of the intervention, exercise therapists and physical therapists test with 50 participants (tested twice, with seven days between the two measurements) in an exercise laboratory with the 10 meter shuttle run test (SRT) of Verschuren. The oxygen uptake and carbon dioxide emissions will be measured via the breath by breath method with a calibrated Cortex Metamax B3.
Step 4: B.1 Create a concept	The researchers have chosen a Research through Design approach. This approach makes it possible to design behaviour-influencing interventions based on an understanding of the target group, the target behaviour and the social and physical context. This increases the perspective on effective interventions, which is reinforced by the participative (co-design) approach.
Step 5: B.2 Explore a concept	During a pilot study, the researchers test whether the intervention is feasible for patients, their caregivers and therapists. The intervention will take place after the completion of the rehabilitation process in the second or first line in order to be able to properly assess the results of the exercise intervention. At the end of the feasibility study, care providers and patients in focus groups will be asked how they experienced the giving and receiving of the intervention.
Step 6: B.2 Explore a concept	After the feasibility study, the researchers conducted a pilot with an experimental design. The results of this study showed no effect, it appears that people have stopped walking. Not walking distance, not fatigue, not fear of falling.
Step 7: A.3 Deliver theory or knowledge	The researchers shared their results and research project on a congress and via a digital newsletter. There is also a project website.
Step 8: B.3 Deliver a concept	The researchers try to stimulate the company partner to further develop the accelerometer. Thereby the researchers tried to implement this accelerometer in the education of the UAS, but without results.

Case 8

Research Pathway Model case 8

Research contexts/ Research activities	Theoretical context	Conceptual context	Real-life practice context
Create	A.1 Create theoretical understanding	B.1 Create a concept	C.1 Create real-life understanding
Explore	A.2 Explore theory or a concept in a controlled situation	B.2 Explore a concept	C.2 Explore a solution in real-life practice
Deliver	A.3 Deliver theory or knowledge	B.3 Deliver a concept	C.3 Deliver change in real-life practice

Research pathway and activity description Research Pathway Model case 8

Research goal	Example
Step 1: C.1 Create real-life understanding	End-users have been interviewed in group sessions and face-to-face interviews in Cyprus, Norway and the Netherlands. The central aims of these interviews were to understand the user needs, in order to be able to define the end-user requirements.
Step 2: B.1 Create a concept	Based on the interviews a design was made and a mock-up version of the tool was created.
Step 3: A.2 Explore theory or a concept in a controlled situation	The mock-up version was tested in a controlled trial.
Step 4: B.1 Create a concept	Based on the controlled trial, the design was adjusted and further developed.
Step 5: B.2 Explore a concept	The prototype was tested in a pilot in three countries.
Step 6: B.1 Create a concept	Based on the field trials in the pilot the prototype was adjusted.
Step 7: B.2 Explore a concept	The tool was tested for six weeks in the homes of end-users in three countries.
Step 8: B.3 Deliver a concept	Based on the field trials in the pilot a final prototype is delivered. A business partner is project leader of this project, and will decide about the further development.
Step 9: A.3 Deliver theory or knowledge	The Dutch researchers have published a paper about the co-design

Case 9

Research Pathway Model case 9

Research contexts/ Research activities	Theoretical context	Conceptual context	Real-life practice context
Create	A.1 Create theoretical understanding	B.1 Create a concept	C.1 Create real-life understanding
Explore	A.2 Explore theory or a concept in a controlled situation	B.2 Explore a concept	C.2 Explore a solution in real-life practice
Deliver	A.3 Deliver theory or knowledge	B.3 Deliver a concept	C.3 Deliver change in real-life practice

Research pathway and activity description Research Pathway Model case 9

Research goal	Example
Step 1: A.1 Create theoretical understanding	The research project is part of a self-management programme. Knowledge of other projects within the programme was the incentive to start with this project. At the beginning of the project, the researchers realized that there was a gap in knowledge about the effect of technological interventions, but during the project this topic became more popular in literature.
Step 2: B.1 Create a concept	Another researcher in the self-management programme had developed an action plan for self-management for COPD patients. The action plan was evaluated and proved effective. The upcoming attention for technological tools caused the researchers to create a digital tool instead of a paper version.
Step 3: B.1 Create a concept	In focus groups with patients and healthcare providers, the researcher started exploring with a need analysis what the expectations of the technology are for the target group. Based on this focus group, it was decided to continue the investigation into the tool.
Step 4: A.1 Create theoretical understanding	A Delphi study with international experts, inside and outside the network of the researchers, was part of the study for the understanding of relevant self-management behaviour and the best chance of influence this behaviour. The researchers tried to reach a consensus with the expert group on promising behaviour. Since the Delphi study the researchers create theoretical understanding about changing behaviour and decided to use the Behaviour Change Wheel.
Step 5: C.1 Create real-life understanding	The researchers conducted a current practice analysis among patients via a focus group to clarify the problem in their home situation. To obtain an idea of what care providers offer in self-management support, various care providers were approached and interviewed via the researchers' own network.
Step 6: A.2 Explore theory or a concept in a controlled situation	Based on literature, the researchers created a timeline to explore the phase in the illness of the patient in which self-management behaviour is needed.
Step 7: B.2 Explore a concept	A test version of the app has been made, a working app for the patient without a version for caregivers. The researchers first wanted to evaluate the app for user-friendliness among patients. It is still an iterative process, in which user-friendliness is evaluated and improvements are made based on the usability testing.
Step 8: B.2 Explore a concept	The researchers started an early feasibility study to see how care providers fit the app in their working method.
Step 9: A.3 Deliver theory or knowledge	The research project was followed up by a PhD project. All studies are or will be published in national and international papers and collected in a manuscript.
Step 10: B.3 Deliver a concept	Based on the study, the idea for the 'Long-attack app' came into being. With the idea for this app, the researcher, together with her colleague and co-promoter, has won the Ureka Mega Challenge 2017. This prize has led to the Long Attack app being realized. The biggest challenge to deliver the concept is to find collaboration with a business partner who is able to launch the app on the market.

Appendix 4. Interview schedule

Protocol for case study interviews with most involved researchers of projects

Introduction

The research is about the implementation of innovations in the field of self-management in healthcare based on practice-based research at universities of applied sciences in the Netherlands. To answer the question, I interview various researchers from the University of Applied Science in Utrecht and actors who are associated with one of the nine selected research projects. The interviews are about the impact of the research results within the network built up before, during and after the research project.

The interview is recorded. Immediately after the interview, it is put in a safe environment (One Drive). We therefore delete the recording from the recording device. Because I will be using the interview for research, I would like to ask to agree with the recording on tape.

During the interview, I would like to hear what your opinion and your own vision is. So there are no right or wrong answers.

STAGE 0: Introduction

1. For how long have you been involved in the project?
2. What is your role in this project?
3. What was the reason to start this project?
4. In which stage is the project at this moment? When did the project end? Is there a follow up project active at this moment?
 - a. If there is a follow up project, are you involved in this project?

Topic 1: Stakeholders

1. What was your role as in the team and the research process?
2. Who were the main researchers?
3. What was the role of collaborators in the research process (both academic and industrial)?
 - a. Which actors played a role in the initial phase? What was their role like?
 - b. Which actors played a role during the project? What was their role like?
 - c. Which actors played a role after the end of the project? What was their role like?
 - d. Are there actors / roles that you have missed during the process, but that you had needed in retrospect?
 - e. What was the role of the end user? Was there any interaction with potential users of the research during the research processes?
 - f. What was the role of the healthcare organization?
 - g. What was the role of stakeholders such as the government, health insurers, etc.?

Topic 2: Research process

I would like to gain insight into how the research process evolved and what different stakeholders contribute to the implementation towards the (healthcare) practice. I assume that impact can already be in small things and can also take place at the beginning of the project, for example through the way in which a consortium is established, the way of working during a joint session, etc.

1. What research activities were done in the initial phase of the research project
 - a. Which actors played a role in the initial phase in the impact of the innovation?
 - b. What was the focus of the project activities in the initial phase?
 - c. What activities focused on the delivery of the innovation in the initial phase?
2. What research activities were done during research project
 - a. Which actors played a role during the project in the impact of the innovation?
 - b. What was the focus of the project activities during the project?
 - c. What activities focused on the delivery of the innovation during the project?
3. What research activities were done at the end of the research project
 - a. Which actors played a role at the end of the project in the impact of the innovation?
 - b. What was the focus of the project activities at the end of the project?
 - c. What activities focused on the delivery of the innovation at the end of the project?

Topic 3: Innovation success

1. Do you think X is a successful innovation?
2. What is a successful innovation in your opinion?
3. How do you see the role of the research team in the implementation of the innovation?

Topic 4: Business model

1. Before, during and / or after the project, has a business model been considered with which this innovation can be brought to the market?
2. Has there been any discussion with end users about how this innovation could be funded?
3. Have there been discussions with healthcare organizations about how this innovation could be financed?
4. Did the research team have consulted entrepreneurs, subsidy providers, governments and / or health insurers about how this innovation could be financed?

Topic 5: Further development

1. Can this innovation also be adapted to the relevant healthcare institution or user?
2. Who is responsible for the further development of the innovation?
3. How will the innovation be developed after the subsidy ends?

Other general questions

1. Is it possible to speak with one of the consortium partners involved to speak to about your research?
2. Are there other questions we should have asked or things that you want to talk about?
3. Are you happy for us to contact you to follow up on details arising from the case study research?